Published in final edited form as:

Photochem Photobiol Sci. 2015 August ; 14(8): 1476–1491. doi:10.1039/c4pp00495g.

The role of photodynamic therapy in overcoming cancer drug resistance

Bryan Q. Spring¹, Imran Rizvi^{1,2}, Nan Xu^{1,3}, and Tayyaba Hasan^{1,4,5}

¹Wellman Center for Photomedicine, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, United States.

²Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts, United States.

³Department of Dermatology, Shanghai East Hospital and Tongji University, Shanghai, China.

⁴Division of Health Sciences and Technology, Harvard University and Massachusetts Institute of Technology, Cambridge, Massachusetts, United States.

Abstract

Many modalities of cancer therapy induce mechanisms of treatment resistance and escape pathways during chronic treatments, including photodynamic therapy (PDT). It is conceivable that resistance induced by one treatment might be overcome by another treatment. Emerging evidence suggests that the unique mechanisms of tumor cell and microenvironment damage produced by PDT could be utilized to overcome cancer drug resistance, to mitigate the compensatory induction of survival pathways and even to re-sensitize resistant cells to standard therapies. Approaches that capture the unique features of PDT, therefore, offer promising factors for increasing the efficacy of a broad range of therapeutic modalities. Here, we highlight key preclinical findings utilizing PDT to overcome classical drug resistance or escape pathways and thus enhance the efficacy of many pharmaceuticals, possibly explaining the clinical observations of the PDT response to otherwise treatment-resistant diseases. With the development of nanotechnology, it is possible that light activation may be used not only to damage and sensitize tumors but also to enable controlled drug release to inhibit escape pathways that may lead to resistance or cell proliferation.

Some major challenges in oncology include treatment toxicity and drug-resistance associated with advanced stage diseases that cannot be completely removed by surgical resection. Because many patients present with local infiltrates and distant metastases, systemic chemotherapy has become an essential partner to surgery and radiotherapy for extending patient survival. Despite tremendous advances in each of these modes of cancer therapy, refractory disease and recurrence remain common. In fact, even patients who have a complete clinical response to the frontline therapies often suffer a relapse with the emergence of lethal, drug-resistant disease—stemming in part from microscopic deposits of surviving cancer cells that escape treatment by various mechanisms. For example, this is common for malignancies of the ovary¹ and the brain².

⁵To whom correspondence should be addressed. thasan@mgh.harvard.edu.

Drug-resistance stems from both intrinsic and acquired mechanisms. These mechanisms include alterations in the drug target, increased drug efflux, and the activation of signaling pathways that promote the repair of damaged cellular components and that suppress cell death³. Many of these classical mechanisms of resistance influence both chemotherapy drugs and small-molecule inhibitors; thus, drug resistance has proven to be a tremendous challenge for gaining improvements using combinations of traditional agents. Compensatory signaling is also a common mode of resistance to molecular-targeted therapeutics, where the cancer cell uses alternative pathways to compensate for the inhibition of a given pathway³. These adaptive processes are influenced by the tumor microenvironment⁴, which can help to create a milieu conducive to resistance and escape. The epithelial-mesenchymal transition (EMT) program^{4,5}, as well as the cancer stem-like cell phenotype⁶, are known to promote metastasis as well as resistance to cell death with decreased sensitivity to a variety of treatment modalities. For instance, cancer stem-like cells highly express drug transporters⁶, are quiescent, and therefore inherently less sensitive to DNA damage⁶, while also possessing enhanced capacities for DNA damage repair⁷. The mesenchymal phenotype⁵ can be induced by cellular, molecular, or physical cues^{5,8,9} in the microenvironment and promotes cell motility, survival, and escape from localized stresses^{4,10}, as well as resistance to conventional agents^{11–14}. The EMT is an important developmental program in cancer invasion and metastasis and can generate the cancer stem-like cell phenotype, suggesting a plasticity among cancer cell subpopulations¹⁵.

Therefore, an emerging concept in oncology is that many cancer therapies actually induce drug resistance as well as enhanced invasiveness and metastasis, which may explain why clinical trials of novel drugs all too often report gains in local tumor control without a significant impact on overall survival (as postulated by Pàez-Ribes *et al.* in regards to antiangiogenic agents¹⁶). That is, increased local invasion and metastasis compensate for local tumor control. For example, this concept is now the subject of several thought-provoking research and perspective articles regarding how best to inhibit tumor escape and progression in response to antiangiogenic therapy^{16–19}. These findings point to the importance of utilizing mechanistically distinct, non-overlapping combination therapies to mop up mechanisms of treatment escape during each cycle of treatment. The combinations of therapeutic modalities should ideally also have non-overlapping toxicities. Dose-limiting toxicities exist for all therapies, such that combining agents with overlapping toxicities can be intolerable. If successful, rationally designed combination therapies offer great promise for reducing toxicity and for enabling the use of multiple treatment cycles to control local tumor growth, whilst suppressing the emergence of drug resistance and invasion. This development may be a key for achieving higher success rates in the clinic to impact patient survival.

In this Perspective, we begin by briefly introducing the principles of PDT. The following sections summarize the unique properties of PDT that overcome classical mechanisms of cancer drug resistance—including the reversal of chemoresistance and the sensitization of tumors to molecular targeted agents—and show how harnessing these distinctive features can make pharmaceuticals work better, while also reducing toxicity. In many cases, provided the mechanistic interactions are appropriately matched, the pharmaceutical-based therapy might in turn enhance PDT. The following discussion also introduces some concepts related

to resistance to PDT itself, but it is not meant to be a comprehensive review of these mechanisms, which we anticipate will be covered in other articles. Throughout, we highlight several important examples of how the photodynamic effect induces mechanisms of physical damage to multiple cellular and tumor compartments—leading to distinct cell death signaling pathways, re-sensitization of drug-resistant cells, and disruption of the tumor microenvironment. We also discuss the emergence of optically active nanomaterials and how light activation can be harnessed for both PDT and for stimulating tumor-confined, controlled drug release such that the drug is at the “right place at the right time”. This development enables a precise control of the interactive combination therapies, where PDT enhances the efficacy of a drug, and likewise, the drug mops up the mechanisms of escape from PDT. Finally, we discuss the prospects for the broader clinical translation of PDT based on these unique advantages.

Principles of PDT

PDT is a photochemistry-based therapeutic modality in which a light-activatable chemical (photosensitizer, PS) is energized by light (600–800 nm) to produce cytotoxic molecular species via an electron transfer to biological substrates (type I photosensitization), and potentially by the indirect excitation of molecular oxygen, or by the direct energy transfer to molecular oxygen (type II photosensitization)²⁰. The principal feature of PDT is its intrinsic dual selectivity—both the PS and light must be present for photodamage—and, therefore, its ability to address highly localized tissue damage and also the absence of toxicity outside of the illumination field. Chromophores with absorption wavelengths beyond ~750 nm (<160 kJ·mol⁻¹) lack sufficient energy for electronic coupling to facilitate the production of excited-state singlet oxygen species—a highly reactive oxygen species and key mediator of photodamage. For example, the S₁–T₁ electronic state energy gap must be at least 94 kJ·mol⁻¹, and ideally more than 157 kJ·mol⁻¹, to generate the major transitions of molecular oxygen²¹. Like with radiation therapy, PDT requires a threshold concentration of these toxic species, which minimizes the damage to surrounding tissues as long as there is a differential in PS concentration between the tumor and the surrounding tissue. The differential tumor uptake of the PS can be accomplished by approaches for passive, targeted, and target-activatable delivery^{22–24}. A number of PSs are already in clinical use or in clinical trials to treat cancer patients, including the hematoporphyrin derivative HpD (Photofrin), a first generation PS with prolonged skin phototoxicity approved by the US Food and Drug Administration for the palliative treatment of obstructive lung and esophageal cancers. Second-generation PSs, many with improved pharmacokinetics and reduced skin photosensitivity, include aminolevulinic acid (ALA; a pro-PS that cancer cells assemble into protoporphyrin IX), benzoporphyrin derivative (BPD), 5-ethylamino-9-diethyl-aminobenzo[a]phenothiazinium chloride (EtNBS), silicon phthalocyanine (Pc4), m-tetrahydroxyphenylchlorin (mTHPC), mesochlorin e₆ (Mce₆), and mono-L-aspartyl chlorin e₆ (NPe₆). Many other PSs exist with unique properties and are in various stages of development, and are reviewed elsewhere^{20,23}.

Unique Mechanisms of PDT: Cell Death Pathways, Direct Damage to Proteins Responsible for Classical Drug Resistance, and Enhanced Drug Delivery

Photodamage of antiapoptotic proteins

At the level of molecular cell biology, PDT induces mechanisms of cell death that depend on the subcellular localization of the particular PS, as well as on the photodynamic dose^{25,26}. Most PSs localize to cellular organelle membranes due to the common core, hydrophobic aromatic ring structure upon photodamage to these intracellular membranes, including their protein components. It is also possible to use molecular-targeted PDT to selectively damage specific proteins²⁷. Cellular photodamage can lead to cell death via any of the normal modes—necrosis, autophagy, or apoptosis. However, autophagy often plays a protective role in promoting cell survival after sub-lethal PDT²⁶. On the other hand, protective autophagy may also play an important role in antigen presentation for PDT-stimulation of an anti-tumor immune response²⁶. The two most well studied modes of PDT are the photodamage of lysosomes (lyso-PDT) and mitochondria (mito-PDT). Lyso-PDT leads to the spillage of proteases (e.g., cathepsins) into the cytosol, which in turn leads to cleavage (via the released lysosomal proteases rather than caspase-8) and activation of the proapoptotic factor BID (tBID)²⁸. A potential advantage of lyso-PDT is that lysosomal damage might circumvent autophagic protection²⁹. PSs that localize to the mitochondria and endoplasmic reticulum selectively damage antiapoptotic proteins of the BCL-2 family, which are then trafficked to the outer mitochondrial membrane³⁰, while the proapoptotic proteins are predominately cytosolic³¹, such as BAX, and are left intact. Following the loss of antiapoptotic proteins, tBID helps insert BAX into the outer mitochondrial membrane to stimulate cytochrome c release, which in turn activates effector caspases that drive the cell along an irreversible path to apoptosis³². Thus, lysosomal and mitochondrial photodamage can tip the balance of the apoptosis pathway towards pro-apoptosis. This mechanism of inducing apoptosis bypasses many of the checkpoints that account for resistance to radio- and chemotherapy (Figure 1). In fact, the overexpression of antiapoptotic BCL-2 family proteins, such as BCL-2 and BCL-X_L, is one of the major mechanisms of classical drug resistance³³.

Photosensitizer localization and impact on treatment outcome

By utilizing combinations of PSs that localize to different cellular compartments, Villanueva et al.^{34,35} and others^{36–38} demonstrated the enhanced efficacy and synergistic induction of cell death in cancer cell cultures. In an elegant study, Kessel and colleagues demonstrated that a sequential low dose of lyso- plus a low dose of mito-PDT in cancer cell cultures could achieve synergistic cancer cell photokilling via enhancement of the proapoptotic signaling, whereas the reverse sequence was less effective³⁸ (Table 1). Cincotta *et al.* showed a synergistic enhancement *in vivo* in a subcutaneous mouse model of large fibrosarcoma tumors (~1 cm at the start-of-treatment) when EtNBS-PDT was combined with BPD-PDT to achieve a 95% tumor reduction and 76% cure rate, whereas each mode of PDT alone was ineffective³⁹. The authors attributed this enhancement to different localizations of the two PSs within the tumor compartments (e.g., vascular versus tumor cell)³⁹. It is also conceivable that a significant component of the robust response comes from the different

subcellular localizations of the two PS—with EtNBS localizing to the lysosome (lyso-PDT) and BPD to the mitochondria (mito-PDT)³⁹—such that the same low dose lyso- plus low dose mito-PDT sequence was used as proposed by Kessel.

Photodamage of drug-efflux pumps

ATP-binding cassette (ABC) transporter proteins, including ABCB1 (MDR1, P-glycoprotein), ABCC1 (MRP1), and ABCG2, constitute another class of proteins involved in classical drug resistance, by increasing the cellular efflux and extracellular sequestration of many drugs⁴⁰, including some PSs as a mechanism for PDT resistance (discussed below in *Mechanisms of Resistance to PDT*). In a pioneering study, Goler-Baron and Assaraf demonstrated the photodestruction of ABCG2-rich extracellular vesicles associated with cancer cell drug efflux and sequestration⁴¹. This approach releases photosensitive drugs concentrated into these vesicles, such that they can reach their intracellular targets⁴¹. Inspired by this report, and as another example of the direct photodamage to proteins involved in drug resistance, our group found that PDT could be applied to directly damage ABCG2 (Huang-Chiao Huang *et al.* unpublished data).

Stimulation of anti-tumor immunity

A tremendous advantage of PDT is that certain regimens stimulate anti-tumor immunity, either using PDT alone^{42,43} or in combination with adjuvant immunostimulatory agents⁴⁴. In contrast to the immunosuppressive effects of traditional therapies, low-dose PDT regimens can induce anti-tumor immunity, and these regimens can be combined with high-dose PDT to achieve local tumor control with the immune suppression of distant disease⁴³. As reviewed in depth elsewhere⁴⁴, the mechanisms of PDT-enhanced anti-tumor immunity are under investigation. In brief, the potential mechanisms involve the acute inflammatory response following PDT, which might increase the presentation of tumor antigens to activate dendritic cells, and their homing in to regional and peripheral lymph nodes—ultimately stimulating CD8⁺ cytotoxic T cells and natural killer cells, accompanied by immune memory and the suppression of any subsequent tumor growth upon rechallenge^{43,44}.

Photodamage of the tumor microenvironment

A unique property of PDT is that the PS–light interval (the time interval between PS administration and the start of photoirradiation) can be exploited to target various tumor compartments. It is possible to induce photodamage to the tumor microvasculature, the parenchyma, and the stroma—or all of these compartments simultaneously—depending on the pharmacokinetics of the particular PS. For instance, the near infrared (NIR) photosensitizer BPD enables the selective damage to microvessels at early time points (~15 minutes) and damage to both the cancer cells and microvasculature at later time points (60–90 minutes post-injection)^{45–47}. The US Food and Drug Administration has approved BPD-PDT for the clinical treatment of macular degeneration, which is the major cause of blindness in older adults. For this application, BPD-PDT selectively destroys choroidal neovascularity associated with the disease, while sparing the overlying neurosensory retina to preserve visual acuity⁴⁸.

Photodynamic enhancement of drug delivery

Finally, PDT can be utilized to enhance drug delivery. First, low-dose vascular PDT can be used to transiently enhance blood vessel permeability, enabling an increased delivery of macromolecular and nanoparticle drug payloads to the tumor^{49–51}. Although this phenomenon has been known for over a decade, more recently it has been termed “super-enhanced permeability and retention (SUPR)”, in reference to further boosting the enhanced permeability and retention (EPR) effect well known in tumor biology⁵². The mechanism of increased tumor vessel leakiness is attributed to the formation of endothelial intercellular gaps, which might be induced via endothelial cell microtubule depolymerization following vascular photosensitization⁵⁰. An early demonstration of this concept by Henderson’s group showed dramatic enhancements in the delivery of fluorescent microspheres (0.1–2 μm) and of a liposome-encapsulated formulation of doxorubicin (Doxil) to subcutaneous tumors⁴⁹. This strategy has also been applied to enhance oncolytic virus accumulation in subcutaneous tumor models⁵¹.

Secondly, PDT can be applied to facilitate the cytosolic delivery of macromolecular drugs that normally cannot enter cells. Photochemical internalization (PCI) is a drug delivery method featuring endocytic escape—pioneered by Berg and colleagues based on the same principles as PDT for the controlled delivery of novel therapeutic agents that normally cannot access their intracellular targets⁵³. For PCI, the PS is used not only to elicit cytotoxic and vascular effects but also to photochemically rupture endocytic vesicles of the targeted cells to enable the photoinduced release of endocytosed therapeutic agents. The therapeutic agent is then released to interact with intracellular targets, rather than being subjected to lysosomal proteolysis and degradation. PCI has been demonstrated to facilitate the intracellular delivery of a variety of macromolecules that do not otherwise readily enter cells, including type I ribosome-inactivating proteins (RIPs), RIP-based immunotoxins, genes, and some chemotherapeutic agents⁵³. PCI can be applied using PSs that are not ABCG2 substrates⁵⁴ and to kill cancer stem-like cells^{55,56}.

Mechanisms of Resistance to PDT

Multidrug resistance (MDR)—where cells with intrinsic or acquired resistance to a single drug also show cross-resistance to other structurally and mechanistically unrelated drugs—is often accompanied by an increased expression of the ABC drug efflux transporters (introduced in *Unique Mechanisms of PDT*). Thus, many studies examining the mechanisms of resistance to PDT have probed for changes in PS cellular uptake, efflux, and localization. Hypoxia, stress responses and antioxidant enzymes are all possible mechanisms of resistance as reviewed previously⁵⁷. Although hypoxia is a general limitation of PDT and many other therapies that depend on oxygen-mediated mechanisms (e.g., radiation), certain PSs can impart photodamage in environments of low molecular oxygen via type I photosensitization. For example, EtNBS-PDT has been applied to selectively kill hypoxic regions of tumor nodules grown as 3D cancer cell cultures⁵⁸. PDT-induced tissue hypoxia as a result of vascular damage and photochemical oxygen consumption may also contribute to the appearance of resistant cells. Although the mechanisms of PDT-resistance have yet to be fully elucidated, and nuances exist due to the diverse mechanisms of PDT and the properties

of the individual PSs, the evidence thus far points to the involvement of a mix of novel mechanisms, as well as some elements of classical MDR. Overall, PDT does not induce significant cross-resistance to other modes of therapy and in fact has been shown to reverse the MDR phenotype⁵⁷. This may well be the case, since PDT has not been used so far as a multiple administration modality and the problem of PDT-induced resistance may emerge only once it has been administered several times, as is typically done in chemotherapy and radiation therapy.

Novel resistance mechanisms

The seminal studies of PDT-resistance pointed to novel mechanisms involving changes in mitochondria structure, function, and number^{59,60}. In 1991, Luna and Gomer used chronic HpD-PDT to induce a PDT-resistant phenotype of mouse RIF-1 (radiation-induced fibrosarcoma) cells⁵⁹. The RIF-1 cells were subjected to 10 cycles of HpD-PDT, with each cycle administered at a dose that kills 99.9% of the wild-type cells using clonogenic assays⁵⁹. The resulting cells were ~1.8-fold less sensitive to HpD-PDT, comparing the dose needed to kill 90% of the parental versus the resistant cell lines^{59,60}. After this exhaustive selection protocol, alterations in the number, structure, and function of the mitochondria were apparent, whereas the uptake, efflux, and localization of the PS were not affected^{60,61}. Changes in the basal levels of antioxidant enzymes, and reduced glutathione and stress responses were also minimal, and hence ruled out as a contributing factor⁵⁹. Interestingly, cross-resistance was observed towards exogenous PpIX-PDT, but not towards ALA-induced PpIX-PDT⁶⁰. Casas *et al.* later developed ALA-induced PpIX-PDT resistant lines with 4- to 7-fold resistance, but these cell lines were not cross-resistant to BPD-PDT⁶². Again, the mechanism was distinct from MDR and the resistant cells had increased numbers of mitochondria, as well as increased protein content with reduced PpIX per protein, although, cellular PpIX production was similar for the parental and resistant cell lines⁶². These data highlight the distinct mechanisms of resistance to PDT, in contrast to classical MDR, and to a lack of cross-resistance amongst different PSs. This contrasts strongly with the classical MDR, in which resistance to one drug leads to resistance to a broad spectrum of drugs.

About a third of the PDT-resistant cell lines (resistant to PDT with a specific PS, but not necessarily to PDT with other PSs) have been found to be cross-resistant to standard chemotherapies⁶². For instance, the HpD-PDT resistant RIF-1 cells are ~1.6-fold cross-resistant to cisplatin chemotherapy⁶¹. However, these levels of PDT- and chemoresistance (~2–7-fold) resulting from mitochondrial alterations are both much lower than the resistance levels commonly observed in drug-resistant cell lines (~10–100-fold^{63,64}). An interesting observation is that the HpD-PDT-resistant RIF-1 cells are several thousand-fold less efficient in tumor initiation than the parental line⁵⁹. In contrast, drug-resistance is often associated with the enrichment of the cancer stem-like cell population (often defined functionally as tumor-initiating cells based on their enhanced tumorigenic potential) that initiates tumors in mice with several orders-of-magnitude fewer cells than normally required⁶. It will be interesting to study the tumor initiation capacity of other PDT-resistant cell lines, as well as biomarkers of the stemness and MDR.

Classical resistance mechanisms that impact PDT efficacy

A second class of PDT resistance does relate to classical MDR mediated by increased PS efflux^{65,66}. Although HpD is not a substrate of ATP transporters (explaining why this mechanism of resistance was not apparent in prior studies with HpD), many second-generation PS have been shown to be effluxed by ABCG2 (introduced above; but, not by other transporters)^{65,67}. Resistance to photocytotoxicity via PS efflux is 4–30-fold, depending on the PS and the treatment protocol⁶⁵. This level of resistance approaches levels commonly found for classical MDR, in contrast to the resistant phenotype with mitochondrial changes discussed above. Thus, a concern is that PS uptake may be insufficient in certain tumor cell side populations (i.e., stem-like cells) that often express higher levels of ABCG2⁶⁷. This effect is reversible using ABCG2 inhibitors⁶⁶; however, none are presently approved for clinical use due to adverse pharmacokinetic interactions with standard chemotherapies⁴⁰. In addition, there are concerns about the systemic use of these inhibitors, which could adversely affect transporter-mediated protection of the central nervous system (e.g., the blood brain barrier) and protection of the body broadly via the excretion of toxins (e.g., from the liver and kidneys). As an alternative to ABC transporter inhibitors, and as already mentioned, PDT can actually be used to directly damage the ABCG2 transporter (Huang-Chiao Huang *et al.* unpublished data), to downregulate cancer cell stemness (and ABCG2 expression)⁶⁸ and to rupture ABC transporter-rich extracellular vesicles, releasing high payloads of sequestered drugs⁴¹. These effects can be used for “chemosensitization” to enhance the effects of standard chemotherapy, as discussed more broadly in the next section.

Reversal of Chemoresistance and the Potentiation of Chemotherapy

Dose-limiting toxicities and the development of treatment resistance limit the utility and efficacy of conventional chemotherapeutic agents. Despite efforts to combine different classes of chemotherapy agents with varying doses and schedules, the clinical response is often not durable and produces only marginal survival benefit with a poor quality of life⁶⁹. For example, FOLFIRINOX, a combination of 4 chemotherapy drugs, generated excitement by extending patient survival of pancreatic cancer by ~4 months compared to standard gemcitabine chemotherapy, but only patients with a good performance status qualify for the treatment regimen due to its increased toxicity⁷⁰. Broadly, the response to chemotherapy is often transient, and patients who develop chemoresistance have a dismal prognosis with little hope for effective treatment of their disease. Overcoming the resistance mechanisms that lead to treatment failure is of critical importance to improving cancer-related outcomes.

The distinct mechanisms of PDT synergize with chemotherapeutics and targeted biologics, and can reverse chemoresistance^{23,71–74}, and PDT has non-overlapping toxicities with these therapies. This section highlights key evidence of chemosensitization and examines the design of PDT-based combinations with chemotherapy. One crucial observation thus far is that successful implementation of PDT-enhanced chemotherapy requires a critical understanding of the biological targets, the specific chemotherapy agents, and the PS being considered for the combination regimen. The following examples highlight the sequence dependent effects among other complexities. Several combinations of antineoplastic drugs

(e.g., cisplatin, carboplatin, doxorubicin, mitomycin C, and methotrexate) and PSs (e.g., HpD, indocyanine G, Mce₆, mTHPC, ALA, and BPD) have been tested with varying results⁷⁴. The therapeutic interaction in these studies ranged from synergistic to antagonistic, depending on the specific PS and chemotherapeutic combination that was evaluated, the treatment sequence, and schedule, as well as on the tumor type⁷⁴.

Early results from preclinical models

Early investigations by Nahabedian *et al.* tested HpD-PDT in combination with cisplatin or doxorubicin in RIF-1 and EMT-6 tumor mouse xenograft models. EMT-6 tumors were only moderately sensitive to cisplatin or doxorubicin alone at the doses evaluated in the study. HpD-PDT significantly enhanced the efficacy of doxorubicin in EMT-6 tumors, but showed no significant additional anti-tumor effect in combination with cisplatin⁷⁴. Canti and co-workers evaluated the same chemotherapeutic agents followed by aluminum phthalocyanine-PDT in L1210 leukemia and P388 tumor-bearing mice and found an additive effect⁷⁴. A study by Baas and colleagues highlighted the importance of the treatment schedule in designing PDT-based combinations. The authors evaluated the effect of combining HpD-PDT and mitomycin C in subcutaneous RIF-1 tumors. The mitomycin C was administered either 15 minutes before or immediately after illumination of the tumors. Mitomycin C given prior to HpD-PDT significantly increased the delay in tumor growth compared to the monotherapies, whereas there was no enhancement when mitomycin C was given after the illumination⁷⁴. A study by Kopecek *et al.* emphasized the importance of proper dosing to assess the therapeutic interaction between the two modalities. The authors evaluated a combination involving doxorubicin and Mce₆-PDT in OVCAR3 human ovarian epithelial carcinoma cells in vitro. Mce₆-PDT and doxorubicin acted additively when each treatment was administered above a dose that was 50% effective (ED₅₀)⁷⁴. The combination was synergistic when the treatments were given at 50% of their ED₅₀ values⁷⁴. Another class of chemotherapeutics includes agents such as lonidamine and levamisole, which are inhibitors of cellular energy metabolism and have a powerful inhibitory effect on oxygen consumption, aerobic glycolysis, and lactate transport in neoplastic cells. These metabolic inhibitors showed a potent anti-tumor effect when combined with ALA-induced PpIX-PDT⁷⁴. Nonaka and colleagues investigated a combination involving cisplatin and HpD-PDT in L5178 mouse lymphoma cells and found a synergistic enhancement of apoptotic cell death due, in part, to the cooperative induction of caspases 3 activity by both PDT and cisplatin⁷⁴.

Evidence of re-sensitization and synergy in chemoresistant models

PDT is effective against a number of chemoresistant tumor cell lines: gemcitabine-resistant pancreatic cancer cell lines⁷⁵, platinum-resistant ovarian cancer cells⁷⁶, head and neck cancer stem-like cells⁶⁸, and temozolomide-resistant glioblastoma stem-like cells (Spring and Watanabe *et al.* unpublished data). PDT re-sensitizes chemoresistant ovarian cancer cell lines and patient-derived primary cultures to standard chemotherapy⁷³, as well as chemoresistant glioblastoma stem-like cells (Spring and Watanabe *et al.* unpublished data). The efficacy of PDT against gemcitabine-resistant pancreatic adenocarcinoma cells is instructive to consider as an example. The refractory nature of pancreatic adenocarcinoma to chemotherapeutics like gemcitabine—as well as its characteristic desmoplastic stroma—led to the hypothesis of insufficient drug delivery and stromal depletion (antistromal therapy) as

an exciting avenue for enhancing drug delivery via penetration of the hypovascular stroma⁷⁷. However, the pivotal clinical trial (NCT01130142) testing inhibition of the Hedgehog signaling pathway to elicit the stromal depletion of pancreatic tumors was stopped due to a difference in survival favoring the placebo plus gemcitabine arm, which had a lower rate of progressive disease than the stromal depleting saridegib plus gemcitabine arm (Infinity Pharmaceuticals, Inc., press release). Although the stroma undoubtedly plays a major role in malignant progression, recent papers have elucidated the role of the stroma in restraining, rather than promoting, tumor progression^{78,79}. PDT addresses perhaps a more fundamental challenge—pancreatic cancer cells are innately unresponsive to many drugs, even when barriers to delivery are not present. The study by Celli *et al.* showed that PDT is effective against a panel of gemcitabine-resistant cell lines by altering the balance of the pro- and antiapoptotic factors towards a proapoptosis state, and this approach works even for cells that become even more resistant to gemcitabine treatment when grown in contact with an extracellular matrix (Figure 2). Therefore, PDT has the potential to be a valuable mechanism-based adjuvant to lower the doses and associated toxicities with the gemcitabine treatment of pancreatic cancer.

Using patient tissue and cell lines, a study by Duska and colleagues demonstrated that photoimmunotherapy (PIT), which uses immunoconjugates to deliver the PS for enhanced PDT selectivity, reverses chemoresistance and synergistically enhances treatment efficacy⁷³ (Figure 3). The authors specifically investigated a combination of chlorin_{e6} (c_{e6})-mediated PIT and cisplatin, a chemotherapeutic agent commonly used to manage many cancers including ovarian cancer. To increase PDT selectivity, c_{e6} was conjugated to the F(ab')₂ fragment of OC-125, a murine monoclonal antibody that recognizes the cell surface antigen CA125, which is overexpressed in 85% of nonmucinous epithelial ovarian carcinomas. The potential of PIT in combination with cisplatin to potentiate the toxicity in tumors was evaluated in five human ovarian and breast cancer cell lines, as well as in tumor samples obtained from 14 patients with ovarian cancer who underwent primary cytoreductive surgery. The combination produced a significant reduction in tumor viability, relative to the monotherapies, and a synergistic enhancement of cisplatin efficacy was found in the cisplatin-resistant samples⁷³. In contrast, the effect was additive in the cisplatin-sensitive samples⁷³.

Sequence-dependence and structural impacts on drug penetration

The results from our group revealed a sequence-dependent synergistic enhancement of carboplatin efficacy with BPD-PDT in a three-dimensional (3D) culture model of micrometastatic ovarian cancer. Treatment with BPD-PDT ($1.25 \mu\text{M} \times \text{J}\cdot\text{cm}^{-2}$) prior to low-dose carboplatin ($40 \text{ mg}\cdot\text{m}^{-2}$) produced a synergistic reduction in the residual tumor volume, compared with PDT alone or carboplatin alone⁷². The reverse sequence, BPD-PDT after low-dose carboplatin, was not synergistic⁷². The explanation for this sequence-dependent synergism may lie in the mechanistic differences, and in cooperation at the subcellular and tumor architecture levels between BPD-PDT and carboplatin. BPD-PDT confers cytotoxicity in part by stimulating mitochondrial-induced apoptosis, which sensitizes the cells to subsequent nuclear damage and apoptotic signaling initiated by carboplatin (Figure

1). BPD-PDT also decreases the size of residual ovarian tumors and disrupts nodular architecture⁷², which are key barriers to the efficacy of platinum-based agents.

Emerging Concepts

An emerging consideration for the inclusion of PDT as part of the regular armamentarium for combination therapies is the potential for PDT to overcome resistance mechanisms conferred by cues from physical factors (such as flow-induced stress)⁸ and communication with stromal partners. Endothelial cells, for example, are emerging as increasingly important drivers of tumor biology and response to treatment, including radiation therapy and chemotherapy^{80,81}. EMT status is an additional critical determinant of the response to chemotherapy and targeted inhibitors^{11–14}. Early evidence suggests that PDT may overcome the chemoresistance conferred by endothelial cells and may be agnostic to EMT status (Rizvi *et al.* unpublished data).

Sensitization to Molecular Targeted Therapies

While PDT successfully kills most tumor cells, like any therapy, it also instigates molecular responses that provide growth and survival support to remaining cells. Like radiation therapy, PDT is a finite treatment that induces acute stress, accompanied by bursts in molecular signaling transduction pathways in response to this damage⁸² (L. Z. Zheng *et al.*, 2009, AACR-NCI-EORTC International Conference, abstract). These signaling events support resistance to cell death via a variety of mechanisms and are mediated largely by secreted factors, cell-surface receptor tyrosine kinases (RTKs), and the intracellular modulators of signaling networks. From another perspective, PDT sensitizes tumors to inhibition of these bursts in molecular signaling pathways. Combination therapies that mop up these signaling events can enhance local tumor control, while also preventing increased invasion and metastasis. Herein, we highlight a few molecular signaling pathways that have been exploited to enhance the outcome of PDT.

Sensitization to antiapoptotic complex inhibitors

Pioneering studies by Gomer and colleagues investigated the effects of PDT on molecular survival signaling cascades^{83–85}, ushering in a new area of research to understand survival signaling and optimal methods to mop up this signaling in the context of PDT. For instance, one early observation was that PDT induces the upregulation of heat shock protein (HSP) expression⁸³ concomitant with increased survivin activity⁸⁵, a member of the inhibitor of apoptosis (IAP) family. The IAP family uniquely forms a final checkpoint that can impart chemo-, radio-, and PDT-resistance to apoptosis by directly inhibiting the effector caspases (caspases 3 and 7). The upregulation of HSP-90 assists survivin activity and thereby the inhibition of apoptosis⁸⁵. This effect was successfully suppressed by interfering with the HSP-survivin complex to enhance PDT cell killing using a derivative of the antibiotic geldanamycin⁸⁵.

Sensitization to kinase inhibitors

Further lines of investigation involved the secreted factors and RTKs. The epidermal growth factor receptor (EGFR) is a RTK that regulates a number of critical cellular functions,

including proliferation, differentiation, motility, and survival via complex signaling cascades. Increased EGFR activity promotes cell cycle progression (G1 to S phase), causing disproportionate cell proliferation. EGFR overexpression is often associated with an increased aggressive or invasive phenotype and a poor prognosis in multiple cancers, including ovarian cancer. Many approaches for the targeted inhibition of the EGFR have been evaluated, but as with chemotherapy, the clinical response has shown limited durability, significant toxicities, and, at best, modest improvements in patient survival. Cetuximab is a chimeric monoclonal antibody approved by the US Food and Drug Administration for the treatment of metastatic colorectal cancer and head and neck cancer. Cetuximab specifically recognizes the EGFR and competes with the native ligand, EGF, to interrupt normal cell proliferation pathways and to induce G1 arrest. To overcome the dose-limiting toxicities and frequent relapse associated with this approach, our group evaluated the effects BPD-PDT combined with cetuximab on acute tumor reduction and survival enhancement in a mouse model for advanced stage human epithelial ovarian cancer⁷¹. The combination treatment produced both a synergistic reduction in mean tumor burden and a synergistic enhancement of the median survival relative to the monotherapies, with no significant increase in toxicity⁷¹. A possible explanation for this interaction involves a cetuximab-mediated blockade of EGFR activity, which prevents cells that overexpress EGFR from aberrantly entering the S phase, thereby inhibiting unregulated progression through the cell cycle and leading to the cytostatic inhibition of tumor growth. Upon inhibition of the EGFR, tumor cells that are highly dependent on these proliferation signals become particularly vulnerable to a secondary insult from a mechanistically non-overlapping treatment. Without this properly timed and rationally-selected insult, the tumor cells eventually develop compensatory pathways to overcome these inhibitory effects and escape cell-cycle arrest, which leads to treatment failure⁸⁶. BPD-PDT complements the cytostatic effects of cetuximab by photochemically triggering apoptosis, in part by inducing mitochondria-mediated apoptosis (Figure 1). Furthermore, a study by Gilaberte et al.⁸⁷ found that EGFR expression is correlated with the resistance to PDT with methyl-aminolevulinic acid (MAL-PDT) in analyses of PDT-resistant squamous cell carcinoma (SCC) cells and tumor biopsies from patients with persistent SCC following MAL-PDT.

Sensitization to antiangiogenic therapy

Vascular endothelial growth factor (VEGF), is a well-studied secreted factor involved in tumor angiogenesis, growth, and survival post-PDT^{84,88–91}. VEGF and its RTKs (e.g., VEGFR2) represent key targets for antiangiogenic therapy; also, the upregulation of VEGF signaling has been observed in response to radiotherapy⁹², chemotherapy⁹³, cytoreductive surgery⁹⁴, and PDT^{84,88}. Ferrario *et al.* introduced the concept of combining antiangiogenic therapy with PDT to improve therapeutic effectiveness motivated by their observation that HpD-PDT induces the expression of hypoxia-inducible factor-1 α (HIF-1 α) and VEGF in mouse mammary carcinomas⁸⁴. Our group found the p38 mitogen-activated protein kinase (MAPK) induction of VEGF by BPD-PDT, without the involvement of HIF-1 α , in an orthotopic mouse model of prostate cancer⁸⁸. The control of local tumor growth and reduced metastasis was observed when combining BPD-PDT with antiangiogenic agents in the same mouse model⁹⁵. We also developed *in vivo* hyperspectral fluorescence molecular imaging to longitudinally monitor the treatment-induced changes in VEGF expression, and found a

burst in tumor VEGF secretion immediately post-BPD-PDT (peaking at ~6–24 hours) in subcutaneous pancreatic and prostate xenograft tumor models⁸². This latter finding points to the importance of the timing of molecular signaling inhibition after PDT.

Due to the bursts in tumor molecular signaling following PDT, the spatiotemporal dynamics of molecular targeted inhibitor delivery become critical. Thus, an attractive area of development is to create optically active nanoparticles that support light-activated drug release in concert with PDT⁹⁶, such that the drug is present at the “right time and right place” to mop up dynamic survival signaling factors. A major contribution towards this goal is the porphyrin-liposome hybrids, termed porphyrosomes, developed by Zheng, Lovell, and colleagues^{97,98}. Porphyrin-phospholipid liposomes, for example, undergo reversible photopermeabilization under NIR irradiation and have been demonstrated to enable the spatial control of drug release (e.g., doxorubicin)⁹⁸. In addition, this concept has been developed for photo-induced gene transfer using optically active micelles⁹⁹.

Overcoming molecular signaling pathway co-activation, compensation and cross-talk

Our group is investigating the use of the light-triggered release of multikinase inhibitors in combination with PDT using a single nanoconstruct. This work addresses a dilemma in oncology—many molecular signaling pathways cross-talk, are co-activated in response to treatment, and compensate for the loss of a given pathway^{77,86,100}. VEGF, hepatocyte growth factor/scatter factor (HGF/SF), and their RTKs (VEGFR and MET, respectively) are prime examples of tumor signaling pathways that collaborate to promote treatment escape. Cancer cell MET signaling promotes the EMT and cancer cell stemness¹⁰¹, as well as tumor growth, invasion, and metastasis^{18,102,103}. Moreover, MET signaling is upregulated in response to anti-VEGF therapy in a number of cell types (e.g., both cancer cells and vascular cells) and comprises a prominent escape mechanism from antiangiogenic treatments¹⁸. When the tumor vasculature is pruned by anti-VEGF therapy, the hypoxic tumor microenvironment stimulates MET expression^{10,16–18}. Sennino et al. elegantly demonstrated that the concurrent inhibition of the VEGF and MET signaling pathways results in enhanced antiangiogenic effects to control tumor growth, while also mitigating cancer cell migration and invasive tumor growth along functional blood vessels or via lymphatic routes^{10,18,104}. We recently found that pancreatic cancer cells transiently upregulate MET signaling in response to PDT¹⁰⁵, which motivates a three-way interactive therapy that utilizes optically active nanoparticles for PDT with simultaneous drug release to enhance the efficacy of cancer cell death, prolonged local tumor control, and the suppression of metastatic escape (Spring, Sears and Zheng *et al.* unpublished data).

Prospects for Broader Clinical Translation of PDT

The complementarity of PDT with other standard therapies for cancer management suggests a broader use than is currently the case. This may be because, as for any therapeutic modality, there are concerns about the clinical use of PDT. The most common concern is the limited penetration of the far red and NIR light into tissue (attenuation depth of 1–4 mm in most tissues, but with photodamage reaching beyond the attenuation depth: up to ~1 cm¹⁰⁶), whereas X-ray radiation penetrates much deeper (>10 cm), albeit with very high doses

deposited near the tissue surface¹⁰⁶. Thus, the reasoning is that PDT cannot be applied beyond the skin. However, in practice, fiber optic light conduits enable clinical applications of PDT using interstitial fiber placement directly in deep seeded tumors, and, furthermore, the placement of multiple fibers is possible for treating large tumors. In our own clinical experience (a phase I/II clinical trial, VERTPAC), BPD-PDT produced a 1–4 cm zone of tumor necrosis (correlating with the administered light dose) with a 100% patient response rate for light delivered via optical fibers positioned percutaneously within locally advanced pancreatic adenocarcinoma tumors under computed tomography guidance¹⁰⁷ (Figure 4). PDT is also possible intraoperatively as an immediate follow-up to surgical debulking, as performed in the brain¹⁰⁸, as well as in the pleural¹⁰⁹ and peritoneal¹¹⁰ cavities. The fact is that the PDT of metastases and deep tumors is practical in clinics with modern technology. Light transport is made efficient in lumens, surgical beds, and cavities using diffusing tip fibers and scattering media (intralipid emulsion) to spread the light over large areas. For example, preclinical studies have demonstrated photodynamic tumor destruction in hepatic, pelvic, subgastric, diaphragmatic, spleen, and bowel sites with the peritoneal cavity¹¹¹, and that light delivery is feasible for the PDT of cancer deposits in the following anatomical sites: the esophagus^{20,23}, bladder^{20,23}, oral cavity and larynx¹¹², brain¹⁰⁸, bone¹¹³, lungs^{20,23,109}, pancreas¹⁰⁷, and those studding the peritoneal organs¹¹⁰. Finally, clinical trials have demonstrated the feasibility, safety, and efficacy for the photodynamic treatment of primary tumors in the pancreas¹⁰⁷, locally malignant glioblastoma multiforme in the brain¹⁰⁸, and disseminated, metastatic tumor deposits spread throughout the pleural (resulting from non-small-cell lung cancer)¹⁰⁹ and peritoneal (resulting from ovarian cancer as well as malignancies of the gastrointestinal tract) cavities¹¹⁰.

A second common misconception is the concern regarding heat generation during light irradiation. The light irradiances used for PDT ($\sim 0.1 \text{ W}\cdot\text{cm}^{-2}$) are an order-of-magnitude lower than those generally needed for laser coagulation and photothermal effects ($\sim 1 \text{ W}\cdot\text{cm}^{-2}$), although, a common mistake in the literature is to attribute photodynamic tumor destruction to photothermal effects under low irradiance continuous wave excitation. For example, photothermal cancer cell killing using IRDye700-antibody conjugates (and an irradiance of $\sim 0.003 \text{ W}\cdot\text{cm}^{-2}$) has been reported as a new modality termed photoimmunotherapy (PIT)¹¹⁴; however, IRDye700 is a silicon phthalocyanine, and phthalocyanines are well known to produce singlet oxygen¹¹⁵. In fact, silicon phthalocyanine Pc 4 is in clinical trials as a promising PDT agent¹¹⁶. Furthermore, phthalocyanine-antibody conjugates for cancer PIT were reported more than a decade ago as efficient PDT agents^{117,118}, with the general concept of PIT as a molecular-targeted approach to PDT being introduced over 30 years ago¹¹⁹.

The major concern among those with clinical experience is the complex dosimetry of PDT. Heroic efforts have been made to ensure uniform light dosimetry by placing photodetectors into the patient during treatment for online monitoring (T. C. Zhu et al., 2013, Proceedings of SPIE, abstract). This enables the clinician to monitor photodeposition in real time. However, concerns still remain since additional factors, such as PS concentration and oxygen perfusion, influence the ultimate photodynamic dose. These are ongoing dilemmas, no different from radiation therapy, and are under intensive development. In the near term, utilizing PDT as an adjuvant alleviates many of these concerns surrounding incomplete

treatment. Finally, improving the selectivity of PS accumulation within cancer cells is also a key for overcoming the toxicities when using PDT for wide-field treatments. As dosimetry and delivery technologies advance, we anticipate that PDT will be more widely adopted and become a core component of the armamentarium of cancer therapies.

Summary and Perspective

In conclusion, PDT has demonstrated promising clinical results for the treatment of cancer patients despite often being given as a last resort after all other options have failed. The use of PDT to address otherwise refractory disease has benefitted numerous patients worldwide—largely to provide palliation of advanced stage disease, but also to obtain durable cures of some early stage cancers (e.g., >90% cure rate of oral and larynx cancers including tumors unresponsive to radiotherapy¹¹²). PDT has unique mechanisms of action that: (1) reverse chemoresistance and sensitize tumors to molecular inhibitors; (2) modulate vascular permeability for enhanced drug delivery and/or to induce vascular occlusion to starve tumors of nutrients; and, (3) stimulate anti-tumor immunity. Resistance to PDT itself is possible but minimal cross-resistance results with other PSs and other modes of therapy, thus making it a legitimate partner, along with current conventional modalities for cancer treatment.

It is our view that, like other cancer therapies, PDT might best be utilized to potentiate a number of other modalities as part of novel frontline combination therapies. Rarely will a single treatment modality be curative. To date, there have been no clinical studies that have explored the use of PDT to potentiate chemotherapy, radiotherapy, or molecular inhibitors although there are several key publications in preclinical models to suggest the value of such a trial. Furthermore, multiple rationales motivate the integration of PDT into the clinical workflow. First, an abundance of preclinical evidence supports the use of PDT to overcome drug resistance. Second, the vast majority of PSs for PDT can be utilized for fluorescence-guided surgery¹²⁰ with follow-up PDT of the surgical bed. For instance, PS fluorescence-guided surgery with follow-up PDT doubled patient survival of glioblastoma multiforme versus standard surgery with follow-up radiotherapy (1 year versus 5.7 months) in a randomized, single center phase III trial. Third, for unresectable tumors, PDT can be applied using interstitial fibers to reach virtually any region in the body. As already mentioned, a phase I/II clinical trial demonstrated that PDT-induced tumor necrosis is feasible and safe in locally advanced, unresectable pancreatic adenocarcinoma tumors¹⁰⁷. Advances like these may pave the way forward to embark on more complex clinical trials incorporating combination regimens.

The major limitation of PDT, as for any cytotoxic modality, is the activation of tumor survival signaling pathways that promote treatment escape. That is, incomplete treatment carries the risks of actually stimulating invasion and metastasis. Nanotechnology-based drug delivery vehicles are now emerging for spatiotemporally synchronized PDT and for the release of potent inhibitors of these molecular signaling events. These advanced drug delivery systems could be used to maximize efficacy per treatment cycle, whilst thwarting survival mechanisms. The second major obstacle is alleviating the precision of light delivery needed to achieve effective PDT with minimal toxicity to off-target tissues. In contrast to traditional therapies, PDT carries an intrinsic dual selectivity for target lesions and,

therefore, minimal toxicity. Nevertheless, dose-limiting toxicities to sensitive tissues has been observed in clinical trials of wide-field PDT of disseminated metastases¹¹⁰. To address this challenge, a recent advance in the field is to utilize tumor biochemistry for PS activation only in the tumor, enabling fluorescence detection and tumor-confined PDT^{22,24}. This approach, combined with the targeting of cell surface molecules overexpressed by cancer cells, enables the monitoring and selective destruction of disseminated, microscopic tumors²⁴ (Figure 5). These activatable PDT agents alleviate the need for precise light delivery in PDT, which should help clinicians use PDT more broadly in the clinic.

Presently, cancer patients endure toxicities associated with high intensity dose schedules needed for optimal chemotherapy, radiation, and molecular inhibitor therapy. This use of high intensity dosimetry with poor efficacy per treatment cycle causes immense suffering, including: cardiovascular problems, hypertension, delayed wound healing, nausea, rash, diarrhea, hair loss, radiation scars, liver damage, gastrointestinal perforations, fatigue, immunosuppression, neurological damage, and more. PDT can be used to potentiate a number of molecular inhibitors and chemotherapy agents that have significant toxicities as single agents or additive toxicities when combined with other toxic drugs. On the other hand, chemotherapy and molecular inhibitor drugs potentiate PDT by enabling systemic effects to mop up residual tumor cells and survival signaling mechanisms. Combining these is mutually beneficial, and when done well, will enable the use of lower dosages of toxic drugs whilst maximizing impact per treatment cycle.

Acknowledgements

This work was supported by National Institutes of Health Grants R01-CA156177, R01-CA158415, R01-CA160998 and P01-CA084203 (to T.H.); and, K99-CA175292 (to I.R.).

References

1. Bast RC, Hennessy B, Mills GB. The biology of ovarian cancer: new opportunities for translation. *Nat Rev Cancer*. 2009; 9:415–428. [PubMed: 19461667]
2. Milano MT, et al. Patterns and timing of recurrence after temozolomide-based chemoradiation for glioblastoma. *Int. J. Radiat. Oncol. Biol. Phys.* 2010; 78:1147–1155. [PubMed: 20207495]
3. Holohan C, Van Schaeybroeck S, Longley DB, Johnston PG. Cancer drug resistance: an evolving paradigm. *Nat Rev Cancer*. 2013; 13:714–726. [PubMed: 24060863]
4. Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. *Cell*. 2011; 144:646–674. [PubMed: 21376230]
5. Epithelial-mesenchymal transitions in development and disease. 2009; 139:871–890.
6. Jordan CT, Guzman ML, Noble M. Cancer Stem Cells. *N Engl J Med*. 2006; 355:1253–1261. [PubMed: 16990388]
7. Bao S, et al. Glioma stem cells promote radioresistance by preferential activation of the DNA damage response. *Nature*. 2006; 444:756–760. [PubMed: 17051156]
8. Rizvi I, et al. Flow induces epithelial-mesenchymal transition, cellular heterogeneity and biomarker modulation in 3D ovarian cancer nodules. *Proc. Natl. Acad. Sci. U.S.A.* 2013; 110:E1974–E1983. [PubMed: 23645635]
9. Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. *J. Clin. Invest.* 2009; 119:1420–1428. [PubMed: 19487818]
10. Sennino B, et al. Suppression of tumor invasion and metastasis by concurrent inhibition of c-Met and VEGF signaling in pancreatic neuroendocrine tumors. *Cancer Discov.* 2012; 2:270–287. [PubMed: 22585997]

11. Thomson S, et al. Epithelial to mesenchymal transition is a determinant of sensitivity of non-small-cell lung carcinoma cell lines and xenografts to epidermal growth factor receptor inhibition. *Cancer Research*. 2005; 65:9455–9462. [PubMed: 16230409]
12. Haslehurst AM, et al. EMT transcription factors snail and slug directly contribute to cisplatin resistance in ovarian cancer. *BMC Cancer*. 2012; 12:91. [PubMed: 22429801]
13. Izumiya M, et al. Chemoresistance is associated with cancer stem cell-like properties and epithelial-to-mesenchymal transition in pancreatic cancer cells. *Anticancer Res*. 2012; 32:3847–3853. [PubMed: 22993328]
14. Du F, et al. Acquisition of paclitaxel resistance via PI3K- dependent epithelial- mesenchymal transition in A2780 human ovarian cancer cells. *Oncology Reports*. 2013; 30:1113–1118. [PubMed: 23807572]
15. Mani SA, et al. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell*. 2008; 133:704–715. [PubMed: 18485877]
16. Pàez-Ribes M, et al. Antiangiogenic Therapy Elicits Malignant Progression of Tumors to Increased Local Invasion and Distant Metastasis. *Cancer Cell*. 2009; 15:220–231. [PubMed: 19249680]
17. Pennacchietti S, et al. Hypoxia promotes invasive growth by transcriptional activation of the met protooncogene. *Cancer Cell*. 2003; 3:347–361. [PubMed: 12726861]
18. Sennino B, McDonald DM. Controlling escape from angiogenesis inhibitors. *Nat Rev Cancer*. 2012; 12:699–709. [PubMed: 23001349]
19. Jain RK. Antiangiogenesis Strategies Revisited: From Starving Tumors to Alleviating Hypoxia. *Cancer Cell*. 2014; 26:605–622. [PubMed: 25517747]
20. Dolmans DEJGJ, Fukumura D, Jain RK. TIMELINE: Photodynamic therapy for cancer. *Nat Rev Cancer*. 2003; 3:380–387. [PubMed: 12724736]
21. Schweitzer C, Schmidt R. Physical mechanisms of generation and deactivation of singlet oxygen. *Chem. Rev*. 2003
22. Lovell JF, Liu TWB, Chen J, Zheng G. Activatable photosensitizers for imaging and therapy. *Chem. Rev*. 2010; 110:2839–2857. [PubMed: 20104890]
23. Celli JP, et al. Imaging and photodynamic therapy: mechanisms, monitoring, and optimization. *Chem. Rev*. 2010; 110:2795–2838. [PubMed: 20353192]
24. Spring B, et al. Selective treatment and monitoring of disseminated cancer micrometastases in vivo using dual-function, activatable immunoconjugates. *Proc. Natl. Acad. Sci. U.S.A.* 2014; 111:E933–E942. [PubMed: 24572574]
25. Agarwal ML, et al. Photodynamic therapy induces rapid cell death by apoptosis in L5178Y mouse lymphoma cells. *Cancer Research*. 1991; 51:5993–5996. [PubMed: 1933862]
26. Kessel D. Death pathways associated with photodynamic therapy. *Med Laser Appl*. 2006; 21:219–224. [PubMed: 19890442]
27. Rahmzadeh R, et al. Ki-67 as a molecular target for therapy in an in vitro three-dimensional model for ovarian cancer. *Cancer Research*. 2010; 70:9234–9242. [PubMed: 21045152]
28. Wan Q, Liu L, Xing D, Chen Q. Bid Is Required in NPe6-PDT-induced Apoptosis. *Photochem Photobiol*. 2007; 84:250–257. [PubMed: 18173728]
29. Reiners JJ, Agostinis P, Berg K, Oleinick NL, Kessel D. Assessing autophagy in the context of photodynamic therapy. *Autophagy*. 2010; 6:7–18. [PubMed: 19855190]
30. Lindsay J, Esposti MD, Gilmore AP. Bcl-2 proteins and mitochondria--specificity in membrane targeting for death. *Biochim. Biophys. Acta*. 2011; 1813:532–539. [PubMed: 21056595]
31. Schellenberg B, et al. Bax exists in a dynamic equilibrium between the cytosol and mitochondria to control apoptotic priming. *Mol. Cell*. 2013; 49:959–971. [PubMed: 23375500]
32. Kessel D, Luo Y. Photodynamic therapy: A mitochondrial inducer of apoptosis. *Cell Death Differ*. 1999; 6:28–35. [PubMed: 10200545]
33. Reed JC, et al. BCL-2 family proteins: Regulators of cell death involved in the pathogenesis of cancer and resistance to therapy. *J. Cell. Biochem*. 1996; 60:23–32. [PubMed: 8825412]
34. Villanueva A, Stockert JC, Cañete M, Acedo P. A new protocol in photodynamic therapy: enhanced tumour cell death by combining two different photosensitizers. *Photochem. Photobiol. Sci*. 2010; 9:295–297. [PubMed: 20221454]

35. Acedo P, Stockert JC, ete MCN, Villanueva A. Two combined photosensitizers: a goal for more effective photodynamic therapy of cancer. 2014; 5 e1122-12.
36. Schneider-Yin X, et al. Hypericin and 5-aminolevulinic acid-induced protoporphyrin IX induce enhanced phototoxicity in human endometrial cancer cells with non-coherent white light. *Photodiagnosis and Photodynamic Therapy*. 2009; 6:12–18. [PubMed: 19447367]
37. Gyenge EB, et al. Photodynamic mechanisms induced by a combination of hypericin and a chlorin based-photosensitizer in head and neck squamous cell carcinoma cells. *Photochem Photobiol*. 2013; 89:150–162. [PubMed: 22882495]
38. Kessel D, Reiners JJ. Enhanced Efficacy of Photodynamic Therapy via a Sequential Targeting Protocol. *Photochem Photobiol*. 2014; 90:889–895. [PubMed: 24617972]
39. Cincotta L, Szeto D, Lampros E, Hasan T, Cincotta AH. Benzophenothiazine and Benzoporphyrin Derivative Combination Phototherapy Effectively Eradicates Large Murine Sarcomas. *Photochem Photobiol*. 1996; 63:229–237. [PubMed: 8657737]
40. Gottesman MM, Fojo T, Bates SE. Multidrug resistance in cancer: role of ATP[ndash]dependent transporters. *Nat Rev Cancer*. 2002; 2:48–58. [PubMed: 11902585]
41. Goler-Baron V, Assaraf YG. Overcoming multidrug resistance via photodestruction of ABCG2-rich extracellular vesicles sequestering photosensitive chemotherapeutics. *PLoS ONE*. 2012; 7:e35487. [PubMed: 22530032]
42. Gollnick SO, Vaughan L, Henderson BW. Generation of effective antitumor vaccines using photodynamic therapy. *Cancer Research*. 2002; 62:1604–1608. [PubMed: 11912128]
43. Shams M, Owczarczak B, Manderscheid-Kern P, Bellnier DA, Gollnick SO. Development of photodynamic therapy regimens that control primary tumor growth and inhibit secondary disease. *Cancer Immunol. Immunother*. 2014 **Advance online publication**,
44. Castano AP, Mroz P, Hamblin MR. Photodynamic therapy and anti-tumour immunity. *Nat Rev Cancer*. 2006; 6:535–545. [PubMed: 16794636]
45. Chen B, Pogue BW, Hoopes PJ, Hasan T. Combining vascular and cellular targeting regimens enhances the efficacy of photodynamic therapy. *Int. J. Radiat. Oncol. Biol. Phys*. 2005; 61:1216–1226. [PubMed: 15752904]
46. Fingar VH, et al. Analysis of acute vascular damage after photodynamic therapy using benzoporphyrin derivative (BPD). *Br J Cancer*. 1999; 79:1702–1708. [PubMed: 10206280]
47. Kurohane K, et al. Photodynamic therapy targeted to tumor-induced angiogenic vessels. *Cancer Letters*. 2001; 167:49–56. [PubMed: 11323098]
48. Schmidt-Erfurth U, Hasan T. Mechanisms of Action of Photodynamic Therapy with Verteporfin for the Treatment of Age-Related Macular Degeneration. *Survey of Ophthalmology*. 2000; 45:195–214. [PubMed: 11094244]
49. Snyder JW, Greco WR, Bellnier DA, Vaughan L, Henderson BW. Photodynamic therapy: a means to enhanced drug delivery to tumors. *Cancer Research*. 2003; 63:8126–8131. [PubMed: 14678965]
50. Chen B, Pogue BW, Luna JM, Hardman RL, Hoopes PJ. Tumor vascular permeabilization by vascular-targeting photosensitization: effects, mechanism, and therapeutic implications. *Clin. Cancer Res*. 2006; 12:917–923. [PubMed: 16467106]
51. Gil M, et al. Photodynamic therapy augments the efficacy of oncolytic vaccinia virus against primary and metastatic tumours in mice. *Br J Cancer*. 2011; 105:1512–1521. [PubMed: 21989183]
52. Sano K, Nakajima T, Choyke PL, Kobayashi H. Markedly enhanced permeability and retention effects induced by photo-immunotherapy of tumors. *ACS Nano*. 2013; 7:717–724. [PubMed: 23214407]
53. Norum O-J, Selbo PK, Weyergang A, Giercksky K-E, Berg K. Photochemical internalization (PCI) in cancer therapy: From bench towards bedside medicine. *Journal of Photochemistry and Photobiology B: Biology*. 2009; 96:83–92.
54. Selbo PK, et al. Strongly amphiphilic photosensitizers are not substrates of the cancer stem cell marker ABCG2 and provides specific and efficient light-triggered drug delivery of an EGFR-targeted cytotoxic drug. *J Control Release*. 2012; 159:197–203. [PubMed: 22349185]
55. Bostad M, et al. Photochemical internalization (PCI) of immunotoxins targeting CD133 is specific and highly potent at femtomolar levels in cells with cancer stem cell properties. *J Control Release*. 2013; 168:317–326. [PubMed: 23567040]

56. Bostad M, et al. Light-triggered, efficient cytosolic release of IM7-saporin targeting the putative cancer stem cell marker CD44 by photochemical internalization. *Mol. Pharm.* 2014; 11:2764–2776. [PubMed: 24960585]
57. Casas A, Di Venosa G, Batlle A. Mechanisms of resistance to photodynamic therapy. *Curr. Med. Chem.* 2011; 18:2486–2515. [PubMed: 21568910]
58. Evans CL, et al. Killing hypoxic cell populations in a 3D tumor model with EtNBS-PDT. *PLoS ONE.* 2011; 6:e23434. [PubMed: 21876751]
59. Luna MC, Gomer CJ. Isolation and initial characterization of mouse tumor cells resistant to porphyrin-mediated photodynamic therapy. *Cancer Research.* 1991; 51:4243–4249. [PubMed: 1831066]
60. Wilson BC, Olivo M, Singh G. Subcellular Localization of Photofrin and Aminolevulinic Acid and Photodynamic Cross-Resistance in Vitro in Radiation-Induced Fibrosarcoma Cells Sensitive or Resistant to Photofrin-Mediated Photodynamic Therapy. *Photochem Photobiol.* 1997; 65:166–176. [PubMed: 9066298]
61. Moorehead RA, Armstrong SG, Wilson BC, Singh G. Cross-Resistance to Cisplatin in Cells Resistant to Photofrin-mediated Photodynamic Therapy. *Cancer Research.* 1994; 54:2556–2559. [PubMed: 7909492]
62. Casas A, et al. Tumor cell lines resistant to ALA-mediated photodynamic therapy and possible tools to target surviving cells. *Int. J. Oncol.* 2006; 29:397–405. [PubMed: 16820882]
63. Duan Z, Lamendola DE, Duan Y, Yusuf RZ, Seiden MV. Description of paclitaxel resistance-associated genes in ovarian and breast cancer cell lines. *Cancer Chemother Pharmacol.* 2004; 55:277–285. [PubMed: 15565326]
64. Roberts D, et al. Identification of genes associated with platinum drug sensitivity and resistance in human ovarian cancer cells. *Br J Cancer.* 2005; 92:1149–1158. [PubMed: 15726096]
65. Robey RW, Steadman K, Polgar O, Bates SE. ABCG2-mediated transport of photosensitizers: potential impact on photodynamic therapy. *Cancer Biol. Ther.* 2005; 4:187–194. [PubMed: 15684613]
66. Liu W, et al. The Tyrosine Kinase Inhibitor Imatinib Mesylate Enhances the Efficacy of Photodynamic Therapy by Inhibiting ABCG2. *Clin. Cancer Res.* 2007; 13:2463–2470. [PubMed: 17438106]
67. Morgan J, Jackson JD, Zheng X, Pandey SK, Pandey RK. Substrate affinity of photosensitizers derived from chlorophyll-a: the ABCG2 transporter affects the phototoxic response of side population stem cell-like cancer cells to photodynamic therapy. *Mol. Pharm.* 2010; 7:1789–1804. [PubMed: 20684544]
68. Yu C-H, Yu C-C. Photodynamic therapy with 5-aminolevulinic acid (ALA) impairs tumor initiating and chemo-resistance property in head and neck cancer-derived cancer stem cells. *PLoS ONE.* 2014; 9:e87129. [PubMed: 24475244]
69. Yap TA, Carden CP, Kaye SB. Beyond chemotherapy: targeted therapies in ovarian cancer. *Nat Rev Cancer.* 2009; 9:167–181. [PubMed: 19238149]
70. Conroy T, et al. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med.* 2011; 364:1817–1825. [PubMed: 21561347]
71. del Carmen MG, et al. Synergism of epidermal growth factor receptor-targeted immunotherapy with photodynamic treatment of ovarian cancer in vivo. *J. Natl. Cancer Inst.* 2005; 97:1516–1524. [PubMed: 16234565]
72. Rizvi I, et al. Synergistic enhancement of carboplatin efficacy with photodynamic therapy in a three-dimensional model for micrometastatic ovarian cancer. *Cancer Research.* 2010; 70:9319–9328. [PubMed: 21062986]
73. Duska LR, Hamblin MR, Miller JL, Hasan T. Combination photoimmunotherapy and cisplatin: effects on human ovarian cancer ex vivo. *J. Natl. Cancer Inst.* 1999; 91:1557–1563. [PubMed: 10491432]
74. Combination of Photodynamic Therapy with Anti-Cancer Agents. 2008; 15:1655–1673.
75. Celli JP, Solban N, Liang A, Pereira SP, Hasan T. Verteporfin-based photodynamic therapy overcomes gemcitabine insensitivity in a panel of pancreatic cancer cell lines. *Lasers Surg. Med.* 2011; 43:565–574. [PubMed: 22057484]

76. Goff BA, Bamberg M, Hasan T. Photoimmunotherapy of human ovarian carcinoma cells ex vivo. *Cancer Research*. 1991; 51:4762–4767. [PubMed: 1716512]
77. Stommel JM, et al. Coactivation of Receptor Tyrosine Kinases Affects the Response of Tumor Cells to Targeted Therapies. *Science*. 2007; 318:287–290. [PubMed: 17872411]
78. Rhim AD, et al. Stromal elements act to restrain, rather than support, pancreatic ductal adenocarcinoma. *Cancer Cell*. 2014; 25:735–747. [PubMed: 24856585]
79. Özdemir BC, et al. Depletion of carcinoma-associated fibroblasts and fibrosis induces immunosuppression and accelerates pancreas cancer with reduced survival. *Cancer Cell*. 2014; 25:719–734. [PubMed: 24856586]
80. Upreti M, et al. Tumor-Endothelial Cell Three-dimensional Spheroids: New Aspects to Enhance Radiation and Drug Therapeutics. *Translational Oncology*. 2011; 4:365–376. [PubMed: 22191001]
81. Franses JW, Baker AB, Chitalia VC, Edelman ER. Stromal endothelial cells directly influence cancer progression. *Sci Transl Med*. 2011; 3:66a5.
82. Chang SK, Rizvi I, Solban N. In vivo optical molecular imaging of vascular endothelial growth factor for monitoring cancer treatment. *Clin. Cancer Res*. 2008; 14:4146–4153. [PubMed: 18593993]
83. Gomer CJ, et al. Photodynamic Therapy-mediated Oxidative Stress Can Induce Expression of Heat Shock Proteins. *Cancer Research*. 1996; 56:2355–2360. [PubMed: 8625311]
84. Ferrario A, et al. Antiangiogenic treatment enhances photodynamic therapy responsiveness in a mouse mammary carcinoma. *Cancer Research*. 2000; 60:4066–4069. [PubMed: 10945611]
85. Ferrario A, Rucker N, Wong S, Luna M, Gomer CJ. Survivin, a Member of the Inhibitor of Apoptosis Family, Is Induced by Photodynamic Therapy and Is a Target for Improving Treatment Response. *Cancer Research*. 2007; 67:4989–4995. [PubMed: 17510430]
86. Wheeler DL, Dunn EF, Harari PM. Understanding resistance to EGFR inhibitors-impact on future treatment strategies. *Nat Rev Clin Oncol*. 2010; 7:493–507. [PubMed: 20551942]
87. Gilaberte Y, et al. Cellular intrinsic factors involved in the resistance of squamous cell carcinoma to photodynamic therapy. *J Invest Dermatol*. 2014; 134:2428–2437. [PubMed: 24717244]
88. Solban N, et al. Mechanistic investigation and implications of photodynamic therapy induction of vascular endothelial growth factor in prostate cancer. *Cancer Research*. 2006; 66:5633–5640. [PubMed: 16740700]
89. Nowak-Sliwinska P, van Beijnum JR, van Berkel M, van den Bergh H, Griffioen AW. Vascular regrowth following photodynamic therapy in the chicken embryo chorioallantoic membrane. *Angiogenesis*. 2010; 13:281–292. [PubMed: 20842454]
90. Weiss A, et al. Low-dose angiostatic tyrosine kinase inhibitors improve photodynamic therapy for cancer: lack of vascular normalization. *Journal of Cellular and Molecular Medicine*. 2014; 18:480–491. [PubMed: 24450440]
91. Nowak-Sliwinska P, et al. Angiostatic kinase inhibitors to sustain photodynamic angio- occlusion. *Journal of Cellular and Molecular Medicine*. 2012; 16:1553–1562. [PubMed: 21880113]
92. Gorski DH, et al. Blockade of the vascular endothelial growth factor stress response increases the antitumor effects of ionizing radiation. *Cancer Research*. 1999; 59:3374–3378. [PubMed: 10416597]
93. Tran J, et al. A role for survivin in chemoresistance of endothelial cells mediated by VEGF. *Proceedings of the National Academy of Sciences*. 2002; 99:4349–4354.
94. Justinger C, et al. Increased growth factor expression after hepatic and pancreatic resection. *Oncology Reports*. 2008; 20:1527–1531. [PubMed: 19020737]
95. Kosharsky B, et al. A mechanism-based combination therapy reduces local tumor growth and metastasis in an orthotopic model of prostate cancer. *Cancer Research*. 2006; 66:10953–10958. [PubMed: 17108133]
96. Huang H-C, Hasan T. The ‘Nano’ World in Photodynamic Therapy. *Austin J Nanomed Nanotechnol*. 2014; 2:4.
97. Lovell JF, et al. Porphysome nanovesicles generated by porphyrin bilayers for use as multimodal biophotonic contrast agents. *Nat Mater*. 2011; 10:324–332. [PubMed: 21423187]

98. Carter KA, et al. Porphyrin-phospholipid liposomes permeabilized by near-infrared light. *Nat Comms.* 2014; 5:3546.
99. Nomoto T, et al. Three-layered polyplex micelle as a multifunctional nanocarrier platform for light-induced systemic gene transfer. *Nat Comms.* 2014; 5:3545.
100. Wilson TR, et al. Widespread potential for growth-factor-driven resistance to anticancer kinase inhibitors. *Nature.* 2012; 487:505–509. [PubMed: 22763448]
101. Hage C, et al. The novel c-Met inhibitor cabozantinib overcomes gemcitabine resistance and stem cell signaling in pancreatic cancer. *Cell Death Dis.* 2013; 4:e627. [PubMed: 23661005]
102. Gherardi E, Birchmeier W, Birchmeier C, Vande Woude G. Targeting MET in cancer: rationale and progress. *Nat Rev Cancer.* 2012; 12:89–103. [PubMed: 22270953]
103. Peters S, Adjei AA. MET: a promising anticancer therapeutic target. *Nat Rev Clin Oncol.* 2012; 9:314–326. [PubMed: 22566105]
104. Sennino B, Ishiguro-Oonuma T, Schriver BJ, Christensen JG, McDonald DM. Inhibition of c-Met reduces lymphatic metastasis in RIP-Tag2 transgenic mice. *Cancer Research.* 2013; 73:3692–3703. [PubMed: 23576559]
105. Zheng LZ, et al. Abstract A127: Combination therapy targeting EGFR/MET crosstalk using nanotechnology improves photodynamic therapy treatment of pancreatic cancer. *Molecular Cancer Therapeutics.* 2009; 8:A127–A127.
106. Bernstein, M.; Berger, MS. *Neuro-oncology: The Essentials.* 2011. Thieme
107. Huggett MT, et al. Phase I/II study of verteporfin photodynamic therapy in locally advanced pancreatic cancer. *Br J Cancer.* 2014; 110:1698–1704. [PubMed: 24569464]
108. Eljamel MS, Goodman C, Moseley H. ALA and Photofrin fluorescence-guided resection and repetitive PDT in glioblastoma multiforme: a single centre Phase III randomised controlled trial. *Lasers Med Sci.* 2008; 23:361–367. [PubMed: 17926079]
109. Friedberg JS, et al. Phase II trial of pleural photodynamic therapy and surgery for patients with non-small-cell lung cancer with pleural spread. *J. Clin. Oncol.* 2004; 22:2192–2201. [PubMed: 15169808]
110. Hahn SM, et al. A phase II trial of intraperitoneal photodynamic therapy for patients with peritoneal carcinomatosis and sarcomatosis. *Clin. Cancer Res.* 2006; 12:2517–2525. [PubMed: 16638861]
111. Molpus KL, et al. Intraperitoneal photodynamic therapy of human epithelial ovarian carcinomatosis in a xenograft murine model. *Cancer Research.* 1996; 56:1075–1082. [PubMed: 8640764]
112. Biel MA. Photodynamic Therapy Treatment of Early Oral and Laryngeal Cancers. *Photochem Photobiol.* 2007; 83:1063–1068. [PubMed: 17880501]
113. Lo VCK, et al. The benefits of photodynamic therapy on vertebral bone are maintained and enhanced by combination treatment with bisphosphonates and radiation therapy. *J. Orthop. Res.* 2013; 31:1398–1405. [PubMed: 23625821]
114. Mitsunaga M, et al. Cancer cell-selective in vivo near infrared photoimmunotherapy targeting specific membrane molecules. *Nat Med.* 2011; 17:1685–1691. [PubMed: 22057348]
115. Doane TL, Chuang C-H, Chomas A, Burda C. Photophysics of silicon phthalocyanines in aqueous media. *Chem. Eur. J. of Chem. Phys.* 2013; 14:321–330.
116. Miller JD, et al. Photodynamic therapy with the phthalocyanine photosensitizer Pc 4: the case experience with preclinical mechanistic and early clinical-translational studies. *Toxicol. Appl. Pharmacol.* 2007; 224:290–299. [PubMed: 17397888]
117. Vrouenraets MB, et al. Targeting of Aluminum (III) Phthalocyanine Tetrasulfonate by Use of Internalizing Monoclonal Antibodies Improved Efficacy in Photodynamic Therapy. *Cancer Research.* 2001; 61:1970–1975. [PubMed: 11280754]
118. Vrouenraets MB, et al. Comparison of aluminium (III) phthalocyanine tetrasulfonate- and meta-tetrahydroxyphenylchlorin- monoclonal antibody conjugates for their efficacy in photodynamic therapy in vitro. *Int. J. Cancer.* 2002; 98:793–798. [PubMed: 11920653]
119. Mew D, Wat CK, Towers GH, Levy JG. Photoimmunotherapy: treatment of animal tumors with tumor-specific monoclonal antibody-hematoporphyrin conjugates. *J. Immunol.* 1983; 130:1473–1477. [PubMed: 6185591]

120. Stummer W, et al. Fluorescence-guided surgery with 5-aminolevulinic acid for resection of malignant glioma: a randomised controlled multicentre phase III trial. *Lancet Oncol.* 2006; 7:392–401. [PubMed: 16648043]
121. Rizvi I, et al. Photoimmunotherapy and irradiance modulation reduce chemotherapy cycles and toxicity in a murine model for ovarian carcinomatosis: perspective and results. *Isr. J. Chem.* 2012; 52:776–787. [PubMed: 23626376]
122. Meirelles K, et al. Human ovarian cancer stem/progenitor cells are stimulated by doxorubicin but inhibited by Mullerian inhibiting substance. *Proc. Natl. Acad. Sci. U.S.A.* 2012; 109:2358–2363. [PubMed: 22308459]

Figure 1.

Overview of the unique mechanisms of PDT-induced apoptosis. PDT directly damages antiapoptotic factors and the drug efflux pumps involved in classical drug resistance. The antiapoptotic BCL-2 family of proteins (e.g., BCL-2 and BCL-X_L) reside on the outer mitochondrial membrane and prevent mitochondria-mediated apoptosis by inhibiting the oligomerization and activation of the proapoptotic family members (e.g., BAX and BAK)³⁰. Proapoptotic proteins like BAX are in dynamic equilibrium between the cytosol and mitochondrial outer membrane but they are largely within the cytosol of healthy cells³¹. Therefore, mito-PDT is observed to predominately destroy antiapoptotic factors. Lyso-PDT induces the release of lysosomal proteases into the cytosol that can cleave BID (independent of caspase 8) to form truncated BID (tBID), which translocates to the mitochondria to promote the oligomerization of BAX and BAK. BAX and BAK oligomers form pore complexes that release cytochrome c and SMAC (second mitochondrion-derived activator of caspases) from the mitochondrial intermembrane space. Once released into the cytoplasm, cytochrome c forms a complex with apoptotic protease-activating factor 1 and procaspase 9, called the apoptosome, to activate caspase 9. SMAC, once released into the cytosol, promotes caspase activation through binding with IAPs (inhibitor of apoptosis proteins) and blocks their antiapoptotic activity. Once activated, the effector caspases (e.g., caspase 3 and caspase 7) carry out cellular degradation process to execute the apoptotic program.

Figure 2.

Cancer cells that are unresponsive to sustained gemcitabine chemotherapy are sensitive to BPD-PDT. (A) A panel of pancreatic adenocarcinoma cell lines contain gemcitabine unresponsive populations (17%–33%) even at extreme gemcitabine doses (up to 1 mM), while moderate BPD-PDT doses (1–6 J·cm⁻²·mM, where the units reflect the product of the light dose and the PS concentration, e.g., 10 J·cm⁻² × 0.25 mM BPD = 2.5 J·cm⁻²·mM) produce nearly complete cancer cell death. (B) BPD-PDT decreases BCL-X_L and increases the ratio of BAX-to-BCL-X_L toward a proapoptotic balance (data are the results from the quantification of western blots). (C) Insensitivity to gemcitabine (*top*), but not to BPD-PDT

(*bottom*), is increased in cells that are adherent to Matrigel basement membrane relative to traditional tissue culture (TC) conditions (NT indicates the no treatment control). Collectively, these results indicate the ability of PDT to bypass intracellular and extracellular cues, leading to gemcitabine resistance and indicate the emerging role of PDT for pancreatic cancer treatment. Adapted from Celli et al. (2011)⁷⁵.

Figure 3.

Photodynamic therapy reverses chemoresistance and synergizes with chemotherapy to destroy platinum-resistant disease. One-fold increase of cytotoxicity following photoimmunotherapy (antibody-PS conjugates) in combination with cisplatin (platinum chemotherapy)—versus cisplatin alone—in cisplatin-resistant (●) and cisplatin-sensitive (■) patient-derived samples and cell line cultures. A total of 19 solid tumor and/or ascites samples were collected from 14 ovarian cancer patients (ages 37–80, stages 1C–4), and 5 cancer cell line cultures were also included. Cisplatin resistance versus sensitivity refers to whether the patient had disease progression or recurrence within 6 months of platinum chemotherapy. Photoimmunotherapy induces a 12.9× enhancement in cytotoxicity against platinum resistant primary cultures (ranging from 1.5–52×) versus 1.8× for platinum sensitive cells. The asterisk indicates $P < 0.05$. Adapted from Duska et al. (1999)⁷³.

Figure 4.

PDT of locally advanced, inoperable pancreatic adenocarcinoma in humans. (A) Contrast-enhanced computed tomography (CT) scans from a patient undergoing BPD-PDT. The images show (*left*) a low attenuation mass in the head of the pancreas prior to treatment, (*center*) placement of a percutaneous needle for fiber optic light delivery into the tumor, and (*right*) a 2.67 cm³ zone of tumor necrosis 5 days post-PDT. (B) CT scans from a patient who qualified for and underwent a successful Whipple's tumor resection following PDT. The pre-PDT image (*left*) shows the tumor abutting the superior mesenteric artery (SMA; arrow); thus, this tumor was inoperable at presentation. Four weeks after PDT, the (*right*) follow-up CT scan for the same patient shows tumor reduction and minimal involvement with the SMA such that surgical resection could then be performed safely. Adapted from Huggett et al. (2014)¹⁰⁷.

Figure 5.

Concepts of tumor-targeted, activatable photoimmunotherapy (taPIT). (A) Activatable immunoconjugates for taPIT are comprised of multiple self-quenching, photocytotoxic chromophores conjugated to antibodies that target and neutralize the key molecules involved in tumorigenesis (e.g., EGFR). (B) Cellular activation of the immunoconjugates via receptor-mediated endocytosis and lysosomal degradation. (C) taPIT concept, in which the immunoconjugates accumulate selectively within the tumor nodules, are activated by cellular processing, and they inhibit molecular signaling and impart selective cytotoxicity to neoplasms upon irradiation while sparing neighboring vital tissues. (D) *Ex vivo* whole mount immunofluorescence image of a micrometastasis, where an anti-human cytokeratin antibody has been applied to visualize the human epithelial cancer cells (orange); an anti-mouse CD31 antibody labels the endothelial cells (green) and immunoconjugates are taken up and activated by tumor cells *in vivo* (red). taPIT enables the safe use of 50× photodynamic dose (PS × light dose) versus “always-on”, unconjugated BPD, and 17× photodynamic dose versus “always-on” PIT (using cetuximab-ce₆ conjugates) in a mouse model of peritoneal disseminated micrometastatic epithelial ovarian cancer²⁴. A single cycle of taPIT plus chemotherapy reduces the micrometastatic burden by 97% versus 3% for chemotherapy alone in the same mouse model^{24,121}, using human chemoresistant OVCAR5 cells^{64,122}. Wide-field taPIT was accomplished by administering scattering media

(Intralipid) to the peritoneal cavity and NIR laser light via a cylindrically diffusing fiber optic tip. Adapted from Spring et al. (2014)²⁴.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Cancer cell colony formation (murine hepatoma 1c1c7 cells) following lysosomal (NPe6, 660 nm)- plus mitochondrial (BPD, 690 nm)-PDT, versus the reverse sequence. Adapted from Kessel and Reiners Jr. (2014)³⁸.

PDT regimen	Clonogenicity (%)
No treatment control	100 ± 3
Low-dose mito-PDT	83 ± 5
Low-dose lyso-PDT	95 ± 2
Low-dose lyso-PDT ⇒ low-dose mito-PDT	17 ± 3 ^{* ‡}
Low-dose mito-PDT ⇒ low-dose lyso-PDT	58 ± 4 [*]

Data represent average ± SD.

^{*} Statistically different from controls;

[‡] statistically different from the result obtained with a reverse order of irradiation (P < 0.05).