Published in final edited form as:

Med Sci Sports Exerc. 2014 February ; 46(2): 232–238. doi:10.1249/MSS.0b013e3182a54d85.

One Bout of Exercise Alters Free-Living Postprandial Glycemia in Type 2 Diabetes

Douglas J. Oberlin¹, Catherine R. Mikus⁵, Monica L. Kearney¹, Pamela S. Hinton¹, Camila Manrique³, Heather J. Leidy¹, Jill A. Kanaley¹, R. Scott Rector^{1,2,4}, and John P. Thyfault^{1,2,4}

¹Department of Nutrition and Exercise Physiology, and Internal Medicine, University of Missouri, Columbia, MO

²Division of Gastroenterology and Hepatology, University of Missouri, Columbia, MO

³Division of Endocrinology, University of Missouri, Columbia, MO

⁴Harry S Truman Memorial VA Hospital, Columbia, MO

⁵Division of Cardiology, Duke University Medical Center, Durham, NC

Abstract

PURPOSE—Elevated postprandial glycemic excursions (PPG) are significant risk factors for cardiovascular disease in type 2 diabetes patients. Here we tested if and for how many meals a single bout of exercise would reduce PPG responses to subsequent meals in type 2 diabetes (T2D) patients using continuous glucose monitors (CGMS).

METHODS—We recruited 9 sedentary (<30 minutes/week of exercise) individuals with T2D (BMI: 36.0 ± 1.1 kg/m²; age 60.3 ± 1.0 years; HbA1c: 6.3 ± 0.2 %). The subjects consumed a eucaloric diet (51% carbohydrate, 31% fat, 18% protein) consisting of 3 meals, identical in composition, over a 2-day period while wearing CGMS in two different conditions (exercise (EX; one 60 minute bout at 60–75% of heart rate reserve performed prior to breakfast) vs. a sedentary (SED) condition). We quantified 24-h average glucose, PPG-AUC (4 h glucose AUC following meals) and PPG-2 h (2 hour post-prandial glucose).

RESULTS—EX significantly reduced average [glucose] during the first 24 hour period ($p=0.03$). EX caused a reduction in PPG-AUC ($p=0.02$) for all of the meals over the two days (main effect between conditions). Comparison between the EX and SED conditions at each meal revealed that EX reduced PPG-AUC following the second meal of day 1 (lunch) ($p=0.04$). PPG-2 h was not significantly different between EX and SED.

CONCLUSION—Although a single EX bout does lower 24-h average [glucose], it only significantly lowered PPG-AUC at the second meal following the bout suggesting that daily exercise may be needed to most effectively improve PPG at the advent of exercise training in T2D patients.

Corresponding Author: John P. Thyfault, PhD Associate Professor Departments of Nutrition and Exercise Physiology and Medicine - Division of Gastroenterology and Hepatology University of Missouri Columbia, MO 65201 Office phone #: 573-814-6000 x53742 Fax #: 573-882-0185 thyfaultj@missouri.edu.

Conflicts of Interest

The authors have no conflicts of interest to report.

Keywords

post-prandial glucose; glycemic control; exercise; type 2 diabetes; continuous glucose monitoring

Introduction

Hyperglycemia is linked to increased risk for cardiovascular disease (CVD) as well as all-cause mortality; thus, improved glycemic control is a critical target for diabetes management (6, 33). Individuals with type 2 diabetes (T2D) are, therefore, given the goal of maintaining glycated hemoglobin (HbA1c), a measure of long term glycemic control, below 7% (2, 31, 39). However, while HbA1c is a good indicator of average blood glucose levels over several months, it does not necessarily reflect the magnitude of changes in glucose levels over shorter periods of time, such as over the course of a day when meals of varying macro-nutrient content are consumed (19, 26, 27). During the course of a day, blood glucose levels can rise and fall several times depending upon frequency of feeding and type of food that is consumed as well as the level of physical activity (19, 26-28, 41). Fasting glucose levels, another common indicator of glycemic control, also fails to assess changes in glucose in response to feeding. Therefore, both HbA1c and fasting glucose may not adequately capture changes in blood glucose concentrations experienced over a typical day (19, 28, 41). Recently it has been reported that post-prandial (or post glucose challenge) glucose excursions (PPG) may be more tightly linked to risk for CVD than HbA1c or fasting glucose levels (6, 33). Therefore, therapies that limit the magnitude of PPG should also lower risk for CVD (4, 6, 7, 21, 33).

Exercise is a powerful method to improve long term glycemic control (3, 11, 14, 15, 18, 30, 32, 34, 35). It has been demonstrated that exercise can 1) act in an acute manner to increase glucose uptake in the absence of insulin, and 2) acutely improve insulin-stimulated glucose uptake in skeletal muscle (30). Moreover, exercise transiently improves insulin sensitivity for a prolonged period of time, suggesting that PPG may also be improved for several meals after one bout of exercise depending on the health status of the population (15, 16, 22). We have recently shown that 7 days of exercise training improves post-prandial glucose levels measured by a continuous glucose monitoring system (CGMS) in free living humans with T2D (24), however, the design did not allow us to determine the residual effects of each exercise bout on PPG. Due to the known effects of exercise to transiently improve skeletal muscle insulin sensitivity, we next questioned if, and for how many subsequent meals (identical in composition and caloric load), one bout of exercise would improve PPG measured by CGMS in free-living individuals with T2D. Importantly, PPG is driven by not just muscle insulin sensitivity but also by hepatic insulin sensitivity, pancreatic β -cell function, and other physiological responses. Although other labs have shown reduced concentrations in PPG and post-prandial insulin at a specific time points post-prandially (2.5 hours) (40), the total area under the glucose curve has not been assessed, nor did subjects consume the same foods at each meal following the exercise bout. We hypothesized that a single, morning exercise session would only improve PPG area under the curve for meals consumed during the same day (meals 1-3), and PPG area under the curve would subsequently return to pre-exercise levels on day 2 (meals 4-6). Our hypothesis for a one

day effect was based on 1). previous data showing one bout of exercise may not improve insulin sensitivity for as long as duration in T2D as in health cohorts, 2.) and that subsequent meals would replete muscle glycogen thus reducing the positive effects of exercise on PPG (9, 15, 22). If proven correct, this hypothesis would indicate that exercise on a daily basis may be needed to most effectively improve PPG in individuals with T2D, at least in the initial stages of training.

Materials and Methods

Subjects

Sedentary and low activity individuals with T2D were recruited in and around Columbia, MO, who on average, took fewer than 6,000 steps per day (pedometer) and did not participate in any formal exercise program (>30 minutes of planned exercise 2 times a week). Pedometers were worn over a 3-day period to measure daily step count. All subjects were non-smokers with a BMI between 30 and 42 kg/m² who did not have orthopedic limitations to exercise safely on a treadmill and stationary bike. They were weight stable ($\pm 5\%$) and had no changes in medication for at least 3 months before entering the study. In addition, the subjects had controlled diabetes with HbA1c < 7.5% with no insulin use, and no reported advanced retinopathy or neuropathy. Other exclusion criteria included breakfast skippers, pregnancy, sleep perturbations, night shift workers, people who have recently traveled across more than two time zones, or individuals with irregular daily schedules to avoid odd sleep or eating patterns which may have conflicted with the study design. All women enrolled in the study were post-menopausal. All subjects signed an informed consent which was approved by the University of Missouri Institutional Review Board.

After the consent meeting, the subjects came to the exercise physiology lab during the morning after a 10 hour fast for measurement of height, weight, blood pressure, fasting blood glucose, glycated hemoglobin (HbA1c) and blood lipids (total cholesterol, LDL, HDL, and triglycerides). HbA1c was analyzed on a Siemens DCA Vantage analyzer using blood drawn in a heparin tube. Blood lipids (cholesterol, triglycerides, HDL, LDL, and total cholesterol) and fasting glucose were measured by a commercial laboratory as done previously (24). Subjects were then given a diet log and a pedometer to measure dietary consumption and daily steps, respectively, over a 3 day period. On another visit, the subjects had their body composition measured via dual-energy X-ray absorptiometry [DEXA; Hologic QDR 4500A (Bedford, MA)]. The subjects then performed an exercise stress test to determine maximal oxygen consumption ($\text{VO}_{2\text{peak}}$), maximal heart rate, and to screen for any potential cardiac abnormalities via EKG. The exercise stress test was performed on a treadmill using a standardized Bruce protocol as performed previously (24). During the test, respiratory gases were measured by a metabolic cart (Parvo Medics True One 2400 Metabolic Measurement System), and cardiovascular function was monitored using a 12 lead EKG (Quinton Q-Stress v3.5 Exercise Test Monitor). Criteria for a maximal test were three of the following: volitional exhaustion, perceived exertion of 17 or greater, respiratory exchange ratio of greater than 1.0, or a leveling off or slight decrease in oxygen consumption. The EKG data from each exercise stress test were reviewed by a cardiologist to ensure that the participants could safely participate in an exercise session. There was a

five to fifteen day washout after the $\text{VO}_{2\text{peak}}$ test before the subjects began the study protocol.

Study Design

The study incorporated a randomized cross-over design and consisted of measuring minute to minute interstitial glucose concentrations by CGMS during a 2 day sedentary condition (subjects maintained normal daily activities) and another 2 day condition in which the subjects exercised during the morning of the first day (followed by maintaining normal daily activities). Therefore, subjects served as their own controls. We have previously used CGMS to measure glycemic control in free living subjects who are healthy as well as those with T2D (24, 25). During each condition, the subjects consumed the study diet which consisted of 3 meals a day representing breakfast (8 am), lunch (1 pm), and dinner (6 pm) for 4 days. The first two days of the diet were to acclimate the subject to the new diet. The following 2 days of the standard diet coincided with CGMS measurements to ensure that any changes to glycemic control were due to exercise and not due to dietary changes. Figure 1 shows the 4-day study period which the subjects repeated twice, once performing an EX bout in the morning prior to breakfast and once while remaining sedentary. During the sedentary condition the subjects continued their typical (sedentary) physical activity, which was verified using a Walk 4 Life Duo pedometer (Plainfield, IL) and an accelerometer (Body Media Sense Wear armband body monitoring system; Pittsburgh, PA). Both pedometer and accelerometer were used primarily to assess physical activity within subjects, between conditions. Validation studies on the accuracy of sensewear to measure energy expenditure have provided mixed results (12, 17), however, the device allowed a basis of comparison between the two conditions and showed that exercise increased energy expenditure. There was a 5-15 day washout period between the 2 conditions of the study.

Exercise Testing and Exercise Session

Graded exercise tests to measure $\text{VO}_{2\text{peak}}$ were performed on a treadmill using a Bruce Protocol as done previously (23). The exercise (EX) bout consisted of 60 minutes of aerobic exercise broken into three 20-minute sections starting at approximately 6:30 AM. This included 20 minutes on a treadmill, 20 minutes on a stationary cycle, and another 20 minutes on a treadmill. The exercise session was broken into these segments because the subjects could not do 60 minutes on the treadmill at this intensity without needing to sit down. The exercise intensity was within five beats per minute of 60% of HRR for the duration of the exercise bout (as determined from a previous graded exercise stress test). Intensity was controlled during the exercise session by adjusting speed or grade on the treadmill or adjusting resistance on the stationary cycle, to maintain the target heart rate throughout the entire exercise bout. The exercise prescription falls in line with the American Diabetes Association and American College of Sports Medicine dual recommendation of 150 minutes per week at an intensity of 40-60% $\text{VO}_{2\text{peak}}$ for patients with T2D (1). In addition, the same exercise prescription was used in a previous study from our laboratory which showed a decrease in PPG during 5-7 days of exercise training in previously sedentary subjects with T2D (24). Subjects were instructed to take their medications as prescribed throughout both conditions, and were monitored post exercise to avoid hypoglycemia. Thus, in this study we

wanted to determine if and for how many meals one bout of exercise prescribed at the same intensity and duration would improve post-prandial glycemic responses to subsequent meals.

Study Diet

As stated previously, the subjects consumed the same study diet in both phases (Table 1). The diet was prepared and packed out by study staff. The subjects were instructed to eat the meals at the same times each day and to allow 5 hours between meals. Every meal had the same nutrient composition, caloric content, and contained the exact same food items prepared as either breakfast, lunch, or dinner. Breakfast was a potato hash with seasoned ground beef topped with salsa and cheese served with buttered toast, applesauce and a juice drink. Lunch was mini cheeseburgers with salsa mixed into the patties and baked french fries served with a side of applesauce and a juice drink. Dinner was a mini-meatloaf with salsa and cheese baked in and mashed potatoes served with garlic toast, applesauce, and a juice drink. Subject were instructed to leave uneaten foods in containers when returning them to the lab, allowing us to assess food compliance to a certain degree. True compliance to the diet was not directly assessed as we did not monitor them eating the meals directly.

The macronutrient distribution based on energy content was 51.4% carbohydrate, 30.9% fat, and 17.8% protein for the total energy content for each meal (Table 1). The glycemic load of each meal was approximately 55. The daily energy requirement for each subject was estimated using the Harris-Benedict equation and also quantified using a three-day dietary record completed by the subject. The three day mean was then averaged with the Harris-Benedict estimate to determine individual energy requirements, and avoid deficits due to under-reporting in the dietary records. From this information, the subjects were provided a diet containing 1600, 1800, 2000, 2200, or 2400 kcals per day, whichever kcal level was within 100 kcals of their predicted requirements. For example, if a person was estimated at 2063 kcal, they would receive the 2000 kcal diet while an individual estimated at 2115 would receive a 2200 kcal diet. The diet was designed to simulate a typical American diet and provide consistent diet composition between meals and across subjects; it was not meant to serve as an intervention, although it did differ from their normal routine.

The subjects were given a log sheet to track when they ate their meals. They were instructed to eat all of the portions within a 15 to 20 minute timeframe. Subjects also noted when they went to bed at night and when they got up in the morning. No additional calories were given to replace those lost in the exercise session, which created a deficit. Although this is a known confounder, replacing the calories would have lead to a mismatch of macronutrients between conditions, which would have also been a confounding variable. Therefore, the meals were kept constant through both conditions.

Postprandial glycemic control

A Medtronic iPro CGMS (Minneapolis, MN) monitor was attached to the subject's abdomen with a probe inserted beneath the skin, and the monitor was attached and taped down with Smith & Nephew IV3000 adhesive pads the night before the first measurement day. The CGMS was then worn for 2 consecutive days. While the CGMS was worn, the subjects recorded four daily blood glucose levels with an Accu-Chek Compact Plus glucometer

which were then used to calibrate the CGMS. We used the CGMS glucose concentration data to quantify 24 h average glucose for each day in each condition. We also used the glucose concentrations along with subject recorded meal times to calculate the PPGAUC for each of the 6 meals [quantified as the 4-h glucose area under the curve (AUC) response following each meal]. Blood glucose values recorded every 15 minutes during the 4 hour period were used to calculate total AUC. We also quantified 2-hour post-prandial glucose concentration for each of the 6 meals (PPG-2h) because it is a strong predictor of cardiovascular events (6).

Physical activity monitoring

Subjects wore pedometers and BodyMedia SenseWear Pro II (Pittsburgh, PA) arm band accelerometers during both phases of the study. Data from the accelerometers (estimated energy expenditure) and pedometers (daily steps) were collected.

Statistical analysis

Statistical analysis was performed with SPSS Version 15 (Armonk, NY) on PPG-AUC, PPG-2 h, average 24 hour glucose concentration, estimated caloric expenditure from accelerometer, and daily steps using a two-way repeated measures ANOVA. The two levels were condition (EX or SED) and meal (1-6) for all but the average 24 hour glucose, steps, and energy expenditure, for which the two levels were condition and day (1 or 2). Post-hoc analyses were conducted using Fisher-LSD. The level of statistical significance was set at a P value of 0.05, and the data are reported as means $\pm$ SE.

Results

Nine subjects completed the study. Baseline anthropometric characteristics, energy intake, blood chemistry, and aerobic fitness data are shown in Table 2. Medication usage for the subjects were the following: 6/9 were taking metformin (biguanides), 9/9 were taking statins, 6/9 were taking ACE inhibitors, and 1/9 were taking angiotensin receptor blockers, and 3/9 were taking diuretics.

Figure 1 depicts the order of events for the study. The exercise sessions were all supervised and were performed at an intensity of 60% HRR (similar to 60% $\text{VO}_{2\text{peak}}$) for 60 minutes. All subjects performed 60 minutes of exercise at an average percent heart rate reserve of 58.1 ± 5.5 . Table 3 shows that the subjects had a statistically significant increase in both steps and energy expenditure on EX day.

Postprandial Glycemic Control

EX lowered average glucose concentration significantly over the first 24-h period compared to SED (EX: 5.98 ± 0.049 vs. SED: 6.62 ± 0.73 mmol/L; $p=0.038$), but differences were not observed in day 2 (EX: 6.59 ± 0.78 vs. SED: 6.62 ± 0.53 mmol/L; $p=0.3$). Figure 2 shows the average glucose responses measured every 5 minutes by CGMS during the entire first 24-h of EX or SED conditions, demonstrating lower circulating glucose levels in the EX vs. SED condition.

Repeated measures ANOVA revealed a main effect of EX to lower PPG-AUC across all 6 meals compared to SED ($p=0.015$) (Figure 3a); this lower PPGAUC was primarily due to the effects seen during day 1. The post hoc analysis between EX and SED at each of the 6 meals revealed that only meal 2 was significantly lower in the EX compared to SED condition (14% lower; $p=0.04$) (Figure 3a). Although other meals showed trends to be lower in the EX vs. SED condition (Meal 1 EX is 15% lower than SED ($p=0.08$); Meal 3 EX is 12.8% lower than SED ($p=0.11$); Meal 6 EX is 19% lower than SED ($p=0.16$)) they were not statistically different. Additionally, there was also a main-effect for the 2-h PPG to be lower during the exercise compared to the sedentary condition ($p=0.005$) (Figure 3b). However, none of the meals were different between EX and SED after post hoc analysis.

Discussion

The current study determined if and for how many subsequent meals one bout of exercise improved PPG in individuals with T2D. Importantly, this was the first study that fed 6 controlled meals (same type and energy content of food at each meal) over a two day period allowing us to ensure that EX induced differences in PPG were not impacted by type or quantity of food. We found that a single bout of exercise significantly lowered PPGAUC responses to the second meal following the exercise bout by 15%, while a similar, but non-significant effect was also shown at the first (15%) and third (12%) meal to also be lowered. These changes in PPG resulted in a significant 10% reduction in average blood glucose concentrations among the subjects during the first 24-h after the exercise session, similar to findings by other groups (13, 20); however, the exercise induced impact on average 24-h glucose concentration was no longer present on day 2. Because the effects of exercise were undetectable by the second day, these data would suggest that exercise should be performed every day to have an optimal effect on postprandial glucose responses at least at the initiation of an exercise program for individuals with T2D. Of note, a previous study from van Dijk and colleagues did find that that one 60 min bout of exercise improved 24 h glucose values on the second day, but key differences between the studies are noted below.

A recent study by Van Dijk et al. (40) showed that a single exercise bout (30 min) each day was sufficient to lower the 2.5 h glucose concentration after breakfast in individuals with T2D. The same study also showed that one 60 min exercise bout lowered post 2.5 hour breakfast glucose on both the day of the bout and the following morning. Thus, they showed that one exercise bout that was the same volume of ours did impact postprandial glucose values on the second day. Key differences exist between the studies which may have impacted the outcomes. First, as already mentioned, our study used controlled meals comprised of the same foods and energy content for each meal limiting the impact that macronutrient content or caloric loads could have altered glycemic responses. Differences in the glycemic control of subjects could have also impacted results. Our T2D patients had good glycemic control ($HbA1c=6.3\%$) compared to the subjects in the van Dijk study ($HbA1c=7.2\%$) (40) suggesting that our subjects may have not had as much room for improvement. In addition, our subjects were primarily only taking metformin as glucose lowering medication, whereas the study from van Dijk used some individuals who were on metformin and some who were also receiving insulin therapy which may have positively interacted with exercise responses and provided even greater improvements in glycemia.

Finally, the timing of the exercise bout could have also impacted results. Our subjects performed an early morning exercise bout and then consumed meal 1 approximately 2 hours later, while the van Dijk study performed exercise after breakfast leaving only two meals to be consumed during the post exercise period of day 1.

While the current study demonstrates that a single exercise bout transiently improves PPG, it is not known whether prolonged exercise training would result in more persistent effects in type 2 diabetics following each exercise bout. In an acute training study, the effects observed are most likely due to either a change in energy homeostasis, or due directly to skeletal muscle contractions which result in a period of enhanced insulin sensitivity that has been documented to last for up to 48 hours (38). However, following prolonged exercise training, many other adaptations may contribute to witnessed improvements in glycemic control (8), such as increased GLUT-4 content (10), increased capillary density in skeletal muscle and improved vascular function (29), and altered body composition (8). Although these adaptations may allow a greater volume of glucose to be taken up by the working tissue, it is unknown whether they significantly change the duration of improvement in skeletal muscle insulin sensitivity and glucose uptake resulting in prolonged reductions in PPG following an exercise bout. Furthermore, exercise training induced adaptations that improve control of hepatic glucose output (18) and pancreatic β -cell function (36) could also play a role in potentiating the lasting impact of an exercise bout on glycemic control during subsequent postprandial periods. These concepts deserve further study.

In the current study, we attempted to control as many variables as possible without keeping subjects in a lab setting for the duration of the study. The number of steps per day and estimated caloric expenditure were both monitored throughout the exercise and sedentary conditions. The significant difference in steps per day and estimated caloric expenditure between the first day of the exercised (including the exercise session) and the first day of the sedentary condition confirms that the subjects did not reduce steps or activity following the exercise bout to compensate for the exercise session in the morning. As previously stated, the diet provided in this study maintained the same food items at every meal, reducing the likelihood of differences in macronutrient content or glycemic load affecting the changes seen in glycemic control. This is a unique feature of this study and we are not aware of a previous study utilizing this approach. Further, since the diet was provided and constant throughout the study, the observed changes that were seen in glycemic control were most likely due to the increased physical activity from the exercise session. However, a limitation of controlling the diet was that the subjects were not re-fed calories to compensate for the energy expended during the exercise bout. This confounds whether the effects seen in the current study were effects of muscle contraction, energy deficit, or a combination of both (5, 37).

In conclusion, the current study demonstrated that a single exercise bout is sufficient to reduce 24 hour average blood glucose concentrations during the first day (day of the exercise bout) as well as post-prandial glycemic AUC for a single meal (meal 2) in individuals with T2D. This study confirms other recent findings which have shown a single bout of endurance exercise or short duration, high intensity exercise reduces average blood glucose concentrations in individuals with T2D for 24 hours and also reduces post-prandial

glucose concentrations (13, 40). Our findings suggest that for optimal blood glucose control it is important for individuals with T2D to perform exercise on a daily basis, particularly at the onset of exercise training.

Acknowledgments

Thank you to JA Fletcher, LJ Boyle, Peggy Nigh, and RA Silverstein for their contributions to performing the tests and preparing the foods in the current study and to Dr. Kevin Dellsperger for reading EKGs. Dr. R. Scott Rector is supported by a VA CDA2 Award (VA-CDA-IK2 BX001299-01) and Dr. John Thyfault is supported by NIH R01DK088940. No author has any personal or financial conflict of interest. This study was funded by the Department of Nutrition and Exercise Physiology at the University of Missouri. The results of the current study do not constitute endorsement by the American College of Sports Medicine.

References

1. American College of Sports Medicine and the American Diabetes Association: joint position statement. Exercise and type 2 diabetes. *Medicine and science in sports and exercise*. 2010; 42(12): 2282–303. [PubMed: 21084931]
2. Barr RG, Nathan DM, Meigs JB, Singer DE. Tests of glycemia for the diagnosis of type 2 diabetes mellitus. *Ann Intern Med*. 2002; 137(4):263–72. [PubMed: 12186517]
3. Black SE, Mitchell E, Freedson PS, Chipkin SR, Braun B. Improved insulin action following short-term exercise training: role of energy and carbohydrate balance. *J Appl Physiol*. 2005; 99(6):2285–93. [PubMed: 16081626]
4. Bonora E, Muggeo M. Postprandial blood glucose as a risk factor for cardiovascular disease in Type II diabetes: the epidemiological evidence. *Diabetologia*. 2001; 44(12):2107–14. [PubMed: 11793012]
5. Braun B, Brooks GA. Critical importance of controlling energy status to understand the effects of “exercise” on metabolism. *Exercise and sport sciences reviews*. 2008; 36(1):2–4. [PubMed: 18156946]
6. Cavalot F, Pagliarino A, Valle M, et al. Postprandial blood glucose predicts cardiovascular events and all-cause mortality in type 2 diabetes in a 14-year follow-up: lessons from the San Luigi Gonzaga Diabetes Study. *Diabetes care*. 2011; 34(10):2237–43. [PubMed: 21949221]
7. Cederberg H, Saukkonen T, Laakso M, et al. Postchallenge glucose, A1C, and fasting glucose as predictors of type 2 diabetes and cardiovascular disease: a 10-year prospective cohort study. *Diabetes care*. 2010; 33(9):2077–83. [PubMed: 20573752]
8. Colberg SR, Albright AL, Blissmer BJ, et al. Exercise and type 2 diabetes: American College of Sports Medicine and the American Diabetes Association: joint position statement. *Exercise and type 2 diabetes. Medicine and science in sports and exercise*. 2010; 42(12):2282–303. [PubMed: 21084931]
9. Cusi K, Maezono K, Osman A, et al. Insulin resistance differentially affects the PI 3- kinase- and MAP kinase-mediated signaling in human muscle. *J Clin Invest*. 2000; 105(3):311–20. [PubMed: 10675357]
10. Dela F, Ploug T, Handberg A, et al. Physical training increases muscle GLUT4 protein and mRNA in patients with NIDDM. *Diabetes*. 1994; 43(7):862–5. [PubMed: 8013748]
11. Frosig C, Richter EA. Improved insulin sensitivity after exercise: focus on insulin signaling. *Obesity (Silver Spring)*. 2009; 17(Suppl 3):S15–20. [PubMed: 19927140]
12. Fruin ML, Rankin JW. Validity of a multi-sensor armband in estimating rest and exercise energy expenditure. *Medicine and science in sports and exercise*. 2004; 36(6):1063–9. [PubMed: 15179178]
13. Gillen JB, Little JP, Punthakee Z, Tarnopolsky MA, Riddell MC, Gibala MJ. Acute high-intensity interval exercise reduces the postprandial glucose response and prevalence of hyperglycaemia in patients with type 2 diabetes. *Diabetes Obes Metab*. 2012; 14(6):575–7. [PubMed: 22268455]
14. Hawley JA. Exercise as a therapeutic intervention for the prevention and treatment of insulin resistance. *Diabetes Metab Res Rev*. 2004; 20(5):383–93. [PubMed: 15343584]

15. Henriksen EJ. Invited Review: Effects of acute exercise and exercise training on insulin resistance. *J Appl Physiol.* 2002; 93(2):788–96. [PubMed: 12133893]
16. Holloszy JO. A forty-year memoir of research on the regulation of glucose transport into muscle. *Am J Physiol Endocrinol Metab.* 2003; 284(3):E453–67. [PubMed: 12556346]
17. Johannsen DL, Calabro MA, Stewart J, Franke W, Rood JC, Welk GJ. Accuracy of armband monitors for measuring daily energy expenditure in healthy adults. *Medicine and science in sports and exercise.* 2010; 42(11):2134–40. [PubMed: 20386334]
18. Kirwan JP, Solomon TP, Wojta DM, Staten MA, Holloszy JO. Effects of 7 days of exercise training on insulin sensitivity and responsiveness in type 2 diabetes mellitus. *Am J Physiol Endocrinol Metab.* 2009; 297(1):E151–6. [PubMed: 19383872]
19. Kuenen JC, Borg R, Kuik DJ, et al. Does glucose variability influence the relationship between mean plasma glucose and HbA1c levels in type 1 and type 2 diabetic patients? *Diabetes care.* 2011; 34(8):1843–7. [PubMed: 21700921]
20. Manders RJ, Van Dijk JW, van Loon LJ. Low-intensity exercise reduces the prevalence of hyperglycemia in type 2 diabetes. *Medicine and science in sports and exercise.* 2010; 42(2):219–25. [PubMed: 19927038]
21. Meigs JB, Nathan DM, D'Agostino RB Sr, Wilson PW. Fasting and postchallenge glycemia and cardiovascular disease risk: the Framingham Offspring Study. *Diabetes care.* 2002; 25(10):1845–50. [PubMed: 12351489]
22. Mikines KJ, Sonne B, Farrell PA, Tronier B, Galbo H. Effect of physical exercise on sensitivity and responsiveness to insulin in humans. *The American journal of physiology.* 1988; 254(3 Pt 1):E248–59. [PubMed: 3126668]
23. Mikus CR, Boyle LJ, Borengasser SJ, et al. Simvastatin impairs exercise training adaptations. *Journal of the American College of Cardiology.* In press.
24. Mikus CR, Oberlin DJ, Libla J, Boyle LJ, Thyfault JP. Glycaemic control is improved by 7 days of aerobic exercise training in patients with type 2 diabetes. *Diabetologia.* 2012; 55(5):1417–23. [PubMed: 22311420]
25. Mikus CR, Oberlin DJ, Libla JL, Taylor AM, Booth FW, Thyfault JP. Lowering physical activity impairs glycemic control in healthy volunteers. *Medicine and science in sports and exercise.* 2012; 44(2):225–31. [PubMed: 21716152]
26. Monnier L, Colette C, Dunseath GJ, Owens DR. The loss of postprandial glycemic control precedes stepwise deterioration of fasting with worsening diabetes. *Diabetes care.* 2007; 30(2):263–9. [PubMed: 17259492]
27. Monnier L, Lapinski H, Colette C. Contributions of fasting and postprandial plasma glucose increments to the overall diurnal hyperglycemia of type 2 diabetic patients: variations with increasing levels of HbA(1c). *Diabetes care.* 2003; 26(3):881–5. [PubMed: 12610053]
28. Nichols GA, Hillier TA, Brown JB. Progression from newly acquired impaired fasting glucose to type 2 diabetes. *Diabetes care.* 2007; 30(2):228–33. [PubMed: 17259486]
29. Padilla J, Simmons GH, Bender SB, Arce-Esquivel AA, Whyte JJ, Laughlin MH. Vascular effects of exercise: endothelial adaptations beyond active muscle beds. *Physiology.* 2011; 26(3):132–45. [PubMed: 21670160]
30. Richter EA, Mikines KJ, Galbo H, Kiens B. Effect of exercise on insulin action in human skeletal muscle. *J Appl Physiol.* 1989; 66(2):876–85. [PubMed: 2496078]
31. Riddle MC, Rosenstock J, Gerich J. The treat-to-target trial: randomized addition of glargine or human NPH insulin to oral therapy of type 2 diabetic patients. *Diabetes care.* 2003; 26(11):3080–6. [PubMed: 14578243]
32. Rockl KS, Witczak CA, Goodyear LJ. Signaling mechanisms in skeletal muscle: acute responses and chronic adaptations to exercise. *IUBMB Life.* 2008; 60(3):145–53. [PubMed: 18380005]
33. Saydah SH, Miret M, Sung J, Varas C, Gause D, Brancati FL. Postchallenge hyperglycemia and mortality in a national sample of U.S. adults. *Diabetes care.* 2001; 24(8):1397–402. [PubMed: 11473076]
34. Sigal RJ, Kenny GP, Boule NG, et al. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med.* 2007; 147(6):357–69. [PubMed: 17876019]

35. Sigal RJ, Kenny GP, Wasserman DH, Castaneda-Sceppa C. Physical activity/exercise and type 2 diabetes. *Diabetes care*. 2004; 27(10):2518–39. [PubMed: 15451933]
36. Solomon TP, Haus JM, Kelly KR, Rocco M, Kashyap SR, Kirwan JP. Diabetes care. 2010; Improved pancreatic beta-cell function in type 2 diabetic patients after lifestyle-induced weight loss is related to glucose-dependent insulinotropic polypeptide. 33(7):1561–6. [PubMed: 20200305]
37. Stephens BR, Granados K, Zderic TW, Hamilton MT, Braun B. Effects of 1 day of inactivity on insulin action in healthy men and women: interaction with energy intake. *Metabolism: clinical and experimental*. 2011; 60(7):941–9. [PubMed: 21067784]
38. Thyfault JP. Setting the stage: possible mechanisms by which acute contraction restores insulin sensitivity in muscle. *American journal of physiology. Regulatory, integrative and comparative physiology*. 2008; 294(4):R1103–10.
39. Umpierre D, Ribeiro PA, Kramer CK, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA : the journal of the American Medical Association*. 2011; 305(17):1790–9. [PubMed: 21540423]
40. van Dijk JW, Tummers K, Stehouwer CD, Hartgens F, van Loon LJ. Exercise therapy in type 2 diabetes: is daily exercise required to optimize glycemic control? *Diabetes care*. 2012; 35(5):948–54. [PubMed: 22399700]
41. Wolever TM, Chiasson JL, Csima A, et al. Variation of postprandial plasma glucose, palatability, and symptoms associated with a standardized mixed test meal versus 75 g oral glucose. *Diabetes care*. 1998; 21(3):336–40. [PubMed: 9540012]

Figure 1.

Study design for the exercise (EX) vs. sedentary (SED) phases of the study. Subjects repeated two separate phases in which they wore CGMS and consumed provided meals 3 times per day. In both phases, CGMS was placed on the evening of the second wash-in day. On the morning of day 1, subjects either performed EX or remained SED. Glucose levels were measured by CGMS and 3 prepared meals were consumed per day (6 meals total) over next 2 days.

Figure 2.

Glucose measured every 5 minutes by CGMS during the first 24 h period. Timepoints for breakfast (8 am), lunch (1 pm), and dinner (6 pm) are approximate times when subjects consumed the 3 meals.

Figure 3.

A) Postprandial glucose area under the curve (PPG-AUC) responses to 6 meals under EX and SED conditions: (†) Main effect of EX to lower PPG-AUC ($P=0.015$) across all meals compared to SED; (*) EX significantly lowered PPG-AUC response at meal 2 ($P=0.04$) compared to SED. Data is shown as means $\pm$ SE. **B)** PPG-2h responses to 6 meals following EX vs. SED conditions: (†) Main effect of EX to lower PPG-AUC ($P=0.005$) across all meals. Data is shown as means $\pm$ SE

Table 1

Composition of subject meals for a 2000 kcal diet

Food item	Amount in meal (g)	Kcals	CHO (g)	Fat (g)	Protein (g)
White sandwich bread	52.00	137.00	28.00	1.00	4.00
Potatoes	140.00	105.95	24.59	0.00	1.89
Salsa	33.00	8.00	2.00	0.00	0.00
Ground beef, 93/7	101.00	147.89	0.00	7.21	20.74
Salted butter, light	14.00	45.00	0.00	5.00	0.00
Olive oil, extra virgin	4.75	42.00	0.00	4.67	0.00
Applesauce	98.00	68.44	17.11	0.00	0.00
Cheese, medium cheddar	14.00	57.00	0.00	5.00	3.00
Fruit juice	125.00	56.00	14.00	0.00	0.00
Totals	581.75	667.28	85.70	22.88	29.63

Foods and quantities in the study diet for a 2000 kcal/per day diet. This table shows the amount of each food item in one meal for the study diet at the 2000 kcal level. The amount of food was adjusted for each calorie level to achieve 200 kcal differences.

Table 2

Baseline subject characteristics

Anthropometric Characteristics	
Age (years)	60.3 ± 1.0
Sex	5 Females / 4 Males
Body Mass Index (kg/m ²)	36.0 ± 1.1
Body Fat (%)	39.6 ± 1.9
Total Cholesterol (mg/dl)	172.9 ± 15.8
LDL (mg/dl)	88.8 ± 12.6
HDL (mg/dl)	48.1 ± 2.7
Triglycerides (mg/dl)	179.7 ± 23.6
Fasting glucose (mmol/l)	6.5 ± 0.6
Hemoglobin A1c (%)	6.3 ± 0.2
Relative VO _{2peak} (ml/kg/min)	20.4 ± 0.7
Average daily caloric intake (self-reported, kcals)	1937.8 ± 176.5

Anthropometric characteristics of the subjects. Baseline characteristics including lipid profile, glucose, and hemoglobin A1c results, metabolic cart readouts from an exercise stress test, and the average daily caloric intake of the subjects for the study participants are shown. Data is shown as means ± SE.

Table 3

Subject activity

Activity Data		
Variable	Sedentary	Exercised
Accelerometer Energy Expenditure (kcal/day)	2343.0 $\pm$ 148.2	2659.2 $\pm$ 50.7 *
Pedometer Steps	3621.1 $\pm$ 391.3	5489.5 $\pm$ 845.0

Physical activity levels. Average daily steps and energy expenditure during EX and SED conditions.

Data is shown as means $\pm$ SE.

* indicates a statistically significant difference ($p < 0.05$).