Published in final edited form as:

J Psychosom Res. 2015 September ; 79(3): 243–245. doi:10.1016/j.jpsychores.2015.06.004.

The Development of Depressive Symptoms During Medical Internship Stress Predicts Worsening Vascular Function

Jess G. Fiedorowicz, M.D., Ph.D.^{a,b,c}, Vicki L. Ellingrod, Pharm. D.^{d,e}, Mariana J. Kaplan, M.D.^g, and Srijan Sen, M.D., Ph.D.^{e,f,h}

^aDepartment of Psychiatry, The University of Iowa, Iowa City, IA

^bDepartment of Internal Medicine, Carver College of Medicine, The University of Iowa, Iowa City, IA

^cDepartment of Epidemiology, College of Public Health, The University of Iowa, Iowa City, IA

^dCollege of Pharmacy, University of Michigan, Ann Arbor, MI

^eDepartment of Psychiatry, University of Michigan, Ann Arbor, MI

^fMolecular and Behavioral Neuroscience Institute, Department of Psychiatry, University of Michigan, Ann Arbor, MI

^gSystemic Autoimmunity Branch, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health, Bethesda, MD

Abstract

Objective—We sought to prospectively determine whether the onset of internship stress and any subsequent depression alters physiological markers of early vascular disease

Methods—We explored potential mechanisms linking stress and depression to vascular disease in a prospective cohort of 37 participants exposed to medical internship stress, an established precipitant of depressive symptomatology.

Results—Change in depressive symptom score from baseline over one year of internship stress was inversely correlated with change in the reactive hyperemia index (RHI), a measure of peripheral endothelial function ($r=0.41$, $p=0.01$). The change in depressive symptoms in the first six months of internship was similarly related to change in RHI over one year ($r=0.38$, $p=0.02$). While the development of depressive symptoms did not significantly impact changes in endothelial progenitor cells (EPCs), EPCs did significantly decrease with the year of internship stress (11.9 to 3.4 cells/ml blood; $p=0.01$).

Conclusion—Endothelial function may be a critical link between stress, depression, and cardiovascular disease and a feasible surrogate outcome for prospective studies.

^hCorresponding author (Dr. Sen): srijan@umich.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

biological makers; cardiovascular diseases; endothelial function; major depressive disorder

1. INTRODUCTION

Depression has been consistently linked with up to a two-fold greater risk of cardiovascular morbidity and mortality [1, 2]. This large association between depression and heart disease has been replicated in representative samples worldwide [1] and independent of a variety of confounding variables [3]. The development and persistence of depressive symptoms have even been purported to exert adverse vascular effects [4].

Understanding the mechanisms linking depressive symptomatology and cardiovascular function holds promise for reducing this excess morbidity and mortality. Animal models of stress and depression have demonstrated related impairments in endothelial function [5, 6]. Endothelial dysfunction precedes the development of atherosclerotic lesions and thus represents an important intermediate phenotype for the study of physiological mechanisms contributing to cardiovascular disease. Depression has been associated with impaired endothelial function and lower levels of endothelial progenitor cells (EPCs) [7–15]. All but one of these studies utilized cross-sectional designs, with the sole exception following a high risk cohort based on family history [7]. Prospective study is essential to establish a temporal relationship between change in depressive symptomatology and vascular outcomes. We sought to study the relationship between depressive symptoms and changes in cardiovascular function in those at risk due to environmental stress.

Medical internship, the first year of professional medical training, represents a unique situation where we can accurately predict a dramatic increase in stress. The Intern Health Study is a prospective cohort study that follows medical trainees from the low stress period prior to internship through the stressful internship year. In early cohorts of the project, we found that rates of depression increased from 4% before internship to 25% during internship [16]. Here, using medical internship as a model, we seek to determine whether the onset of internship stress and any subsequent depression alters physiological markers of early vascular disease: EPCs and endothelial function.

2. METHODS**2.1. Sample**

We recruited medical interns who were participating in the Intern Health Study, a prospective cohort designed to assess predictors for the development of depressive symptoms during internship, which has been described in detail elsewhere [16, 17]. Fifty-three study participants who graduated from the University of Michigan and University of Iowa medical schools in 2012 and 2013, and matched to stay at these institutions for internship were invited to take part in this arm of the Intern Health Study, designed to assess vascular function, with 37 choosing to participate (70% participation rate). The study was

approved the University of Michigan and the University of Iowa Institutional Review Boards and all persons participating in this study arm provided written informed consent.

2.2. Outcome Assessment

Depressive symptoms were measured using the 9-item self-report Patient Health Questionnaire (PHQ-9), at baseline under lower stress conditions and then at 3-month intervals during the stressful internship year [18]. A PHQ-9 score >10 has good sensitivity and specificity for a diagnosis of major depressive disorder [16]. Endothelial function was assessed non-invasively, with participants seated, using finger plethysmography with the EndoPAT2000 (Itamar Medical, Caesarea, Israel). A reactive hyperemia index (RHI), which is predictive of subsequent cardiovascular events [19], was determined by assessing vasodilation following a 5-minute occlusion of the brachial artery and comparing pulse-wave readings through pneumatic finger probes before and after occlusion relative to an unoccluded control arm. To measure EPCs, flow cytometry was utilized to count CD34+ and CD133+ labeled cells in the peripheral blood cells using established methods [8]. A decrease in EPCs has been linked to an impaired ability to repair arteries and increased risk of myocardial infarctions and strokes [20]. Endothelial function and EPCs were assessed at baseline and 12 months.

2.3. Statistical Analysis

SPSS Version 21 (IBM Corp., Armonk, NY) was used for the analysis reported herein. The distribution of the variables of interest were examined using histograms (PHQ-9, RHI, EPC) and tested for any deviation from normality using the Kolmogorov-Smirnov test. Depressive symptoms were treated as a continuous variable for all analyses. We also report the proportion of participants with a PHQ-9>10 to provide an estimate the proportion of participants likely to have developed major depressive disorder. Paired *t*-tests were utilized to assess for significant changes in depressive symptom score, RHI, and EPCs from the baseline to the internship time points. Change in depressive symptoms was calculated by taking the average of the PHQ-9 scores at 3-, 6-, 9-, and 12-months and subtracting this from the baseline PHQ-9. Two subjects did not complete within-internship PHQ-9 assessments and could not be included in analyses involving depressive symptom change. Pearson's correlation was used to examine the relationship between change in depressive symptoms, as measured by the PHQ-9, and changes in RHI and EPC levels. The relationship between RHI change and mean PHQ-9 change was graphed using SigmaPlot (Cystat Software, San Jose, CA). To clarify temporality, the relationship between change in depression scores from baseline to 6 months and change in RHI from baseline to 12 months was assessed. In a sensitivity analysis to assess for potential confounding, linear regression models which adjusted for age and sex were performed.

3. RESULTS

Sociodemographic and clinical characteristics of the sample are highlighted in Table 1. The sample included 14 women and 23 men pursuing training in a range of medical specialties with a mean age of 27.0 years. The most common specialties pursued were internal medicine (38%) and pediatrics (16%). Mean PHQ-9 score increased from a mean of 2.2

before internship to 4.7 during internship ($p<0.001$). No participants met PHQ-9 criteria for depression (PHQ-9 >10) at baseline, while 26% met criteria at least once during internship.

Over the course of internship, RHI did not change significantly (pre-internship stress RHI 1.81, post-internship stress RHI 2.00, $p=0.16$, where an RHI <1.67 is indicative of endothelial dysfunction) and baseline RHI did not predict depression during internship ($p=0.96$). Change in depressive symptoms was strongly and inversely correlated with change in RHI ($r=-0.41$, $p=0.01$). Changes in depression preceding the endothelial function assessment (baseline to 6 months) were similarly related to change in RHI over the intern year (baseline to 12 months) ($r=-0.38$, $p=0.02$). EPC levels dramatically decreased during internship from 11.9 cells/ml blood to 3.4 cells/ml blood ($t=2.60$, $p=0.01$). The change in depression scores under internship stress was not significantly related to the change in EPC levels ($p=0.96$). Estimates were not substantively altered in multivariate models controlling for age and sex.

4. DISCUSSION

Our finding of a temporal relationship between depression and endothelial dysfunction in a healthy, young cohort provides further evidence of a causal relationship between depression and cardiovascular health, suggesting vascular dysfunction may be a key link. Although there have been several case-control studies supporting a relationship between mood disorders and vascular function, there has been limited prospective study to establish temporality. In one prospective study of adolescent and young adult females with a family history of mood disorders, Tomfohr et al. similarly found an inverse relationship between depression and subsequent endothelial functioning [7]. Our cohort was at risk of depression due to occupational stress rather than family history and yielded similar findings, suggesting that the temporal relationship between depression and vasculopathy may be generalizable. We also demonstrated that internship stress, inferred from changes observed over the course of internship, independent of depression, was associated with substantial decreases in EPCs, which play a role in vessel repair and regeneration and whose decline may be an early harbinger of vascular disease [21]. EPC levels may be affected earlier in the progression from stress to vascular disease than endothelial function and lower levels of EPCs may impede recovery from any vascular insults. If this hypothesis is correct, we would predict that longer term prospective study of similar intensity stressors would lead to corresponding decreases in endothelial function with stress, regardless of depressive symptoms.

Limitations of the current study primarily involve the small samples size. We had a relatively high participation rate for a study involving physicians and the sample was representative of the larger cohort, even demonstrating a nearly identical proportion with significant depressive symptoms based on the PHQ-9 threshold [16]. The sample size also precluded any assessment of potential mediators or moderators, such as diet or inflammation, and limited the ability to assess confounding although the prospective design allowed individual participants to serve as their own matched control and improved statistical efficiency. Three or more assessments of RHI would have allowed a more refined assessment of the temporal relationship between depression and this outcome and more definitely determine the causal relationship between the two variables. The generalizability

of these findings to those who develop depressive symptoms due to other stressors or at other ages invites further investigation of this phenomenon.

This study prospectively demonstrates that the development of depression exerts physiological effects on vascular function in young adults with severe occupational stress. Such physiological effects could explain the dose-response between depressive symptom burden and vascular outcomes [22, 23]. The neurobiological underpinnings of depression may most plausibly mediate any such vascular effects through alterations in the hypothalamic-pituitary-adrenal axis, the autonomic nervous system, and inflammation. Further prospective studies, with repeated assessments, will be critical to identify the most clinically relevant mediators of depression on endothelial function and potential targets for intervention. Moderators such as sex may also be examined. The results reported herein would suggest that one year of follow-up may provide an adequate timeline to reliably characterize such relationships, in the setting of a sustained, chronic stressor such as medical internship.

ACKNOWLEDGEMENTS

The authors thank Srilakshmi Yalavarthi, M.S. who completed the EPC experiments. We are grateful to interns who participated in the study. Funding was provided by the NIMH (R01 MH101459, K23 MH095109) and the Michigan Institute for Clinical and Health Research (UL1RR024986). Dr. Fiedorowicz is funded by the NIMH (K23 MH083695) and the NHLBI (P01HL014388). Dr. Ellingrod is also funded by the NIMH (R01 MH082784). These funding organizations played no role in the design and conduct of the study, collection, management, analysis, and interpretation of the data, preparation and review of the manuscript or the decision to submit the manuscript for publication. This article was prepared while Dr. Kaplan was employed at the University of Michigan. The opinions expressed in this article are the author's own and do not reflect the view of the National Institutes of Health, the Department of Health and Human Services, or the United States government.

REFERENCES

1. Ormel J, Von Korff M, Burger H, Scott K, Demyttenaere K, Huang YQ, et al. Mental disorders among persons with heart disease - results from World Mental Health surveys. General hospital psychiatry. 2007; 29:325–334. [PubMed: 17591509]
2. Osby U, Brandt L, Correia N, Ekblom A, Sparen P. Excess mortality in bipolar and unipolar disorder in Sweden. Archives of general psychiatry. 2001; 58:844–850. [PubMed: 11545667]
3. Fiedorowicz JG, He J, Merikangas KR. The association between mood and anxiety disorders with vascular diseases and risk factors in a nationally representative sample. J Psychosom Res. 2011; 70:145–154. [PubMed: 21262417]
4. Fiedorowicz JG. Depression and cardiovascular disease: an update on how course of illness may influence risk. Current psychiatry reports. 2014; 16:492. [PubMed: 25163592]
5. Stanley SC, Brooks SD, Butcher JT, d'Audiffret AC, Frisbee SJ, Frisbee JC. Protective effect of sex on chronic stress- and depressive behavior-induced vascular dysfunction in BALB/cJ mice. Journal of applied physiology. 2014; 117:959–970. [PubMed: 25123201]
6. Bouzinova EV, Norregaard R, Boedtkjer DM, Razgovorova IA, Moeller AM, Kudryavtseva O, et al. Association between endothelial dysfunction and depression-like symptoms in chronic mild stress model of depression. Psychosomatic medicine. 2014; 76:268–276. [PubMed: 24804883]
7. Tomfohr LM, Murphy ML, Miller GE, Puterman E. Multiwave associations between depressive symptoms and endothelial function in adolescent and young adult females. Psychosomatic medicine. 2011; 73:456–461. [PubMed: 21715299]
8. Dome P, Teleki Z, Rihmer Z, Peter L, Dobos J, Kenessey I, et al. Circulating endothelial progenitor cells and depression: a possible novel link between heart and soul. Molecular psychiatry. 2009; 14:523–531. [PubMed: 18180758]

9. Harris KF, Matthews KA, Sutton-Tyrrell K, Kuller LH. Associations between psychological traits and endothelial function in postmenopausal women. *Psychosomatic medicine*. 2003; 65:402–409. [PubMed: 12764213]
10. Sherwood A, Hinderliter AL, Watkins LL, Waugh RA, Blumenthal JA. Impaired endothelial function in coronary heart disease patients with depressive symptomatology. *Journal of the American College of Cardiology*. 2005; 46:656–659. [PubMed: 16098431]
11. Pizzi C, Manzoli L, Mancini S, Costa GM. Analysis of potential predictors of depression among coronary heart disease risk factors including heart rate variability, markers of inflammation, and endothelial function. *European heart journal*. 2008; 29:1110–1117. [PubMed: 18400765]
12. Cooper DC, Milic MS, Tafur JR, Mills PJ, Bardwell WA, Ziegler MG, et al. Adverse impact of mood on flow-mediated dilation. *Psychosomatic medicine*. 2010; 72:122–127. [PubMed: 20100885]
13. Chen H, Zhang L, Zhang M, Song X, Zhang H, Liu Y, et al. Relationship of depression, stress and endothelial function in stable angina patients. *Physiology & behavior*. 2013; 118:152–158. [PubMed: 23688945]
14. Tomfohr LM, Martin TM, Miller GE. Symptoms of depression and impaired endothelial function in healthy adolescent women. *Journal of behavioral medicine*. 2008; 31:137–143. [PubMed: 18165894]
15. Osika W, Montgomery SM, Dangardt F, Wahrborg P, Gan LM, Tideman E, et al. Anger, depression and anxiety associated with endothelial function in childhood and adolescence. *Archives of disease in childhood*. 2011; 96:38–43. [PubMed: 19822537]
16. Sen S, Kranzler HR, Krystal JH, Speller H, Chan G, Gelernter J, et al. A prospective cohort study investigating factors associated with depression during medical internship. *Archives of general psychiatry*. 2010; 67:557–565. [PubMed: 20368500]
17. Grant F, Guille C, Sen S. Well-being and the risk of depression under stress. *PloS one*. 2013; 8:e67395. [PubMed: 23840872]
18. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire*. *JAMA : the journal of the American Medical Association*. 1999; 282:1737–1744. [PubMed: 10568646]
19. Rubinshtein R, Kuvlin JT, Soffler M, Lennon RJ, Lavi S, Nelson RE, et al. Assessment of endothelial function by non-invasive peripheral arterial tonometry predicts late cardiovascular adverse events. *European heart journal*. 2010; 31:1142–1148. [PubMed: 20181680]
20. Shantsila E, Watson T, Lip GY. Endothelial progenitor cells in cardiovascular disorders. *Journal of the American College of Cardiology*. 2007; 49:741–752. [PubMed: 17306702]
21. Yoder MC. Human endothelial progenitor cells. *Cold Spring Harbor perspectives in medicine*. 2012; 2:a006692. [PubMed: 22762017]
22. Fiedorowicz JG, Jancic D, Potash JB, Butcher B, Coryell WH. Vascular mortality in participants of a bipolar genomics study. *Psychosomatics*. 2014; 55:485–490. [PubMed: 24746452]
23. Sodhi SK, Linder J, Chenard CA, Miller del D, Haynes WG, Fiedorowicz JG. Evidence for accelerated vascular aging in bipolar disorder. *Journal of psychosomatic research*. 2012; 73:175–179. [PubMed: 22850256]

HIGHLIGHTS

We assessed vascular function in a prospective cohort of medical interns.

The development of depressive symptoms predicted worse endothelial function.

Endothelial progenitor cells were lower after one year of chronic stress.

Endothelial function may link stress, depression, and cardiovascular disease.

Figure 1. Depression worsens endothelial function

This scatterplot (N=37) demonstrates the relationship between change in depressive symptoms from baseline to the mean depressive symptoms score across the 3, 6, 9, and 12 month timepoints, as measured by the PHQ-9, and the observed changes in endothelial function, based on the Reactive Hyperemia Index (RHI), from baseline to 12 months. Participants with the greatest increases in depressive symptoms, as measured by the PHQ-9, had the greatest decreases in endothelial function as quantified by the RHI ($r=0.41$, $p=0.01$). Female participants are indicated with open circles and male participants with closed circles.

Table 1

Demographic characteristics of study participants.

Variable	Full Sample (%)	Female (%)	Male (%)
Gender	37	14 (37.8)	23 (62.2)
Age, y			
25	7 (18.9)	2 (14.3)	5 (21.7)
26–30	27 (73.0)	12 (85.7)	15 (65.3)
31–35	2 (5.4)	0 (0.0)	2 (8.6)
>36	1 (2.7)	0 (0.0)	1 (4.3)
Married/Engaged			
Yes	9 (24.4)	4 (28.6)	5 (21.7)
No	28 (75.6)	10 (71.4)	18 (78.3)
Family History of Depression			
Yes	20 (54.1)	8 (57.1)	12 (52.2)
No	17 (45.9)	6 (42.9)	11 (47.8)
Medical Specialty			
Internal Medicine	9 (24.4)	2 (14.3)	7 (30.4)
Other	15 (40.5)	6 (42.9)	9 (39.1)
Pediatrics	7 (18.9)	4 (28.6)	3 (13.0)
Emergency Medicine	2 (5.4)	0 (0.0)	2 (8.6)
Transitional Year	2 (5.4)	1 (7.4)	1 (4.3)
Internal medicine/pediatrics	2 (5.4)	1 (7.4)	1 (4.3)
Baseline PHQ	1.92	2.29	1.70
Mean Internship PHQ	4.39	4.82	4.11