Published in final edited form as:

Curr Hypertens Rep. 2015 February ; 17(2): 12. doi:10.1007/s11906-014-0523-z.

The Conundrum of Arterial Stiffness, Elevated blood pressure, and Aging

Majd AlGhatrif^{1,2} and Edward G. Lakatta¹

¹Laboratory of Cardiovascular Science, NIA, NIH, Baltimore, MD

²Department of Medicine, Johns Hopkins School of Medicine, Baltimore, MD

Abstract

Isolated systolic hypertension is a major health burden that is expanding with the aging of our population. There is evidence that central arterial stiffness contributes to the rise in systolic blood pressure (SBP); at the same time central arterial stiffening is accelerated in patients with increased (SBP). This bidirectional relationship created a controversy in the field on whether arterial stiffness leads to hypertension or vice versa. Given the profound interdependency of arterial stiffness and blood pressure, this question seems intrinsically challenging, or probably naïve. The aorta's function of dampening the pulsatile flow generated by the heart is optimal a range of distending pressure that secures the required distal flow, keeps the aorta in an optimal mechanical conformation, and minimizes cardiac work. This homeostasis is disturbed by age-associated, minute alterations in aortic hemodynamic and mechanical properties that induce short- and long-term alterations in each other. Hence, it is impossible to detect an "initial insult" at an epidemiological level. Earlier manifestations of these alterations are observed in young adulthood with a sharp decline in aortic strain and distensibility accompanied by an increase in diastolic blood pressure. Subsequently aortic mechanical reserve is exhausted, and aortic remodeling with wall stiffening and dilatation ensue. These two phenomena affect pulse pressure in opposite directions and different magnitudes. With early remodeling there is an increase in pulse pressure, due to the dominance of arterial wall stiffness, which in turn accelerates aortic wall stiffness and dilation. With advanced remodeling, which appears to be greater in men, the effect of diameter becomes more pronounced and partially offsets the effect of wall stiffness leading to plateauing in pulse pressure in men, and slower increase in PP than that of wall stiffness in women. The complex nature of hemodynamic changes with aging makes the "one-size-fits-all" approach suboptimal, and urges for therapies that addresses the vascular profile that underlies a given blood pressure, rather than the blood pressure values themselves.

Keywords

aging; hypertension; pulse wave velocity; aortic diameter; aortic remodeling

Introduction

Hypertension is an escalating major global health problem with an 8 out of 10 people are likely to have hypertension in their lifetime¹. It affects more than 40% of the world's population aged 25 and older, and nearly 80% of people aged 75 and older; it is responsible for about 13% of the total of all deaths worldwide². The direct and indirect cost of hypertension in the United States is estimated at \$46 billion and is projected to increase to \$274 billion by 2030³. Despite the great efforts to control hypertension, the number of subjects with uncontrolled hypertension increased from 800 million in 1988 to nearly 1 billion in 2008². This increase is mainly due to the aging of our population with the percentage of the people aged 65 and older rising, from 5% in 1950 to 8% in 2010, with a projected increase to 16% in 2050, accounting for 1.5 billion people⁴. This implies a vast expansion of an already costly and fatal epidemic of hypertension in the next few decades that warrants great efforts to prevent and treat this condition.

A deeper understanding of the pathophysiology that underlies hypertension is essential for any breakthrough in this direction. The mainstay approach to control blood pressure is to target mean blood pressure (MBP) by affecting heart rate, stroke volume, and peripheral vascular resistance; this approach is ineffective, if not harmful, when the issue is actually an elevation in pulse pressure (PP) rather than MBP⁵. This is the case in isolated systolic hypertension (ISH), the dominant form of hypertension in both young and old individuals with exponentially increasing dominance with advancing age^{6,7}.

While central arterial stiffness with aging is strongly implicated in the pathogenesis of ISH, controversy exists regarding the direction and nature of this relationship, with evidence supporting a role of arterial stiffness in the initiation and progression of ISH, and vice versa. Here, we aim to elucidate the conundrum of arterial stiffness, elevated blood pressure and aging by reviewing the physiological function of the aorta, and the epidemiological evidence of age-associated alterations in aortic properties and blood pressure parameters.

The “Purpose” of the Aorta: An evolutionary perspective

In early stages of life, eukaryotes evolved in a physically stationary, chemically renewable, microenvironment. The evolution of multicellular organisms with increasing complexity required solving the problem of maintaining the characteristics of microenvironments for the individual cells that are mostly not in direct contact with the external environment. This was an easier task in early forms of multicellular organisms such as the hydra, with a cylindrical body consisting of 2-cell thick wall enclaving a gastrovascular cavity connected with the external world; the peduncle of the hydra provides random contractions “beats” shaking the intracavitary fluid, while the epithelial flagellated cells simply move the fluids around⁸.

A dedicated circulatory system evolved with the evolution of more complex organisms in which internal and external environments are separated by an epithelium. While the heart generates pulsatile flow, it is the aorta that receives this pulsation and, via a Windkessel effect, dampens it into an almost continuous distal flow securing a physically stable environment for cells in the periphery⁹. The aorta provides elasticity over a physiological perfusing pressure range of a given organism in order to achieve dampening of pulsatility in

the physiological pressure range, and tensile strength to contain the stroke volume and avoid rupture at higher pressure.

Having an aortae composed of fibrillin microfibrils, along with other extra cellular matrix components, such as collagen, provided the elasticity required to minimize pulsatility in animal with low-pressurized, open circulation¹⁰. The appearance of vertebrates, however, with a closed circulation resulted in increased peripheral resistance and higher mean arterial pressure¹¹. Since the fibrillin microfibrils alone become very stiff in the context of newly evolved organisms requiring higher perfusion pressure, an increase in pulsatility might have imposed a selection pressure for the evolution of another extra-cellular matrix element required to provide elasticity at higher pressures; this element is tropoelastin¹⁰. In contrary to other ECM components, tropoelastin evolution coincided with the evolution of the highly-pressurized systems indicating its critical role in modulating pulsatile hemodynamics in this range.

The Cross-Talk between the Heart, Central Arteries, Peripheral Arteries, and the Microcirculation

The evolution of elastin is an example of cardiac-central arterial-peripheral arterial crosstalk at an evolutionary level; the peripheral modification accompanying the evolution of a closed-circulation required a central arterial modification to adapt to/optimize the new peripheral hemodynamics, while maintaining optimal cardiac work.

Similar to many phylogenic evolutionary phenomena, ontogenic cross-talk between the different components of the vascular bed is recapitulated in fetal development; the shift from low-pressurized, to high-pressurized circulation due to peripheral changes around birth triggers the production of elastin in central arteries¹⁰. This cross-talk represents a complex control-circuit including: cardiac output, elastin-collagen recruitment, vascular smooth muscle tone, and endothelial function, to count few. This control-circuit assures optimal distal flow, arterial wall stress, and cardiac loading in different physiological states i.e. exercise and pregnancy. One limitation in this control-circuit is the non-renewable status of elastin, which is only produced around birth and goes through slow turnover with aging¹². This, along with additional alterations to other components of the circuit that might arise with aging, convert this elegant control-circuit into a vicious cycle of drastic hemodynamic alteration. Hence, age-associated alterations to the structure and function of elastin are implicated in the increased central pulsatility that accompanies advancing age. These alterations are associated with activation of a proinflammatory profile leading to fibrotic and calcification processes leading to further increase in arterial wall stiffness (Figure1).

Age-Associated Changes in Aortic Properties and Blood Pressure in Humans

Epidemiological studies have been pursued to describe changes in arterial stiffness with aging and to answer the question of whether central arterial stiffness is a cause or an effect of elevated systolic and pulse blood pressure. One of the difficulties in addressing this issue relates to a degree of ambiguity and restrictions in these terms. This question might be better

articulated if we expand these terms and rename “arterial stiffness” as “arterial mechanical alterations”, and “elevated blood pressure” as “hemodynamic alterations”; then, it bears apparent that arterial mechanical properties and hemodynamics are inseparable and the question on what starts first, mechanical or hemodynamic alterations appears to be somewhat naïve.

In the beginning... Early alterations of arterial mechanics and the rise in diastolic blood pressure with Aging

The initial evidence of age-associated arterial mechanical alterations is observed by the third decade of age with sharp declines in aortic strain, the difference between aortic systolic and diastolic diameter relative to the diastolic diameter, and in aortic distensibility, i.e. aortic strain divided by pulse pressure; analysis of cardiac magnetic resonance (CMR) imaging of 111 healthy participants have shown that nearly 80% of the total decline of aortic strain occurs before the 5th decade of age¹³ (Figure 2). Preliminary analyses further examination the decline in arterial strain and distensibility in the SardiNIA project, a large cohort study of 6000 community-dwelling participant from Sardinia, Italy, have shown that the earlier decline in common carotid arterial strain and distensibility is not associated with a decreased “restricted” systolic diameter, but rather with an increase in diastolic diameter, independent of changes in distending blood pressure¹⁴. This later phenomena is consistent with an initial impaired diastolic recoil and not a restrictive process, commonly inferred from the notion that increased fibrosis and calcification are among the main stiffening processes.

During the same stages of life, as central arterial strain is becoming reduced, there is an increase in diastolic blood pressure^{15, 5}, which could be associated with progression of increased endothelial dysfunction with aging, leading to increased peripheral vascular resistance. While such changes are not very well studied in normal human subjects, those with essential hypertension demonstrate eutrophic, inward remodeling of small arteries^{16,17}. It is not clear whether the increase in diastolic alters the optimal conformation of the arterial wall making it prone to increased hemodynamic stress, and to additional mechanical alterations beyond those resulting from simply stretching elastin fibers at higher pressures and shifting the load to the stiffer collagen fibers.

Beyond the 5th Decade of Age: Central Arterial Mechanics and the Pulsatile Hemodynamic

Dramatic hemodynamic alterations ensue beyond the 5th decade of life, as increases in SBP and PP become the hallmarks of arterial aging⁵. Major histological and subsequent mechanical alterations in the central arterial wall are observed at this stage. These are characterized by a proinflammatory profile, metabolic stress, elastin fragmentation and calcification, collagen deposition, and vascular smooth muscle cells (VSMCs) stiffness¹⁸, reduced responsiveness to vasodilators, and increased proliferation and migration¹⁹ (Figure 3); among these factors, the effects of changes in elastin²⁰ and central arterial VSMC stiffness^{20–22} on arterial wall stiffness are more pronounced. These changes result in

mechanical alterations that manifest in a dramatic increase in PWV, the best currently available measurement of arterial stiffness.

An experimental animal model has shown that an increase in PWV occurs without any increase in blood pressure in nonhuman primates fed with diet high in fat and sucrose²³. Similarly Dahl salt-sensitive hypertensive rats develop an increase in PWV that precedes the increase in blood pressure in response to salt loading²⁴. Although these experimental animal models did not address the issue of aging, however, in reality, they do support the notion that the increase in PWV occurs independent of pressure changes.

Having said that, PWV and blood pressure parameters have been shown to be strongly associated in cross-sectional studies²⁵. A study of 483 participants evaluated at 2 visits about 6 years apart has shown that subjects with uncontrolled hypertension have greater increase in PWV compared to those with normal blood pressure or those with controlled hypertension²⁶. Data from the Caerphilly Prospective Study have shown that the cumulative effect of PP and HR also predicts PWV at 20-year, however, unfortunately this study did not have baseline PWV limiting the interpretation of these results²⁷. On the other hand, in a prospective study of 259 subjects who underwent repeated follow up for 16 years, PWV increase over time was observed independent of changes in mean arterial pressure due to treatment²⁸, it is worth noting in this study that those with AGTR1 genotype 1166C was associated with greater increase in PWV, independent of blood pressure or antihypertensive treatment after the age of 55²⁹.

The Framingham Heart Study, using data from Cycle 7 to predict SBP and PWV in Cycle 8, has shown that higher PWV in Cycle 7 was associated with higher SBP in Cycle 8³⁰; however, the opposite was not true SBP at Cycle 7 was not associated with higher PWV in Cycle 8. It is noteworthy, however, that one of the shortcomings of such design, with a relatively short follow-up time of 7 years, is that it ignores that the magnitude and the direction of association between SBP and PWV could change with aging and differ by gender; merely adjusting for these variables does not inform whether these associations differed between the different categories of these variable. Hence, the findings reflect that the average associations over the age spectrum studied for both genders.

New longitudinal perspective on the conundrum of arterial wall stiffness, blood pressure, and aging: A vicious cycle between teammates that eventually diverge

Indications of a vicious cycle between arterial stiffness and systolic blood pressure

The BLSA is a cohort study of community-swelling populations with extended follow-up time (1988–2013) and multiple repeated measures of PWV and BP. Earlier analyses from the BLSA have shown that greater PWV was associated with larger increase in SBP with aging, and predicted the incidence of hypertension³¹. Recent analysis from the BLSA, however, using linear mixed effects models, shed light on the vicious cycle, showing that higher SBP, in a dose-dependent fashion, is associated also associate with a greater rate of

increase in PWV overtime; this association was more pronounced in men with accelerating rates of increase in PWV at higher SBP with advancing age³².

Dissociation between PWV and SBP trajectories: When the teammates diverge

New insight on the longitudinal changes in PWV and SBP parameters came from the SardiNIA data with examination of the concurrent trajectories of repeated measures of PWV and SBP; using linear mixed effects models allows the examination of whether the longitudinal changes of these parameters over time vary by starting age¹⁵. This analysis demonstrated a striking dissociation in the trajectories of these parameters with advancing age; a dissociation more pronounced in men than women¹⁵. Figures 4A and B illustrate the cross-sectional differences “beginning of the splines” and the longitudinal changes (slopes of the splines) with aging (Rates of changes are illustrated in the lower panels) of PWV and SBP in both men and women. In men (Figure 4A) PWV increased with age at rates that increase linearly with advancing age; however, although cross-sectional SBP continues to increase, the longitudinal rates of change, while initially increasing, they begin to decline with time, thus the rates of change in SBP diverge from those of PWV by the fifth decade. A similar, but a less dramatic, divergence is observed in women (Figure 4B); while, PWV showed the same pattern of longitudinal changes in men with linearly increasing rates of change with advancing age, SBP increased longitudinally at a steady, rather than increasing, rates throughout the age range studied. Of note, the longitudinal associations between changes in PWV and SBP, which would have demonstrated the vicious cycle, were not evident in the SardiNIA study in part due to the conservative statistical approach that was implemented. Preliminary analyses from the BLSA using the same approach of examining the concurrent trajectories showed a similar pattern of dissociations between PWV trajectories and those of SBP and PP, which were more pronounced in men than women³³.

Epidemiological insight into the discrepancy between the SardiNIA and BLSA findings and the established views on SBP increase with aging

The longitudinal plateauing and then decrease in SBP among men with advancing age found in the SardiNIA¹⁵ and the BLSA data (preliminary analysis)³³ seem to contradict the conventional wisdom of an ongoing increase in SBP with aging; this apparent contradiction brings the attention to the impact of the analytical approaches to longitudinal data on the results that one would get. First, a pronounced limitation of any epidemiological study aimed at examining the trajectories of a parameter, SBP in this case, over a broad age range is that it is close to impossible to have a cohort that enters the study at the same young age and then to be followed all the way to an advanced age. Hence, large epidemiological studies implement a design in which participants enter studies at different ages and then be followed for a specific time; then the whole data is analyzed to produce the trajectories over a wide age range by compiling the trajectories of the different “sub-cohorts” studied. The main reason for constructing longitudinal data is to avoid the selection biases that are implicated in cross-sectional studies, especially at older age where those who are enrolled have SBP that are usually consistent with a survival/functional status advantage. This fact is often overlooked and, when constructing these trajectories, the average values of a blood pressures at a given age are derived from all participants, those who were just enrolled at that age and those who enrolled earlier and has been followed up till that age. Hence, any

longitudinal changes could be contaminated with the cross-sectional differences and the established trajectories would represent the average of these two forces.

The analytical approach implemented in analyzing the SardiNIA and the BLSA data using linear mixed effects models with Age expressed as Entry Age “cross-sectional differences” and Followup time “longitudinal changes”, allows separate assessment of the age-related cross-sectional differences and longitudinal changes of SBP, and beyond that to assess whether the longitudinal changes vary with age. This approach allows the revelation of “pure” longitudinal changes, while showing the age-related cross-sectional differences. A study that attempted reconstructing the life course trajectory of SBP using series of eight different longitudinal cohorts that each covered a segment of human lifespan showed similar results to that shown in the SardiNIA and the BLSA supporting this explanation. In addition, if the SardiNIA data, for example, is reanalyzed ignoring the cross-sectional differences and focusing on the follow up SBP values alone (the end point of the splines in the upper panels of Figure 4), the results show slowing of SBP increase, rather than a decline, which is almost identical to those from the Framingham Heart Study, in which the cross-sectional differences and the longitudinal changes in SBP were averages. The facts that the steeper rates of decline in SBP was not monotonic but was accentuated with advancing age, was different by gender, independent of antihypertensive treatment, and that it was associated with an escalating rates of increase in PWV, all make it unlikely to be a manifestation of regression to the mean, first-visit white coat effect, or secular changes, although one can’t totally exclude such effects.

A potential physiological explanation for the dissociation between PWV and the decline in SBP with aging in men

Dissociations between the rates at which PWV and SBP change over time bring our attention again to terminology; “Arterial stiffness” usually implies an increase in arterial opposition to flow i.e. characteristics (Z_c) impedance. However, Z_c , per the water hammer equation, is a function of both PWV and aortic diameter squared³⁴. Hence, an explanation for the dissociation between SBP and PWV longitudinal trajectories in men is an increase in aortic diameter. Preliminary analysis from the BLSA show a greater increase in aortic root dilatation with increasing age in men than in women³³. The net effect of increasing PWV and aortic diameter approximated by applying the water hammer equation is, in fact, a less pronounced increase in calculated Z_c in men despite their more pronounced increase in PWV which was offset by the greater increase in diameter³³. While these might explain how PWV and SBP/PP trajectories would diverge, the role of wave reflection in this dissociation is not well clear and it is worth further examination.

SUMMARY

The hypertension field has been struggling to better understand the complex relationship between arterial mechanical and hemodynamic alterations and fixated on answering the question: “which is the culprit?” The failure to reach an unequivocal answer is not surprising and is probably a reflection of the naivety of the question is. In health “homeostasis”, a functional crosstalk between central and peripheral segments of the circulation is required for optimal operation. Once this homeostasis is broken, for any reason, a vicious cycle of

minute alterations in central arterial mechanical and hemodynamics starts and propagates, leading to the dramatic changes in arterial properties observed with aging. Thus, in this paradigm, it is close to impossible to detect the “initial” minute alteration and point to it as the “culprit”.

However, the data discussed above suggest that the vicious cycle evolves into different stages the hemodynamic and mechanical patterns of which depends on the net effect of multiple factors that are interdependent and changes at different rates and directions with aging over a lifetime. In early adulthood, the heart, the aorta, and the peripheral circulation system form an “optimal” setup; the perfusing pressure needed puts the aorta in an optimal conformation for best Windkessel effect that in turn provides optimal pulsatility of the system. In the first phase of the vicious cycle, there are early alterations in arterial strain and distensibility associated with a rise in diastolic and mean blood pressure. By the 5th decade, arterial strain and distensibility reach a nadir while diastolic and mean blood pressures reach a peak and start a downward trend; these changes lag about a decade in women than men. In the second phase, aortic structural remodeling is more pronounced with an increase in arterial wall stiffness and aortic dilatation; remodeling progresses at a greater rate in men than women with a more pronounced dilatation and an increase in wall stiffness. Since the main manifestations of aortic remodeling, i.e. wall stiffness and lumen dilatation, affect aortic characteristic impedance, a major determinant of pulsatility, in opposite directions and at different rates, the net effect of these changes would vary across the age spectrum. Initially, they result in increased impedance and widening pulse pressure, which interplay with wall stiffness and lumen dilatation in a vicious cycle. Finally, given the greater increase in arterial stiffness and wall dilatation in men than women and that diameter imposes a stronger effect on impedance, the increase in pulse pressure observed in the previous phase is attenuated in men as result of a decline in SBP that parallels the ongoing decline in of DBP, leading to a plateau in PP. In women, on the other hand, the lower rate of remodeling slows the increase in SBP but does not result in a decline like the case in men; with DBP continuing to drop, pulse pressure continue to increase with age.

Given this extreme complexity, any efforts directed to treating or preventing the increase in blood pressure would be infertile until major efforts are committed to further explore the underpinning pathophysiology of this complex relationship, starting in young adulthood, before these alterations reach the clinical threshold, and to develop stage/process-specific interventions rather than the “one-size fits all” approach that dominate the field of hypertension.

References

1. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jones DW, Materson BJ, Oparril S, Wright JT, Roccella EJ. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003; 289:2560–72.
2. WHO. Raised blood pressure. [cited 2014 Aug 21]; Available from: http://www.who.int/gho/ncd/risk_factors/blood_pressure_prevalence_text/en/
3. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, Dai S, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit Ja, Howard VJ, Huffman MD, Judd SE, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Mackey RH, Magid DJ, Marcus

- GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Neumar RW, Nichol G, Pandey DK, Paynter NP, Reeves MJ, Sorlie PD, Stein J, Towfighi A, Turan TN, Virani SS, Wong ND, Woo D, Turner MB. Heart disease and stroke statistics--2014 update: a report from the American Heart Association. *Circulation*. 2014; 129:e28–e292.
4. World Population Aging: Clocks Illustrate Growth in Population Under Age 5 and Over Age 65 [Internet]. [cited 2014 Nov 5]; Available from: <http://www.prb.org/Publications/Articles/2011/agingpopulationclocks.aspx>
 5. Franklin SS, Gustin W, Wong ND, Larson MG, Weber MA, Kannel WB, Levy D. Hemodynamic Patterns of Age-Related Changes in Blood Pressure : The Framingham Heart Study. *Circulation*. 1997; 96:308–315. [PubMed: 9236450]
 6. Franklin SS, Jacobs MJ, Wong ND, L'Italien GJ, Lapuerta P. Predominance of Isolated Systolic Hypertension Among Middle-Aged and Elderly US Hypertensives : Analysis Based on National Health and Nutrition Examination Survey (NHANES) III. *Hypertension*. 2001; 37:869–874. [PubMed: 11244010]
 7. Grebla RC, Rodriguez CJ, Borrell LN, Pickering TG. Prevalence and determinants of isolated systolic hypertension among young adults: the 1999–2004 US National Health And Nutrition Examination Survey. *J Hypertens*. 2010; 28:15–23. [PubMed: 19730124]
 8. Shimizu H, Fujisawa T. Peduncle of Hydra and the heart of higher organisms share a common ancestral origin. *Genesis*. 2003; 36:182–6.
 9. Lakatta EG. Arterial and Cardiac Aging: Major Shareholders in Cardiovascular Disease Enterprises: Part I: Aging Arteries: A “Set Up” for Vascular Disease. *Circulation*. 2003; 107:139–146. [PubMed: 12515756]
 10. Faury G. Function-structure relationship of elastic arteries in evolution: from microfibrils to elastin and elastic fibres. *Pathol Biol (Paris)*. 2001; 49:310–25. [PubMed: 11428167]
 11. Schulte K, Kunter U, Moeller MJ. The evolution of blood pressure and the rise of mankind. *Nephrol Dial Transplant*. 2014; 1–11.
 12. Watanabe M, Sawai T, Nagura H, Suyama K. Age-related alteration of cross-linking amino acids of elastin in human aorta. *Tohoku J Exp Med*. 1996; 180:115–30. [PubMed: 9111761]
 13. Redheuil A, Yu W-C, Wu CO, Mousseaux E, de Cesare A, Yan R, Kachenoura N, Bluemke D, Lima JAC. Reduced ascending aortic strain and distensibility: earliest manifestations of vascular aging in humans. *Hypertension*. 2010; 55:319–26. [PubMed: 20065154]
 14. AlGhatrif M, Scuteri A, Morrell C, Canepa M, Strait J, Tarasov K, Orru' M, Schlessinger D, Lakatta E. GENDER DIFFERENCES IN EARLY MANIFESTATIONS OF ARTERIAL STIFFNESS: RESULTS FROM THE SARDINIA PROJECT. *J Am Coll Cardiol*. 2014; 63:A2037.
 15. Scuteri A, Morrell CH, Orru M, Strait JB, Tarasov KV, Ferrelli LAP, Loi F, Pilia MG, Delitala A, Spurgeon H, Najjar SS, AlGhatrif M, Lakatta EG. Longitudinal Perspective on the Conundrum of Central Arterial Stiffness, Blood Pressure, and Aging. *Hypertension*. 2014 This paper is the first to examine the concurrent longitudinal trajectories of pulse wave velocity and the different blood pressure parameters in a large community dwelling population. It showed dissociation in the trajectories of PWV and PP/SBP with advancing age that is more pronounced in men.
 16. Rizzoni D, Muiesan ML, Porteri E, De Ciuceis C, Boari GEM, Salvetti M, Paini A, Rosei EA. Interrelationships between macro and microvascular structure and function. *Artery Res*. 2010; 4:114–117.
 17. Mulvany MJ. Small artery remodelling in hypertension. *Basic Clin Pharmacol Toxicol*. 2012; 110:49–55. [PubMed: 21733124]
 18. Sehgel NL, Zhu Y, Sun Z, Trzeciakowski JP, Hong Z, Hunter WC, Vatner DE, Meininger GA, Vatner SF. Increased vascular smooth muscle cell stiffness: a novel mechanism for aortic stiffness in hypertension. *Am J Physiol Heart Circ Physiol*. 2013; 305:H1281–7. [PubMed: 23709594]
 19. Lakatta EG, Wang M, Najjar SS. Arterial aging and subclinical arterial disease are fundamentally intertwined at macroscopic and molecular levels. *Med Clin North Am*. 2009; 93:583–604. [PubMed: 19427493]

20. Valentín A, Humphrey JD, Holzapfel GA. A multi-layered computational model of coupled elastin degradation, vasoactive dysfunction, and collagenous stiffening in aortic aging. *Ann Biomed Eng.* 2011; 39:2027–45. [PubMed: 21380570]
21. McEniery CM, Qasem A, Schmitt M, Avolio AP, Cockcroft JR, Wilkinson IB. Endothelin-1 regulates arterial pulse wave velocity in vivo. *J Am Coll Cardiol.* 2003; 42:1975–81. [PubMed: 14662262]
22. Yin FC, Weisfeldt ML, Milnor WR. Role of aortic input impedance in the decreased cardiovascular response to exercise with aging in dogs. *J Clin Invest.* 1981; 68:28–38. [PubMed: 7251864]
23. Mattison JA, Wang M, Bernier M, Zhang J, Park S-S, Maudsley S, An SS, Santhanam L, Martin B, Faulkner S, Morrell C, Baur JA, Peshkin L, Sosnowska D, Csiszar A, Herbert RL, Tilmont EM, Ungvari Z, Pearson KJ, Lakatta EG, de Cabo R. Resveratrol prevents high fat/sucrose diet-induced central arterial wall inflammation and stiffening in nonhuman primates. *Cell Metab.* 2014; 20:183–90. [PubMed: 24882067]
24. Herrera VL, Decano JL, Giordano N, Moran AM, Ruiz-Opazo N. Aortic and carotid arterial stiffness and epigenetic regulator gene expression changes precede blood pressure rise in stroke-prone dahl salt-sensitive hypertensive rats. *PLoS One.* 2014; 9:e107888. [PubMed: 25229245]
25. Cecelja M, Chowienczyk P. Dissociation of aortic pulse wave velocity with risk factors for cardiovascular disease other than hypertension: a systematic review. *Hypertension.* 2009; 54:1328–1336. [PubMed: 19884567]
26. Benetos A. Determinants of Accelerated Progression of Arterial Stiffness in Normotensive Subjects and in Treated Hypertensive Subjects Over a 6-Year Period. *Circulation.* 2002; 105:1202–1207. [PubMed: 11889014]
27. McEniery CM, Spratt M, Munnery M, Yarnell J, Lowe GD, Rumley A, Gallacher J, Ben-Shlomo Y, Cockcroft JR, Wilkinson IB. An Analysis of Prospective Risk Factors for Aortic Stiffness in Men 20-Year Follow-Up From the Caerphilly Prospective Study. *Hypertension.* 2010; 56:36–43. This paper showed that the cumulative effect of the product of heart rate and pulse pressure over twenty years was associated with greater PWV at the end of the follow up.
28. Benetos A, Giron A, Joly L, Temmar M, Nzietchueng R, Pannier B, Bean K, Thomas F, Labat C, Lacolley P. Influence of the AGTR1 A1166C genotype on the progression of arterial stiffness: A 16-year longitudinal study. *Am J Hypertens.* 2013; 26:1421–7. [PubMed: 23942655]
29. Moens AL, Leyton-Mange JS, Niu X, Yang R, Cingolani O, Arkenbout EK, Champion HC, Bedja D, Gabrielson KL, Chen J, Xia Y, Hale AB, Channon KM, Halushka MK, Barker N, Wuyts FL, Kaminski PM, Wolin MS, Kass DA, Barouch LAC-P. Adverse ventricular remodeling and exacerbated NOS uncoupling from pressure-overload in mice lacking the beta3-adrenoreceptor. *J Mol Cell Cardiol.* 2009; 47:576–585. [PubMed: 19766235]
30. Kaess BM, Rong J, Larson MG, Hamburg NM, Vita JA, Levy D, Benjamin EJ, Vasan RS, Mitchell GF. Aortic stiffness, blood pressure progression, and incident hypertension. *JAMA.* 2012; 308:875–81. This paper showed that arterial stiffness at baseline was associated with the higher blood pressure at follow up, while the opposite was not true in this cohort. [PubMed: 22948697]
31. Najjar SS, Scuteri A, Shetty V, Wright JG, Muller DC, Fleg JL, Spurgeon HP, Ferrucci L, Lakatta EG. Pulse wave velocity is an independent predictor of the longitudinal increase in systolic blood pressure and of incident hypertension in the Baltimore Longitudinal Study of Aging. *J Am Coll Cardiol.* 2008; 51:1377–83. This is first study to show that in a community dwelling population, arterial stiffness was associated with greater increase in systolic blood pressure and the incidence of hypertension. [PubMed: 18387440]
32. AlGhatrif M, Strait JB, Morrell CH, Canepa M, Wright J, Elango P, Scuteri A, Najjar SS, Ferrucci L, Lakatta EG. Longitudinal Trajectories of Arterial Stiffness and the Role of Blood Pressure: The Baltimore Longitudinal Study of Aging. *Hypertension.* 2013; 62:934–951. This is the first analysis to show the longitudinal changes in PWV with aging in a community dwelling population and that traditional cardiovascular risk factors, except SBP and waist circumference, were not associated with the longitudinal increase in arterial stiffness. [PubMed: 24001897]
33. AlGhatrif M, Strait JB, Morrell CH, Canepa M, Wright J, Elango P, Scuteri A, Najjar SS, Ferrucci L, Lakatta EG. Attenuated Aortic Dilatation, Not Increased Wall Stiffness Best Explains the Rise

in Pulse Pressure in Women With Aging: Results From the Baltimore Longitudinal Study of Aging. *Circulation*. 2013; 128:A18061.

34. Mitchell GF, Gudnason V, Launer LJ, Aspelund T, Harris TB. Hemodynamics of increased pulse pressure in older women in the community-based Age, Gene/Environment Susceptibility-Reykjavik Study. *Hypertension*. 2008; 51:1123–1128. [PubMed: 18259005]
35. Safar, ME.; O'Rourke, MF.; Frohlich, ED., editors. *Blood Pressure and Arterial Wall Mechanics in Cardiovascular Diseases*. London: Springer London; 2014.

Figure 1. A schematic model illustrating the mechanism of TGF-β1 activation with loss of elasticity

TGF-β1 is secreted in a large latent complex, consisting of TGF-β1 associated with the latency associated peptide (LAP) and the latent TGF-β1-binding protein (LTBP-1). This structure links VSMCs to elastic fibers. In normal-functioning elastic arteries, VSMCs contraction pulls the whole compound with elastic fiber connected to it. With the loss of elasticity due to calcification, amyloidization, and other factors, the elastic fiber is less mobile and traction forces exerted by the stiffer VSMCs is transferred to LAP potentially resulting in activation of TGF-β1 and subsequent fibrosis. (Adapter from Blood Pressure and Arterial Wall Mechanics in Cardiovascular Diseases, Safar ME, O'Rourke MF, Frohlich ED, London, 2014)³⁵

Figure 2. Early age-related changes in aortic distensibility precede the changes in PWV
 Aortic distensibility (triangles) decreases sharply between the 3rd and 5th decade of life after which there is a sharp rise in aortic pulse wave velocity (modified from Redheuil et al, Hypertension, 2010)¹³

Figure 3. Conceptual model of arterial aging

Age-associated cellular disorders and cumulative mechanical stress lead to a state of chronic inflammation, elastin degradation, and endothelial and VSMC dysfunction. These processes interact and lead to arterial wall calcification, fibrosis, amyloid deposition, VSMCs proliferation, and increased intimal medial thickness. These structural changes lead to functional alterations resulting in widened pulse pressure. (Adapter from Blood Pressure and Arterial Wall Mechanics in Cardiovascular Diseases, Safar ME, O'Rourke MF, Frohlich ED, London, 2014)³⁵

Figure 4. Linear mixed-effects models predicted PWV and SBP values illustrating gender-specific cross-sectional differences “beginning of the splines” and the longitudinal changes (slopes of the splines) with aging (Rates of changes are illustrated in the lower panels) in men (A) and women (B) from the SardiNIA project. (Adapted with modification from Scuteri et al, Hypertension, 2014)¹⁵