CORRESPONDENCE

The Treatment of Displaced Intra-articular Distal Radius Fractures in Elderly Patients

A Randomized Multi-center Study (ORCHID) of Open Reduction and Volar Locking Plate Fixation Versus Closed Reduction and Cast Immobilization

by PD Dr. med. Christoph Bartl, PD Dr. med. Dirk Stengel MSc.,
Dr. sc. hum. Thomas Bruckner, Prof. Dr. med. Florian Gebhard in issue 46/2014

Prof. Dr. med. Lukas Prantl
Dr. med. Silvan Klein
Dr. med. Sebastian Geis
PD Dr. med. Jürgen Dolderer
Universitätsklinikum Regensburg
Zentrum für Plastische-, Hand- und
Wiederherstellungschirurgie
lukas.prantl@ukr.de

Prof. Dr. phil. Michael Koller
Universitätsklinikum Regensburg
Zentrum für klinische Studien
michael.koller@ukr.de

Prof. Dr. med. Michael Nerlich
Dr. med. Markus Loibl
Universitätsklinikum Regensburg
Klinik für Unfallchirurgie
michael.nerlich@ukr.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

Duration of Immobilization to Be Factored in

The authors' results show the continued need for a critical evaluation by means of randomized controlled trials of the use of volar locking plate fixation, even though today this technique has been widely adopted in clinical practice (1).

In our letter, we like to highlight an important aspect which has only been briefly touched in the study. We think that the type of aftercare provided was among the key factor influencing the treatment outcomes achieved. In the Methods section of the article, the aftercare is described as consisting of two weeks of postoperative physiotherapy according to local standards (1). However, we would have been interested in information about how long the periods of immobilization were in the various centers. While some hospitals altogether avoid postoperative immobilization after volar locking plate fixation, other centers prescribe postoperative immobilization periods of several weeks (2). Likewise, there are several distinct immobilization regimes used together with the standard casting techniques for this injury (3). Prolonged periods of immobilization frequently increase the need for physiotherapy after cast removal to such an extent that two weeks, the time limit stated in the article, are not enough (4). Thus, different periods of immobilization within and between the treatment arms could act as a confounder for the level of recovery, at least at the follow-up study visits at 3 and 6 months (3).

These differences between aftercare regimes are of interest from the perspective of resource utilization and against the background of the epidemiological trend described by the authors. DOI: 10.3238/arztebl.2015.0486a

REFERENCES

1. Bartl C, Stengel D, Bruckner T, Gebhard F and the ORCHID Study Group: The treatment of displaced intra-articular distal radius fractures in elderly patients—a randomized multi-center study (ORCHID) of open reduction and volar locking plate fixation versus closed reduction and cast immobilization. *Dtsch Arztebl Int* 2014; 111: 779–87.
2. Klein SM, Prantl L, Koller M, et al.: Evidence based postoperative treatment of distal radius fractures following internal locking plate fixation. *Acta Chir Orthop Traumatol Cech* 2015; 82: 33–40.
3. Millet PJ, Rushton N: Early mobilization in the treatment of Colles' fracture: a 3 year prospective study. *Injury* 1995; 26: 671–5.
4. Cyr LM, Ross RG: How controlled stress affects healing tissues. *J Hand Ther* 1998; 11: 125–30.

Age Should Be no Obstacle to Treatment

While in the last century reduction loss and its consequences for the patient (renewed reduction, pain, complex regional pain syndrome [CRPS]...) were common problems, these complications occur only very rarely today thanks to the surgical management of unstable distal radius fractures with volar locking plate fixation (1, 2). The sometimes catastrophic outcomes in patients who developed a complex regional pain syndrome are almost unknown to younger colleagues; the ability to prevent this complication is a blessing for patients and this treatment should be made available for "older patients ≥ 65 years of age" as well. We enrolled 762 patients in our prospective study; stable fractures (282 patients) were treated with standard casting techniques, while unstable fractures (480 patients) were treated with volar locking plate fixation—none of the patients treated with this approach developed a complex regional pain syndrome (3).

The authors' conclusion that the standard casting techniques represent a valid treatment concept for the primary management of unstable C2 and C3 fractures in elderly patients does not make sense to us, given a conversion rate of 41% due to secondary reduction loss, especially when the distribution pattern of the fractures in the group treated with standard casting techniques is taken into consideration (4). It is likely that secondary slipping occurred mostly along with unstable fractures—as expected—and that therefore surgical treatment was ultimately required. Under this assumption, we are approaching—with 37 converted patients and 38 patients with C2/C3 fractures treated with standard casting techniques—a conversion rate of almost 100%. My interpretation of the data: Elderly patients with unstable distal radius fracture should be immediately treated with locking plate fixation, while stable fractures should be treated with standard casting techniques. With this approach, predictable

complications can be prevented: despite economic constraints, elderly patients, too, have the right to receive care that quickly restores hand function.

DOI: 10.3238/arztebl.2015.0486b

REFERENCES

1. Dresing K, Stürmer KM, et al.: Deutsche Gesellschaft für Unfallchirurgie: Leitlinien Distale Radiusfraktur. <http://www.awmf.org/leitlinien/detail/II/012-015.html>
2. Stanos SP, Harden RN, Wagner-Raphael L, Saltz SL: A prospective clinical model for investigating the development of CRPS. in: Complex regional pain syndrome: progress in pain research and management. vol. 22. IASP Press, Seattle; 2001: 151–64.
3. Maier KJ: Behandlungsergebnisse nach palmarer, optional variabel winkelstabiler Plattenosteosynthese distaler Radiusfrakturen mit einer 2,7 mm Platte. Dissertation zum Erwerb des Doktorgrades in der Medizin an der Med. Fakultät der LMU München.
4. Bartl C, Stengel D, Bruckner T, Gebhard F and the ORCHID Study Group: The treatment of displaced intra-articular distal radius fractures in elderly patients—a randomized multi-center study (ORCHID) of open reduction and volar locking plate fixation versus closed reduction and cast immobilization. *Dtsch Arztebl Int* 2014; 111: 779–87.

Dr. med. Klaus-Jürgen Maier
Unfallchirurgie, RoMed Klinik Bad Aibling
klaus-juergen.maier@ro-med.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Take Courage to Perform Open Articular Surface Repair

The authors do not explicitly point out the need for a preoperative CT scan as a prerequisite for optimum surgery planning. The reduction of the articular surface achieved with the procedure can only be documented by postoperative computed tomography and these scans can help surgeons to learn from the outcomes of the procedures they have performed.

While intraoperative fluoroscopy can control the correct angular alignment of the main fragments and the position of the locking plate after reduction, it cannot be used to verify the articular congruency and the absence of steps in the articular surface. Therefore, intra-articular inspection of the joint surface is required and the fragments that bear areas of articular surface must be reduced and secured under direct visualization. With very small fragments, this can only be achieved using thin K-wires. The best outcome requires an articular surface without any steps; this is crucial to restore good load-bearing capacity of the radiocarpal joint. Patient satisfaction despite articular surface incongruity should not blind surgeons to the fact that significantly better outcomes would have been achieved with open step-free reduction. Unfortunately, this technique greatly prolongs operating times.

DOI: 10.3238/arztebl.2015.0487a

REFERENCES

1. Bartl C, Stengel D, Bruckner T, Gebhard F and the ORCHID Study Group: The treatment of displaced intra-articular distal radius fractures in elderly patients—a randomized multi-center study (ORCHID) of open reduction and volar locking plate fixation

versus closed reduction and cast immobilization. *Dtsch Arztebl Int* 2014; 111: 779–87.

Dr. med. Martin Artmann

Traunstein
art-hof@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We like to thank our colleagues Prantl et al., Artmann and Maier for their constructive criticism. With regard to aftercare, active physiotherapy started after two weeks in the surgical treatment arm. Patients began with finger movements on the first postoperative day and a removable dorsal plaster splint was applied for two weeks. All patients were shown how to perform an exercise program at home and instructions for independent training were handed out. In the plaster cast treatment arm, active physiotherapy started at six weeks. At 3 months the surgical treatment arm showed superior wrist mobility; it is likely that the earlier start of physiotherapy/home program significantly contributed to this outcome. At 12 months, both treatment arms showed comparable wrist mobility and the number of physiotherapy prescriptions had no direct influence on the range of motion achieved.

A preoperative CT scan was performed in addition to standard radiographic studies to optimize treatment planning, where necessary. Postoperatively, no routine CT scan was performed. At 12 months, the superior restoration of radiologic anatomy achieved with locking plate fixation compared with plaster cast treatment (1) was not associated with significantly better wrist mobility and daily-living function. To answer the question whether the superior restoration of joint anatomy will ultimately translate into long-term gains in wrist function and patient satisfaction, a long-term follow-up study is required.

In our study, the rate of complications and severe adverse events was lower than the one reported in the literature, even though in the ORCHID study the procedures were performed by surgeons of all training levels. Remarkably, only one case of CRPS (Sudeck's atrophy) was observed after plaster cast treatment. A positive correlation between higher-degree damage to the articular surface and metaphysis and an increase in the rate of secondary conversion to surgical treatment was found. In our opinion, not only the vast majority of non-displaced distal radius fractures but most displaced fractures where it is possible to largely stabilize the reduction with cast immobilization can be managed with plaster cast treatment until healing is complete (2, 3).

Since over the last decades a significant increase in the average activity levels of elderly patients has been observed, a patient's individual functional requirements should guide the decision on which treatment to select to manage a wrist fracture.

DOI: 10.3238/arztebl.2015.0487b

REFERENCES

1. Diaz-Garcia RJ, Oda T, Shauver MJ, Chung KC: A systematic review of outcomes and complications of treating unstable distal radius fractures in the elderly. *J Hand Surg Am* 2011; 36: 824–35.
2. Arora R, Lutz M, Deml C, Krappinger D, Haug L, Gabl M: A prospective randomized trial comparing nonoperative treatment with volar locking plate fixation for displaced and unstable distal radial fractures in patients sixty-five years of age and older. *J Bone Joint Surg Am* 2011; 93: 2146–53.
3. Bartl C, Stengel D, Bruckner T, Gebhard F and the ORCHID Study Group: The treatment of displaced intra-articular distal radius fractures in elderly patients—a randomized multi-center study (ORCHID) of open reduction and volar locking plate fixation versus closed reduction and cast immobilization. *Dtsch Arztebl Int* 2014; 111: 779–87.

PD Dr. med. Christoph Bartl

Klinik für Unfall-, Hand-,
Plastische und Wiederherstellungschirurgie
Universität Ulm
chbartl@gmx.de

PD Dr. med. Dirk Stengel MSc

Zentrum für klinische Forschung
Unfallkrankenhaus Berlin

Dr. sc. hum. Thomas Bruckner

Institut für Medizinische Biometrie und Informatik (IMBI)
Universität Heidelberg

Prof. Dr. med. Florian Gebhard

Klinik für Unfall-, Hand-,
Plastische und Wiederherstellungschirurgie
Universität Ulm

Conflict of interest statement

PD Dr. Stengel has received payments for consultation, reimbursement of congress attendance fees, and honoraria for preparation of scientific meetings from Biomet, Stryker, and the AO Foundation.

He has received reimbursement of travel or accommodation costs from Biomet, Stryker, the AO Foundation, the German Trauma Society (Deutsche Gesellschaft für Unfallchirurgie e.V., DGU), and the German Social Accident Insurance (Deutsche Gesetzliche Unfallversicherung, DGU).

PD Dr. Bartl, Dr. Bruckner and Prof. Gebhard declare that no conflict of interest exists.