Published in final edited form as:

J Periodontol Res. 2012 April ; 47(2): 236–242. doi:10.1111/j.1600-0765.2011.01427.x.

***Porphyromonas gingivalis* Promotes Monocyte Migration by Activating MMP-9**

Jing Zhou¹, Jingmei Zhang², and Julie Chao²

¹Department of Pediatric Dentistry and Orthodontics, Medical University of South Carolina, Charleston, SC, USA

²Department of Molecular Biology and Biochemistry, Medical University of South Carolina, Charleston, SC, USA

Abstract

Objectives—The migration of monocytes into local environment is crucial for its maturation into macrophages or osteoclasts in the pathogenesis of periodontal disease. The objective of this study was to investigate the role and mechanisms mediated by *Porphyromonas gingivalis* (*P. gingivalis*) in promoting the migration of monocytes through regulating matrix metalloproteinase (MMP)-9 and tissue inhibitor of metalloproteinase (TIMP)-1 expression.

Methods—Human THP1 monocytes were treated with culture supernatant derived from *P. gingivalis* (ATCC 33277) for 24 hours. Zymography, Western blot analysis and quantitative polymerase chain reactions were performed to analyze protein and mRNA levels of MMP-9. Protein and mRNA levels of TIMP-1 from monocytes treated with or without *P. gingivalis* were determined as well. Transwell migration assay was carried out to analyze the effect of *P. gingivalis* on the migration of human peripheral blood CD14⁺ monocytes. MMP inhibitor (GM6001) and proteinase inhibitor (Leupeptin) were used to determine the role of MMP-9 in *P. gingivalis* supernatant- and Lipopolysaccharide (LPS)-induced monocyte migration.

Results—In zymography and Western blot, an 82-kDa band of active MMP-9 emerged in *P. gingivalis*-treated monocyte culture media in a dose-dependent manner in addition to the MMP-9 proenzyme (92 kDa) band expressed in control cell culture media. *P. gingivalis* supernatant increased both the protein and mRNA levels of MMP-9 and TIMP-1. *P. gingivalis* supernatant, but not its LPS, increased the migratory ability of CD14⁺ monocytes. The increased migratory ability of *P. gingivalis*-treated monocytes was partially inhibited by leupeptin (200 µg/ml) and completely antagonized by the MMP inhibitor GM6001 (100 nM). LPS of *P. gingivalis* increased protein and mRNA levels of MMP-9 in monocytes, but had no effect on the migratory ability or MMP-9 activation.

Conclusion—*P. gingivalis* supernatant increased the migratory ability of monocytes, in part, by increasing activation and expression of MMP-9.

Keywords

Porphyromonas gingivalis; monocytes; migration; MMP-9

Introduction

Recent case-control studies and longitudinal studies have established the association between chronic periodontal disease and cardiovascular disease (1). This relationship could be mediated by inflammatory responses and cellular events induced by periodontal pathogens (2). *Porphyromonas gingivalis* (*P. gingivalis*), a major periodontal pathogen, was shown to be present in both periodontal pockets and atherosclerotic plaques (3, 4).

Monocytes are a circulating white blood cell and part of the human body's immune system. Activation of immune cells by periodontal pathogens, e.g., *P. gingivalis*, is an important feature of chronic inflammatory periodontal disease. The migration and localization of monocytes has been recognized to be a crucial step of its differentiation and maturation into macrophages and osteoclasts, which play important roles in the development of local inflammation and connective tissue destruction (5, 6). Previous studies have shown that *P. gingivalis* modulates adhesion, chemotaxis, and migration of monocytes and its derived cells by regulating the levels of cytokines (7–10), chemoattractants (11–13), cytosolic proteins (14, 15), and matrix metalloproteinases (MMPs) (16).

The migratory abilities of monocytes and derived cells are partially accomplished through the regulation of MMPs and tissue inhibitors of metalloproteinases (TIMPs) (17–19). MMPs are a family of zinc-dependent endopeptidases responsible for the degradation of multiple extracellular matrix (ECM) components (20). MMPs are secreted in a latent proform by multiple host structural cells and immune cells. The activation of MMPs involves the loss of a propeptide (20). TIMPs function as inhibitors of MMPs by forming monocovalent complexes with MMPs, thus blocking the access of substrates to the MMP catalytic site (20). Numerous MMPs and TIMPs are expressed in peripheral blood monocytes (21) including MMP-9 and TIMP-1. MMP-9, also known as gelatinase-B, cleaves denatured collagen, in particular, type IV collagen, which constitutes the major component of the basement membranes (22). This cleavage helps the white blood cells that play a role in immune responses, such as monocytes, to enter and leave the blood circulations. Previous studies suggest that MMP-9 and TIMP-1 are involved in the migration and localization of monocytes and their differential cells (16, 18, 19, 23). However, how MMPs are regulated by periodontal pathogen in monocytes and how this regulation affects the migratory ability of monocyte have not been fully elicited.

Therefore, the objective of this study was to investigate the role and mechanisms mediated by *P. gingivalis* in promoting the migration of monocytes through regulating MMP and TIMP expression. It was hypothesized that *P. gingivalis* induces monocyte migration via regulating matrix metalloproteinase (MMP)-9 and tissue inhibitor of metalloproteinase (TIMP)-1. Understanding how periodontal pathogens alternate the function of host immune cells in the pathogenesis of periodontal disease may help explain the association of periodontal disease and other systemic conditions such as cardiovascular disease.

Material and Methods

Bacteria growth and the collection of the culture supernatant

P. gingivalis supernatant has been used in previous studies as a stimulus to mimic the toxins released from this bacterium in the periodontal pocket (24–26). *P. gingivalis* ATCC 33277 supernatant was provided by Dr. A. Progulske-Fox (University of Florida College of Dentistry, Gainesville, FL, USA). *P. gingivalis* ATCC 33277 was cultured in supplemented brain heart infusion growth media as described previously (25). The collected supernatant was filtered twice through 2 µm membranes to remove larger cell remnants and then stored at –20°C until utilized. The lipopolysaccharide (LPS) of *P. gingivalis* ATCC 33277 was also provided by Dr. A. Progulske-Fox. The methodology for isolation and purification of LPS from *P. gingivalis* was described in a previous paper (27).

Cell Culture

THP1 human monocytic leukemia cell line was utilized to study the effect of *P. gingivalis* on intact human monocytes (28). Monocytes isolated from mononuclear fractions of peripheral blood of healthy donors and commercially available human CD14+ monocytes were also used in this study. Monocytes were cultured at 37°C and 5% CO₂ in low glucose (1g/l) α-minimal essential medium (MEM) media (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum, 4mM L-glutamine, 100 U/ml penicillin, 50 µg/ml gentamicin, and 2.5 µg/ml fungizone.

THP1 cells were cultured with *P. gingivalis* supernatant or LPS. Monocytes were collected from the culture flask, centrifuged, and resuspended in serum-free α-MEM media at a concentration of 50,000 cells/ml. The cells were then distributed in 6-well culture plates and cultured for 24 hours with *P. gingivalis* supernatant (0.5–5% v/v) and *P. gingivalis* LPS (100 µg/ml). Cells cultured in serum-free medium were used as experimental controls. The conditioned culture media were collected and concentrated for zymography and Western blot analysis.

Gelatin zymography

The conditioned THP1 media was mixed with non-reducing loading buffer and resolved at 150 V in 10% sodium dodecylsulphate–polyacrylamide gel electrophoresis (SDS–PAGE) gels copolymerized with 1 mg/ml gelatin. The gels were then washed with a washing solution containing 2.5% Triton X-100 for three 20-minute cycles. The gels were then incubated in fresh developing buffer containing 50 mM Tris pH 7.5, 3mM NaN₃, 5mM CaCl₂, and 1 µM ZnCl₂ at 37°C overnight and stained with Coomassie blue to visualize the proteolytic bands. The experiment was repeated three times.

Western blot

An equal amount of total protein of untreated and *P. gingivalis* treated cell culture media was resolved in 10% SDS–PAGE gels at 150 V. The proteins on the gels were transferred to polyvinylidene difluoride membrane (Millipore, Bedford, MA, USA) at 0.3 A for 1 hour using blotting buffer (25 mM Tris-HCl, pH 8.3, 192 mM glycine, and 10% methanol). The membranes were then incubated in 5% milk in phosphate-buffered saline (PBS) solution

with 0.1% Tween-20 (pH 7.4) for 1 hour to block non-specific binding. The membranes were then incubated with primary antibodies at 4°C overnight. Antibodies used in this study were monoclonal anti-MMP-9 (1:1000, Fitzgerald, Concord, MA, USA) and polyclonal antibody 2315 (5 µg/ml, a gift from Dr. L.J. Windsor, Indianapolis, IN, USA) for TIMP-1. The membranes were then washed three times with PBS with Tween and incubated with horseradish peroxidase-conjugated secondary antibodies for 1 hour at room temperature. Chemiluminescence was detected by an ECL-Plus kit (Perkin-Elmer, Foster City, CA, USA). The Western blot analyses were repeated three to five times.

Real-time PCR

Total RNA was isolated from cultured THP1 cells with TRIzol reagent (Invitrogen) following the manufacturer's protocol. Total RNA was reverse-transcribed using the High-Capacity cDNA Reverse Transcription Kit. Real-time polymerase chain reactions (PCR) was performed with the TaqMan Gene Expression Assay and was normalized against 18S RNA using an ABI 7300 real-time PCR System (Applied Biosystems, Foster City, CA, USA). The primers Hs00234579_ml and Hs171558_ml were used for the detection of MMP-9 and TIMP-1 expression, respectively. The experiments were repeated three or more times.

Cell Migration assay

Commercially available human CD14+ peripheral monocytes were used for monocyte migration studies due to the fact that THP1 cells do not attach to the culture wells. Monocyte migration assay was performed using modified Boyden Chamber methods as previously described (29). Briefly, peripheral blood CD14+ monocytes (Lonza, Allendale, NJ, USA) were cultured in α -MEM with 10% FBS and 10ng/ml of Macrophage Colony-Stimulating Factor (M-CSF) for 5 days before being used in the transwell migration assay. Cells were collected by trypsin treatment (0.25%), centrifugation, and then placed in the upper chambers of transwell plates (10^5 cells/well in 250 µl α -MEM supplied with 10ng/ml M-CSF). The lower chambers were filled with control media (500 µl α -MEM) and different treatment media: *Pg* (α -MEM + 1% *P. gingivalis* supernatant), *Pg* LPS (α -MEM + 100 µg/ml *P. gingivalis* LPS), boiled *Pg* (α -MEM + 1% denatured *P. gingivalis* supernatant), *Pg* + leupeptin (α -MEM + 1% *P. gingivalis* supernatant + 200 µg/ml leupeptin), or *Pg* + GM6001 (α -MEM + 1% *P. gingivalis* supernatant + 200 nM GM6001). The cells were cultured in 24 well transwell plates for 16 hours at 37°C. The upper wells were wiped by cotton tips, then fixed with 3.7% formalin overnight. The wells were then rinsed with 1× PBS twice and stained with crystal violet for 20 minutes. Migrated cells were counted under four random 40× microscope fields. The means of the cell numbers in these fields were compared between groups. The experiments were repeated three times.

Statistical analysis

All data are presented as means $\pm$ SE. Comparison between groups was made using one-way ANOVA with the Fisher multiple comparison test. A probability value of $P < 0.05$ was considered statistically significant. All cell culture experiments were performed in triplicates on at least three separate occasions.

Results

P. gingivalis supernatant induces MMP-9 activation

A proteolytic band of approximately 92 kDa of MMP-9 was detected in the untreated THP1 conditioned media (Fig. 1A). It was observed that with the addition of 0.5%, 1%, and 2.5% of *P. gingivalis* supernatant in the media, the 92-kDa band weakened and a band with molecular weight of approximately 82 kDa emerged (Fig. 1A) in a dose-dependent manner. With increased concentrations of *P. gingivalis* supernatant (5%), both bands weakened indicated either the degradation of host MMPs or cell depression by bacterial supernatant. There is an enhanced 92-kDa band but no 82-kDa band in the *P. gingivalis* LPS-treated THP1 cell culture media. The results indicate a dose-dependent conversion of pro-MMP-9 to active MMP-9.

P. gingivalis supernatant or LPS treatment increased the protein levels of MMP-9 and TIMP-1 in monocytes

The Western blot confirmed the findings in gelatin zymography. Latent MMP-9 was detected in untreated THP1 culture media, *P. gingivalis* LPS-treated THP1 conditioned media, and *P. gingivalis* supernatant (0.5–2.5%)-treated THP1 conditioned media (Fig. 1B). Active form of MMP-9 (82 kDa) was only detected in *P. gingivalis* supernatant (0.5–2.5%)-treated THP1 conditioned media (Fig. 1B). Treatment with a higher concentration of *P. gingivalis* supernatant (5%) resulted in a decreased detectable MMP-9 amount (Fig. 1). This could be the result of either the direct degradation of MMP-9 by bacterial proteinases, or host cell depression by high concentrations of bacterial supernatant. Therefore, 1% *P. gingivalis* supernatant was utilized in the cell migration analysis. *P. gingivalis* LPS increased the expression of latent MMP-9 (92 kDa) but not its active form (82 kDa). Both *P. gingivalis* supernatant 1% and LPS (100 µg/ml) increased the expression of TIMP-1 (Fig. 2A).

P. gingivalis supernatant or LPS treatment increased mRNA levels of MMP-9 and TIMP-1 in monocytes

The mRNA expression levels of MMP-9 and TIMP-1 from untreated THP1 cells and THP1 cells treated with *P. gingivalis* supernatant or LPS for 24 hours were determined. Both treated and untreated THP1 cells were shown to express mRNA for MMP-9 (Fig. 3) and TIMP-1 (Fig. 2B). *P. gingivalis* supernatant or its purified LPS increased the mRNA expression of MMP-9 (Fig. 3) and TIMP-1 (Fig. 2B). However, the mRNA level of MMP-9 (Fig. 3) was increased at a much higher level than that of TIMP-1 (Fig. 2B) from *P. gingivalis*-treated THP1 cells.

P. gingivalis supernatant increased the migration of human peripheral monocytes

Treatment with 1% *P. gingivalis* supernatant increased the migration of CD14+ monocytes into the lower chamber (Fig. 4). Neither *P. gingivalis* LPS nor boiled *P. gingivalis* supernatant increased the amount of migrated CD14+ monocytes. The cysteine and serine proteinase inhibitor leupeptin (200 µg/ml) partially reduced the cell migratory ability of the *P. gingivalis* supernatant-treated monocytes. The MMP inhibitor GM6001 (200 nM)

completely antagonized the increased cell migratory ability of *P. gingivalis* supernatant-treated CD14+ monocytes.

Discussion

It was revealed in our study that *P. gingivalis* supernatant increased the migratory ability of CD14+ monocytes *in vitro*. This effect was accompanied by the activation of MMP-9 from the *P. gingivalis* supernatant treated monocytes (in both THP1 cells (Fig. 1) and isolated human peripheral monocytes (data not shown)). The increased migration of *P. gingivalis*-treated monocytes was partially reduced by a cysteine and serine proteinase inhibitor, and completely antagonized by an MMP inhibitor (Fig. 4). These results suggest that *P. gingivalis* increased the migratory ability of human monocytes through the activation and expression of MMP-9, partially mediated by the bacterial proteinases. The remaining increased migratory ability of monocyte may be induced by other protein components in the *P. gingivalis* supernatant, such as the fimbriae (30). This hypothesis is also indirectly supported by the fact that neither *P. gingivalis* LPS nor boiled *P. gingivalis* supernatant increased the migratory ability of the monocytes. These findings help us to understand how periodontal pathogens promote peripheral blood monocytes migration into the local environment of periodontitis. Furthermore, these findings together with previous publications on how *P. gingivalis* regulates the proadhesive pathway of monocytes (30–33), provide insight in the association between *P. gingivalis* and atherosclerotic vascular disease.

This study also demonstrated that *P. gingivalis* upregulated both MMP-9 and TIMP-1 at protein and mRNA levels. TIMP-1 is a major type of tissue inhibitors of MMPs that is synthesized by most cells (20). The balance of MMPs and TIMPs in the local environment is important in maintaining the collagen turn over rate and disturbances of this balance may result in excessive degradation of the ECM (20). In the current study, the levels of MMP-9 and TIMP-1 were upregulated at different levels, resulting in a high MMP-9/TIMP-1 ratio. This result agreed with a previous study, which reported a high MMP-9/TIMP-1 ratio induced by *P. gingivalis* (live strains and LPS) from dendritic cells (16). The high MMP/TIMP ratio implies an imbalance of MMP functions in the inflammatory conditions such as periodontal disease and atherosclerotic vascular disease.

The activation of the MMPs is an essential step in regulating its biological function (20). Proteolytic activation of MMPs was determined by an approximately 10-kDa reduction in molecular weight. Studies have shown that proteinases expressed by periodontal pathogens can activate latent MMPs (34, 35) and may play additional roles in initiating or accelerating extracellular matrix degradation in periodontal tissues (36). Pro-MMP-1 and pro-MMP-8 can be activated by supra- and subgingival plaque extracts (37). Pro-MMP-9 can be activated by bacteria and their outer membrane extracts and cell suspension (22). Pro-MMP-2 can be activated by gingival crevicular fluid and *P. gingivalis* supernatant (38, 39). *In vitro*, *P. gingivalis* proteinases can activate latent MMP-1, MMP-2, MMP-3, MMP-8, MMP-9, as well as catalyze the superactivation of MMP-1 by MMP-3 (34, 35, 38, 40). Activation of these MMPs has been demonstrated to result from an initial hydrolysis within the propeptide of the MMPs. For the first time, our study demonstrates that *P. gingivalis* activates MMP-9 expressed in monocytes (Fig. 1). This activation of MMP-9, in part,

contributes to the increased migratory ability of human monocytes. Activated forms of MMP-9 were detected in differentiated monocytes and were suggested to play an important role in promoting monocyte invasiveness and its differentiation into macrophages (23).

Previous studies have shown that *P. gingivalis* and its components regulate the levels of cytokines from immune cells via different mechanisms (41, 42). *P. gingivalis* fimbrial protein molecules regulate monocyte chemoattractant protein-1 expression in endothelial cells (11–13). These molecules may play an important role in structurally facilitating the attachment of bacterial protein to host cells such as erythrocytes and monocytes (30, 31, 43). Live *P. gingivalis* and its LPS or fimbriin induce different cytokines and chemokines by mouse macrophages through different ligand pathways (42, 44). *P. gingivalis* LPS also induces calprotectin (a cytosolic protein) and multiple co-stimulatory molecules in human monocytes (14, 15, 45), as well as upregulates MMP-9 and TIMP-1 expression in human dendritic cells (16). Our study suggests that *P. gingivalis* LPS regulates MMP-9 and TIMP-1 expression in monocytes, but the *P. gingivalis* proteinases may play a crucial role in the activation of MMP-9.

Cysteine proteinases (gingipains) account for the major proteolytic activity of *P. gingivalis*. They play various roles in the pathogenesis of periodontal disease, including the development of edema, neutrophil infiltration, and bleeding (46). Gingipains exhibit enzymatic activity against many host proteins such as immunoglobulins, extracellular matrix proteins, iron transporting factors and inhibitors, and proteins involved in the coagulation, complement, and kallikrein-kinin cascades, as well as host MMPs and their activators (25, 34, 35, 46, 47). Gingipains also regulate monocyte chemoattractant protein-1 expression in endothelial cells (48). Our findings suggest that gingipains may increase the migratory ability of monocytes by activating MMP-9. Further studies are indicated to identify the role of gingipains in regulating the localization and maturation of monocytes.

In summary, the findings of this study suggest that activation of MMP-9 is a crucial step in promoting monocyte migration induced by *P. gingivalis*. This study provided insights into the role and mechanisms of the periodontal pathogen *Porphyromonas gingivalis* in the development of periodontal disease and its systemic complications.

Acknowledgements

This study was conducted with support from the National Institutes of Health, National Center of Research Resources Grant Number P20 RR01796, South Carolina COBRE for Oral Health Research. The authors greatly acknowledge Dr. Ann Progulski-Fox for kindly providing the *P. gingivalis* supernatant and LPS. The authors also greatly acknowledge Dr. L. Jack Windsor for providing the anti-TIMP-1 polyclonal antibody.

References

1. Beck JD, Offenbacher S, Williams R, Gibbs P, Garcia R. Periodontitis: a risk factor for coronary heart disease? *Ann Periodontol*. 1998; 3:127–141. [PubMed: 9722697]
2. Kuramitsu HK, Qi M, Kang IC, Chen W. Role for periodontal bacteria in cardiovascular diseases. *Ann Periodontol*. 2001; 6:41–47. [PubMed: 11887470]
3. Gaetti-Jardim E Jr, Marcelino SL, Feitosa AC, Romito GA, Avila-Campos MJ. Quantitative detection of periodontopathic bacteria in atherosclerotic plaques from coronary arteries. *J Med Microbiol*. 2009; 58:1568–1575. [PubMed: 19679682]

4. Mahendra J, Mahendra L, Kurian VM, Jaishankar K, Mythilli R. 16S rRNA-based detection of oral pathogens in coronary atherosclerotic plaque. *Indian J Dent Res.* 2010; 21:248–252. [PubMed: 20657096]
5. Okada H, Silverman MS. Chemotactic activity in periodontal disease. I. The role of complement in monocyte chemotaxis. *J Periodontal Res.* 1979; 14:20–25. [PubMed: 153959]
6. Pradeep AR, Daisy H, Hadge P. Gingival crevicular fluid levels of monocyte chemoattractant protein-1 in periodontal health and disease. *Arch Oral Biol.* 2009; 54:503–509. [PubMed: 19286166]
7. Barksby HE, Nile CJ, Jaedicke KM, Taylor JJ, Preshaw PM. Differential expression of immunoregulatory genes in monocytes in response to *Porphyromonas gingivalis* and *Escherichia coli* lipopolysaccharide. *Clin Exp Immunol.* 2009; 156:479–487. [PubMed: 19438601]
8. Eskin MA, Hajishengallis G, Kinane DF. Differential activation of human gingival epithelial cells and monocytes by *Porphyromonas gingivalis* fimbriae. *Infect Immun.* 2007; 75:892–898. [PubMed: 17118977]
9. Ogawa T, Uchida H, Hamada S. *Porphyromonas gingivalis* fimbriae and their synthetic peptides induce proinflammatory cytokines in human peripheral blood monocyte cultures. *FEMS Microbiol Lett.* 1994; 116:237–242. [PubMed: 7908651]
10. Shimauchi H, Ogawa T, Okuda K, Kusumoto Y, Okada H. Autoregulatory effect of interleukin-10 on proinflammatory cytokine production by *Porphyromonas gingivalis* lipopolysaccharide-tolerant human monocytes. *Infect Immun.* 1999; 67:2153–2159. [PubMed: 10225868]
11. Choi EK, Park SA, Oh WM, et al. Mechanisms of *Porphyromonas gingivalis*-induced monocyte chemoattractant protein-1 expression in endothelial cells. *FEMS Immunol Med Microbiol.* 2005; 44:51–58. [PubMed: 15780578]
12. Kang IC, Kuramitsu HK. Induction of monocyte chemoattractant protein-1 by *Porphyromonas gingivalis* in human endothelial cells. *FEMS Immunol Med Microbiol.* 2002; 34:311–317. [PubMed: 12443832]
13. Kobayashi-Sakamoto M, Isogai E, Hirose K. *Porphyromonas gingivalis* modulates the production of interleukin 8 and monocyte chemotactic protein 1 in human vascular endothelial cells. *Curr Microbiol.* 2003; 46:109–114. [PubMed: 12520365]
14. Kido J, Hayashi N, Kataoka M, Nagata T. Calprotectin expression in human monocytes: induction by *porphyromonas gingivalis* lipopolysaccharide, tumor necrosis factor- α , and interleukin-1 β . *J Periodontol.* 2005; 76:437–442. [PubMed: 15857079]
15. Suryono, Kido J, Hayashi N, Kataoka M, Nagata T. Effect of *Porphyromonas gingivalis* lipopolysaccharide, tumor necrosis factor- α , and interleukin-1 β on calprotectin release in human monocytes. *J Periodontol.* 2003; 74:1719–1724. [PubMed: 14974811]
16. Jotwani R, Eswaran SV, Moonga S, Cutler CW. MMP-9/TIMP-1 imbalance induced in human dendritic cells by *Porphyromonas gingivalis*. *FEMS Immunol Med Microbiol.* 2009; 58:314–321. [PubMed: 20030715]
17. Baratelli FE, Heuze-Vourc'h N, Krysan K, et al. Prostaglandin E2-dependent enhancement of tissue inhibitors of metalloproteinases-1 production limits dendritic cell migration through extracellular matrix. *J Immunol.* 2004; 173:5458–5466. [PubMed: 15494493]
18. Osman M, Tortorella M, Londei M, Quarantino S. Expression of matrix metalloproteinases and tissue inhibitors of metalloproteinases define the migratory characteristics of human monocyte-derived dendritic cells. *Immunology.* 2002; 105:73–82. [PubMed: 11849317]
19. Wagsater D, Zhu C, Björck HM, Eriksson P. Effects of PDGF-C and PDGF-D on monocyte migration and MMP-2 and MMP-9 expression. *Atherosclerosis.* 2009; 202:415–423. [PubMed: 18573494]
20. Birkedal-Hansen H, Moore WG, Bodden MK, et al. Matrix metalloproteinases: a review. *Crit Rev Oral Biol Med.* 1993; 4:197–250. [PubMed: 8435466]
21. Newby AC. Metalloproteinase expression in monocytes and macrophages and its relationship to atherosclerotic plaque instability. *Arterioscler Thromb Vasc Biol.* 2008; 28:2108–2114. [PubMed: 18772495]
22. Fravallo P, Menard C, Bonnaure-Mallet M. Effect of *Porphyromonas gingivalis* on epithelial cell MMP-9 type IV collagenase production. *Infect Immun.* 1996; 64:4940–4945. [PubMed: 8945530]

23. Zhou J, Zhu P, Jiang JL, et al. Involvement of CD147 in overexpression of MMP-2 and MMP-9 and enhancement of invasive potential of PMA-differentiated THP-1. *BMC Cell Biol.* 2005; 6:25. [PubMed: 15904490]
24. Zhou J, Olson BL, Windsor LJ. Nicotine increases the collagen-degrading ability of human gingival fibroblasts. *J Periodontal Res.* 2007; 42:228–235. [PubMed: 17451542]
25. Zhou J, Windsor LJ. Porphyromonas gingivalis affects host collagen degradation by affecting expression, activation, and inhibition of matrix metalloproteinases. *J Periodontal Res.* 2006; 41:47–54. [PubMed: 16409255]
26. Zhou J, Windsor LJ. Heterogeneity in the collagen-degrading ability of Porphyromonas gingivalis-stimulated human gingival fibroblasts. *J Periodontal Res.* 2007; 42:77–84. [PubMed: 17214643]
27. Shah HN, Gharbia SE, Progulske-Fox A, Brocklehurst K. Evidence for independent molecular identity and functional interaction of the haemagglutinin and cysteine proteinase (gingivain) of Porphyromonas gingivalis. *J Med Microbiol.* 1992; 36:239–244. [PubMed: 1313881]
28. Auwerx J. The human leukemia cell line, THP-1: a multifaceted model for the study of monocyte-macrophage differentiation. *Experientia.* 1991; 47:22–31. [PubMed: 1999239]
29. Huang X, Griffiths M, Wu J, Farese RV Jr, Sheppard D. Normal development, wound healing, and adenovirus susceptibility in beta5-deficient mice. *Mol Cell Biol.* 2000; 20:755–759. [PubMed: 10629031]
30. Hajishengallis G, Wang M, Harokopakis E, Triantafilou M, Triantafilou K. Porphyromonas gingivalis fimbriae proactively modulate beta2 integrin adhesive activity and promote binding to and internalization by macrophages. *Infect Immun.* 2006; 74:5658–5666. [PubMed: 16988241]
31. Harokopakis E, Albzreh MH, Martin MH, Hajishengallis G. TLR2 transmodulates monocyte adhesion and transmigration via Rac1- and PI3K-mediated inside-out signaling in response to Porphyromonas gingivalis fimbriae. *J Immunol.* 2006; 176:7645–7656. [PubMed: 16751412]
32. Roth GA, Ankersmit HJ, Brown VB, Papapanou PN, Schmidt AM, Lalla E. Porphyromonas gingivalis infection and cell death in human aortic endothelial cells. *FEMS Microbiol Lett.* 2007; 272:106–113. [PubMed: 17459112]
33. Roth GA, Moser B, Roth-Walter F, et al. Infection with a periodontal pathogen increases mononuclear cell adhesion to human aortic endothelial cells. *Atherosclerosis.* 2007; 190:271–281. [PubMed: 16620832]
34. Sorsa T, Ingman T, Suomalainen K, et al. Identification of proteases from periodontopathogenic bacteria as activators of latent human neutrophil and fibroblast-type interstitial collagenases. *Infect Immun.* 1992; 60:4491–4495. [PubMed: 1398963]
35. Uitto VJ, Larjava H, Heino J, Sorsa T. A protease of Bacteroides gingivalis degrades cell surface and matrix glycoproteins of cultured gingival fibroblasts and induces secretion of collagenase and plasminogen activator. *Infect Immun.* 1989; 57:213–218. [PubMed: 2535833]
36. Birkedal-Hansen H. Role of matrix metalloproteinases in human periodontal diseases. *J Periodontol.* 1993; 64:474–484. [PubMed: 8315570]
37. Ding Y, Haapasalo M, Kerosuo E, Lounatmaa K, Kotiranta A, Sorsa T. Release and activation of human neutrophil matrix metallo- and serine proteinases during phagocytosis of Fusobacterium nucleatum, Porphyromonas gingivalis and Treponema denticola. *J Clin Periodontol.* 1997; 24:237–248. [PubMed: 9144046]
38. Grayson R, Douglas CW, Heath J, Rawlinson A, Evans GS. Activation of human matrix metalloproteinase 2 by gingival crevicular fluid and Porphyromonas gingivalis. *J Clin Periodontol.* 2003; 30:542–550. [PubMed: 12795793]
39. Pattamapun K, Tiranathanagul S, Yongchaitrakul T, Kuwatanasuchat J, Pavasant P. Activation of MMP-2 by Porphyromonas gingivalis in human periodontal ligament cells. *J Periodontal Res.* 2003; 38:115–121. [PubMed: 12608904]
40. DeCarlo AA Jr, Windsor LJ, Bodden MK, Harber GJ, Birkedal-Hansen B, Birkedal-Hansen H. Activation and novel processing of matrix metalloproteinases by a thiol-proteinase from the oral anaerobe Porphyromonas gingivalis. *J Dent Res.* 1997; 76:1260–1270. [PubMed: 9168859]
41. Zhou Q, Amar S. Identification of proteins differentially expressed in human monocytes exposed to Porphyromonas gingivalis and its purified components by high-throughput immunoblotting. *Infect Immun.* 2006; 74:1204–1214. [PubMed: 16428770]

42. Hajishengallis G, Tapping RI, Harokopakis E, et al. Differential interactions of fimbriae and lipopolysaccharide from *Porphyromonas gingivalis* with the Toll-like receptor 2-centred pattern recognition apparatus. *Cell Microbiol.* 2006; 8:1557–1570. [PubMed: 16984411]
43. Ogawa T, Hamada S. Hemagglutinating and chemotactic properties of synthetic peptide segments of fimbrial protein from *Porphyromonas gingivalis*. *Infect Immun.* 1994; 62:3305–3310. [PubMed: 7913697]
44. Zhou Q, Desta T, Fenton M, Graves DT, Amar S. Cytokine profiling of macrophages exposed to *Porphyromonas gingivalis*, its lipopolysaccharide, or its FimA protein. *Infect Immun.* 2005; 73:935–943. [PubMed: 15664935]
45. Mahanonda R, Pothiraksanon P, Sa-Ard-Iam N, et al. The effects of *Porphyromonas gingivalis* LPS and *Actinobacillus actinomycetemcomitans* LPS on human dendritic cells in vitro, and in a mouse model in vivo. *Asian Pac J Allergy Immunol.* 2006; 24:223–228. [PubMed: 17348245]
46. Travis J, Pike R, Imamura T, Potempa J. *Porphyromonas gingivalis* proteinases as virulence factors in the development of periodontitis. *J Periodontal Res.* 1997; 32:120–125. [PubMed: 9085221]
47. Imamura T, Potempa J, Pike RN, Travis J. Dependence of vascular permeability enhancement on cysteine proteinases in vesicles of *Porphyromonas gingivalis*. *Infect Immun.* 1995; 63:1999–2003. [PubMed: 7729914]
48. Nassar H, Chou HH, Khlgtian M, Gibson FC 3rd, Van Dyke TE, Genco CA. Role for fimbriae and lysine-specific cysteine proteinase gingipain K in expression of interleukin-8 and monocyte chemoattractant protein in *Porphyromonas gingivalis*-infected endothelial cells. *Infect Immun.* 2002; 70:268–276. [PubMed: 11748192]

Fig. 1. MMP-9 expression and activation in THP1 monocytes cultured with or without *P. gingivalis* supernatant in different concentrations. Cell culture media were collected and concentrated for zymography (A) and Western blot analysis (B) after 24 hours of incubation. 1. Control THP1 culture media, 2. THP1 treated with 100 μ g/ml of *P. gingivalis* LPS, 3. THP1 treated with 0.5% *P. gingivalis* supernatant, 4. THP1 treated with 1% *P. gingivalis* supernatant, 5. THP1 treated with 2.5% *P. gingivalis* supernatant, 6. THP1 treated with 5% *P. gingivalis* supernatant.

Fig. 2.

Protein and relative mRNA levels of TIMP-1 from THP1 monocytes after 24 hours of culture with or without *P. gingivalis* supernatant. (A). Western Blot analysis. (B). Real-time PCR analysis. Control: THP1 monocytes cultured in α -MEM media only. *Pg*: THP1 monocytes treated with 1% *P. gingivalis* supernatant. *Pg* LPS: THP1 monocytes treated with 100 μ g/ml of *P. gingivalis* LPS.

Fig. 3.
Relative mRNA levels (by real-time PCR analysis) of MMP-9 from THP1 monocytes after 24 hours of culture with or without *P. gingivalis* supernatant. Control: THP1 monocytes cultured in α -MEM media only. *Pg*: THP1 monocytes treated with 1% *P. gingivalis* supernatant. *Pg* LPS: THP1 monocytes treated with 100 μ g/ml of *P. gingivalis* LPS.

Fig. 4.

Transwell migration assay for human peripheral blood CD14⁺ monocytes. CD14⁺ monocytes were collected and placed in the upper chambers of transwell plates (10^5 cells/well in 250 μ l α -MEM supplied with 10ng/ml M-CSF). The lower chambers were filled with control media (500 μ l α -MEM) and different treatment media: *Pg* (α -MEM + 1% *P. gingivalis* supernatant), *Pg* LPS (α -MEM + 100 μ g/ml *P. gingivalis* LPS), boiled *Pg* (α -MEM + 1% denatured *P. gingivalis* supernatant), *Pg* + leupeptin (α -MEM + 1% *P. gingivalis* supernatant + 200 μ g/ml leupeptin), or *Pg* + GM6001 (α -MEM + 1% *P.*

gingivalis supernatant + 200 nM GM6001). Migrated cells were stained and counted under four random 40× microscope fields after 16 hours of incubation. The pictures were taken in 20× field.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript