Published in final edited form as:

Toxicol Appl Pharmacol. 2011 September 1; 255(2): 184–192. doi:10.1016/j.taap.2011.06.013.

Exposure to Diesel Exhaust Up-regulates iNOS Expression in ApoE Knockout Mice

Ni Bai^{1,2}, Takashi Kido², Terrance J. Kavanagh³, Joel D. Kaufman³, Michael E. Rosenfeld³, Cornelis van Breemen¹, and Stephan F. van Eeden²

¹Department of Anesthesiology, Pharmacology & Therapeutics, University of British Columbia, Vancouver, BC, Canada

²The James Hogg Research Centre, Providence Heart and Lung Institute, St. Paul's Hospital, University of British Columbia, Vancouver, BC, Canada

³Department of Occupational and Environmental Health, University of Washington, Seattle, WA, USA

Abstract

Traffic related particulate matter air pollution is a risk factor for cardiovascular events; however, the biological mechanisms are unclear. We hypothesize that diesel exhaust (DE) inhalation induces up-regulation of inducible nitric oxide synthase (iNOS), which is known to contribute to vascular dysfunction, progression of atherosclerosis and ultimately cardiovascular morbidity and mortality.

Methods—ApoE knockout mice (30-week) were exposed to DE (at 200 μ g/m³ of particulate matter) or filtered-air (control) for 7 weeks (6h/day, 5days/week). iNOS expression in the blood vessels and heart was evaluated by immunohistochemistry and western blotting analysis. To examine iNOS activity, thoracic aortae were mounted in a wire myograph, and vasoconstriction stimulated by phenylephrine (PE) was measured with and without the presence of the specific inhibitor for iNOS (1400W). NF- κ B (p65) activity was examined by ELISA. The mRNA expression of iNOS and NF- κ B (p65) was determined by real-time PCR.

Results—DE exposure significantly enhanced iNOS expression in the thoracic aorta (4-fold) and heart (1.5 fold). DE exposure significantly attenuated PE-stimulated vasoconstriction by ~20%, which was partly reversed by 1400W. The mRNA expression of iNOS and NF- κ B was significantly augmented after DE exposure. NF- κ B activity was enhanced 2-fold after DE inhalation, and the augmented NF- κ B activity was positively correlated with iNOS expression ($R^2=0.5998$).

Corresponding author: Stephan van Eeden, MD, PhD, **Mailing address:** 1081 Burrard Street, Vancouver, BC, Canada, V6Z1Y6, The James Hogg Research Centre, Providence Heart and Lung Institute, St. Paul's Hospital, University of British Columbia, Vancouver, BC, Canada, Stephan.vanEeden@hli.ubc.ca, **Phone:** 1-604-806-8346, **Fax:** 1-604-806-8351.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusions—We show that exposure to DE increases iNOS expression and activity possibly via NF- κ B-mediated pathway. We suspect that DE exposure-caused up-regulation of iNOS contributes to vascular dysfunction and atherogenesis, which could ultimately lead to urban air pollution-associated cardiovascular morbidity and mortality.

Keywords

Diesel exhaust; iNOS; NF- κ B; Atherosclerosis; ApoE knock out mice

1. Introduction

Numerous epidemiological studies have demonstrated an association between exposure to particulate matter air pollution with diameter less than 10 μ m (or called PM₁₀) and increased cardiovascular morbidity and mortality (Kaiser, 2000; Peters *et al.*, 2004; Pope, III *et al.*, 2004; Samet *et al.*, 2000; Miller *et al.*, 2007). Furthermore, recent studies have also shown that reducing PM₁₀ levels results in a declined in cardiovascular deaths by 10.3% (Clancy *et al.*, 2002; Laden *et al.*, 2006), suggesting a causal effect of PM₁₀ and cardiovascular mortality. It has been well established that deposition of airborne particles in the lung provokes low-grade alveolar inflammation with a secondary systemic inflammatory response resulting in downstream cardiovascular dysfunction. Atherosclerosis has been recognized as a chronic inflammatory disorder of blood vessels involving vascular, metabolic, and immune systems (Brook *et al.*, 2004; Ross, 1999). We previously showed that exposure to PM₁₀ caused progression of atherosclerosis of coronary arteries and aorta in rabbits that naturally developed atherosclerosis (Suwa *et al.*, 2002). This sentinel finding was confirmed in ApoE knockout mice (Sun *et al.*, 2005). In addition, human study also showed an association between progression of carotid atherosclerosis and the levels of ambient air pollution (Kunzli *et al.*, 2005, 2010). This progression of atherosclerosis induced by PM₁₀ exposure could contribute to the increased cardiovascular morbidity and mortality. The observations of PM₁₀-induced atherosclerosis progression are very compelling; however, the mechanisms whereby PM₁₀ exposure causes progression in atherosclerosis, have not been fully elucidated.

Diesel exhaust (DE) is a mixture of fine particles and gases, and represents a useful model of traffic-related air pollutants, which accounts for up to 90% of the fine particulate mass in ambient air of many major cities, such as London (Maheswaran *et al.*, 2005; Salvi *et al.*, 1999). A recent study demonstrated an association between progression of atherosclerosis and living near major roads (Kunzli *et al.*, 2010). In addition, evidence has shown that 6% of coronary heart disease deaths are linked to traffic-related pollution (Hoek *et al.*, 2002; Maheswaran *et al.*, 2005). Therefore, we selected to use DE to explore the underlying mechanisms of atherogenesis induced by ambient particulate matter exposure, using ApoE knockout mice that spontaneously develop atherosclerosis.

Inducible nitric oxide synthase (iNOS) is up-regulated in response to inflammatory cytokines as part of host defense responses (Stuehr *et al.*, 1987), and generates 100–1000 fold more nitric oxide (NO) than does endothelium NOS (eNOS) (micromolar vs. nanomolar levels). NO derived from eNOS plays an important role in protecting vasculature from

inflammation and atherosclerosis. However, excessive NO production from iNOS has detrimental effects on cardiovascular function. The large amount of locally released NO has been linked to the generation of harmful oxidative products, such as peroxynitrite, which is implicated in iNOS-mediated development of atherosclerosis (Ito *et al.*, 2000; Sun *et al.*, 2005; White *et al.*, 1994). iNOS is undetectable under normal physiological conditions, but its expression can be detected in macrophages, endothelial cells and smooth muscle cells of human atherosclerotic plaques (Wilcox *et al.*, 1997). The notion that iNOS plays a causative role in the progression of atherosclerosis is supported by the observation that atherosclerotic lesions were diminished in iNOS/ApoE double knockout mice, compared with ApoE knockout mice (Detmers *et al.*, 2000; Kuhlencordt *et al.*, 2001). iNOS overexpression was shown to be responsible for DE-induced lung inflammation (Ito *et al.*, 2000; Takano *et al.*, 1999), while iNOS knockout mice had a significant reduction of cytokines in the lung after exposure to ambient particles (Becher *et al.*, 2007). iNOS expression is up-regulated by NF- κ B, which is sensitive to inflammation and oxidative stress stimulation (Nathan *et al.*, 1994). We recently have shown that DE exposure accelerates the progression of atherosclerosis in ApoE mice (Hirano *et al.*, 2003). In this study, we test our hypothesis that DE exposure up-regulates iNOS expression and activity in vasculature via NF- κ B-mediated pathway.

2. Material and methods

2.1. DE generation

Characteristics of the exposure system have previously been described (Gould *et al.*, 2008). Briefly, DE was derived from a 2002 model turbocharged direct-injection 5.9-L Cummins B-series engine (6BT5.9G6; Cummins, Inc., Columbus, IN) in a generator set. Load was maintained at 75% of rated capacity, using a load-adjusting load bank (Simplex, Springfield, IL) throughout the exposures. We used No.2 undyed, on-highway fuel and Valvoline 15W-40 crankcase oil. All dilution air for the system was passed through HEPA and carbon filters, permitting a filtered air control exposure option with very low particulate and gaseous organic pollutant levels. The air entering the exposure room was conditioned to 18°C and 60% relative humidity. During exposures, DE concentrations were continuously measured and maintained at steady concentrations using a feedback controller monitoring fine particulate levels. Multistage samples collected on a micro-orifice uniform deposition impactor (MOUDI; MSP, Shoreview, MN) indicated a mass median diameter of 0.104 μ m.

2.2. Animals, exposure protocol, and sample collections

Male ApoE knockout mice were housed in a temperature- and humidity-controlled environment with a 12-h light/dark cycle with free access to water and standard rodent chow. At the age of 30 weeks, these mice were moved to a “Biozone” facility adjacent to the exposure chamber where exposure was controlled by opening or closing a valve to animal cages resulting in minimal stress for animals during the exposure period. We exposed ApoE knockout mice (10 mice/group) to DE for 7 weeks (5 days/week, 6 hrs/day) at the concentration of 200 μ g/m³ of particulate. Exposing mice to filtered air was the control. Animal procedures were approved by the Animal Care and Use Committee of the University of Washington.

After exposure, sodium pentobarbital (100mg/kg, Abbott Laboratories, IL) and heparin sulfate (500U/kg) were administered intraperitoneally. Upon the loss of all reflexes, the thoracic aorta, aortic root, and heart were carefully dissected from their connective tissues and kept at appropriate conditions until assay.

2.3. Measurement of vascular tone and iNOS activity

Thoracic aorta was carefully cleaned off all connective tissues without damaging the endothelium, and placed in ice-cold physiological salt solution (PSS). The vessels were cut to 2mm rings and mounted on a wire myograph (Model 610M; Danish Myo Technology, Aarhus, Denmark). Each vessel was bathed in oxygenated PSS at 37°C for an hour during which the resting tension was gradually increased to 6mN with three changes of PSS at 10minute intervals followed by stabilizing the vessels at resting tension (6mN) for 30min. Thereafter the vessels were stimulated with 80 mM KCl twice. Smooth muscle contractility was studied by the addition of cumulative concentrations of phenylephrine (PE, 1nM-10µM). To examine the impact of DE on iNOS activity, 1400W, a specific inhibitor for iNOS, was used. Vessels were incubated with 1400W (10µM) for 30min followed by addition of cumulative concentrations of PE (1nM-10µM). iNOS activity was determined by the fractional changes of maximal PE constriction with and without the presence of 1400W.

2.4. Immunohistochemistry of analysis of macrophages, smooth muscle cells, the expression of iNOS, CD36, and nitrotyrosine

Thoracic aorta tissue adjacent to those used for functional studies, aortic root sections contained three complete valve leaflets, and the heart were fixed with 10% neutral formalin for 24 hours, then embedded in paraffin. Sections (5 m) were deparaffinized in xylene and hydrated by passing through a series of graded alcohol. Thereafter, these sections were treated with citrate buffer (Invitrogen) to unmask the antigenic sites, and incubated with 10% goat serum at room temperature for 30min to block nonspecific binding proteins, followed by incubation with specific primary antibodies to: iNOS (1:100, Santa Cruz); macrophage (F4/80; 1:50, AbD Serotec); CD36 (1:50, Santa Cruz); smooth muscle cells (α -actin; 1:600, Abcam); and nitrotyrosine (1:400, Upstate Biotechnology) at 4°C overnight. Negative controls were included with non-immune isotype antibody or omission of the primary antibody. Subsequently, sections were incubated with biotinylated goat anti rabbit IgG (1:800, Vector Laboratories) at room temperature for 30min, followed by avidin-biotin conjugated alkaline phosphatase and Vector red (Vector Laboratories) to detect the antigen-antibody complexes.

Random images were captured by a spot digital camera (Nikon, Japan). We used Image Pro Plus software and a colour segmentation method to identify positive staining of the antigens of interest, and examined without knowledge of experimental groups. The volume fraction (V/v%) of specific staining was determined and normalized to the area of atherosclerotic plaque (aorta roots), and the thickness of vascular wall (thoracic aorta), respectively.

2.5. Western blotting analysis of iNOS expression in the heart

Frozen heart tissues were homogenized in 10 volumes of ice-cold RIPA buffer (Sigma) supplemented with protease inhibitor cocktail (Roche Diagnostics). Supernatant was

obtained following centrifugation of homogenates at 12,000 rpm for 15 min (4°C). The protein concentrations of supernatant were quantified by Bradford method using Bio-Rad DC Protein Assay kit according to the manufacturer's instructions (Biorad). Same amount of protein samples (40µg/lane) from different mice were separated on 7.5% SDS-PAGE gel, subsequently electrotransferred to polyvinylidene difluoride membranes (Biorad) and blocked with 5% BSA dissolved in 0.1% TBS-Tween 20 for 60min at room temperature. The membrane was incubated with specific primary antibody for iNOS (1:500, Santa Cruz) at 4°C overnight, followed by 1-hour incubation with horseradish peroxidase-conjugated goat anti-rabbit IgG (Santa Cruz). Immunoreactive bands were visualized by enhanced chemiluminescence kit (Amersham Life Sciences). Densitometry was carried out using Image J (NIH), and the results were expressed in optical intensity per mg protein.

2.6. ELISA of NF-κB (p65) activity

Nuclear extracts from frozen heart tissues were obtained using nuclear extract kit (Cayman Chemical). Protein concentrations of nuclear extract were quantified. NF-κB (p65) activity in the nuclear extracts from heart tissues was measured by enzyme-linked immunosorbent assay (ELISA) using NF-κB (p65) transcription factor kit according to the manufacturer's instructions (Cayman Chemical). This kit is a non-radioactive, sensitive method for detecting NF-κB (p65) DNA binding activity in nuclear extracts as a replacement for the radioactive electrophoretic mobility shift assay. Results were expressed as absorbance (450nm) per mg protein.

2.7. Real-Time Reverse Transcription Polymerase Chain Reaction (RT-PCR) of iNOS and NF-κB mRNA expression

Total RNA was extracted from heart tissues using RNeasy fibrous tissue mini kit (Qiagen). RNA concentrations were measured by Nanodrop (Thermo Scientific). Reverse transcription was performed using SuperScript III first-strand synthesis system for RT-PCR (Invitrogen). The same amount of RNA was loaded in triplicates for each assay, and RT-PCR was performed using TaqMan universal PCR master mix and Taqman gene expression assay system (Applied Biosystems) according to the manufacturer's instructions. The mRNA expression of iNOS, NF-κB, β-actin, and hypoxanthine phosphoribosyltransferase-1 (HPRT1) was determined by RT-PCR using ABI Prism 7900HT sequence detection system (Applied Biosystems). Gene expression values were calculated based on the comparative threshold cycle method, normalized to the expression values of β-actin and HPRT1, and displayed as ratio relative to β-actin.

2.9. Solutions and chemicals

The PSS consisted of the following (in mM): NaCl 119, KCl 4.7, KH₂PO₄ 1.18, NaHCO₃ 24, MgSO₄·7H₂O 1.17, CaCl₂ 1.6, glucose 5.5 and EDTA 0.026. All reagents were purchased from Sigma.

2.10. Statistical analysis

Data are shown as mean±SEM. The statistical significance was evaluated using the unpaired Student's t test for simple comparison between two values. Non-parametric analysis

(Kruskal-Wallis test) was used to analyze data that were not normally distributed and Bonferroni correction was used for multiple comparisons. Linear regression modeling was used to assess the relationship between parameters. In all experiments, n equals the number of mice from which samples were obtained. $P < 0.05$ was considered to be significant.

3. Results

3.1. iNOS expression in the thoracic aorta

Immunohistochemical staining of thoracic aorta sections demonstrates that iNOS was predominantly expressed by smooth muscle cells and macrophages in the vessels (Fig 1A,B,C). Immunohistochemistry analysis shows that iNOS expression was significantly elevated in the thoracic aorta after DE exposure ($11.9 \pm 3.2\%$ vs. $40.2 \pm 6.5\%$; V/v% of staining in filtered air vs. DE; $p < 0.002$) (Fig 1D,E,F).

3.2. iNOS expression in heart tissues

Western blotting analysis shows that DE exposure significantly augmented iNOS expression in heart tissues ($0.7 \pm 0.1\%$ vs. $1.0 \pm 0.1\%$; filtered air vs. DE; $p < 0.03$) (Fig 2A,B).

3.3. Similarly localized expression of iNOS with CD36 and nitrotyrosine in aortic root lesions

iNOS was predominantly expressed by macrophages in aortic root sections (Fig 3B,C). Serial sections of aortic root from the same mouse shows that the expression of iNOS, CD36 and nitrotyrosine was similarly localized and predominantly expressed by foam cells (Fig 3D,E,F). In addition, iNOS expression was positively associated with CD36 expression ($R^2 = 0.4580$, $P < 0.01$) (Fig 3G), and CD36 expression correlated with nitrotyrosine formation ($R^2 = 0.4629$, $P < 0.01$) (Fig 3H), suggesting a close relationship between iNOS and these oxidative stress markers.

3.4. iNOS activity in the thoracic aorta

PE-stimulated vasoconstriction of the thoracic aorta was significantly suppressed after DE exposure ($9.5 \pm 0.6\text{mN}$ vs. $7.4 \pm 0.6\text{mN}$, filtered air vs. DE; $p < 0.02$) (Fig 4A,B). In the presence of 1400W, the fractional change of PE-elicited maximal constriction was significantly increased in DE exposure group, compared with the control ($5.2 \pm 3.0\%$ vs. $24.1 \pm 7.6\%$; filtered air vs. DE; $p < 0.04$) (Fig 4C,D,E), suggesting an increase in iNOS activity after DE exposure. To verify whether the reduced vasoconstriction was due to morphological abnormalities, we stained the thoracic aorta adjacent to the segments for functional studies with hematoxylin and eosin, and examined the thickness and the structure of the vessels using morphologic analysis. We found no difference in the thickness and structure of the vessels after DE exposure (data not shown).

3.5. The mRNA expression of iNOS and NF- κ B (p65)

The mRNA expression of both iNOS ($76 \pm 3\%$ vs $90 \pm 6\%$; filtered air vs. DE; $p < 0.004$) (Fig 5A) and NF- κ B ($83 \pm 1\%$ vs $88 \pm 3\%$; filtered air vs. DE; $p < 0.01$) (Fig 5B) were significantly higher in heart tissues after DE exposure.

3.6. NF- κ B (p65) activity

To assess whether iNOS was regulated at the transcription level via NF- κ B activation, we extracted nuclear protein from frozen hearts, and NF- κ B activity was evaluated using ELISA. DE exposure up-regulated NF- κ B (p65) activity (0.014 ± 0.0004 vs. 0.029 ± 0.006 absorbance of 450nm/mg protein; filtered air vs. DE; $p < 0.01$) (Fig 6A). iNOS expression was positively correlated with NF- κ B activity ($R^2 = 0.5998$, $p < 0.0005$) (Fig 6B), indicating that DE exposure up-regulated iNOS expression at the transcriptional level via NF- κ B activation (p65).

4. Discussion

Studies from our (Bai *et al.*, 2011; Suwa *et al.*, 2002) and other (Sun *et al.*, 2002; Kunzli *et al.*, 2005, 2010), laboratories have shown that exposure to PM₁₀ that was derived from different sources is associated with development of atherosclerosis. In this study, we show that exposure to DE, (urban particulate matter) increased iNOS expression in the thoracic aorta and heart (Fig 1,2). DE inhalation also enhanced iNOS activity in blood vessels, which resulted in decreased vascular contractility (Fig 4). In this study, we show that iNOS was expressed by smooth muscle cells, macrophages and cardiac myocytes, and the expression of iNOS, CD36 and nitrotyrosine was similarly localized in aortic lesions (Fig 3D,E,F). The positive association between enhanced NF- κ B activity and iNOS expression (Fig 6B) suggests that activation of transcription factor NF- κ B played a key role in this enhanced iNOS activity. The augmented mRNA expression of iNOS and NF- κ B (Fig 6) also indicates that iNOS was up-regulated at both transcription and translation levels. To our knowledge, this is first study demonstrating that DE exposure up-regulates iNOS activity and expression in the vasculature and heart, and we speculate that oxidative stress induced NF- κ B activation contributes to the increased iNOS expression and activity.

4.1. The level of DE exposure

In this study, we exposed ApoE knockout mice fed with regular chow to DE for 7 weeks using a well-controlled inhalation system that mimics real-world exposure. Our average DE concentration throughout a 24-hour period is around $35 \mu\text{g}/\text{m}^3$ (averaged concentration by 30h/week exposure of DE at the concentration of $200 \mu\text{g}/\text{m}^3$ for 7 weeks), which is environmentally relevant and within the National Ambient Air Quality Standard (Brook *et al.*, 2004).

4.2. DE exposure and up-regulation of iNOS

Nitric oxide (NO) is an essential biological mediator protecting the vessels from inflammation and atherosclerosis; however, overproduction of NO catalyzed by iNOS is cytotoxic. iNOS is not expressed under normal physiological conditions, but can be up-regulated in response to pro-inflammatory mediators, and oxidative stress stimulation. The up-regulation of iNOS is implicated in the pathogenesis of a number of cardiovascular diseases, including the progression in atherosclerosis (Cromheeke *et al.*, 1999). In this study, we show that DE exposure increased iNOS activity (Fig 4), and enhanced iNOS expression in blood vessels (Fig 1) and heart (Fig 2) at both protein and mRNA levels (Fig 6A). We also found a positive correlation between iNOS expression with foam cell formation and the

number of smooth muscle cells in aortic lesions (Supplementary Fig A.1). These associations suggest that iNOS contributes to the progression of atherosclerosis induced by DE inhalation.

Similar to the finding of Sun and colleagues in which ApoE knockout mice were exposed to concentrated ambient particles for 6 months (Sun *et al.*, 2005), we found that iNOS expression was up-regulated at both protein and mRNA levels after exposure to DE for a shorter exposure period (7 weeks vs. 6 months), suggesting that the constituents of DE may be more active, compared with concentrated ambient particles (Hirano, Furuyama, Koike, and Kobayashi, 2003b). Sun and colleagues found that exposure to concentrated ambient particles enhanced vasoconstriction to phenylephrine, which likely resulted from reduced NO production or endothelial dysfunction. In contrast, we found suppressed vasoconstriction to phenylephrine, which was likely due to excessive NO production derived from iNOS, because the inhibition of iNOS enhanced the constriction (Fig 4D,E).

Diesel exhaust is a complex mixture of gases and diesel particulate matter. The gas-phase pollutants include ozone, sulphur dioxide, carbon monoxide (CO), and nitrogen oxides (NO and NO₂), and the particulate components consist of a central carbon core nucleus with an estimated 18,000 combustion products adsorbed, including organic chemicals, such as polycyclic aromatic hydrocarbons, and transition metals. The composition of DE (gas- and particulate-phase) is different from other ambient particulate (such as concentrated ambient particles) exposed in different studies, and the variation may explain the discrepancy of vascular effects. In our study, the concentrations of CO and NO were lower than those in other studies (Campen *et al.*, 2010; Quan *et al.*, 2010), despite that NO in the DE exposure group (1515.80ppb) was higher than filtered air group (38.63ppb) in the present study. Our study did not address the differential effect of DE composition on vascular function, because a number of studies have examined the impact of different composition of gasoline and diesel exhaust on cardiovascular function. For example, Campen and colleagues reported that CO and NO, but not NO₂, had greater effect on ET-1 mRNA expression in thoracic aorta (Campen *et al.*, 2010b). Furthermore, organic compounds are not only potential occupational carcinogens (Mauderly *et al.*, 2008), but can also generate oxidative stress in rat heart microvessel endothelial cells (Hirano *et al.*, 2003), and induce pro-inflammatory responses in human aortic endothelial cells (Li *et al.*, 2010), which may contribute to myocardial ischemia associated with non-particulate matter component of DE (Campen *et al.*, 2005).

We found no impairment of ACh-stimulated endothelium-dependent relaxation (unpublished data). These data indicate that effect on smooth muscle cells is prominent after DE inhalation for 7 weeks. Sun and colleagues also observed that mice fed with regular chow did not develop endothelial dysfunction after exposure to concentrated ambient particles (supporting our finding), suggesting that high fat diet-mediated modifications on vasculature are important elements and may have synergistic effect on ambient particle exposure-induced vascular dysfunction. Data from a recent human study have implied that increased lipid levels contribute to atherosclerosis associated with ambient particle exposure (Bröske *et al.*, 2011)

iNOS activation is mainly regulated at the transcriptional level by NF- κ B (Taylor *et al.*, 1998). In this study, we found that NF- κ B activity and mRNA expression were significantly increased (Fig 5B & Fig 6A). The positive correlation between iNOS and increased NF- κ B activity (Fig 6B) suggests that NF- κ B plays an important role in the up-regulation of iNOS following DE exposure. NF- κ B is known to regulate the transcription of various genes participating in oxidative stress responses. NF- κ B dimers (p50/65) are sequestered in the cytoplasm by a family of inhibitors, called inhibitors of κ B (I κ B). Activation of NF- κ B is initiated by degradation of I κ B proteins triggered by oxidative stress. NF- κ B complex enters into the nucleus in which it can turn on the expression of specific genes by binding to their promoters. The promoter of the murine gene coding for iNOS contains the binding sites for NF- κ B, which are sensitive to oxidative stress stimulation (Xie *et al.*, 1993). It was reported that exposure to ambient particles caused oxidative stress-induced NF- κ B activation in epithelial cells (Shukla *et al.*, 2000). Exposure to diesel exhaust particles was found to cause ROS generation in epithelial cells, alveolar macrophages, blood vessels, endothelial cells, and cardiac myocytes (Ikeda *et al.*, 1995; Ito *et al.*, 2000; Okayama *et al.*, 2006; Shukla *et al.*, 2000; Bai *et al.*, 2001). We recently demonstrated that DE inhalation increased systemic and tissue (aortic root) oxidative stress (Bai *et al.*, 2011). In the present study, we show that CD36 and nitrotyrosine are concomitantly up-regulated with iNOS after DE inhalation (Fig 3), suggesting that ROS could play a key role in the activation of NF- κ B, whereby up-regulates iNOS expression and activation. ROS generation may arise directly from the surface of ambient particles (Brown *et al.*, 2000), soluble compounds (e.g., transition metals) (Donaldson *et al.*, 2001; Li *et al.*, 2002), or organic compounds (Baulig *et al.*, 2003; Hirano *et al.*, 2003; Sagai *et al.*, 1993). In addition, direct interactions between DE, airway epithelial cells and alveolar macrophages may also contribute to ROS generation (Bai *et al.*, 2007; Rudell *et al.*, 1999).

4.3. iNOS and Atherosclerosis

The notion that overexpression of iNOS has causal effect on the development of atherosclerosis has previously been established by showing that atherosclerotic lesions were attenuated when iNOS was knocked out in ApoE mice (Detmers *et al.*, 2000; Kuhlencordt *et al.*, 2001). iNOS is capable of generating superoxide (Xia *et al.*, 1997; Xia *et al.*, 1998), which can rapidly deplete endothelium-derived NO, attenuate eNOS activity and expression, leading to endothelial dysfunction and development of atherosclerosis. It is well known that oxidative modification of low-density lipoprotein (LDL) accelerates lipoprotein-uptake by macrophages and the formation of foam cells (Krieger *et al.*, 1993), which is the characteristic component of atherosclerotic lesions and landmark of development of atherosclerosis. CD36 is one of the major macrophage scavenger receptor that uptakes oxidized LDL (oxLDL), and is also known as oxLDL receptor (Endemann *et al.*, 1993). We recently reported that DE inhalation increased CD36 expression (Bai *et al.*, 2011). CD36 is up-regulated by proatherogenic cytokines and oxidized lipid. In this study we show that the expression of iNOS and CD36 was co-localized in atherosclerotic lesions (Fig 3). The positive correlation between iNOS and CD36 (Fig 3G) suggests that iNOS, as a free radical generator or a surrogated marker for ROS, may contribute to the augmented CD36 expression leading to the foam formation and progression of atherosclerosis. In addition, up-regulation of iNOS also contributes to the generation of peroxynitrite, which is a potent

oxidant and implicated in the pathogenesis of several cardiovascular diseases, including atherosclerosis, myocardial infarction, and heart failure (Graham *et al.*, 1993).

5. Conclusions

In summary, we show that DE inhalation causes significant up-regulation of iNOS expression and activity in blood vessels and heart. Our data indicate that the up-regulation of iNOS was mediated by the activation of NF- κ B. This study provides some novel insights into the mechanisms underlying DE inhalation-induced progression of atherosclerosis, including increased lipid deposition, increased macrophages and smooth muscle cells in the lesion (Bai *et al.* 2011). iNOS may be a potential target to decrease the impact of air pollution on cardiovascular morbidity and mortality.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH grants P50ES015915 (MER, TJK, JDK, SVE), R01ES13434 (MER), K24ES013195 (JDK), the Heart and Stroke Foundation of Canada (SVE, NB), and Michael Smith Foundation for Health Research (NB). SVE is a Senior Scholar with the Michael Smith Foundation for Health Research and CIHR/GSK professor in Chronic Obstructive Pulmonary Disease.

Abbreviations

CO	Carbon Monoxide
DE	Diesel Exhaust
ELISA	Enzyme-Linked Immunosorbent Assay
GM-CSF	Granulocyte-Macrophage Colony-Stimulating Factor
HPRT1	Hypoxanthine Phosphoribosyltransferase-1
iNOS	Inducible Nitric Oxide Synthase
IL-1	Interleukin-1
MIP	Macrophage Inflammatory Protein
NF-κB	Nuclear Factor-kappa B
NO	Nitric Oxide
NT	Nitrotyrosine
OxLDL	Oxidized Low-Density-Lipoprotein
PE	Phenylephrine
PM₁₀	Particulate matter air pollution with the diameter less than 10 μ m
ROS	Reactive Oxidative Species
RT-PCR	Real Time Reverse Transcription Polymerase Chain Reaction

TNF

Tumour Necrosis Factor

References

1. Bai N, Khazaei M, van Eeden SF, Laher I. The pharmacology of particulate matter air pollution-induced cardiovascular dysfunction. *Pharmacol.Ther.* 2007; 113:16–29. [PubMed: 16920197]
2. Bai N, Takashi K, Suzuki H, Yang G, Kavanagh TJ, Kaufman JD, Rosenfeld ME, van Breemen C, van Eeden SF. Changes in atherosclerotic plaques induced by inhalation of diesel exhaust. *Atherosclerosis*. 2011 (in press).
3. Baulig A, Garlatti M, Bonvallot V, Marchand A, Barouki R, Marano F, Baeza-Squiban A. Involvement of reactive oxygen species in the metabolic pathways triggered by diesel exhaust particles in human airway epithelial cells. *Am.J.Physiol Lung Cell Mol.Physiol.* 2003; 285:L671–L679. [PubMed: 12730081]
4. Becher R, Bucht A, Ovrevik J, Hongslo JK, Dahlman HJ, Samuelsen JT, Schwarze PE. Involvement of NADPH oxidase and iNOS in rodent pulmonary cytokine responses to urban air and mineral particles. *Inhal.Toxicol.* 2007; 19:645–655. [PubMed: 17510837]
5. Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, Luepker R, Mittleman M, Samet J, Smith SC Jr, Tager I. Air pollution and cardiovascular disease: a statement for healthcare professionals from the Expert Panel on Population and Prevention Science of the American Heart Association. *Circulation.* 2004; 109:2655–2671. [PubMed: 15173049]
6. Brown DM, Stone V, Findlay P, Macnee W, Donaldson K. Increased inflammation and intracellular calcium caused by ultrafine carbon black is independent of transition metals or other soluble components. *Occup.Environ.Med.* 2000; 57:685–691. [PubMed: 10984341]
7. Campen MJ, Babu NS, Helms GA, Pett S, Wernly J, Mehran R, McDonald JD. Nonparticulate components of diesel exhaust promote constriction in coronary arteries from ApoE^{−/−} mice. *Toxicol.Sci.* 2005; 88:95–102. [PubMed: 16093524]
8. Campen MJ, Lund AK, Doyle-Eisele ML, McDonald JD, Knuckles TL, Rohr AC, Knipping EM, Mauderly JL. A comparison of vascular effects from complex and individual air pollutants indicates a role for monoxide gases and volatile hydrocarbons. *Environ.Health Perspect.* 2010b; 118:921–927. [PubMed: 20197249]
9. Clancy L, Goodman P, Sinclair H, Dockery DW. Effect of air-pollution control on death rates in Dublin, Ireland: an intervention study. *Lancet.* 2002; 360:1210–1214. [PubMed: 12401247]
10. Cromheeke KM, Kockx MM, De Meyer GR, Bosmans JM, Bult H, Beelaerts WJ, Vrints CJ, Herman AG. Inducible nitric oxide synthase colocalizes with signs of lipid oxidation/peroxidation in human atherosclerotic plaques. *Cardiovasc.Res.* 1999; 43:744–754. [PubMed: 10690346]
11. Detmers PA, Hernandez M, Mudgett J, Hassing H, Burton C, Mundt S, Chun S, Fletcher D, Card DJ, Lisnock J, Weikel R, Bergstrom JD, Shevell DE, Hermanowski-Vosatka A, Sparrow CP, Chao YS, Rader DJ, Wright SD, Pure E. Deficiency in inducible nitric oxide synthase results in reduced atherosclerosis in apolipoprotein E-deficient mice. *J.Immunol.* 2000; 165:3430–3435. [PubMed: 10975863]
12. Donaldson K, MacNee W. Potential mechanisms of adverse pulmonary and cardiovascular effects of particulate air pollution (PM10). *Int.J.Hyg.Environ.Health.* 2001; 203:411–415. [PubMed: 11556145]
13. Endemann G, Stanton LW, Madden KS, Bryant CM, White RT, Protter AA. CD36 is a receptor for oxidized low density lipoprotein. *J.Biol.Chem.* 1993; 268:11811–11816. [PubMed: 7685021]
14. Gould T, Larson T, Stewart J, Kaufman JD, Slater D, McEwen N. A controlled inhalation diesel exhaust exposure facility with dynamic feedback control of PM concentration. *Inhal.Toxicol.* 2008; 20:49–52. [PubMed: 18236222]
15. Graham A, Hogg N, Kalyanaraman B, O'Leary V, rley-Usmar V, Moncada S. Peroxynitrite modification of low-density lipoprotein leads to recognition by the macrophage scavenger receptor. *FEBS Lett.* 1993; 330:181–185. [PubMed: 8365489]

16. Hirano S, Furuyama A, Koike E, Kobayashi T. Oxidative-stress potency of organic extracts of diesel exhaust and urban fine particles in rat heart microvessel endothelial cells. *Toxicology*. 2003; 187:161–170. [PubMed: 12699905]
17. Hoek G, Brunekreef B, Goldbohm S, Fischer P, van den Brandt PA. Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet*. 2002; 360:1203–1209. [PubMed: 12401246]
18. Ikeda M, Suzuki M, Watarai K, Sagai M, Tomita T. Impairment of endothelium-dependent relaxation by diesel exhaust particles in rat thoracic aorta. *Jpn.J.Pharmacol*. 1995; 68:183–189. [PubMed: 7563975]
19. Ito T, Ikeda M, Yamasaki H, Sagai M, Tomita T. Peroxynitrite formation by diesel exhaust particles in alveolar cells: Links to pulmonary inflammation. *Environ.Toxicol.Pharmacol*. 2000; 9:1–8. [PubMed: 11137462]
20. Kaiser J. Air pollution. Evidence mounts that tiny particles can kill. *Science*. 2000; 289:22–23. [PubMed: 10928919]
21. Krieger M, Acton S, Ashkenas J, Pearson A, Penman M, Resnick D. Molecular flypaper, host defense, and atherosclerosis. Structure, binding properties, and functions of macrophage scavenger receptors. *J.Biol.Chem*. 1993; 268:4569–4572. [PubMed: 8383115]
22. Kuhlencordt PJ, Chen J, Han F, Astern J, Huang PL. Genetic deficiency of inducible nitric oxide synthase reduces atherosclerosis and lowers plasma lipid peroxides in apolipoprotein E-knockout mice. *Circulation*. 2001; 103:3099–3104. [PubMed: 11425775]
23. Kunzli N, Jerrett M, Garcia-Esteban R, Basagana X, Beckermann B, Gilliland F, Medina M, Peters J, Hodis HN, Mack WJ. Ambient air pollution and the progression of atherosclerosis in adults. *PLoS.One*. 2010; 5:e9096. [PubMed: 20161713]
24. Kunzli N, Jerrett M, Mack WJ, Beckerman B, LaBree L, Gilliland F, Thomas D, Peters J, Hodis HN. Ambient air pollution and atherosclerosis in Los Angeles. *Environ.Health Perspect*. 2005; 113:201–206. [PubMed: 15687058]
25. Laden F, Schwartz J, Speizer FE, Dockery DW. Reduction in fine particulate air pollution and mortality: Extended follow-up of the Harvard Six Cities study. *Am.J.Respir.Crit Care Med*. 2006; 173:667–672. [PubMed: 16424447]
26. Li N, Wang M, Oberley TD, Sempf JM, Nel AE. Comparison of the pro-oxidative and proinflammatory effects of organic diesel exhaust particle chemicals in bronchial epithelial cells and macrophages. *J.Immunol*. 2002; 169:4531–4541. [PubMed: 12370390]
27. Li R, Ning Z, Majumdar R, Cui J, Takabe W, Jen N, Sioutas C, Hsiai T. Ultrafine particles from diesel vehicle emissions at different driving cycles induce differential vascular pro-inflammatory responses: implication of chemical components and NF-kappaB signaling. *Part Fibre.Toxicol*. 2010; 7:6. [PubMed: 20307321]
28. Maheswaran R, Haining RP, Brindley P, Law J, Pearson T, Fryers PR, Wise S, Campbell MJ. Outdoor air pollution, mortality, and hospital admissions from coronary heart disease in Sheffield, UK: a small-area level ecological study. *Eur.Heart J*. 2005; 26:2543–2549. [PubMed: 16166102]
29. Mauderly JL, Chow JC. Health effects of organic aerosols. *Inhal.Toxicol*. 2008; 20:257–288. [PubMed: 18300047]
30. Miller KA, Siscovick DS, Sheppard L, Shepherd K, Sullivan JH, Anderson GL, Kaufman JD. Long-term exposure to air pollution and incidence of cardiovascular events in women. *N.Engl.J.Med*. 2007; 356:447–458. [PubMed: 17267905]
31. Nathan C, Xie QW. Regulation of biosynthesis of nitric oxide. *J.Biol.Chem*. 1994; 269:13725–13728. [PubMed: 7514592]
32. Okayama Y, Kuwahara M, Suzuki AK, Tsubone H. Role of reactive oxygen species on diesel exhaust particle-induced cytotoxicity in rat cardiac myocytes. *J.Toxicol.Environ.Health A*. 2006; 69:1699–1710. [PubMed: 16864420]
33. Peters A, von KS, Heier M, Trentinaglia I, Hormann A, Wichmann HE, Lowel H. Exposure to traffic and the onset of myocardial infarction. *N.Engl.J.Med*. 2004; 351:1721–1730. [PubMed: 15496621]
34. Pope CA III, Burnett RT, Thurston GD, Thun MJ, Calle EE, Krewski D, Godleski JJ. Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological

- evidence of general pathophysiological pathways of disease. *Circulation*. 2004; 109:71–77. [PubMed: 14676145]
35. Quan C, Sun Q, Lippmann M, Chen LC. Comparative effects of inhaled diesel exhaust and ambient fine particles on inflammation, atherosclerosis, and vascular dysfunction. *Inhal.Toxicol*. 2010; 22:738–753. [PubMed: 20462391]
  36. Ross R. Atherosclerosis—an inflammatory disease. *N.Engl.J.Med*. 1999; 340:115–126. [PubMed: 9887164]
  37. Rudell B, Blomberg A, Helleday R, Ledin MC, Lundback B, Stjernberg N, Horstedt P, Sandstrom T. Bronchoalveolar inflammation after exposure to diesel exhaust: comparison between unfiltered and particle trap filtered exhaust. *Occup.Enviroin.Med*. 1999; 56:527–534. [PubMed: 10492649]
  38. Sagai M, Saito H, Ichinose T, Kodama M, Mori Y. Biological effects of diesel exhaust particles. I. In vitro production of superoxide and in vivo toxicity in mouse. *Free Radic.Biol.Med*. 1993; 14:37–47. [PubMed: 8384149]
  39. Salvi SS, Frew A, Holgate S. Is diesel exhaust a cause for increasing allergies? *Clin.Exp.Allergy*. 1999; 29:4–8. [PubMed: 10051695]
  40. Samet JM, Dominici F, Curriero FC, Coursac I, Zeger SL. Fine particulate air pollution and mortality in 20 U.S cities 1987–1994. *N.Engl.J.Med*. 2000; 343:1742–1749. [PubMed: 11114312]
  41. Shukla A, Timblin C, BeruBe K, Gordon T, McKinney W, Driscoll K, Vacek P, Mossman BT. Inhaled particulate matter causes expression of nuclear factor (NF)-kappaB-related genes and oxidant-dependent NF-kappaB activation in vitro. *Am.J.Respir.Cell Mol.Biol*. 2000; 23:182–187. [PubMed: 10919984]
  42. Stuehr DJ, Marletta MA. Induction of nitrite/nitrate synthesis in murine macrophages by BCG infection, lymphokines, or interferon-gamma. *J.Immunol*. 1987; 139:518–525. [PubMed: 3110273]
  43. Sun Q, Wang A, Jin X, Natanzon A, Duquaine D, Brook RD, Aguinaldo JG, Fayad ZA, Fuster V, Lippmann M, Chen LC, Rajagopalan S. Long-term air pollution exposure and acceleration of atherosclerosis and vascular inflammation in an animal model. *JAMA*. 2005; 294:3003–3010. [PubMed: 16414948]
  44. Suwa T, Hogg JC, Quinlan KB, Ohgami A, Vincent R, van Eeden SF. Particulate air pollution induces progression of atherosclerosis. *J.Am.Coll.Cardiol*. 2002; 39:935–942. [PubMed: 11897432]
  45. Takano H, Lim HB, Miyabara Y, Ichinose T, Yoshikawa T, Sagai M. Manipulation of the L-arginine-nitric oxide pathway in airway inflammation induced by diesel exhaust particles in mice. *Toxicology*. 1999; 139:19–26. [PubMed: 10614685]
  46. Taylor BS, de Vera ME, Ganster RW, Wang Q, Shapiro RA, Morris SM Jr, Billiar TR, Geller DA. Multiple NF-kappaB enhancer elements regulate cytokine induction of the human inducible nitric oxide synthase gene. *J.Biol.Chem*. 1998; 273:15148–15156. [PubMed: 9614127]
  47. White CR, Brock TA, Chang LY, Crapo J, Briscoe P, Ku D, Bradley WA, Gianturco SH, Gore J, Freeman BA. Superoxide and peroxynitrite in atherosclerosis. *Proc.Natl.Acad.Sci.U.S.A*. 1994; 91:1044–1048. [PubMed: 8302829]
  48. Wilcox JN, Subramanian RR, Sundell CL, Tracey WR, Pollock JS, Harrison DG, Marsden PA. Expression of multiple isoforms of nitric oxide synthase in normal and atherosclerotic vessels. *Arterioscler.Thromb.Vasc.Biol*. 1997; 17:2479–2488. [PubMed: 9409218]
  49. Xia Y, Roman LJ, Masters BS, Zweier JL. Inducible nitric-oxide synthase generates superoxide from the reductase domain. *J.Biol.Chem*. 1998; 273:22635–22639. [PubMed: 9712892]
  50. Xia Y, Zweier JL. Superoxide and peroxynitrite generation from inducible nitric oxide synthase in macrophages. *Proc.Natl.Acad.Sci.U.S.A*. 1997; 94:6954–6958. [PubMed: 9192673]
  51. Xie QW, Whisnant R, Nathan C. Promoter of the mouse gene encoding calcium-independent nitric oxide synthase confers inducibility by interferon gamma and bacterial lipopolysaccharide. *J.Exp.Med*. 1993; 177:1779–1784. [PubMed: 7684434]

Highlights

Exposed ApoE knockout mice (30-week) to diesel exhaust (DE) for 7 weeks

Examine iNOS expression and activity in the blood vessels and heart.

DE exposure enhanced iNOS protein and mRNA expression in the aorta and heart

iNOS activity was also increased after DE exposure

This up-regulation of iNOS may contribute to vascular dysfunction and atherogenesis.

Figure 1. Immunohistochemical analysis of iNOS expression in the thoracic aorta
 Representative photomicrographs of thoracic aorta sections stained for A) iNOS (400X) (arrow); B) macrophages (F4/80) (400X) (arrowhead); C) smooth muscle cells (α -actin) (400X) (arrow); representative photomicrographs of thoracic aorta sections stained for iNOS exposed to filtered air (D) or DE (E) (600X); F) Exposure to DE increased iNOS expression in the thoracic aorta, n=6 (filtered air) and n=8 (DE), *P<0.002. L: lumen. Values are mean $\pm$ SEM.

Figure 2. Western blotting analysis of iNOS expression in the heart

A) Representative photomicrograph of western blotting of iNOS expression in the heart; B) Western blotting analysis demonstrates that iNOS expression was enhanced following DE exposure, $n=8$, $*P<0.03$. Values are mean $\pm$ SEM.

A **B** **C**

Figure 3. iNOS expression in atherosclerotic plaque in aortic root and its relationship with the expression of CD36 and nitrotyrosine (NT)
 Representative photomicrographs of aortic root sections stained for A) negative control, B) iNOS, and C) macrophages (F4/80) (400X); Representative photomicrographs of aortic root sections showing similarly localized expression of D) iNOS, E) CD36, and F) NT; G) Positive correlation of the expression between iNOS and CD36, $R^2=0.4580$, $P<0.01$; H) Positive correlation of the expression between CD36 and NT, $R^2=0.4629$, $P<0.01$.

Figure 4. Reduced vasoconstriction and increased iNOS activity in the thoracic aorta

A) Dose-response curves of PE-elicited constriction show that exposure to DE attenuated vasoconstriction, $n=9$, $*P<0.02$; B) DE exposure caused significant attenuation of maximum vasoconstriction, compared with the filtered air exposure, $n=9$, $*P<0.02$; C) In the control group, the presence of iNOS blocker (1400W) had no effect on PE-elicited constriction; D) In DE exposure group, the reduced vasoconstriction was partly restored by 1400W; E) Inhibition of iNOS by 1400W caused a significant elevation of maximal vasoconstriction in DE exposure group, $n=9$, $*P<0.04$. Values are mean $\pm$ SEM.

Figure 5. Increased mRNA expression of iNOS and NF- κ B in the heart

A) Increased mRNA expression of iNOS in the heart after DE exposure, n=7 (filtered air) and n=8 (DE), *P<0.004; B) Exposure to DE increased NF- κ B mRNA expression, n=7 (filtered air), n=8 (DE), *P<0.01. Values are mean $\pm$ SEM.

Figure 6. DE exposure increased NF-κB activity of heart nuclear extracts

A) Exposure to DE increased NF-κB activity of heart nuclear extracts, n=9 (filtered air), n=10 (DE), *P<0.01; B) Positive correlation between NF-κB activity and iNOS expression, $R^2=0.5998$, P<0.0005. Values are mean±SEM.