Published in final edited form as:

Curr Opin Cell Biol. 2015 August ; 35: 117–122. doi:10.1016/j.ceb.2015.04.018.

Recycling endosomes

James R Goldenring^{1,2,3,4}

¹ Department of Surgery, Vanderbilt University School of Medicine, Nashville, TN, USA

² Department of Cell & Developmental Biology, Vanderbilt University School of Medicine, Nashville, TN, USA

³ The Epithelial Biology Center, Vanderbilt University School of Medicine, Nashville, TN, USA

⁴ The Nashville VA Medical Center, Nashville, TN, USA

Abstract

The endosomal membrane recycling system represents a dynamic conduit for sorting and re-exporting internalized membrane constituents. The recycling system is composed of multiple tubulovesicular recycling pathways that likely confer distinct trafficking pathways for individual cargoes. In addition, elements of the recycling system are responsible for assembly and maintenance of apical membrane specializations including primary cilia and apical microvilli. The existence of multiple intersecting and diverging recycling tracks likely accounts for specificity in plasma membrane recycling trafficking.

Plasma membrane recycling: a general requirement for surface homeostasis

The presentation of proteins, including receptors, ion channels, ion pumps and adhesion molecules, at the membrane surface determines many of the aspects of differentiated cell physiology. It is increasingly clear that many of these proteins are internalized using a number of different endocytic pathways. Once internalized, membrane proteins must undergo a number of sorting steps that determine their fate including targeting trafficking to degradation pathways or processing for return to the membrane surface through membrane recycling (Figure 1). The list of proteins that are recognized to recycle to the membrane surface has expanded markedly over the past decade. It now appears that membrane recycling can regulate not only surface receptors and ion channels, but also mediates aspects of junctional protein maintenance as well as the maintenance of apical and basolateral specializations in polarized cells. Together these processes place the endosomal recycling system in the center of coordinating decisions in membrane trafficking that can radically affect the cell surface.

Corresponding author: Goldenring, James R (jim.goldenring@vanderbilt.edu).

Conflict of interest The author has declared that no conflict of interest exists.

The recycling endosome is a dynamic structure

Maxfield's original studies on transferrin trafficking established that the membrane recycling system is a highly dynamic and morphologically diverse complex of membranous elements [1]. Recent studies have now focused on how regulators of the recycling system can alter the morphology as well as the function of the recycling system. The most prominent morphological dynamic in the recycling system involves dynamic tubulation of the recycling system membranes. This tubulation is less apparent in fixed cells where the endosomal morphologies are often fragmented by formaldehyde fixation. However, in live cells, the extensive tubulation is readily apparent and highly dynamic [2,3^{*}]. Some of the characteristics of tubulation likely reflect the regulation of the passage of membrane and cargoes through the recycling system. Overexpression of Rab11-Family Interacting Proteins (Rab11-FIPs) can result in dynamic expansion of tubules, while dual overexpression with Rab11a can shift the dynamics towards a more vesicular phenotype, indicating that multiple effectors may be competing for a limiting pool of Rab11a [3^{*}]. KIF13A can associate directly with Rab11a and promotes the formation of tubules in the recycling endosome system [4^{**}]. Recent studies have implicated the microtubule-severing protein, spastin, and ESCRT in scission of recycling system tubules [5^{**}]. Thus, regulation of the tubulation of the Rab11a-containing recycling system utilizes multiple mechanisms for the growth of tubules and their subsequent vesiculation.

Rab8a distributes into tubular and vesicular endosomes that also recruit EHD proteins [6]. Tubulation of the Rab8a-dependent recycling system, which is responsible for recycling of non-clathrin-dependent endocytosed cargoes, such as MHC Class I, is regulated by MICAL-L1 [7,8]. MICAL-L1 recruits both Arf6 and EHD1 to recycling membranes and promotes tubulation [7,9,10^{*},11]. The association of MICAL-L1 with tubular endosomes is regulated by Rab35 [8,12]. These studies suggest that regulation of the morphology of the endosomal recycling membranes may reflect alterations in trafficking patterns. It appears that recycling endosomes demarcated by Rab11a or Rab8a utilize distinguishable mechanisms for their dynamic regulation. It remains to be determined if the morphological alterations in the recycling system are reflected in either the kinetics or the specificity of recycling trafficking. Nevertheless, it seems likely that different sets of proteins associated with Rab11a or Rab8a-containing membranes will govern the formation tubular elements radiating from more central recycling system cisternae.

How many recycling systems are there?

When addressing the question of membrane recycling from endosomes, it is now clear that a number of pathways can lead to recycling of proteins following internalization. It is therefore of importance to ask how many pathways exist for recycling and are they acting separately or in concert? A number of regulatory proteins have been associated with recycling pathways. The greatest amount of literature addresses the role of Rab11a as a regulator of 'slow' recycling pathways that are present in all cells. These Rab11a-dependent slow recycling pathways are involved in a vast array of cell surface proteins from transferrin receptor to ion channels to junctional proteins and integrins. While there appears to be considerable overlap in the Rab11a-mediated recycling membranes, it is increasingly clear

that there may be multiple pathways for both entry and exit from these recycling membranes.

Recent studies in live cells have noted a considerable amount of subcompartmentalization of regions within the Rab11a-containing recycling system [2,3*]. What is even less clear is whether other Rab protein-dependent recycling pathways are truly distinct or are integrated into general recycling endosome systems. Previous investigations had suggested that Rab8a and Rab11a-containing recycling pathways may be distinct, with distinct Rab11a-containing vesicles separate from Rab8a-containing membranes [6]. However, Rab8a and Rab11a both interact with the actin motors Myosin VA and Myosin Vb [6,13*] as well as with the Rab8 GEF, Rabin8 [14]. A coordination of Rab8a and Rab11a with Rabin8 is required for initial lumen formation [15], but lumen formation is also regulated by Rab8a and Rab11a interacting with Myosin Vb [16]. Although their binding sites in Myosin Vb are distinct, there is no definitive evidence that Rab11a and Rab8a can bind simultaneously to Myosin Vb. It therefore remains unclear whether these Rab proteins may occupy parts of the same membrane cisternae or are present on separate membranes that either juxtapose each other during transport along microtubules or gather in 'crowded' regions in the pericentriolar space or in cell extensions. Rab10 appears to occupy domains at the ends of Rab8a-containing tubules [17]. Furthermore, the concept of recycling pathways may be more diverse. Thus, trafficking from the recycling endosome to the plasma membrane may also occur through shunting into autophagy pathways that lead to degradation or recycling through lysosomal exocytosis [18,19]. Finally, it should be noted that roles of Rab8a and Rab11a may likely extend past the recycling system. Thus, lumen formation and primary cilium formation, as will be discussed below, may involve specialized functions of Rab11a and Rab8a that do not necessarily have to be mediated by recycling membranes *per se*.

Defects in apical recycling system function influence polarized apical trafficking

Polarized cells require distinct regulation of trafficking to segregated apical and basolateral domains. Previous studies have established that polarized cells have a distinct Rab11a-containing apical recycling system that is separate from the basolateral recycling system which is not dependent on Rab11a [20,21]. Over the years, there has been much discussion of whether the Rab11a-containing recycling system functions as an intermediate for trafficking of *de novo* synthesized proteins to the apical membrane. Recently, substantial evidence has been reported analyzing the passage of newly synthesized Rhodopsin through the recycling endosome on its way to the plasma membrane [22*]. These studies indicate that the recycling endosome does function as an intermediate trafficking compartment, at least for proteins that normally undergo recycling. Other recent studies have now focused interest in the recycling system as a distinct coordinator of the integrity of apical specializations and polarity. Bryant, *et al.* have established in MDCK cells that Rab11a and Rab8a regulate the development of apical lumens early in the process of polarization [15]. Inhibition of Rab11a or Rab8a interaction with MYO5B leads to aberrant apical lumen formation [16]. Manipulation of the expression or phosphorylation of a number of Rab11 interacting proteins leads to alterations in lumen formation or polarity [23,24].

A critical point of coordinated regulation between Rab11a and Rab8a-dependent trafficking systems exists in Rab11a targeting of Rabin8 to allow activation of Rab8a (Figure 2).

This pathway plays a central role in the formation of primary cilia, where Rab11a-dependent Rabin8 recruitment promotes activation of Rab8a, which is necessary for formation of apical primary cilia [25]. These studies have pointed to a level of coordination of trafficking in the region of the centrioles. The recycling system has often been described as ‘pericentriolar’ and a number of studies have demonstrated the microtubule dependence of recycling system trafficking, but recent investigations indicate that coordination of signaling at the centrosome and trafficking into and out of the recycling system in the pericentriolar region is likely critical for assembly and maintenance of the primary cilium (Figure 2). Loss of casein kinase 1-delta, causes dissociation of Rab11a and Rab8a from the pericentriolar region [26*]. Recent studies have also demonstrated that Rabin8 activity is regulated by ERK1/2 phosphorylation [27**]. While this study did not address an impact of Rabin8 phosphorylation on primary cilia, these results do suggest a possible mechanism for regulation of ciliary growth. Two mother centriole appendage proteins, centriolin and cenexin/ODF2, associate with a Rab11a-GAP protein, Evi5, and influence primary cilium formation and centriolin also impacts the localization of the Rab11a-containing recycling system [28,29**]. While all of these studies indicate that the mother centriole is a central player in coordinating the localization of recycling system machinery involved in primary cilium assembly, the exact ultra-structure of how these processes are coordinated is less clear. The mother centriole may coordinate multiple vesicle trafficking pathways converging along microtubule tracks and/or it may act as the site of vesicle trafficking hand-offs necessary for targeting membrane components into the primary cilium. Still, the mother centriole appears to coordinate the recruitment of SNAP29, EHD1 and EHD3 to the preciliary vesicle early in the process of cilium formation and EHD1 and EHD3 localize with the ciliary pocket [30].

Components of the apical recycling system influence the maintenance of the intestinal enterocyte brush border

Recent investigations have also focused on the role of recycling system components on the process of maintaining apical microvilli in intestinal epithelial cells. Studies on patients with neonatal diarrhea from Microvillus Inclusion Disease (MVID) have led to important insights into how defects in apical trafficking can alter epithelial cell function. MVID patients have secretory diarrhea beginning in the first weeks of life [31] and display significant small intestine enterocyte villus blunting, absence of microvilli, malabsorption of glucose, and defective sodium transport. In 2008 reports from Austria and the United States demonstrated that mutations in the *MYO5B* gene caused MVID [32,33]. The vast majority of MVID patients have disabling mutations in *MYO5B* that disrupt the function of this motor or induce premature termination before the tail domain [32,34]. Loss of functional *MYO5B* leads to delocalization of Rab8a and Rab11a from the apical domain of enterocytes in association with loss of ezrin and deficits in microvillar formation and microvillar trafficking [16,35*]. Findings in cell culture models demonstrated that Rab8a association with *MYO5B* is required for maintenance of apical microvilli [35*,36,37], while the

observation of microvillus inclusions was dependent on disruption of MYO5B association with Rab11a [35*]. Interestingly, Rab8a KO mice also show deficits in microvillus formations in the intestines and these defects are augmented in Rab8a/Rab8b double KO mice [38,39*]. While Rab11a germline KO mice are embryonic lethal [40], targeted KO of Rab11a in intestinal enterocytes leads to short apical microvilli and deficits in TLR9 recycling [41–43].

Interestingly, the formation of microvillus inclusions appears to be due to activation of macropinocytosis [35*], so it appears that Rab11a/MYO5B must be involved in recycling of membranes from macropinosomes. The process of large-scale membrane internalization and recycling may be a major concept in specialized membrane dynamics. A recent study has found that bulk clathrin and dynamin-independent membrane internalization is involved in recycling membrane into synaptic vesicle pools [44**]. Large-scale membrane internalization and processing into regulated recycling membranes has previously been identified for the recycling of H/K-ATPase and Rab11a-containing tubulovesicular membranes in gastric parietal cells [45–47]. Thus, elements of apical recycling from macropinocytosed membranes may be a major pathway for maintenance of the apical surface in polarized epithelial cells.

Recently, mutations in *STX3* (Syntaxin 3) and *STXBP2* (*Munc18-2*) have been implicated as the causative agents for an atypical form of neonatal diarrhea [32,48*,49], which is characterized by loss of apical microvilli along with lateral microvilli formation. The phenotype is notably similar to that observed in the mice with intestinal targeted KO of Rab11a, where lateral microvilli are observed without microvillus inclusions [43]. Interestingly, previous studies have identified STX3 as a critical regulator of Rab11a-dependent apical recycling in gastric parietal cells [50–52]. These rare but devastating neonatal diarrhea syndromes demonstrate that aberrations in apical recycling trafficking can have functional implications for epithelial cell physiology. It is important to note, however, that while MYO5B and STX3 are ubiquitously expressed in multiple epithelial cells, these children manifest their major symptoms focally in the intestine. Thus, some epithelial cells must be able to compensate for loss of function mutations in apical trafficking proteins better than others. It seems likely that unrecognized mutations in other proteins regulating apical membrane trafficking and recycling may be responsible for undiagnosed diarrhea syndromes in neonates.

Conclusions

The foregoing discussion demonstrates that the recycling endosome can no longer be treated as a single organelle, but is rather more appropriately represented as a system of recycling membrane pathways that may be semi-autonomous or at least interconnected. Multiple autonomous trafficking pathways may be operating independently to move specific cargoes along common established micro-tubule-based pathways. Effectors of Rab small GTPases can modulate the dynamics of trafficking and it seems likely that more examples will be identified reflecting the regulation of critical trafficking steps and decision points by second messenger and protein kinase-mediated signaling. In the future, it will be critical to dissect the complexity of trafficking decisions made within the tubular recycling system membranes

and determine how specializations can account for regulation of multiple cargo recycling pathways within specific cell systems.

Acknowledgement

Dr. Goldenring's research is supported by NIH RO1 DK48370 and RO1 DK070856.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

•of special interest

••of outstanding interest

1. Marsh EW, Leopold PL, Jones NL, Maxfield FR. Oligomerized transferrin receptors are selectively retained by a luminal sorting signal in a long-lived endocytic recycling compartment. *J Cell Biol.* 1995; 129:1509–1522. [PubMed: 7790351]
2. Baetz NW, Goldenring JR. Distinct patterns of phosphatidylserine localization within the Rab11a-containing recycling system. *Cell Logist.* 2014; 4:e28680. [PubMed: 25210648]
3. Baetz NW, Goldenring JR. Rab11-Family Interacting Proteins define spatially and temporally distinct regions within the dynamic Rab11a-dependent recycling system. *Mol Biol Cell.* 2013; 24:643–658. [PubMed: 23283983] [This report details live cell imaging studies of the tubular subdomains within the recycling endosome that are defined by individual Rab11-FIP proteins]
4. Delevoe C, Miserey-Lenkei S, Montagnac G, Gilles-Marsens F, Paul-Gilloteaux P, Giordano F, Waharte F, Marks MS, Goud B, Raposo G. Recycling endosome tubule morphogenesis from sorting endosomes requires the kinesin motor KIF13A. *Cell Rep.* 2014; 6:445–454. [PubMed: 24462287] [This report details the direct interaction of Rab11a with a microtubule motor that is involved in the process of tubulation in the recycling endosome]
5. Allison R, Lumb JH, Fassier C, Connell JW, Ten Martin D, Seaman MN, Hazan J, Reid E. An ESCRT-spastin interaction promotes fission of recycling tubules from the endosome. *J Cell Biol.* 2013; 202:527–543. [PubMed: 23897888] [This study describes a critical mechanism for accomplishing fission of tubules emanating from the recycling system]
6. Roland JT, Kenworthy AK, Peranen J, Caplan S, Goldenring JR. Myosin Vb Interacts with Rab8a on a Tubular Network Containing EHD1 and EHD3. *Mol Biol Cell.* 2007; 18:2828–2837. [PubMed: 17507647]
7. Xie S, Naslavsky N, Caplan S. Diacylglycerol kinase α regulates tubular recycling endosome biogenesis and major histocompatibility complex class I recycling. *J Biol Chem.* 2014; 289:31914–31926. [PubMed: 25248744]
8. Rahajeng J, Giridharan SS, Cai B, Naslavsky N, Caplan S. MICALL1 is a tubular endosomal membrane hub that connects Rab35 and Arf6 with Rab8a. *Traffic.* 2012; 13:82–93. [PubMed: 21951725]
9. Reinecke JB, Katafiasz D, Naslavsky N, Caplan S. Regulation of Src trafficking and activation by the endocytic regulatory proteins MICAL-L1 and EHD1. *J Cell Sci.* 2014; 127:1684–1698. [PubMed: 24481818]
10. Cai B, Giridharan SS, Zhang J, Saxena S, Bahl K, Schmidt JA, Sorgen PL, Guo W, Naslavsky N, Caplan S. Differential roles of C-terminal Eps15 homology domain proteins as vesiculators and tubulators of recycling endosomes. *J Biol Chem.* 2013; 288:30172–30180. [PubMed: 24019528] [The study demonstrates the ability of EHD1 to tubulate recycling endosomes containing MICAL-L1]
11. Giridharan SS, Cai B, Vitale N, Naslavsky N, Caplan S. Cooperation of MICAL-L1, syndapin2, and phosphatidic acid in tubular recycling endosome biogenesis. *Mol Biol Cell.* 2013; 24:1776–1790 S1771-1715. [PubMed: 23596323]

12. Kobayashi H, Etoh K, Ohbayashi N, Fukuda M. Rab35 promotes the recruitment of Rab8 Rab13 and Rab36 to recycling endosomes through MICAL-L1 during neurite outgrowth. *Biol Open*. 2014; 3:803–814. [PubMed: 25086062]
- 13•. Lindsay AJ, Jollivet F, Horgan CP, Khan AR, Raposo G, McCaffrey MW, Goud B. Identification and characterization of multiple novel Rab-myosin Va interactions. *Mol Biol Cell*. 2013; 24:3420–3434. [PubMed: 24006491] [These investigations detail interactions of multiple Rab proteins with MYO5A and suggest that MYO5 motors may regulate multiple trafficking steps in cells]
14. Knodler A, Feng S, Zhang J, Zhang X, Das A, Peranen J, Guo W. Coordination of Rab8 and Rab11 in primary ciliogenesis. *Proc Natl Acad Sci U S A*. 2010; 107:6346–6351. [PubMed: 20308558]
15. Bryant DM, Datta A, Rodriguez-Fraticelli AE, Peranen J, Martin-Belmonte F, Mostov KE. A molecular network for de novo generation of the apical surface and lumen. *Nat Cell Biol*. 2010; 12:1035–1045. [PubMed: 20890297]
16. Roland JTE, Bryant DM, Datta A, Itzen A, Mostov KE, Goldenring JR. Rab GTPase-Myo5B complexes control membrane recycling and epithelial polarization. *Proc Natl Acad Sci U S A*. 2011; 108:2789–2794. [PubMed: 21282656]
17. Roland JT, Lapierre LA, Goldenring JR. Alternative splicing in class V myosins determines association with Rab10. *J Biol Chem*. 2009; 284:1213–1223. [PubMed: 19008234]
18. Szatmari Z, Kis V, Lippai M, Hegedus K, Farago T, Lorincz P, Tanaka T, Juhasz G, Sass M. Rab11 facilitates cross-talk between autophagy and endosomal pathway through regulation of Hook localization. *Mol Biol Cell*. 2014; 25:522–531. [PubMed: 24356450]
19. Puri C, Renna M, Bento CF, Moreau K, Rubinsztein DC. ATG16L1 meets ATG9 in recycling endosomes: additional roles for the plasma membrane and endocytosis in autophagosome biogenesis. *Autophagy*. 2014; 10:182–184. [PubMed: 24257061]
20. Wang E, Brown PS, Aroeti B, Chapin SJ, Mostov KE, Dunn KW. Apical and basolateral endocytic pathways of MDCK cells meet in acidic common endosomes distinct from a nearly neutral apical recycling endosome. *Traffic*. 2000; 1:480–493. [PubMed: 11208134]
21. Lapierre LA, Kumar R, Hales CM, Navarre J, Bhartur SG, Burnette JO, Mercer JA, Bahler M, Goldenring JR. Myosin Vb is associated with and regulates plasma membrane recycling systems. *Mol Biol Cell*. 2001; 12:1843–1857. [PubMed: 11408590]
- 22•. Thuenauer R, Hsu YC, Carvajal-Gonzalez JM, Deborde S, Chuang JZ, Romer W, Sonnleitner A, Rodriguez-Boulan E, Sung CH. Four-dimensional live imaging of apical biosynthetic trafficking reveals a post-Golgi sorting role of apical endosomal intermediates. *Proc Natl Acad Sci U S A*. 2014; 111:4127–4132. [PubMed: 24591614] [This study provides convincing evidence for the function of the apical recycling system as an intermediate trafficking point for newly synthesized proteins destined for delivery to the apical membrane]
23. Li D, Mangan A, Cicchini L, Margolis B, Prekeris R. FIP5 phosphorylation during mitosis regulates apical trafficking and lumenogenesis. *EMBO Rep*. 2014; 15:428–437. [PubMed: 24591568]
24. Lapierre LA, Avant KM, Caldwell CM, Oztan A, Apodaca G, Knowles BC, Roland JT, Ducharme NA, Goldenring JR. Phosphorylation of Rab11-FIP2 regulates polarity in MDCK cells. *Mol Biol Cell*. 2012; 23:2302–2318. [PubMed: 22553350]
25. Westlake CJ, Baye LM, Nachury MV, Wright KJ, Ervin KE, Phu L, Chalouni C, Beck JS, Kirkpatrick DS, Slusarski DC, et al. Primary cilia membrane assembly is initiated by Rab11 and transport protein particle II (TRAPP2) complex-dependent trafficking of Rabin8 to the centrosome. *Proc Natl Acad Sci U S A*. 2011; 108:2759–2764. [PubMed: 21273506]
- 26•. Greer YE, Westlake CJ, Gao B, Bharti K, Shiba Y, Xavier CP, Pazour GJ, Yang Y, Rubin JS. Casein kinase 1delta functions at the centrosome and Golgi to promote ciliogenesis. *Mol Biol Cell*. 2014; 25:1629–1640. [PubMed: 24648492] [The report elucidates a novel mechanism for localized kinase regulation of trafficking in the pericentriolar region]
- 27••. Wang J, Ren J, Wu B, Feng S, Cai G, Tuluc F, Peranen J, Guo W. Activation of Rab8 guanine nucleotide exchange factor Rabin8 by ERK1/2 in response to EGF signaling. *Proc Natl Acad Sci U S A*. 2015; 112:148–153. [PubMed: 25535387] [This study reports the phosphorylation and

activation of Rabin8 by ERK1/2 and provides a possible regulatory mechanism for pericentriolar trafficking]

28. Hehnly H, Hung HF, Doxsey S. One among many: ODF2 isoform 9, a.k.a Cenexin-1, is required for ciliogenesis. *Cell Cycle*. 2013; 12:1021. [PubMed: 23511168]
- 29•. Hehnly H, Chen CT, Powers CM, Liu HL, Doxsey S. The centrosome regulates the Rab11-dependent recycling endosome pathway at appendages of the mother centriole. *Curr Biol*. 2012; 22:1944–1950. [PubMed: 22981775] [This investigation details evidence for an association of Rab11a-containing recycling endosomes with the mother centriole]
30. Lu Q, Insinna C, Ott C, Stauffer J, Pintado PA, Rahajeng J, Baxa U, Walia V, Cuenca A, Hwang YS, et al. Early steps in primary cilium assembly require EHD1/EHD3-dependent ciliary vesicle formation. *Nat Cell Biol*. 2015; 17:228–240. [PubMed: 25686250]
31. Davidson GP, Cutz E, Hamilton JR, Gall DG. Familial enteropathy: a syndrome of protracted diarrhea from birth, failure to thrive, and hypoplastic villus atrophy. *Gastroenterology*. 1978; 75:783–790. [PubMed: 100367]
32. Muller T, Hess MW, Schiefermeier N, Pfaller K, Ebner HL, Heinz-Erian P, Ponstingl H, Partsch J, Rollinghoff B, Kohler H, et al. MYO5B mutations cause microvillus inclusion disease and disrupt epithelial cell polarity. *Nat Genet*. 2008; 40:1163–1165. [PubMed: 18724368]
33. Erickson RP, Larson-Thorne K, Valenzuela RK, Whitaker SE, Shub MD. Navajo microvillous inclusion disease is due to a mutation in MYO5B. *Am J Med Genet A*. 2008; 146A:3117–3119. [PubMed: 19006234]
34. Ruemmele FM, Muller T, Schiefermeier N, Ebner HL, Lechner S, Pfaller K, Thoni CE, Goulet O, Lacaille F, Schmitz J, et al. Loss-of-function of MYO5B is the main cause of microvillus inclusion disease: 15 novel mutations and a CaCo-2 RNAi cell model. *Hum Mutat*. 2010; 31:544–551. [PubMed: 20186687]
- 35•. Knowles BC, Roland JT, Krishnan M, Tyska MJ, Lapierre LA, Dickman PS, Goldenring JR, Shub MD. Myosin Vb uncoupling from RAB8A and RAB11A elicits microvillus inclusion disease. *J Clin Invest*. 2014; 124:2947–2962. [PubMed: 24892806] [This report demonstrates that the manifestation of microvillus inclusions in patients with Microvillus Inclusion Disease likely accrues from an inability of enterocytes to recycle apical membranes internalized by macropinocytosis]
36. Thoeni CE, Vogel GF, Tancevski I, Geley S, Lechner S, Pfaller K, Hess MW, Muller T, Janecke AR, Avitzur Y, et al. Microvillus inclusion disease: loss of myosin Vb disrupts intracellular traffic and cell polarity. *Traffic*. 2013; 15:22–42. [PubMed: 24138727]
37. Kravtsov D, Mashukova A, Forteza R, Rodriguez MM, Ameen NA, Salas PJ. Myosin 5b loss of function leads to defects in polarized signalling: implication for Microvillus Inclusion Disease pathogenesis and treatment. *Am J Physiol Gastrointest Liver Physiol*. 2014; 307:G992–G1001. [PubMed: 25258405]
38. Sato T, Mushiaki S, Kato Y, Sato K, Sato M, Takeda N, Ozono K, Miki K, Kubo Y, Tsuji A, et al. The Rab8 GTPase regulates apical protein localization in intestinal cells. *Nature*. 2007; 448:366–369. [PubMed: 17597763]
- 39•. Sato T, Iwano T, Kunii M, Matsuda S, Mizuguchi R, Jung Y, Hagiwara H, Yoshihara Y, Yuzaki M, Harada R, et al. Rab8a and Rab8b are essential for several apical transport pathways but insufficient for ciliogenesis. *J Cell Sci*. 2014; 127:422–431. [PubMed: 24213529] [This paper from the group that originally reported deficits in enterocyte apical microvilli in Rab8a KO mice, demonstrates that dual KO of both Rab8a and Rab8b causes a more severe phenotype]
40. Yu S, Yehia G, Wang J, Stypulkowski E, Sakamori R, Jiang P, Hernandez-Enriquez B, Tran TS, Bonder EM, Guo W, et al. Global ablation of the mouse Rab11a gene impairs early embryogenesis and matrix metalloproteinase secretion. *J Biol Chem*. 2014; 289:32030–32043. [PubMed: 25271168]
41. Yu S, Nie Y, Knowles B, Sakamori R, Stypulkowski E, Patel C, Das S, Douard V, Ferraris RP, Bonder EM, et al. TLR sorting by Rab11 endosomes maintains intestinal epithelial-microbial homeostasis. *EMBO J*. 2014; 33:1882–1895. [PubMed: 25063677]
42. Sobajima T, Yoshimura S, Iwano T, Kunii M, Watanabe M, Atik N, Mushiaki S, Morii E, Koyama Y, Miyoshi E, et al. Rab11a is required for apical protein localisation in the intestine. *Biol Open*. 2014; 4:86–94. [PubMed: 25527643]

43. Knowles BC, Weis VG, Yu S, Roland JT, Williams JA, Alvarado GS, Lapierre LA, Shub MD, Gao N, Goldenring JR. Rab11a regulates Syntaxin 3 localization and microvillus assembly in enterocytes. *J Cell Sci.* 2015; 128:1617–1626. [PubMed: 25673875]
44. Wu Y, O'Toole ET, Girard M, Ritter B, Messa M, Liu X, McPherson PS, Ferguson SM, De Camilli P. A dynamin 1-, dynamin 3- and clathrin-independent pathway of synaptic vesicle recycling mediated by bulk endocytosis. *Elife.* 2014; 3:e01621. [PubMed: 24963135] [This paper suggests a novel pathway for bulk internalization of membrane and reformation of synaptic vesicles. Much of these processes have similarities to recycling of membranes from macropinosomes]
45. Forte TM, Machen TE, Forte JG. Ultrastructural changes in oxyntic cells associated with secretory function: a membrane recycling hypothesis. *Gastroenterology.* 1977; 73:941–955. [PubMed: 902958]
46. Peng X- P, Yao X, Chow D- C, Forte JG, Bennett MK. Association of Syntaxin 3 and vesicle associated (VAMP) + membrane protein with H + /K -ATPase-containing tubulovesicles in gastric parietal cells. *Mol Biol Cell.* 1997; 8:399–407. [PubMed: 9188093]
47. Calhoun BC, Goldenring JR. Two Rab proteins, vesicle-associated membrane protein 2 (VAMP-2) and secretory carrier membrane proteins (SCAMPs), are present on immunoisolated parietal cell tubulovesicles. *Biochem J.* 1997; 325(Pt 2):559–564. [PubMed: 9230141]
- 48•. Wiegerinck CL, Janecke AR, Schneeberger K, Vogel GF, van Haaften-Visser DY, Escher JC, Adam R, Thoni CE, Pfaller K, Jordan AJ, et al. Loss of Syntaxin 3 causes variant Microvillus Inclusion Disease. *Gastroenterology.* 2014; 147:65–68. [PubMed: 24726755] [This report identifies children with neonatal diarrhea due to syntaxin 3 mutations who show a partial loss of polarity defect in enterocytes with the formation of microvilli along the lateral membranes. This report, in addition to findings in Microvillus Inclusion Disease, suggests that defects in apical recycling and trafficking may account for previously undiagnosed neonatal diarrhea syndromes]
49. Stepensky P, Bartram J, Barth TF, Lehmborg K, Walther P, Amann K, Philips AD, Beringer O, Zur Stadt U, Schulz A, et al. Persistent defective membrane trafficking in epithelial cells of patients with familial hemophagocytic lymphohistiocytosis type 5 due to STXB2/MUNC18-2 mutations. *Pediatr Blood Cancer.* 2013; 60:1215–1222. [PubMed: 23382066]
50. Calhoun BC, Lapierre LA, Chew CS, Goldenring JR. Rab11a redistributes to apical secretory canaliculus during stimulation of gastric parietal cells. *Am J Physiol.* 1998; 275:C163–C170. [PubMed: 9688847]
51. Karvar S, Zhu L, Crothers J Jr, Wong W, Turkoz M, Forte JG. Cellular localization and stimulation-associated distribution dynamics of syntaxin-1 and syntaxin-3 in gastric parietal cells. *Traffic.* 2005; 6:654–666. [PubMed: 15998321]
52. Lapierre LA, Avant KM, Caldwell CM, Ham AJ, Hill S, Williams JA, Smolka AJ, Goldenring JR. Characterization of immunoisolated human gastric parietal cells tubulovesicles: identification of regulators of apical recycling. *Am J Physiol Gastrointest Liver Physiol.* 2007; 292:G1249–G1262. [PubMed: 17255364]

Figure 1.

A complex scheme for endosomal pathways leading to membrane recycling. Membrane proteins are internalized by multiple mechanisms including clathrin-dependent endocytosis (CDE), non-clathrin-dependent endocytosis (NCDE) and macropinocytosis (MP). The early endosomes from these pathways are thought to traffic through a common sorting endosome (SE). Proteins can then be shunted for degradation to the lysosome, recycled to the plasma membrane using a rapid recycling pathway (thought to be mediated by Rab4) or processed into slower recycling endosome (RE) pathways (thought to be regulated by Rab11a and Rab8a). While both Rab11a and Rab8a can interact with MYO5B, specific proteins interacting with membranes containing either Rab11a (Rab11-FIPs and KIF13) or Rab8a (MICAL-L1, EHD1 and EHD3) may regulate specific trafficking pathways out of the recycling system.

Figure 2.

Pericentriolar coordination of primary cilium construction by components of recycling endosomes. The mother centriole provides a point of coordination for trafficking through the pericentriolar region. Microtubule-based tracks lead towards and away from the pericentriolar region and provide paths for anterograde and retrograde trafficking to the plasma membrane surface. These trafficking processes are necessary for the assembly and maintenance of apical specializations such as the primary cilium. The mother centriole acts as a coordinating point for the recruitment by Rab11a of the Rab8a GEF, Rabin8, which can in turn activate Rab8a. It is important to note that this coordinated activity in the pericentriolar region likely involves the juxtaposition of distinct endosomal membranes containing either Rab11a or Rab8a.