secondary care health delivery system is well developed and regular follow up is well ensured, we can design a larger study with an inbuilt system to assess the achievement of treatment goals periodically. This mechanism to audit comprehensive healthcare delivery in a condition like Type 2 DM which has such a significant impact on cardiovascular morbidity and mortality will be a huge step towards achieving improved health for our clientele.

REFERENCES

1. Menon AS, Ahluwalia AI. The ABC of diabetes. How many patients are able to achieve the goal laid down by American Diabetes Association? *Med J Armed Forces India*. 2015;71: 132–134.
2. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*. 1998;352:837–853.

3. Casagrande SS, Fradkin JE, Saydah SH, Rust KF, Cowie CC. The prevalence of meeting A1C, blood pressure, and LDL goals among people with Diabetes, 1988–2010. *Diabetes Care*. 2013;36: 2271–2279.

Col J. Muthukrishnan, SM*

Senior Adviser (Med & Endocrinologist), Command Hospital
(Southern Command), Pune 40, India

*Tel.: +91 8698951237.

E-mail address: jmuthukrishnan@hotmail.com

Available online 2 July 2015

DOI of original article: <http://dx.doi.org/10.1016/j.mjafi.2015.06.002>

0377-1237/\$ – see front matter

© 2015, Armed Forces Medical Services (AFMS). All rights reserved.
<http://dx.doi.org/10.1016/j.mjafi.2015.06.003>

Reply

Dear Editor,

We thank the reader for the keen interest he has shown in reading our article. There is lack of data on the effectiveness of therapy for Diabetes in our country. We agree with the reader's contention that larger studies should be planned to get a comprehensive picture of standard of care of Diabetes in our clientele. We feel Armed Forces Medical Services with its stratified structure of providing medical care is ideally suited for conducting the study and could take a lead in this respect.

The study design was formulated before the concept of individualisation of glycemic target was brought in diabetes care. We are not certain whether the outcome would have been any different even if we had used individualised targets.

The necessity for auditing performance is the first step in improving quality of care in chronic non communicable disease like Diabetes and we hope more attention is paid to this aspect of diabetes care.¹

REFERENCE

1. Kapur A, Harries AD. Cohort monitoring: as a tool to improve diabetes care services. *Diab Res Clin Prac*. 2013;102:260–264.

Col A.S. Menon*

Senior Adviser (Medicine & Endocrinology), Command Hospital
(Central Command), Lucknow 226002, India

Surg Capt A.I. Ahluwalia

Senior Adviser (Medicine & Endocrinology), Base Hospital,
Delhi Cantt, India

*Corresponding author.

E-mail address: menonara@yahoo.com

Available online 2 July 2015

0377-1237/\$ – see front matter

© 2015, Armed Forces Medical Services (AFMS). All rights reserved.
<http://dx.doi.org/10.1016/j.mjafi.2015.06.004>

Letter to the Editor

Dear Editor,

I read with extreme interest the original article titled “Major laparoscopic surgery under regional anesthesia: A prospective

feasibility study” by Singh et al. published in *Med J Armed Forces India* 2015; 71:126–131.¹

Laparoscopic surgeries have revolutionized the very outlook of surgeons. Gone are the days of large incisions and

“good exposure” for elective abdominal surgeries. Apart from being more acceptable to the patient from the cosmetic point of view, it also reduces the burden on the health care system by decreasing hospital stay, decreasing the overall cost and time of surgery in addition to abbreviating recovery time to such an extent that Laparoscopic surgeries are now in the domain of “Day Care Surgeries”. Along with the advent and increasing popularity of Laparoscopic surgeries, anaesthesia for the same has also evolved keeping the same objectives of improving patient outcomes in perspective. However, this evolution has given rise to new challenges for the anaesthesiologist. Thus though the use of Propofol for General Anaesthesia was thought to be an improvement for Day Care Surgeries, the quest for better techniques has always been on. The article titled “Major laparoscopic surgery under regional anesthesia: A prospective feasibility study” takes us further on that road of evolution. The earlier studies with regional anaesthesia for laparoscopic surgeries have been done with patients presenting with respiratory and (or) cardiac compromise and a poor risk situation for general anaesthesia.^{2,3} I must congratulate the authors for successful outcomes where many a study including one by myself in 2008 had to be abandoned or had poorer outcomes. A pilot study was done with 15 cases and reported good results with minimal post operative pain.⁴ One of the similar studies which reported good results, however used segmental thoracic spinal anaesthesia, hypothesizing that the “tenth thoracic interspace lies in the centre of the surgical field”.⁵ Though this particular study did not report any adverse event most anaesthesiologists would be wary of attempting a thoracic sub arachnoid block and would prefer the conventional lumbar sub arachnoid block with a Trendelenberg tilt till the desired dermatome block was achieved.^{6,7} A recent study has even compared thoracic vs. lumbar sub arachnoid block and reported better results with the thoracic sub arachnoid block, however with low pressure pneumoperitoneum.⁸

Despite the growing trend towards regional anaesthesia certain issues need consideration. The first being the preferred position of the patient by the surgeon, which is usually Anti-Trendelenberg. This position promotes further venous pooling of blood in a patient where sympatholysis by virtue of the neuraxial block already exists and results in profound hypotension. Also to be considered is the fact that the pneumoperitoneum itself would further decrease the venous return thereby adding to the hypotension. The second issue is that of rising blood CO₂ levels. Though it has been addressed in studies of shorter duration an arterial blood gas analysis is essential in surgeries of longer duration (upto 170 min in the present article).^{9,10} These studies have also reported that the rising CO₂ level was offset by an increased respiratory rate as well as minute ventilation which should definitely add to the discomfort level of an awake patient. Lastly Laparoscopic Surgeries do require muscle relaxation for pneumoperitoneum hence extreme care and caution must be exercised at achieving the right

dermatomal level of block, which has been mentioned as T₄ in some studies.¹⁰

The entire concept of Day Care Laparoscopic Surgery is faster recovery time concomitant with early discharge and independence from the hospital, which is well achieved by the use of regional anaesthesia.

REFERENCES

1. Singh RK, Saini AM, Goel N, Bisht D, Seth A. Major laparoscopic surgery under regional anesthesia: A prospective feasibility study. *Med J Armed Forces India*. 2015;71:126–131.
2. Pursnani KG, Bazza Y, Calleja M, Mughal MM. Laparoscopic cholecystectomy under epidural anesthesia in patients with chronic respiratory disease. *Surg Endosc*. 1998;12:1082–1084.
3. Gramatica L, Brasesco OE, Luna AM, et al. Laparoscopic cholecystectomy performed under regional anesthesia in patients with chronic obstructive pulmonary disease. *Surg Endosc*. 2002;16:472–475.
4. Tzoravaras G, Fafoulakis F, Pratsas K, Georgopoulou S, Stamatou G, Hatzitheofilou C. Laparoscopic cholecystectomy under spinal anesthesia. A pilot study. *Surg Endosc*. 2006;20:580–582.
5. van Zundert AA, Stultiens G, Jakimowicz JJ, et al. Laparoscopic cholecystectomy under segmental thoracic spinal anaesthesia: a feasibility study. *Br Journal of Anaesthesia*. 2007;98:682–686.
6. Tiwari S, Chauhan A, Chatterjee P, Alam MT. Laparoscopic cholecystectomy under spinal anaesthesia: A prospective, randomised study. *J Minim Access Surg*. 2013;9:65–71.
7. Sinha R, Gurwara AK, Gupta SC. Laparoscopic cholecystectomy under spinal anesthesia: a study of 3492 patients. *J Laparoendosc Adv Surg Tech A*. 2009;19:323–327.
8. Imbelloni LE. Spinal anesthesia for laparoscopic cholecystectomy: thoracic vs. Lumbar Technique. *Saudi J Anaesth*. 2014;8:477–483.
9. Subba B, Gupta I, Singh H. Studies of cardiovascular and arterial blood gas changes during carbon dioxide pneumoperitoneum for laparoscopic sterilization under general anaesthesia versus local anaesthesia. *Asia-Oceania J of Obstetrics and Gynaecology*. 1991;17:31–35.
10. Diamant M, Benumof JL, Saidman LJ, Kennedy J, Young P. Laparoscopic sterilization with local anesthesia: complications and blood gas changes. *Anesthesia and Analgesia*. 1977;56:335–337.

Col R.N. Verma

Associate Professor, Dept of Anaesthesia, Critical Care & Pain Management, Armed Forces Medical College, Pune 40, India

E-mail address: docrishiraj68@gmail.com

Available online 2 July 2015

DOI of original article: <http://dx.doi.org/10.1016/j.mjafi.2015.06.006>

0377-1237/\$ – see front matter

© 2015, Armed Forces Medical Services (AFMS). All rights reserved.
<http://dx.doi.org/10.1016/j.mjafi.2015.06.006>