Published in final edited form as:

Neuropharmacology. 2015 October ; 97: 270–274. doi:10.1016/j.neuropharm.2015.05.023.

Effects of orally-bioavailable short-acting kappa opioid receptor-selective antagonist LY2456302 on nicotine withdrawal in mice

K. J. Jackson^a, A. Jackson^b, F. Ivy Carroll^c, and M. I. Damaj^b

^aDepartment of Psychiatry, Virginia Commonwealth University, 800 E. Leigh St., Suite 390A, Richmond, VA 23219, U.S.A.

^bDepartment of Pharmacology/Toxicology, Virginia Commonwealth University, 1220 E. Broad St., MMRB, Box 980613, Richmond, VA 23219, U.S.A

^cResearch Triangle Institute, Research Triangle Park, NC, United States

Abstract

Kappa opioid receptor (KOR) signaling has been implicated in mediating behavioral and biochemical effects associated with drug dependence. The most commonly used KOR antagonists, norbinaltorphimine (norBNI) and (3R)-7-Hydroxy-N{(1S)-1-[(3R,4R)-4-(3-hydroxyphenyl)-3,4-dimethyl-1-piperidiny]methyl}-2-methylpropyl}-1,2,3,4-tetrahydro-3-isoquinoline-carboxamide (JDTic), have provided a wealth of information in this area; however, the delayed onset and long-lasting effects of these antagonists complicate experimental design and interpretation of results, and make them less than ideal for clinical studies. Initial studies with the recently developed KOR antagonist, LY2456302, show that the compound is a short acting, high-affinity, selective KOR antagonist with therapeutic potential for mood disorders and ethanol use in animal models, and is well tolerated in humans. The goal of the current study was to evaluate the effectiveness of LY2456302 in alleviating the nicotine withdrawal syndrome in mice. Mice were chronically treated with nicotine for 14 days and physical and affective nicotine withdrawal signs were measured using a spontaneous nicotine withdrawal model and conditioned place aversion (CPA) following pre-treatment with LY2456302, administered orally. Vehicle treated nicotine withdrawn mice displayed significant anxiety-related behavior, somatic signs, hyperalgesia, and CPA. Similar to previous studies with norBNI and JDTic, LY2456302 alleviated the nicotine withdrawal syndrome, as evidenced by decreased expression of nicotine withdrawal induced anxiety-related behavior, somatic signs, and CPA, and increased hotplate latency in nicotine withdrawn mice following pre-treatment. Given the current results, and with its favorable pharmacokinetic and pharmacodynamic profile, LY2456302 may be a useful therapeutic agent for treatment of multiple aspects of the nicotine withdrawal syndrome.

Corresponding Author: M. Imad Damaj, Ph.D., 1220 East Broad Street, Molecular Medicine Research Building Room 3044, P.O. Box 980613, Richmond, Virginia 23298-0613, Phone: (804) 828-1676, mdamaj@vcu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

There are no conflicts of interest to disclose.

Keywords

LY2456302; kappa opioid receptor; nicotine withdrawal; kappa opioid receptor antagonist; nicotine dependence

1. Introduction

Scientific evidence increasingly supports a role for kappa opioid receptor (KOR) signaling in mediating the behavioral and biochemical effects associated with aversive and depressive-like states, and substance use dependence. The mechanism of KOR involvement in regulating motivational and emotional states has been suggested to involve dynorphin, the endogenous KOR ligand. Rewarding and stressful stimuli increase cyclic adenosine monophosphate response element binding protein (CREB), resulting in increased levels of dynorphin, which have been observed after stress or drug exposure (Nestler, Barrot, DiLeone, Eisch, Gold, & Monteggia, 2002). The increased levels of dynorphin bind to the KOR, resulting in reduced levels of dopamine and a state of anhedonia. Blockade of KOR by antagonists alleviates negative motivational and emotional states through blocking dynorphin interactions with the receptor (Carroll & Carlezon, Jr., 2013). In particular, disruption of KOR function attenuates stress responses, which can serve as an environmental trigger for neuropsychiatric conditions, such as depressive disorders and addiction (Van't Veer & Carlezon, Jr., 2013). The prototypical KOR antagonists, norbinaltorphimine (norBNI) and (3R)-7-Hydroxy-N{(1S)-1-[(3R,4R)-4-(3-hydroxyphenyl)-3,4-dimethyl-1-piperidinyl]methyl}-2-methylpropyl}-1,2,3,4-tetrahydro-3-isoquinoline-carboxamide (JDTic), and the selective KOR agonist, U50,488, have been used to assess KOR involvement in these effects, and have provided much of the current knowledge in this area.

In rodent models of drug withdrawal, norBNI and/or JDTic attenuated nicotine and morphine somatic signs of withdrawal (Tejeda, Natividad, Orfila, Torres, & O'Dell, 2012; Jackson, Carroll, Negus, & Damaj, 2010; Kelsey, Verhaak, & Schierberl, 2015), ethanol and nicotine withdrawal-induced anxiety-related behavior as measured by the elevated plus maze (Schank, Goldstein, Rowe, King, Marusich, Wiley et al., 2012; Valdez & Harshberger, 2012; Gillett, Harshberger, & Valdez, 2013; Jackson et al., 2010), nicotine and morphine withdrawal conditioned place aversion (CPA) (Jackson et al., 2010; Kelsey et al., 2015), and ultrasonic vocalizations associated with ethanol withdrawal (Berger, Williams, McGinnis, & Walker, 2013). JDTic and norBNI also decreased ethanol self-administration in rats (Schank et al., 2012; Walker & Koob, 2008; Walker, Zorrilla, & Koob, 2011). Many of these behaviors were exacerbated by treatment with U50,488 (Valdez & Harshberger, 2012; Berger et al., 2013; Schank et al., 2012; Gillett et al., 2013; Tejeda et al., 2012). Alternatively, administration of the peripheral KOR agonist, ICI 204,448, inhibited nicotine withdrawal induced increases in feeding, metabolism, and locomotor activity in rats (Sudakov, Nazarova, Alekseeva, & Kolpakov, 2014), suggesting that peripheral and central KOR signaling may differentially mediate nicotine withdrawal.

Despite their utility in understanding KOR involvement in addictive behaviors, various confounds affecting experimental design and interpretation of results are evident with these

antagonists. First, the onset of KOR antagonism by norBNI and JDTic can be delayed for hours (Munro, Berry, Van't Veer, Beguin, Carroll, Zhao et al., 2012; Carroll, Thomas, Dykstra, Granger, Allen, Howard et al., 2004). Both compounds also have very long durations of action, which can result in an extended pharmacodynamic effects or increase the potential for undesirable drug-drug interactions. NorBNI and JDTic have been shown to block antinociceptive activity in mice for up to 2 weeks (Carroll et al., 2004; Patkar, Wu, Ganno, Singh, Ross, Rasakham et al., 2013), and significant KOR antagonist activity is detectable in rodents for up to 28 days (Munro et al., 2012; Carroll et al., 2004; Patkar et al., 2013). Such drug properties would also be unfavorable for clinical use. LY2456302, a recently developed, potent, high-affinity selective KOR antagonist, which has a plasma half-life of 2–4 h in rats and mice and receptor occupancy <50% within 48 h after oral administration (Rorick-Kehn, Witkin, Statnick, Eberle, McKinzie, Kahl et al., 2014), may be a more useful tool for assessing the therapeutic potential of KOR modulation as treatment for addictive disorders in preclinical and clinical models.

Although literature is limited, LY2456302 has shown promise in initial preclinical and clinical studies. LY2456302 reversed KOR agonist-induced analgesia and prepulse inhibition in rats, reduced immobility in the mouse forced swim test, and reduced ethanol self-administration in rats, with no significant effects on motor performance, thermoregulation, or dopamine levels in the nucleus accumbens (Rorick-Kehn et al., 2014). In a clinical study, acute and chronic (14 day) administration of LY2456302 was generally well tolerated in healthy participants, and had a favorable pharmacokinetic profile, supporting its development for therapeutic use in psychiatric disorders (e.g., depression, anxiety, substance use) (Lowe, Wong, Witcher, Gonzales, Dickinson, Bell et al., 2014). LY2456302 has recently advanced to phase II clinical trials for augmentation of antidepressant therapy in treatment-resistant depression and is currently being clinically evaluated for use in mood and anxiety spectrum disorders (ClinicalTrials.gov, 2014). Because of its ability to reduce ethanol self-administration, similar to norBNI and JDTic, LY2456302 may also have therapeutic effects on other drug-induced behaviors.

The goal of this study was to evaluate the effectiveness of the novel KOR antagonist, LY2456302, in established models of nicotine withdrawal in mice, and to determine if a short acting KOR antagonist would show the same effects in a nicotine withdrawal study as previously tested longer acting KOR antagonists, JDTic and norBNI (Jackson et al., 2010). Interest in LY2456302 as a potential therapeutic agent in the treatment of nicotine withdrawal is also heightened since the drug is the only KOR to proceed through to a phase II clinical study. Mice were chronically treated with nicotine for two weeks, and spontaneous nicotine withdrawal behaviors were measured 18–24 hr after mini pump removal, and 60 min after pretreatment with orally-administered LY2456302. The ability of LY2456302 to attenuate expression of nicotine withdrawal CPA was also tested.

2. Materials and Methods

2.1. Animals

Naïve male 8–10 week-old ICR mice (Harlan Laboratories; Indianapolis, IN) served as subjects. Mice were housed five per cage in a 21°C humidity controlled facility with ad

libitum access to food and water. The animal facility was approved by the Association for Assessment and Accreditation of Laboratory Animal Care. Experiments were performed during the light cycle and were approved by the Institutional Animal Care and Use Committee of Virginia Commonwealth University and followed the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals.

2.2. Drugs

(-)-Nicotine hydrogen tartrate salt [(-)-1-methyl-2-(3-pyridyl)pyrrolidine (+)-bitartrate salt] was purchased from Sigma-Aldrich Inc. (St. Louis, MO, USA). (S)-3-fluoro-4-(4-((2-(3,5-dimethylphenyl)pyrrolidin-1-yl)methyl)phenoxy) benzamide (LY2456302) was supplied by the National Institute on Drug Abuse (Washington, DC, USA). The doses of LY2456302 used in our study are within the range of doses used to assess *in vivo* KOR effects as reported in the literature (Rorick-Kehn et al., 2014). Nicotine was dissolved in physiological saline (0.9% sodium chloride). The pH of the nicotine solution was checked and neutralized if necessary. LY2456302, was dissolved in water with the addition of 2% methylcellulose. LY2456302 was given to mice orally via gavage at the volume of 10 ml/kg. Drugs were mixed fresh on the day of dosing. All doses are expressed as the free base of the drug.

2.3. Nicotine chronic administration protocol

Mice were anesthetized with sodium pentobarbital (45 mg/kg, intraperitoneal) and implanted subcutaneously (s.c.) with Alzet osmotic mini pumps [(model 2002); Durect Corporation, Cupertino, CA]. Mice received one dose of ketoprofen after the mini pump operation. The concentration of nicotine was adjusted according to animal weight and the mini pump flow rate to deliver 24 mg/kg/day for 14 days. The spontaneous nicotine withdrawal model for induction of nicotine dependence has generally used the 24 mg/kg/day nicotine dose, as this dose was shown to produce a significant withdrawal syndrome in a spontaneous model for all three behavioral tests when compared to a higher dose of 48 mg/kg/day (Damaj, Kao, & Martin, 2003).

2.4. Spontaneous nicotine withdrawal

To assess the effects of acute LY2456302 administration on spontaneous withdrawal from nicotine, mice were infused with nicotine or saline for 14 days. On day 14, mice were anesthetized under isoflurane anesthesia and minipumps were removed in the evening. No analgesic was given after mini pump removal, as this would interfere with hyperalgesia testing. Mice were habituated to the gavage procedure for oral drug administration for 3 days prior to mini pump removal. On day 15, 18–24 hr after mini pump removal, these mice were pretreated orally (p.o.) 60 min prior to testing with either vehicle or LY2456302 (1, 3, and 10 mg/kg, p.o.). Immediately following the 60 min pre-treatment, mice were observed for physical and affective nicotine withdrawal signs as described in Damaj et al (2003). Mice were first evaluated for 5 min in the plus maze test for anxiety-related behavior. Time spent on the open arms of the plus maze was assessed as a measure of anxiety-related response. The number of arm crosses between the open and closed arms was also counted as a measure of locomotor activity. The plus maze assessment was immediately followed by a 20-min observation of somatic signs measured as paw and body tremors, head shakes, backing, jumps, curls, and ptosis. Mice were placed in clear activity cages without bedding

for the observation period. The total number of somatic signs was tallied for each mouse and the average number of somatic signs during the observation period was plotted for each test group. Hyperalgesia was evaluated using the hot plate test immediately following the somatic sign observation period. Mice were placed into a 10-cm wide glass cylinder on a hot plate (Thermojust Apparatus, Richmond, VA) maintained at 52°C. The latency to reaction time (jumping or paw licking) was recorded. The specific testing sequence was chosen based on our prior studies showing that this order of testing reduced within-group variability and produced the most consistent results (Jackson, Martin, Changeux, & Damaj, 2008). All studies were performed by an observer blinded to experimental treatment.

2.5. Nicotine withdrawal in the CPA test

The CPA protocol was conducted over the course of four days in a biased fashion as described in Jackson et al. (2009). Briefly, mice were implanted with 28-day nicotine (36 mg/kg/day) or saline mini pumps 14 days prior to initiation of CPA testing to induce dependence. Infusion continued throughout the duration of testing. The nicotine withdrawal CPA model was developed using the dose of 36 mg/kg/day (Jackson, Kota, Martin, & Damaj, 2009), as this dose produced a more robust CPA than lower nicotine doses. On day 1 of the CPA procedure, after a 5 min habituation period in the center compartment, mice were allowed to roam freely between compartments for 15 minutes to determine baseline responses. The pre-preference score was used to pair each mouse with mecamlamine (3.5 mg/kg, s.c.) to its initially preferred compartment. On days 2 and 3 of CPA training, all mice received injections of saline in the morning and mecamlamine in the afternoon. In addition, 60 min prior to afternoon conditioning animals received LY2456302 (0.3, 1, and 3 mg/kg, p.o.) or vehicle (p.o.). On day 4 in the afternoon, mice moved freely between compartments as on day 1 and time spent in each compartment was recorded for each mouse. The CPA score was calculated as the amount of time spent in the initially preferred compartment at baseline minus the amount of time spent in the initially preferred compartment on test day. A reduction in time spent in the initially preferred compartment was interpreted as CPA.

2.6. Statistical analysis

Statistical analyses of all behavioral data were performed using Prism 6 (GraphPad Software, La Jolla, CA, USA). All behavioral scores were expressed as mean $\pm$ S.E.M. for each group. Statistical analyses of all studies were analyzed using one-way analyses of variance where appropriate, with treatment as the between subject factor. *p* values <0.05 were considered to be statistically significant. Significant results were further analyzed using the Student Neuman-Keuls post-hoc test.

3. Results

3.1 Physical and affective nicotine withdrawal signs are attenuated by KOR antagonist LY2456302

Physical (somatic signs and hyperalgesia) and affective (anxiety-related behavior) signs were measured in mice following 18–24 h withdrawal from chronic nicotine and treatment with either LY2456302 or vehicle. Results are shown in Figure 1. Nicotine withdrawal significantly increased anxiety-related behavior in the plus maze (Figure 1A), increased

expression of somatic withdrawal signs (Figure 1B), and decreased response latencies in the hot-plate test (Figure 1C). Pretreatment with LY2456302 at 3 and 10 mg/kg (p.o.) significantly decreased the expression of anxiety-related behaviors (time in open arms in the plus-maze test) ($[F(5,40)=7.888, p<0.001]$) (Figure 1A). To ensure that the results in the plus maze were not due to changes in locomotor activity, the average total number of arm crosses was analyzed for each group. There were no significant between group differences in this measure (Table 1). In addition, pretreatment with LY2456302 at all three doses (1, 3, and 10 mg/kg, p.o.) dose-dependently decreased the expression of nicotine somatic withdrawal signs ($[F(5,36)=41.26, p<0.0001]$) (Figure 1B). Finally, pretreatment with LY2456302 at 3 and 10 mg/kg (p.o.) decreased the expression of hyperalgesia (hot-plate latency) ($[F(5,36)=4.449, p<0.002]$) (Figure 1C). The highest dose of LY2456302 (10 mg/kg) used in this assessment did not significantly affect behavioral responses in saline-infused mice in any withdrawal test and did not precipitate significant nicotine withdrawal signs in nicotine-dependent mice at 1, 8, or 18 h after administration (data not shown).

3.2. Expression of nicotine withdrawal aversion is blocked by pretreatment with KOR antagonist LY2456302

A place-conditioning procedure was used to measure effects of KOR antagonism on expression of an aversion associated with nicotine withdrawal. Mice receiving chronic infusions of nicotine or saline via mini pump were exposed to conditioning sessions with mecamylamine (3.5 mg/kg, s.c.) or its vehicle, and LY2456302 was administered 60 min prior to testing. Figure 2 shows a significant effect of treatment on CPA [$F(6,35)=8.994; p<0.0001$]. Post hoc testing indicated that mecamylamine (3.5 mg/kg, s.c.) precipitated a significant CPA in chronic nicotine-exposed mice pretreated with vehicle. Pretreatment with LY2456302 (1 and 3 mg/kg, p.o.) prior to the test day blocked expression of the mecamylamine-induced CPA. The highest dose of LY2456302 (3 mg/kg, p.o.) did not produce significant responses in saline-infused mice.

4. Discussion

LY2456302 is a short acting, high-affinity, selective KOR antagonist that has demonstrated therapeutic potential for psychiatric disorders in animal models. Findings from this study show that, like the longer acting prototypical KOR antagonists (norBNI, JDTic), LY2456302 alleviates physical and affective nicotine withdrawal signs in mice, as demonstrated by reduced anxiety-related behavior, somatic signs, and CPA, and increased hot plate latency compared to vehicle treated nicotine withdrawn mice. This is the first study evaluating the behavioral effects of this novel KOR antagonist with nicotine. It is also emphasized that LY2456302 is the only KOR antagonist that has passed through to phase II clinical trial, thus, it is of particular interest to examine the utility of this compound for smoking cessation.

The nicotine withdrawal syndrome is considered to be one of the major causes of high relapse rate in individuals undergoing smoking cessation (Le Foll & Goldberg, 2009). The nicotine withdrawal syndrome in abstinent smokers is comprised of physical, or somatic components, cognitive, and affective components. The somatic signs include bradycardia,

gastrointestinal discomfort, and increased appetite. Affective symptoms primarily include depressed mood including anhedonia, dysphoria, anxiety, irritability, difficulty concentrating, and craving (Hughes, 2007). Although the somatic manifestations of withdrawal from drugs of abuse are certainly unpleasant, escape from affective components of withdrawal may play a more important role in the maintenance of dependence to drugs of abuse, including nicotine (Koob & Le Moal, 2005; Koob & Volkow, 2010). KOR antagonists blocked expression of all physical and affective nicotine withdrawal behaviors measured in our study, though these behaviors are mediated by different nicotinic acetylcholine receptor subtypes and neuronal pathways (Jackson et al., 2008; Jackson et al., 2009). Currently available therapies involve nicotine replacement (e.g., patch, gum, nasal spray), bupropion and varenicline. While the mechanisms of action for the pharmacological agents, bupropion and varenicline are not totally clear, they likely involve interaction with nicotinic receptors. Although nicotinic receptors are the primary site of action for nicotine, KOR antagonists would have a totally different mechanism of action. Previous studies have shown that extracellular levels of dopamine in the nucleus accumbens (NAc) are reduced in adult rats experiencing withdrawal (Markou, 2008). Since the dynorphin/KOR system maintains tonic activity to inhibit dopamine transmission in the NAc, it is possible that the withdrawal signs and dysphoric state associated with nicotine withdrawal are regulated by the dynorphin/KOR system through dopaminergic transmission (Shippenberg, Zapata, & Chefer, 2007). Ideally, KOR antagonists would target both affective and somatic nicotine withdrawal signs in clinical studies, thus alleviating multiple aspects of the nicotine withdrawal syndrome.

KOR signaling has also been implicated in stress-induced reinstatement of drug use. Stress is a risk factor in maintenance of drug addictive behaviors and relapse to drug abuse. In the case of nicotine, exposure to stressors in humans increases smoking and nicotine intake (Todd, 2004; Pomerleau & Pomerleau, 1987), alters smoking topography (e.g., increased cigarette puff rate and inhaled smoke volume) (Rose, Ananda, & Jarvik, 1983), and desire to smoke (Todd, 2004; Perkins & Grobe, 1992). NorBNI blocked stress-induced reinstatement of nicotine conditioned place preference in mice, but not nicotine-primed reinstatement, suggesting that the kappa opioid system is specifically involved in signaling processes that mediate stress induced drug relapse, as opposed to drug-induced relapse (Jackson, McLaughlin, Carroll, & Damaj, 2013). The effect is not specific to nicotine, as similar findings were observed with stress-induced relapse to cocaine and ethanol-induced behaviors in rodents (Carey, Borozny, Aldrich, & McLaughlin, 2007; Beardsley, Howard, Shelton, & Carroll, 2005; McLaughlin, Marton-Popovici, & Chavkin, 2003; Sperling, Gomes, Sypek, Carey, & McLaughlin, 2010; Redila & Chavkin, 2008). Because KOR signaling is specific to stress-induced drug relapse, and spans across multiple drugs of abuse, it would be useful to examine LY2456302 efficacy in this response.

The results of this study show that, similar to the effects observed with norBNI and JDTic (Jackson et al., 2010), pre-treatment with LY2456302 alleviates physical and affective nicotine withdrawal signs in mouse models of nicotine withdrawal. Of importance for a potential therapeutic drug, LY2456302 appears to have low risk for abuse liability, as it does not significantly affect dopamine levels in the nucleus accumbens at behaviorally relevant doses (Rorick-Kehn et al., 2014), and it is generally well tolerated in rodents and humans

(Lowe et al., 2014; Rorick-Kehn et al., 2014). Though few studies have evaluated the behavioral effects of LY2456302, results from the current study support findings from previous studies using rodent models of mood and addictive disorders (Rorick-Kehn et al., 2014), and suggest that LY2456302 may be efficacious in the treatment of behaviors associated with ethanol and nicotine dependence. Because KOR receptors are also involved in stress-induced drug relapse, drug action at these receptors may decrease the likelihood of relapse, craving, and increased cigarette smoking resulting from exposure to stressors. With its favorable pharmacokinetic and pharmacodynamic profile, it would be beneficial to expand studies using LY2456302 to behaviors associated with stress-induced drug relapse, and other drugs of abuse to further elucidate the therapeutic potential of this compound for drug addiction disorders.

Acknowledgements

The authors would like to thank Tie Han for his work in conducting the withdrawal assays. This work was supported the NIDA Medications Development Program and by National Institute on Drug Abuse grants R01DA032933 and N01DA-12-8904 to MID.

References

- Beardsley PM, Howard JL, Shelton KL, Carroll FI. Differential effects of the novel kappa opioid receptor antagonist, JD1c, on reinstatement of cocaine-seeking induced by footshock stressors vs cocaine primes and its antidepressant-like effects in rats. *Psychopharmacology*. 2005; 183(1):118–126. [doi]. Retrieved from PM:16184376. [PubMed: 16184376]
- Berger AL, Williams AM, McGinnis MM, Walker BM. Affective cue-induced escalation of alcohol self-administration and increased 22-kHz ultrasonic vocalizations during alcohol withdrawal: role of kappa-opioid receptors. *Neuropsychopharmacology*. 2013; 38(4):647–654. doi:npp2012229 [pii]; 10.1038/npp.2012.229 [doi]. Retrieved from PM:23212453. [PubMed: 23212453]
- Carey AN, Borozny K, Aldrich JV, McLaughlin JP. Reinstatement of cocaine place-conditioning prevented by the peptide kappa-opioid receptor antagonist aroclon. *Eur J Pharmacol*. 2007; 569(1–2):84–89. doi:S0014-2999(07)00548-1 [pii];10.1016/j.ejphar.2007.05.007 [doi]. Retrieved from PM:17568579. [PubMed: 17568579]
- Carroll FI, Carlezon WA Jr. Development of kappa opioid receptor antagonists. *J Med Chem*. 2013; 56(6):2178–2195. [doi]. Retrieved from PM:23360448. [PubMed: 23360448]
- Carroll I, Thomas JB, Dykstra LA, Granger AL, Allen RM, Howard JL, et al. Pharmacological properties of JD1c: a novel kappa-opioid receptor antagonist. *Eur J Pharmacol*. 2004; 501(1–3): 111–119. doi:S0014-2999(04)00926-4 [pii];10.1016/j.ejphar.2004.08.028 [doi]. Retrieved from PM:15464069. [PubMed: 15464069]
- ClinicalTrials.gov. A service of the U.S. National Institutes of Health: Trial records 1,2,3 for LY2456302. 2014. <https://clinicaltrials.gov/ct2/show?term=LY2456302&rank=1> visited 3 Feb 2015.
- Damaj MI, Kao W, Martin BR. Characterization of spontaneous and precipitated nicotine withdrawal in the mouse. *The Journal of pharmacology and experimental therapeutics*. 2003; 307(2):526–534. Retrieved from PM:12970387. [PubMed: 12970387]
- Gillett K, Harshberger E, Valdez GR. Protracted withdrawal from ethanol and enhanced responsiveness stress: regulation via the dynorphin/kappa opioid receptor system. *Alcohol (Fayetteville, N.Y.)*. 2013; 47(5):359–365. doi:S0741-8329(13)00071-2 [pii];10.1016/j.alcohol.2013.05.001 [doi]. Retrieved from PM:23731692.
- Hughes JR. Effects of abstinence from tobacco: valid symptoms and time course. *Nicotine Tob Res*. 2007; 9(3):315–327. doi:772841444;10.1080/14622200701188919. Retrieved from PM:17365764. [PubMed: 17365764]

- Jackson KJ, Carroll FI, Negus SS, Damaj MI. Effect of the selective kappaopioid receptor antagonist JDTC on nicotine antinociception, reward, and withdrawal in the mouse. *Psychopharmacology*. 2010; 210(2):285–294. Retrieved from PM:20232057. [PubMed: 20232057]
- Jackson KJ, Kota DH, Martin BR, Damaj MI. The role of various nicotinic receptor subunits and factors influencing nicotine conditioned place aversion. *Neuropharmacology*. 2009; 56(6–7):970–974. doi:S0028-3908(09)00044-6;10.1016/j.neuropharm.2009.01.023. Retrieved from PM: 19371584. [PubMed: 19371584]
- Jackson KJ, Martin BR, Changeux JP, Damaj MI. Differential role of nicotinic acetylcholine receptor subunits in physical and affective nicotine withdrawal signs. 2008; 325:302–312.
- Jackson KJ, McLaughlin JP, Carroll FI, Damaj MI. Effects of the kappa opioid receptor antagonist, norbinaltorphimine, on stress and drug-induced reinstatement of nicotine-conditioned place preference in mice. *Psychopharmacology*. 2013; 226(4):763–768. [doi]. Retrieved from PM: 22526543. [PubMed: 22526543]
- Kelsey JE, Verhaak AM, Schierberl KC. The kappa-opioid receptor antagonist, nor-binaltorphimine (nor-BNI), decreases morphine withdrawal and the consequent conditioned place aversion in rats. *Behav Brain Res*. 2015; 283C:16–21. doi:S0166-4328(15)00012-1 [pii];10.1016/j.bbr.2015.01.008 [doi]. Retrieved from PM:25591478. [PubMed: 25591478]
- Koob GF, Le Moal M. Plasticity of reward neurocircuitry and the 'dark side' of drug addiction. *Nature neuroscience*. 2005; 8(11):1442–1444. doi:nn1105-1442 [pii];10.1038/nn1105-1442 [doi]. Retrieved from PM:16251985. [PubMed: 16251985]
- Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacology*. 2010; 35(1):217–238. doi:npp2009110 [pii];10.1038/npp.2009.110 [doi]. Retrieved from PM:19710631. [PubMed: 19710631]
- Le Foll B, Goldberg SR. Effects of nicotine in experimental animals and humans: an update on addictive properties. *Handb.Exp.Pharmacol*. 2009; (192):335–367. Retrieved from PM:19184655. [PubMed: 19184655]
- Lowe SL, Wong CJ, Witcher J, Gonzales CR, Dickinson GL, Bell RL, et al. Safety, tolerability, and pharmacokinetic evaluation of single- and multiple-ascending doses of a novel kappa opioid receptor antagonist LY2456302 and drug interaction with ethanol in healthy subjects. *J Clin Pharmacol*. 2014; 54(9):968–978. [doi]. Retrieved from PM:24619932. [PubMed: 24619932]
- Markou A. Review. Neurobiology of nicotine dependence. *Philosophical transactions of the Royal Society of London.Series B, Biological sciences*. 2008; 363(1507):3159–3168. doi:T037063847WM0768;10.1098/rstb.2008.0095. Retrieved from PM:18640919. [PubMed: 18640919]
- McLaughlin JP, Marton-Popovici M, Chavkin C. Kappa opioid receptor antagonism and prodynorphin gene disruption block stress-induced behavioral responses. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2003; 23(13):5674–5683. doi:23/13/5674 [pii]. Retrieved from PM:12843270. [PubMed: 12843270]
- Munro TA, Berry LM, Van't Veer A, Beguin C, Carroll FI, Zhao Z, et al. Long-acting kappa opioid antagonists nor-BNI, GNTI and JDTC: pharmacokinetics in mice and lipophilicity. *BMC Pharmacol*. 2012; 12:5. doi:1471-2210-12-5 [pii];10.1186/1471-2210-12-5 [doi]. Retrieved from PM:22642416. [PubMed: 22642416]
- Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM. Neurobiology of depression. *Neuron*. 2002; 34(1):13–25. doi:S0896627302006530 [pii]. Retrieved from PM:11931738. [PubMed: 11931738]
- Patkar KA, Wu J, Ganno ML, Singh HD, Ross NC, Rasakham K, et al. Physical presence of nor-binaltorphimine in mouse brain over 21 days after a single administration corresponds to its long-lasting antagonistic effect on kappa-opioid receptors. *The Journal of pharmacology and experimental therapeutics*. 2013; 346(3):545–554. doi:jpet.113.206086 [pii];10.1124/jpet.113.206086 [doi]. Retrieved from PM:23853171. [PubMed: 23853171]
- Perkins KA, Grobe JE. Increased desire to smoke during acute stress. *Br.J Addict*. 1992; 87(7):1037–1040. Retrieved from PM:1643396. [PubMed: 1643396]
- Pomerleau CS, Pomerleau OF. The effects of a psychological stressor on cigarette smoking and subsequent behavioral and physiological responses. *Psychophysiology*. 1987; 24(3):278–285. Retrieved from PM:3602283. [PubMed: 3602283]

- Redila VA, Chavkin C. Stress-induced reinstatement of cocaine seeking is mediated by the kappa opioid system. *Psychopharmacology*. 2008; 200(1):59–70. [doi]. Retrieved from PM:18575850. [PubMed: 18575850]
- Rorick-Kehn LM, Witkin JM, Statnick MA, Eberle EL, McKinzie JH, Kahl SD, et al. LY2456302 is a novel, potent, orally-bioavailable small molecule kappa-selective antagonist with activity in animal models predictive of efficacy in mood and addictive disorders. *Neuropharmacology*. 2014; 77:131–144. doi:S0028-3908(13)00428-0 [pii];10.1016/j.neuropharm.2013.09.021 [doi]. Retrieved from PM:24071566. [PubMed: 24071566]
- Rose JE, Ananda S, Jarvik ME. Cigarette smoking during anxiety-provoking and monotonous tasks. *Addict Behav*. 1983; 8(4):353–359. doi:0306-4603(83)90035-7 [pii]. Retrieved from PM:6677075. [PubMed: 6677075]
- Schank JR, Goldstein AL, Rowe KE, King CE, Marusich JA, Wiley JL, et al. The kappa opioid receptor antagonist JDTC attenuates alcohol seeking and withdrawal anxiety. *Addict Biol*. 2012; 17(3):634–647. [doi]. Retrieved from PM:22515275. [PubMed: 22515275]
- Shippenberg TS, Zapata A, Chefer VI. Dynorphin and the pathophysiology of drug addiction. *Pharmacology & therapeutics*. 2007; 116(2):306–321. doi:S0163-7258(07)00147-7 [pii];10.1016/j.pharmthera.2007.06.011 [doi]. Retrieved from PM:17868902. [PubMed: 17868902]
- Sperling RE, Gomes SM, Sypek EI, Carey AN, McLaughlin JP. Endogenous kappa-opioid mediation of stress-induced potentiation of ethanol-conditioned place preference and self-administration. *Psychopharmacology*. 2010; 210(2):199–209. [doi]. Retrieved from PM:20401606. [PubMed: 20401606]
- Sudakov SK, Nazarova GA, Alekseeva EV, Kolpakov AA. Effect of activated peripheral kappa-opioid receptors on the action of nicotine and its withdrawal in nicotine-dependent rats. *Bull Exp Biol Med*. 2014; 156(5):609–611. [doi]. Retrieved from PM:24770739. [PubMed: 24770739]
- Tejeda HA, Natividad LA, Orfila JE, Torres OV, O'Dell LE. Dysregulation of kappa-opioid receptor systems by chronic nicotine modulate the nicotine withdrawal syndrome in an age-dependent manner. *Psychopharmacology*. 2012; 224(2):289–301. Retrieved from PM:22659976. [PubMed: 22659976]
- Todd M. Daily processes in stress and smoking: effects of negative events, nicotine dependence, and gender. *Psychology of addictive behaviors : journal of the Society of Psychologists in Addictive Behaviors*. 2004; 18(1):31–39. Retrieved from PM:15008683. [PubMed: 15008683]
- Valdez GR, Harshberger E. kappa opioid regulation of anxiety-like behavior during acute ethanol withdrawal. *Pharmacol Biochem Behav*. 2012; 102(1):44–47. doi:S0091-3057(12)00090-1 [pii]; 10.1016/j.pbb.2012.03.019 [doi]. Retrieved from PM:22487769. [PubMed: 22487769]
- Van't Veer A, Carlezon WA Jr. Role of kappa-opioid receptors in stress and anxiety-related behavior. *Psychopharmacology*. 2013; 229(3):435–452. [doi]. Retrieved from PM:23836029. [PubMed: 23836029]
- Walker BM, Koob GF. Pharmacological evidence for a motivational role of kappa-opioid systems in ethanol dependence. *Neuropsychopharmacology*. 2008; 33(3):643–652. doi:1301438 [pii]; 10.1038/sj.npp.1301438 [doi]. Retrieved from PM:17473837. [PubMed: 17473837]
- Walker BM, Zorrilla EP, Koob GF. Systemic kappa-opioid receptor antagonism by nor-binaltorphimine reduces dependence-induced excessive alcohol self-administration in rats. *Addict Biol*. 2011; 16(1):116–119. doi:ADB226 [pii];10.1111/j.1369-1600.2010.00226.x [doi]. Retrieved from PM:2057900. [PubMed: 20579007]

Highlights

- Kappa opioid receptors, in part, mediate mood and substance use disorders.
- A new kappa opioid receptor antagonist, LY2456302, may have therapeutic potential.
- LY2456302 was evaluated for potential to alleviate nicotine withdrawal in mice.
- LY2456302 reduced expression of physical and affective nicotine withdrawal in mice.
- Findings suggest that LY2456302 may be efficacious across multiple drugs of abuse.

Figure 1. LY2456302 attenuates physical and affective nicotine withdrawal signs in mice
 Withdrawal from nicotine induced a A) significant increase in anxiety-related behavior, B) significant increase in somatic signs, and C) a significant decrease in hot plate latency. Compared to vehicle, pretreatment with LY2456302 A) significantly decreased expression of anxiety-related behavior at 3 and 10 mg/kg p.o., B) significantly reduced somatic signs at all doses tested, and C) significantly increased hot plate latency at 3 and 10 mg/kg p.o. in nicotine withdrawal mice. Each point represents the mean $\pm$ S.E.M. of 7–8 mice per group. All doses are expressed in mg/kg. * denotes $p < 0.05$ vs. Sal/Veh group, # denotes $p < 0.05$ vs. Nic/Veh group. Veh, vehicle; Sal, saline; Nic, nicotine; LY, LY2456302

Figure 2. LY2456302 blocks expression of nicotine withdrawal aversion

Mecamylamine (3.5 mg/kg, s.c.) precipitated a significant nicotine withdrawal CPA in chronic nicotine-infused mice. Pretreatment with LY2456302 dose-dependently blocked expression of CPA. Each point represents the mean $\pm$ S.E.M. of 7–8 mice per group. * denotes $p < 0.05$ vs. Sal-Mec/Veh control group, # denotes $p < 0.05$ vs. Nic-Mec/Veh. Veh, vehicle; Mec, mecamylamine; Sal, saline; Nic, nicotine; LY, LY2456302

Table 1
LY2456302 does not significantly alter the average number of arm crosses in the plus maze assessment

Nicotine-withdrawn mice were treated with vehicle or LY2456302 (1, 3, and 10 mg/kg, p.o.), and the average number of crosses between the open and closed arms of the plus maze was counted. Numbers are presented as the total average number of arm crosses $\pm$ SEM for seven to eight mice per group MP, mini pump.

Treatment	Average number of arm crosses $\pm$ SEM
Saline MP-vehicle	6.3 $\pm$ 0.7
Saline MP-LY2456302 (10)	6.4 $\pm$ 0.9
Nicotine MP-vehicle	6.6 $\pm$ 0.6
Nicotine MP-LY2456302 (1)	5.7 $\pm$ 0.7
Nicotine MP-LY2456302 (3)	5.9 $\pm$ 0.3
Nicotine MP-LY2456302 (10)	5.6 $\pm$ 0.4