Rare disease

Cerebellar haemangioblastoma: a rare entity

Mehar Aziz,¹ Kiran Alam,¹ Manoranjan Varshney,¹ Veena Maheshwari,¹ Rana K Sherwani,¹ Kavita Gaur,¹ Vinod Kumar Srivastava²

¹Pathology Department, J. N. Medical College, Aligarh, Uttar Pradesh, India;

²Surgery Department, J. N. Medical College, Aligarh, Uttar Pradesh, India

Correspondence to Dr Kavita Gaur, kavgaur@gmail.com

Summary

A 45-year-old female presented with headache, nausea, vomiting and ataxia of 4-month duration. CT head showed a posterior fossa tumour which was diagnosed as a case of haemangioblastoma on microscopic examination.

BACKGROUND

Haemangioblastoma is a rare central nervous system tumour which often mimics haemangioma and astrocytoma clinically and radiologically.

CASE PRESENTATION

A 45-year-old female presented to neurosurgery outpatient department with a 4-month history of headache, nausea, vomiting and ataxia. General examination was unremarkable; however, systemic examination showed features of cerebellar ataxia.

INVESTIGATIONS

Contrast-enhanced CT scan of head revealed a well-defined cystic density lesion measuring 30×48×25 mm in right cerebellar hemisphere, extending superiorly to vermis. Small mural nodule was noted in the posterior aspect of the lesion. It was causing mass effect in the form of mild compression of the fourth ventricle. After intravenous injection of contrast, the mural nodule showed enhancement whereas the cystic lesion did not show any enhancement (figure 1A,B). Fundus examination, audiometry and

CT abdomen were unremarkable. Molecular genetic testing for von Hippel–Lindau (VHL) disease did not detect any abnormality. There was no family history suggestive of similar disease. Surgical resection of tumour was done. Gross examination showed a greyish soft tissue piece measuring 1.5 cm in diameter which was solid on cut section, and microscopic examination revealed polygonal cells with juxtaposed capillary channels (figure 2). These polygonal cells had clear cytoplasm and minimal nuclear pleomorphism (figure 3A,B). Mitosis was rare. Surrounding tissue had gliosis (figure 3C) and normal brain parenchyma adjacent to tumour was also noticed (figure 3D). Reticulin stain delineates the outlines of vascular architecture (figure 4). On the basis of above findings, a diagnosis of haemangioblastoma was rendered.

DIFFERENTIAL DIAGNOSIS

Differential diagnoses include metastatic tumour especially renal cell carcinoma, clear cell meningioma, pilocytic astrocytoma, clear cell ependymoma, arteriovenous malformation and haemangioma. Our case lacked perivascular pseudorosettes, endothelial proliferation and hypercellularity

Figure 1 (A,B) CT head showing enhancement in the mural nodule while the cystic lesion did not show any enhancement.

Figure 2 Section showing polygonal cells with juxta-posed capillary channels (H&E ×50).

Figure 3 (A) High power view showing clear vacuolated cells (H&E ×500). (B) High power view showing vacuolated cells with juxta-posed blood vessels (H&E ×500). (C) High power view showing gliosis surrounding the lesion (H&E ×500). (D) High power view showing normal brain parenchyma adjacent to the lesion (H&E ×500).

frequent mitosis of clear cell ependymoma. The presence of intimate mixture of vacuolated stromal cells and vascular spaces rule out arteriovenous malformation, haemangioma and clear cell meningioma which may also have vague whorl formation and intranuclear inclusions. Gliosis

surrounding the tumour may sometime mimic pilocytic astrocytoma but presence of prominent vascularity rule it out. Metastatic renal cell carcinoma can be a close mimic, but it is composed of prominent nucleoli, non-uniform chromatin, mitosis and necrosis.

Figure 4 (A,B) Reticulin stain delineating the outline of blood vessel and vacuolated cells (Reti $\times 500$).

TREATMENT

Surgical excision of tumour was done.

OUTCOME AND FOLLOW-UP

Patient was discharged in good conditions, and till date (9 months after surgery) no foci of recurrence is seen. During follow-up visits, ophthalmologic examination, audiometry, blood pressure, ultrasonography of the abdomen and urinary examination for catecholamine metabolites were done and found to be unremarkable.

DISCUSSION

Haemangioblastomas are benign vascular tumours comprising 1–2% of all intracranial neoplasm^{1–3} and about 7.3% of posterior fossa tumours.⁴ They can manifest sporadically or associated with von VHL disease in 20–25% of cases.⁵ When associated with VHL, the mean age at diagnosis is third decade; however, sporadic cases are seen a decade later. It is commonly seen in cerebellum^{6–8}; however, they can be found in brain stem and spinal cord.^{3,9} On the basis of macroscopic examination, they can be divided into simple cystic, macrocystic, solid and microcystic type. Macrocystic type comprises cyst of variable size within a mural nodule as in our case. Clinical features depend upon the anatomic location of tumour. When present in posterior fossa, the patient presents with cerebellar dysfunctions and features of raised intracranial pressure which include headache, nausea, vomiting, ataxia and dizziness. Histopathological examination reveals proliferating blood vessels with stromal cells which are large and vacuolated. Definitive treatment is complete resection of tumour. After complete resection of tumour, recurrence is rare.^{10,11} Prognosis is better in our case as it was not a part of VHL syndrome.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Cushing H**, Bayley B. *Tumours Arising from the Blood Vessels of the Brain: Angiomatous Malformations and Hemangioblastomas*. Springfield IL: Charles C Thomas, 1928.
2. **Neumann HP**, Eggert HR, Weigel K, *et al*. Hemangioblastomas of the central nervous system. A 10-year study with special reference to von Hippel–Lindau syndrome. *J Neurosurg* 1989;**70**:24–30.
3. **Weil RJ**, Vortmeyer AO, Zhuang Z, *et al*. Clinical and molecular analysis of disseminated hemangioblastomatosis of the central nervous system in patients without von Hippel–Lindau disease. Report of four cases. *J Neurosurg* 2002;**96**:775–87.
4. **OLIVECRONA H**. The cerebellar angiorectulomas. *J Neurosurg* 1952;**9**:317–30.
5. **Böhling T**, Plate KH, Haltia MJ, *et al*. Von Hippel–Lindau disease and capillary hemangioblastoma. In: Kleihues P, Caveness WK, eds. *World Health Organization Classification of Tumours. Pathology and Genetics-Tumours of the Nervous System*. Lyon: IARC Press, 2000;223 6.
6. **Fomekong E**, Hernalsteen D, Godfraind C, *et al*. Pituitary stalk hemangioblastoma: the fourth case report and review of the literature. *Clin Neurol Neurosurg* 2007;**109**:292–8.
7. **Rumboldt Z**, Gnjidi Z, Talan-Hranilovi J, *et al*. Intracellar hemangioblastoma: characteristic prominent vessels on MR imaging. *AJR Am J Roentgenol* 2003;**180**:1480–1.
8. **Wasenko JJ**, Rodziewicz GS. Suprasellar hemangioblastoma in Von Hippel–Lindau disease: a case report. *Clin Imaging* 2003;**27**:18–22.
9. **Conway JE**, Chou D, Clatterbuck RE, *et al*. Hemangioblastomas of the central nervous system in von Hippel–Lindau syndrome and sporadic disease. *Neurosurgery* 2001;**48**:55–62.
10. **Kato M**, Ohe N, Okumura A, *et al*. Hemangioblastomatosis of the central nervous system without von Hippel–Lindau disease: a case report. *J Neurooncol* 2005;**72**:267–70.
11. **Reyns N**, Assaker R, Louis E, *et al*. Leptomeningeal hemangioblastomatosis in a case of von Hippel–Lindau disease: case report. *Neurosurgery* 2003;**52**:1212–15.

Learning points

- Haemangioblastoma is a rare posterior fossa tumour.
- One-third cases are associated with VHL disease.
- It is earlier when associated with VHL than in sporadic cases.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Aziz M, Alam K, Varshney M, Maheshwari V, Sherwani RK, Gaur K, Srivastava VK. Cerebellar haemangioblastoma: a rare entity. *BMJ Case Reports* 2011;10.1136/bcr.03.2011.3943, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow