Clinical use of the β_3 adrenoceptor agonist mirabegron in patients with overactive bladder syndrome

Monika Vij and Marcus J. Drake

Ther Adv Urol

2015, Vol. 7(5) 241–248

DOI: 10.1177/
1756287215591763

© The Author(s), 2015.

Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Abstract: Mirabegron is a β_3 adrenoceptor agonist licensed for the treatment of overactive bladder symptoms, such as urinary urgency or urgency incontinence. β_3 adrenoceptor activation causes detrusor muscle relaxation, but mirabegron may also act by binding other targets in the bladder, and it may also reduce activity in sensory nerves. Phase III clinical trials (SCORPIO, ARIES, and CAPRICORN) evaluated mirabegron at various doses, demonstrating reduction from baseline to endpoint in mean incontinence episodes and mean number of micturitions per 24 h (coprimary endpoints), along with health-related quality of life and a range of secondary measures. Efficacy was seen in many patients who had previously discontinued antimuscarinic therapy on the grounds of lack of efficacy or poor tolerability. Treatment emergent adverse effects were documented in a long-term study (TAURUS), mostly being of mild or moderate severity. The most frequent adverse effects were hypertension, dry mouth, constipation, and headache, with a lower incidence of dry mouth than for the antimuscarinic active comparator. Efficacy and safety are not substantially different in older patients. A urodynamic safety study in men showed no consistent effect on voiding function, but a small increase in postvoid residual. Use of mirabegron in combination with α -adrenergic blockers does not appear to increase adverse effects. Dose reduction is needed in people with severe renal failure, or moderate hepatic failure. Dose adjustment is not needed in relation to food intake. Ongoing research is evaluating the potential for combination therapy with antimuscarinics.

Keywords: mirabegron, beta-3 agonist, overactive bladder, LUTS, urinary urgency, urgency urinary incontinence

Introduction

Lower urinary tract symptoms (LUTS) affecting urine storage include urinary urgency, urgency incontinence, increased daytime frequency and nocturia [Abrams *et al.* 2002]. Overactive bladder syndrome (OAB) is defined by the International Continence Society as urinary urgency, with or without urgency incontinence, usually with increased daytime frequency and nocturia in the absence of proven infection or any other pathology [Drake, 2014]. It is a chronic symptom complex that can substantially impair quality of life. The prevalence of LUTS [Coyne *et al.* 2009], OAB and urgency urinary incontinence is substantial, affecting both women and men, and resulting in considerable morbidity, personal cost and health economic burden [Coyne *et al.* 2014; Milsom *et al.* 2014].

Initial management uses conservative measures, such as fluid advice and bladder training [Lucas *et al.* 2012]. Drug therapy using antimuscarinics has long been the mainstay of the pharmacological treatment of OAB [Chapple *et al.* 2008; Kaplan *et al.* 2011; Lucas *et al.* 2012]. However, compliance with these drugs is generally poor [Wagg *et al.* 2012], due to perceived lack of efficacy and potential high rate of side effects. The emergence of new drugs and treatment regimens should improve patient persistence with treatment and acceptability. Development of mirabegron (YM178), a β_3 -adrenoceptor (AR) agonist now offers a new pharmacotherapy option for the treatment of OAB [Maman *et al.* 2014]. Mirabegron is available for clinical use as a modified release film-coated tablet in an oral controlled

Correspondence to:
Marcus J. Drake, DM, MA, FRCS (Urol)
University of Bristol,
Southmead Hospital,
Bristol BS9 1BQ, UK
marcus.drake@bui.ac.uk
Monika Vij, MS, MRCOG
Southmead Hospital,
Bristol, UK

absorption system (OCAS), with two licensed dosage strengths of 25 and 50 mg.

Mechanism of action

There are three subtypes of β -AR, with the β_3 -AR predominating in the detrusor muscle of some species [Ochodnick *et al.* 2013], including the human bladder. β_3 -AR constitutes a high proportion of the β -AR mRNA in the human detrusor [Igawa *et al.* 2010]. Activation of these receptors elicits detrusor relaxation in bladder strips from various species [Takasu *et al.* 2007], through generation of cyclic adenosine monophosphate [Takasu *et al.* 2007; Limberg *et al.* 2010]. β_3 -AR may also be present in other cell types of the bladder, such as urothelium [Otsuka *et al.* 2008; Limberg *et al.* 2010; Kullmann *et al.* 2011] or interstitial cells [Limberg *et al.* 2010]. However, the role of their activation in the urothelium remains to be established [Moro *et al.* 2013].

In animal studies, activation of β_3 -AR gives dose-dependent detrusor relaxation during the storage phase of the micturition cycle. Functionally, it increases intervoid interval and bladder compliance, while preserving voiding function in a rodent model [Sadananda *et al.* 2013]. β_3 -AR are also suggested to inhibit sensory nerve activation, including activity in mechanosensitive A- δ fibers [Aizawa *et al.* 2010], and also reducing autonomous non-voiding contractions [Woods *et al.* 2001; Aizawa *et al.* 2010; Sadananda *et al.* 2013]. β -AR activation can elicit release of nitric oxide by urothelium, giving a potential indirect mechanism which could further contribute to bladder relaxation [Murakami *et al.* 2007; Yamaguchi and Chapple, 2007]. A critical influence of β_3 -ARs in increasing bladder capacity, without changing micturition pressure or residual volume, may be mediated by inhibition of afferent nerve fiber activity [Kanai *et al.* 2011].

Pharmacokinetics

The marketed formulation of mirabegron is in an OCAS, with oral bioavailability in the range of 24–53% [Kashyap and Tyagi, 2013]. The plasma protein binding is 71% [Dickinson *et al.* 2013]. The time to reach maximum concentration after dosing is 3–4 h, and the terminal elimination half-life is approximately 50 h. The absolute bioavailability is dose-dependent and affected by gender [Kashyap and Tyagi, 2013]. Interactions with cytochrome P450 (CYP) metabolism [Lee *et al.* 2013a; Krauwinkel *et al.* 2014] are potentially relevant clinically. For

example, drug–drug interactions may occur with concomitant administration of drugs that exert inducible (rifampin) or inhibitory (ketoconazole) effects on CYP3A4/5 [Kashyap and Tyagi, 2013]. CYP2D6 inhibition by mirabegron might alter the plasma exposure of metoprolol [Kashyap and Tyagi, 2013]. One study has shown that combining mirabegron with an α -adrenergic antagonist (tamsulosin) did not affect safety aspects, notably cardiovascular adverse effects, in 48 healthy men aged 44 years or older [van Gelderen *et al.* 2014].

Mirabegron OCAS tablets show a decrease in mirabegron plasma exposure with food that is independent of dose (50 or 100 mg) or sex, but dependent on meal composition; however, this does not warrant dose adjustment in clinical practice [Lee *et al.* 2013b]. Pharmacokinetic changes observed in subjects with severe renal impairment or moderate hepatic impairment [Dickinson *et al.* 2013] mean that dose reduction is necessary.

Clinical evidence of efficacy

Mirabegron has been studied extensively, with more than 10,000 subjects involved in research studies over approximately 10 years. The safety and efficacy was evaluated in five global trials (two in phase II and three in phase III) comparing various doses of mirabegron, either with placebo or tolterodine.

BLOSSOM was a phase II proof of concept trial conducted in six European countries [Chapple *et al.* 2013a]. The study randomized subjects to mirabegron at doses of 100 or 150 mg twice daily, placebo (twice daily), or tolterodine 4 mg extended release (ER) once daily for 4 weeks. The primary endpoint was change from baseline to end of treatment in mean number of micturition episodes per 24 h. Mirabegron was superior to placebo and tolterodine, with mean reductions in micturition frequency of 2.2, 1.2, and 1.5, respectively. Mirabegron was superior to placebo in terms of mean volume voided, mean number of incontinence episodes, nocturia episodes and urgency episodes in 24 h. There was no significant difference found between 100 and 150 mg twice daily doses.

The DRAGON study was a 12-week phase IIB dose-ranging study conducted predominantly in Europe [Chapple *et al.* 2013b]. A total of 919 patients were randomized in six arms: placebo, mirabegron at doses of 25, 50, 100, or 200 mg

once daily, or tolterodine as an active comparator. A dose-dependent decrease in mean number of micturitions per 24 h was seen with mirabegron, which was significant at the doses of 50 mg and above compared with placebo. Twenty-eight percent of people receiving mirabegron 50 mg were classified as 'responders' in terms of a reduction in voiding frequency (down to eight times daily or less). This compared with 19% for placebo and 19% for tolterodine. There were also significant improvements seen in secondary endpoints, such as mean number of urgency and urgency incontinence episodes, nocturia, and mean voided volume. Responses were evident at 1 week of treatment, with maximum efficacy gained at 8–12 weeks.

The phase III development program comprised three studies: SCORPIO, ARIES, and CAPRICORN. These were large-scale, 12-week, multicenter, randomized, double-blind, parallel-group studies undertaken in Europe, Australia, and North America. They looked at two coprimary efficacy endpoints: the change from baseline to endpoint in mean incontinence episodes and mean number of micturitions per 24 h. The study populations were adults of at least 18 years of age, with at least eight micturitions per 24 h, and at least three urgency episodes in their 3-day bladder diary, with or without urgency incontinence. Different doses of mirabegron were compared with placebo, and an active antimuscarinic comparator in some studies.

SCORPIO [Khullar *et al.* 2013a] recruited 1978 patients with OAB and showed statistically significant improvement with 50 and 100 mg mirabegron in both the measures. Although improvement was also seen in the tolterodine 4 mg ER arm, it was not statistically significant. Significant improvement was also seen in mean number of urgency episodes and mean voided volume with mirabegron 50 mg compared with placebo.

ARIES enrolled 1328 patients with OAB, and significant improvement was seen in both coprimary endpoints, along with mean level of urgency, mean urgency incontinence episodes per 24 h, and mean nocturia episodes [Nitti *et al.* 2013a].

CAPRICORN included 1306 patients who were randomized to placebo, mirabegron 25 mg, or mirabegron 50 mg [Herschorn *et al.* 2013]. Mean reductions in incontinence episodes were 1.36

and 1.38 for mirabegron 25 and 50 mg, compared with 0.96 for the placebo group. Mean reductions in micturitions per 24 h were 1.65 and 1.60 for mirabegron 25 and 50 mg, respectively (1.18 for placebo). Mirabegron 50 mg demonstrated significantly greater improvements *versus* placebo in mean volume voided/micturition, and this was not seen for mirabegron 25 mg [Herschorn *et al.* 2013].

Pooled analysis of the data from the above trials reviewed outcomes of 3542 patients randomized between placebo, mirabegron 50, and mirabegron 100 mg [Nitti *et al.* 2013b]. This showed the reduction in mean number of incontinence episodes per 24 h after 12 weeks of therapy was −1.10, −1.49, and −1.50, respectively. The equivalent reductions for number of micturitions per 24 h were −1.20, −1.75, and −1.74 [Nitti *et al.* 2013b]. The higher dose did not appear to achieve additional benefit. The active doses also showed benefits in 'dry rates' (zero incontinence episodes per 24 h), at 44.1% (mirabegron 50 mg) and 46.4% (100 mg) compared with 37.8% for placebo [Nitti *et al.* 2013b]. Statistically significant change in nocturia was seen, but the extent of clinical benefit is arguable (−0.55 voids per night for the active compound, −0.42 for placebo). Significant difference relative to placebo in these parameters was seen at 4 weeks.

Mirabegron 25 mg and mirabegron 50 mg were also shown to be effective in older patients with OAB (>65 or >75 years), demonstrating improvements in the primary endpoints [Wagg *et al.* 2014].

Considering that antimuscarinic medications have long been the primary drug therapy for OAB, *post hoc* subgroup analysis looked at the effectiveness in treatment-naïve patients and those who had previously discontinued antimuscarinic treatment (either from lack of efficacy, or due to difficulty tolerating the medication) [Khullar *et al.* 2013b]. Response to mirabegron in terms of primary outcome measures was not affected by antecedent medication exposure (for those who discontinued due to poor efficacy), but it did significantly impact on response to the antimuscarinic active comparator. The placebo effect was found to be greater in the treatment-naïve patients. Further information on this will be given by the BEYOND study [ClinicalTrials.gov identifier: NCT01638000], which studied 1870 patients who were not satisfied by antimuscarinic therapy

(on a treatment satisfaction Likert scale), and randomized them to either mirabegron 50 mg or solifenacin 5 mg.

Overall, onset of response appears to be achieved for efficacy and quality of life measures at 1 month [Chapple *et al.* 2014]. Long-term efficacy was evaluated in the TAURUS study, which was a randomized, double-blind parallel group phase III trial, using mirabegron at doses of 50 or 100 mg and tolterodine 4 mg ER as active comparator [Chapple *et al.* 2013c]. In this trial, 2444 patients were recruited. Efficacy endpoints were secondary measures, with improvements in OAB symptoms for both doses (50 and 100 mg) seen by 1 month, using various measures, and maintained throughout the follow-up period.

Health-related quality of life (HRQL) and treatment satisfaction were assessed in the mirabegron trials using various validated OAB-specific and general scales, such as the overactive bladder questionnaire [Coyne *et al.* 2002], the patient perception of bladder condition [Coyne *et al.* 2006], and a treatment satisfaction visual analog scale. Significant improvement was demonstrated from baseline to final visit in these parameters of HRQL in all three of the main phase III studies.

In the DRAGON study [Chapple *et al.* 2013b], International Consultation on Incontinence Questionnaire-Overactive Bladder (ICIQ-OAB) and ICIQ-OAB quality of life questionnaires were used. Patient-reported benefit was also evaluated, with the question ‘has the treatment been of any benefit to you?’ The percentage of responders was 59.0%, 65%, 65.8%, and 70.8% for 25, 50, 100, and 200 mg mirabegron, respectively compared with 51% for placebo and 55% of the tolterodine group.

The studies included male patients, who generally made up between one quarter and one third of the study population in most of the reported studies, but the analyses did not separate findings according to sex.

Safety and tolerability

Mirabegron appears to have acceptable safety and tolerability in all the reported trials. In the BLOSSOM trial [Chapple *et al.* 2013b], the incidence of treatment-emergent adverse effects (TEAEs) was 39.2% with mirabegron compared with 36.4% and 48.4% for placebo and

tolterodine, respectively. Most of the adverse effects were minor or moderate in severity. The most commonly reported side effects were gastrointestinal disorders (13.8%), followed by headache (6.9%). Dizziness and palpitations were more common in mirabegron groups than the tolterodine group. The discontinuation rates due to adverse effects were 4.6% and 7.7% for the mirabegron groups compared with 1.5% in the placebo group and 3.1% in the tolterodine group.

In the DRAGON trial [Chapple *et al.* 2013b], the incidence of serious adverse events was reported as less than 2% of patients across the treatment groups. The most common side effects reported were again gastrointestinal (7.2–8.3% with mirabegron *versus* 5.3% with placebo), including constipation, dry mouth, dyspepsia, and nausea. Incidence of dry mouth was higher with tolterodine (3.5%) than mirabegron (1.8–3.0%). A dose-dependent small rise in pulse rate was also noted with mirabegron; this was not associated with any cardiovascular side effect, with no difference found in electrocardiogram parameters and blood pressure across the treatment groups. Discontinuation due to side effects was low, at 3.0% for placebo (3.0%), 2.4–5.3% for mirabegron and 1.2% for tolterodine. There were no clinically significant episodes of acute retention observed with mirabegron.

Pooled safety data from SCORPIO, ARIES, and CAPRICORN showed that the overall incidence of TEAEs was similar across the treatment groups and there was no evidence of a dose–response relationship among the mirabegron treatment groups for overall rates of TEAEs (mirabegron 25 mg 48.6%, mirabegron 50 mg 47.1%, mirabegron 100 mg 43.3%, total mirabegron 46.0%, placebo 47.7% and tolterodine ER 4 mg 46.7%) [Nitti *et al.* 2013b]. The most common side effects noted in the mirabegron groups were hypertension (7.3%), nasopharyngitis (3.4%), and urinary tract infection (3.0%). Dry mouth occurred more commonly in the tolterodine group (2.0% for the total mirabegron groups *versus* 2.1% for placebo and 10.1% for the tolterodine group). The discontinuation rates were mirabegron 25 mg 3.9%, mirabegron 50 mg 3.9%, mirabegron 100 mg 3.7%, total mirabegron 3.8%, placebo 3.3%, and tolterodine ER 4 mg 4.4%.

In the 12-month TAURUS study, the incidence of TEAEs was similar across the mirabegron 50 mg (59.7%), mirabegron 100 mg (61.3%), and

tolterodine ER 4 mg (62.6%) groups. Most of the side effects were of mild or moderate severity. The most frequent were hypertension, constipation, and headache, occurring at a similar incidence across treatment groups, and dry mouth. Dry mouth had a higher incidence in the tolterodine group (8.6% *versus* mirabegron 2.3–2.8%). Discontinuations due to adverse events were comparable across treatment groups, occurring in 6.4%, 5.9%, and 6.0% of patients receiving mirabegron 50 mg, mirabegron 100 mg, and tolterodine ER 4 mg, respectively.

In older patients, 49% of placebo patients and 55% on active compound reported a TEAE. The three main adverse effects on active compound were hypertension, nasopharyngitis, and urinary tract infection [Wagg *et al.* 2014]. Dry mouth and constipation did not differ from placebo.

For men, in whom bladder outlet obstruction might be a feature due to benign prostate enlargement, the possibility of impairing voiding function as a consequence of administering medications to aid urine storage has to be considered. A urodynamic safety study was undertaken in 200 men receiving mirabegron 50 or 100 mg or placebo for 12 weeks [Nitti *et al.* 2013c]. Maximum flow rate and detrusor pressure at maximum flow were not significantly impaired by active treatment. A small increase in postvoid residual was seen for mirabegron 100 mg. Acute urinary retention occurred in one man on placebo and one on mirabegron 100 mg. The proportion of patients experiencing at least 150 ml change from baseline in postvoid residual volume was lower in the mirabegron groups compared with tolterodine: mirabegron 25 mg (0%), mirabegron 50 mg (0.3%), mirabegron 100 mg (0.4%), placebo (0.7%), and tolterodine ER 4 mg (0.8%) in the pooled analysis of the phase III trials [Nitti *et al.* 2013b].

Mirabegron was shown not to raise intraocular pressure after treatment with 100 mg daily for 8 weeks in healthy volunteers [Novack *et al.* 2013], which is advantageous when considering that closed-angle glaucoma is a contraindication to antimuscarinic therapy.

Combination therapy

Management of LUTS is increasingly reliant on combination pharmaceutical therapy, and the introduction of a new drug class for OAB offers further opportunities for combinations aiming to increase efficacy or reduce side effects. The

SYMPHONY study [Abrams *et al.* 2014] was a dose-ranging phase II study using a range of permutations combining mirabegron with solifenacin. Several combination doses achieved greater improvements in mean voided volume, micturition frequency, and urgency than solifenacin monotherapy. The BESIDE study [ClinicalTrials.gov identifier: NCT01908829] has recently completed recruitment, and was designed to see if adding mirabegron to an antimuscarinic treatment (solifenacin) will be more effective in controlling incontinence than when using the antimuscarinic treatment alone.

In male LUTS, combinations of antimuscarinic medication and α -adrenergic blockers are routinely used in refractory cases (for example, solifenacin and tamsulosin) [van Kerrebroeck *et al.* 2013]. Combining mirabegron with α -adrenergic blockers is likely to be considered in developing the therapeutic area. Efficacy assessments for such combinations have recently been reported in men receiving mirabegron with tamsulosin (0.2 mg), in whom the OAB Symptom Score fell by -2.21 for the combination group and by -0.87 in the tamsulosin monotherapy group [Ichihara *et al.* 2015]. The safety aspects of combining mirabegron with tamsulosin have previously been reported in a separate study [van Gelderen *et al.* 2014].

Conclusion

Mirabegron is a first-in-class β_3 agonist for treatment of OAB in women or men, which appears to have good efficacy and tolerability. It can be used in patients who have discontinued antimuscarinic therapy, or who have contraindications for antimuscarinics. Development of combination therapies may further extend the management options in OAB in the future.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest statement

Marcus J. Drake: Advisory boards, research and speaker bureaux for Allergan, Astellas and Ferring. Monika Vij has no CoI.

References

- Abrams, P., Cardozo, L., Fall, M., Griffiths, D., Rosier, P., Ulmsten, U. *et al.* (2002) The

- standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. *Neurourol Urodyn* 21: 167–178.
- Abrams, P., Kelleher, C., Staskin, D., Griffiths, D., Rosier, P., Ulmsten, U. *et al.* (2014) Combination treatment with mirabegron and solifenacin in patients with overactive bladder: efficacy and safety results from a randomised, double-blind, dose-ranging, phase 2 study (Symphony). *Eur Urol* 67: 577–588.
- Aizawa, N., Igawa, Y., Nishizawa, O. and Wyndaele, J. (2010) Effects of CL316,243, a β_3 -adrenoceptor agonist, and intravesical prostaglandin E2 on the primary bladder afferent activity of the rat. *Neurourol Urodyn* 29: 771–776.
- Chapple, C., Amarenco, G., Lopez Aramburu, M., Everaert, K., Liehne, J., Lucas, M. *et al.* (2013a) A proof-of-concept study: mirabegron, a new therapy for overactive bladder. *Neurourol Urodyn* 32: 1116–1122.
- Chapple, C., Dvorak, V., Radziszewski, P., Van Kerrebroeck, P., Wyndaele, J., Bosman, B. *et al.* (2013b) A phase II dose-ranging study of mirabegron in patients with overactive bladder. *Int Urogynecol J* 24: 1447–1458.
- Chapple, C., Kaplan, S., Mitcheson, D., Klecka, J., Cummings, J., Drogendijk, T. *et al.* (2013c) Randomized double-blind, active-controlled phase 3 study to assess 12-month safety and efficacy of mirabegron, a $\beta(3)$ -adrenoceptor agonist, in overactive bladder. *Eur Urol* 63: 296–305.
- Chapple, C., Khullar, V., Gabriel, Z., Muston, D., Bitoun, C. and Weinstein, D. (2008) The effects of antimuscarinic treatments in overactive bladder: an update of a systematic review and meta-analysis. *Eur Urol* 54: 543–562.
- Chapple, C., Nitti, V., Khullar, V., Wyndaele, J., Herschorn, S., van Kerrebroeck, P. *et al.* (2014) Onset of action of the β_3 -adrenoceptor agonist, mirabegron, in Phase II and III clinical trials in patients with overactive bladder. *World J Urol* 32: 1565–1572.
- Coyne, K., Matza, L., Kopp, Z. and Abrams, P. (2006) The validation of the patient perception of bladder condition (PPBC): a single-item global measure for patients with overactive bladder. *Eur Urol* 49: 1079–1086.
- Coyne, K., Revicki, D., Hunt, T., Corey, R., Stewart, W., Bentkover, J. *et al.* (2002) Psychometric validation of an overactive bladder symptom and health-related quality of life questionnaire: the OAB-q. *Qual Life Res* 11: 563–574.
- Coyne, K., Sexton, C., Thompson, C., Milsom, I., Irwin, D., Kopp, Z. *et al.* (2009) The prevalence of lower urinary tract symptoms (LUTS) in the USA, the UK and Sweden: results from the Epidemiology of LUTS (EpiLUTS) study. *BJU Int* 104: 352–360.
- Coyne, K., Wein, A., Nicholson, S., Kvasz, M., Chen, C. and Milsom, I. (2014) Economic burden of urgency urinary incontinence in the United States: a systematic review. *J Manag Care Pharm* 20: 130–140.
- Dickinson, J., Lewand, M., Sawamoto, T., Krauwinkel, W., Schaddelee, M., Keirns, J. *et al.* (2013) Effect of renal or hepatic impairment on the pharmacokinetics of mirabegron. *Clin Drug Invest* 33: 11–23.
- Drake, M. (2014) Do we need a new definition of the overactive bladder syndrome? ICI-RS 2013. *Neurourol Urodyn* 33: 622–624.
- Herschorn, S., Barkin, J., Castro-Diaz, D., Frankel, J., Espuna-Pons, M., Gousse, A. *et al.* (2013) A phase III, randomized, double-blind, parallel-group, placebo-controlled, multicentre study to assess the efficacy and safety of the $\beta(3)$ adrenoceptor agonist, mirabegron, in patients with symptoms of overactive bladder. *Urology* 82: 313–320.
- Ichihara, K., Masumori, N., Fukuta, F., Tsukamoto, T., Iwasawa, A. and Tanaka, Y. (2015) A randomized controlled study of the efficacy of tamsulosin monotherapy and its combination with mirabegron for overactive bladder induced by benign prostatic obstruction. *J Urol* 193: 921–926.
- Igawa, Y., Aizawa, N. and Homma, Y. (2010) Beta3-adrenoceptor agonists: possible role in the treatment of overactive bladder. *Korean J Urol* 51: 811–818.
- Kanai, A., Wyndaele, J., Andersson, K., Fry, C., Ikeda, Y., Zabbarova, I. *et al.* (2011) Researching bladder afferents-determining the effects of $\beta(3)$ -adrenergic receptor agonists and botulinum toxin type-A. *Neurourol Urodyn* 30: 684–691.
- Kaplan, S., Roehrborn, C., Abrams, P., Chapple, C., Bavendam, T. and Guan, Z. (2011) Antimuscarinics for treatment of storage lower urinary tract symptoms in men: a systematic review. *Int J Clin Pract* 65: 487–507.
- Kashyap, M. and Tyagi, P. (2013) The pharmacokinetic evaluation of mirabegron as an overactive bladder therapy option. *Expert Opin Drug Metab Toxicol* 9: 617–627.
- Khullar, V., Amarenco, G., Angulo, J., Cambroner, J., Høye, K., Milsom, I. *et al.* (2013a) Efficacy and tolerability of mirabegron, a $\beta(3)$ -adrenoceptor agonist, in patients with overactive bladder: results from a randomised European-Australian phase 3 trial. *Eur Urol* 63: 283–295.
- Khullar, V., Cambroner, J., Angulo, J., Wooning, M., Blauwet, M., Dorrepaal, C. *et al.* (2013b) Efficacy of mirabegron in patients with and without prior antimuscarinic therapy for overactive bladder: a post

- hoc analysis of a randomized European-Australian Phase 3 trial. *BMC Urol* 13: 45.
- Krauwinkel, W., Dickinson, J., Schaddelee, M., Meijer, J., Tretter, R., van de Wetering, J. *et al.* (2014) The effect of mirabegron, a potent and selective β_3 -adrenoceptor agonist, on the pharmacokinetics of CYP2D6 substrates desipramine and metoprolol. *Eur J Drug Metab Pharmacokinet* 39: 43–52.
- Kullmann, F., Downs, T., Artim, D., Limberg, B., Shah, M., Contract, D. *et al.* (2011) Urothelial beta-3 adrenergic receptors in the rat bladder. *Neurourol Urodyn* 30: 144–150.
- Lee, J., Moy, S., Meijer, J., Krauwinkel, W., Sawamoto, T., Kerbusch, V. *et al.* (2013a) Role of cytochrome p450 isoenzymes 3A and 2D6 in the in vivo metabolism of mirabegron, a β_3 -adrenoceptor agonist. *Clin Drug Invest* 33: 429–440.
- Lee, J., Zhang, W., Moy, S., Krauwinkel, W., Sawamoto, T., Kerbusch, V. *et al.* (2013b) Effects of food intake on the pharmacokinetic properties of mirabegron oral controlled-absorption system: a single-dose, randomized, crossover study in healthy adults. *Clin Ther* 35: 333–341.
- Limberg, B., Andersson, K., Aura Kullmann, F., Burner, G., de Groat, W. and Rosenbaum, J. (2010) β -Adrenergic receptor subtype expression in myocyte and non-myocyte cells in human female bladder. *Cell Tissue Res* 342: 295–306.
- Lucas, M., Bosch, R., Burkhard, F., Cruz, F., Madden, T., Nambiar, A. *et al.* (2012) EAU Guidelines on Assessment and Nonsurgical Management of Urinary Incontinence. *Eur Urol* 62: 1130–1142.
- Maman, K., Aballea, S., Nazir, J., Desroziars, K., Neine, M., Siddiqui, E. *et al.* (2014) Comparative efficacy and safety of medical treatments for the management of overactive bladder: a systematic literature review and mixed treatment comparison. *Eur Urol* 65: 755–765.
- Milsom, I., Coyne, K., Nicholson, S., Kvasz, M., Chen, C. and Wein, A. (2014) Global prevalence and economic burden of urgency urinary incontinence: a systematic review. *Eur Urol* 65: 79–95.
- Moro, C., Tajouri, L. and Chess-Williams, R. (2013) Adrenoceptor function and expression in bladder urothelium and lamina propria. *Urology* 81: 211 e1–e7.
- Murakami, S., Chapple, C., Akino, H., Sellers, D. and Chess-Williams, R. (2007) The role of the urothelium in mediating bladder responses to isoprenaline. *BJU Int* 99: 669–673.
- Nitti, V., Auerbach, S., Martin, N., Calhoun, A., Lee, M. and Herschorn, S. (2013a) Results of a randomized phase III trial of mirabegron in patients with overactive bladder. *J Urol* 189: 1388–1395.
- Nitti, V., Khullar, V., van Kerrebroeck, P., Herschorn, S., Cambronero, J., Angulo, J. *et al.* (2013b) Mirabegron for the treatment of overactive bladder: a prespecified pooled efficacy analysis and pooled safety analysis of three randomised, double-blind, placebo-controlled, phase III studies. *Int J Clin Pract* 67: 619–632.
- Nitti, V., Rosenberg, S., Mitcheson, D., He, W., Fakhoury, A. and Martin, N. (2013c) Urodynamics and safety of the β_3 -adrenoceptor agonist mirabegron in males with lower urinary tract symptoms and bladder outlet obstruction. *J Urol* 190: 1320–1327.
- Novack, G., Lewis, R., Vogel, R., Sheth, N., Swearingen, D., Rasmussen, S. *et al.* (2013) Randomized, double-masked, placebo-controlled study to assess the ocular safety of mirabegron in healthy volunteers. *J Ocul Pharmacol Ther* 29: 674–680.
- Ochodnick, P., Uvelius, B., Andersson, K. and Michel, M. (2013) Autonomic nervous control of the urinary bladder. *Acta Physiol (Oxf)* 207: 16–33.
- Otsuka, A., Shinbo, H., Matsumoto, R., Kurita, Y. and Ozono, S. (2008) Expression and functional role of β -adrenoceptors in the human urinary bladder urothelium. *Naunyn Schmiedebergs Arch Pharmacol* 377: 473–481.
- Sadananda, P., Drake, M., Paton, J. and Pickering, A. (2013) A functional analysis of the influence of β_3 -adrenoceptors on the rat micturition cycle. *J Pharmacol Exp Ther* 347: 506–515.
- Takasu, T., Ukai, M., Sato, S., Matsui, T., Nagase, I., Maruyama, T. *et al.* (2007) Effect of (R)-2-(2-aminothiazol-4-yl)-4'-[2-[(2-hydroxy-2-phenylethyl) amino]ethyl] acetanilide (YM178), a novel selective beta3-adrenoceptor agonist, on bladder function. *J Pharmacol Exp Ther* 321: 642–647.
- van Gelderen, M., Tretter, R., Meijer, J., Dorrepaal, C., Gangaram-Panday, S., Brooks, A. *et al.* (2014) Absence of clinically relevant cardiovascular interaction upon add-on of mirabegron or tamsulosin to an established tamsulosin or mirabegron treatment in healthy middle-aged to elderly men. *Int J Clin Pharmacol Ther* 52: 693–701.
- van Kerrebroeck, P., Chapple, C., Drogendijk, T., Klaver, M., Sokol, R., Speakman, M. *et al.* (2013) Combination therapy with solifenacin and tamsulosin oral controlled absorption system in a single tablet for lower urinary tract symptoms in men: efficacy and safety results from the randomised controlled NEPTUNE trial. *Eur Urol* 64: 1003–1012.
- Wagg, A., Cardozo, L., Nitti, V., Castro-Diaz, D., Auerbach, S., Blauwet, M. *et al.* (2014) The

Visit SAGE journals online
<http://tau.sagepub.com>

SAGE journals

efficacy and tolerability of the β 3-adrenoceptor agonist mirabegron for the treatment of symptoms of overactive bladder in older patients. *Age Ageing* 43: 666–675.

Wagg, A., Compion, G., Fahey, A and Siddiqui, E. (2012) Persistence with prescribed antimuscarinic therapy for overactive bladder: a UK experience. *BJU Int* 110: 1767–1774.

Woods, M., Carson, N., Norton, N., Sheldon, J. and Argentieri, T. (2001) Efficacy of the β 3-adrenergic receptor agonist CL-316243 on experimental bladder hyperreflexia and detrusor instability in the rat. *J Urol* 166: 1142–1147.

Yamaguchi, O. and Chapple, C. (2007) β 3-adrenoceptors in urinary bladder. *Neurourol Urodyn* 26: 752–756.