Published in final edited form as:

J Rheumatol. 2015 September ; 42(9): 1685–1693. doi:10.3899/jrheum.150071.

No association between daily walking and structural changes in people at risk of or with mild knee osteoarthritis. Prospective data from the Multicenter Osteoarthritis Study

Britt Elin Øiestad, Emily Quinn, Daniel White, Frank Roemer, Ali Guermazi, Michael Nevitt, Neil A. Segal, Cora E. Lewis, and David T. Felson

Department of Orthopedics, Oslo University Hospital, Oslo, Norway

Clinical Epidemiology Research and Training Unit, Boston University School of Medicine, Boston, MA, US

Data Coordinating Center, Boston University School of Public Health, Boston, MA US

Quantitative Imaging Center, Department of Radiology, Boston University School of Medicine, Boston, US

University of California, San Francisco, San Francisco, CA, US

Abstract

Corresponding author address/ request for reprint to: Britt Elin Øiestad, Oslo and Akershus University College of Applied Sciences, P.O. Box 4 St. Olavs plass, N-0130 OSLO, Norway, Britt-Elin.Oiestad@hioa.no, Phone: 0047-92802089.

B.E. Øiestad, PT, PhD, Department of Orthopedics, Oslo University Hospital, Oslo, Norway

E. Quinn, MSc, Data Coordinating Center, Boston University School of Public Health, Boston, MA US

D. White, PT, ScD, MSc, Department of Physical Therapy, University of Delaware, Newark, DE, US and Clinical Epidemiology Research and Training Unit, Boston University School of Medicine, Boston, MA, US

F. Roemer, MD, PhD, Quantitative Imaging Center, Department of Radiology, Boston University School of Medicine, Boston, US and Department of Radiology, University of Erlangen-Nuremberg, Erlangen, Germany

A. Guermazi, MD, PhD, Quantitative Imaging Center, Department of Radiology, Boston University School of Medicine, Boston, US

M. Nevitt, PhD, University of California, San Francisco, San Francisco, CA, US

N. A. Segal, MD, MS, The University of Kansas, Kansas City, KS, US

C. E. Lewis, MD, MSPH, Division of Preventive Medicine, University of Alabama at Birmingham, Birmingham, AL, US

D. T. Felson, MD, MPH, Clinical Epidemiology Research and Training Unit, Boston University School of Medicine, Boston, MA, US and Arthritis Research UK Epidemiology Unit, University of Manchester, Manchester, UK

Author contribution

Britt Elin Øiestad contributed with paper idea, analysis and interpretation of data, and has written the manuscript.

Emily Quinn contributed with analysis and interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

Daniel White contributed with acquisition of data, interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

Frank Roemer contributed with acquisition of data, interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

Ali Guermazi contributed with acquisition of data, interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

Michael Nevitt contributed with acquisition of data, interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

Neil Segal contributed with interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

Cora E. Lewis contributed with acquisition of data, interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

David Felson contributed with paper idea, acquisition of data, analysis and interpretation of data, revising the manuscript critically for important intellectual content, and approved the version to be submitted.

We have no financial conflicts of interests

Objective—We investigated the association between objectively measured daily walking and knee structural change, defined either as radiographic worsening or as cartilage loss, in people at risk of or with knee osteoarthritis (OA).

Methods—Participants from the Multicenter Osteoarthritis (MOST) study with Kellgren and Lawrence (KL) grades 0–2 and daily walking (measured with the StepWatch) at the 60-month visit, were included. Participants had fixed flexion weight bearing radiographs and knee magnetic resonance images (MRIs) at 60 and 84 months. Radiographic worsening was read in both knees using the OARSI grading, and MRIs were read for one knee using WOMBS semiquantitative scoring. Odds ratios (OR) and 95% confidence intervals (CI) were calculated comparing those in the middle tertile against the lowest and highest tertiles of daily walking using logistic regression models and generalized estimating equations. Data on walking with moderate to vigorous intensity (minutes with >100 steps/min/day) was associated to structural change using multivariate and logistic regression models.

Results—The 1179 study participants (59% females) were 67.0 (± 7.6) years, with a mean ($\pm$ SD) body mass index of 29.8 (± 5.3) kg/m² who walked 6981 (± 2630) steps/day. After adjusting for confounders, we found no significant associations between daily walking and radiographic worsening or cartilage loss. More time spent walking at a moderate to vigorous intensity was not associated with either radiographic worsening or cartilage loss.

Conclusion—Results from the MOST study indicated no association between daily walking and structural changes over two years in people at risk of or with mild knee OA.

Key Indexing Terms

physical activity; osteoarthritis; structural changes; MRI

INTRODUCTION

The role of physical activity in the development and progression of knee osteoarthritis (OA) is controversial. Some studies suggest that physical activity at both high and low levels increase the risk of developing structural disease from repetitive compressive forces(1, 2). Others conclude that physical activity is protective against disease by promoting joint health(3–5).

One reason for the conflicting findings of prior studies examining physical activity and structural change in the knee joint may be the assumption of a unidirectional effect of physical activity on risk of structural change, rather than adverse effects of high and low levels of activity. Very low levels of walking may result in limited joint compression, which may provide insufficient stimulation of cartilage cellular and matrix components. For instance, a sedentary lifestyle in animal models has been shown to lower proteoglycan content and synovial fluid volume(6), which further led to cartilage fibrillation, ulceration, and erosion of the extracellular matrix(7). At the opposite extreme, high levels of walking may lead to detrimental mechanical stress on joint tissues. Excessive mechanical stress may shift the chondrocyte to more catabolic activity(7). Also, little is known about the risk of moderate to vigorous physical activity (MVPA) often characterized as walking at a frequency of >100 steps/minute(8, 9) on cartilage loss in people at risk of or with mild knee

OA(10). Walking is the most common type of physical activity for older adults(11), however, the association of walking with structural changes in the knee joint as seen on radiograph and MRI is not well understood.

The aim of the present study was to examine the association of objectively measured daily walking with structural change two years later in people at risk of or with mild knee OA. We hypothesized that individuals in the lowest and highest tertiles of daily walking would have a higher degree of radiographic worsening than those in the middle tertile. Furthermore, we hypothesized that individuals in the lowest and highest tertiles of daily walking would have a higher degree of cartilage loss, than people in the middle tertile. In addition, we hypothesized that no time (minutes) spent in walking at MVPA intensity (> 100 steps/minute) would be associated with a higher risk of structural change.

MATERIALS AND METHODS

The Multicenter Osteoarthritis (MOST) study is a prospective cohort study of men and women between 50–79 years of age at baseline, at risk of knee OA (i.e. overweight, obese, a history of knee injury, or with frequent knee pain), or with established knee OA. The study participants were from Birmingham, Alabama and Iowa City, Iowa in the US, and were contacted by advertisements and mass mailings of study brochures in the first step and screened by telephone for eligibility in the next step(12). The study started in 2003 and the study participants were interviewed by telephone and attended clinical visits at the time points zero (baseline), 30, 60, and 84 months (12, 13). The visits included examination of background variables such as age, height, weight, self-reported physical activity level, knee extensor muscle strength, self-reported physical function and pain, radiographic evaluation, and MRI.

For inclusion in our study, we required that participants attended the 60-month visit when daily walking was measured for the first time using a StepWatch, and the 84 month visit, the first follow-up visit after StepWatch data were collected. Subjects had to have radiographic and/or MRI data at the 60- and 84-month visits and at least 3 days of step data. Also, we included those who at the 60-month visit had knee radiographs that showed Kellgren and Lawrence (KL) grades < 3 in at least one knee. Study participants who had KL grade 3 or 4 or who had undergone total knee replacement at the 30- or 60-month visits were excluded. We have previously written that identifying risk factors for progressive OA among those with more advanced disease is formidably difficult due to collider selection bias and other methodological challenges(14, 15).

Assessments

Radiographic assessment—At the 60- and 84-month visits, standing posterior-anterior and lateral radiographs were taken using SynaFlexer frame for standardized positioning (Synarc Inc., San Francisco, CA). The radiographs were graded according to the KL classification system (grade 0–4)(16) (posteroanterior view), and the Osteoarthritis Research Society International (OARSI) atlas(17) (posteroanterior and lateral view) by a musculoskeletal radiologist and a rheumatologist at Boston University blinded to clinical and MRI data. Discrepancies were solved in a group of three readers (including DTF). Inter-

rater reliability results between $k=0.77$ – 0.80 have previously been published(18) for radiographic scoring of MOST data. Radiographic worsening in the tibiofemoral joint from 60 to 84 months was defined as any increase in OARSI grade, e.g. increase within grade (0.5 grade); or one or more grade increase(19, 20). The 60 and 84-month radiographs were read paired and compared for radiographic change(19). Full limb radiographs of both legs were obtained at the 60-month visit(18).

MRI assessment—MRIs were obtained of both knees at the 60- and 84-month visits using a 1.0 Tesla MRI system (OrthOne, GE Medical Systems, Wilmington, MA, USA), and read for one randomly selected knee per participant. A circumferential transmit-receive extremity coil was used. Sagittal and axial fat-suppressed fast spin-echo proton density-weighted sequences (repetition time=5800/2500 ms; time to echo=35 ms; slice thickness=3 mm; field of view=14 cm; matrix=288×192 pixels), and coronal Short Tau Inversion Recovery (STIR) sequence (repetition time=7820 ms, time to echo=15 ms, slice thickness=3 mm, field of view=14 cm, matrix=256×256 pixels) were used for the MRI examination(21). The MRIs were read paired by two experienced readers (AG, FWR) for one randomly selected knee for each participant using the Whole-Organ MRI Score (WORMS), including a total of 14 subregions(22). Cartilage loss was graded using WORMS from 0–6(22): 0= normal thickness and no signal; 1= normal thickness, but increased signal on proton density-weighted images; 2= partial thickness focal defect <1 cm in greatest width; 2.5= full thickness focal defect <1 cm in greatest width; 3= multiple areas of partial-thickness, intermixed with areas of normal thickness, or a grade 2.0 wider than 1 cm, but >75% of the region; 4=diffuse (>75% of the region) partial-thickness loss; 5= multiple areas of full-thickness loss or a grade 2.5 lesion wider than 1 cm, but <75% of the region; 6= diffuse (>75% of the region). Cartilage loss vs. no change between 60–84 months was defined on the basis of a comparison between the 60 and 84 months images of five subregions in the lateral and medial regions. The five medial and lateral subregions included femur central and posterior, and tibia anterior, central, and posterior. The change variables were defined as no change in the WORMS score, half-grade change, one grade change, and more than one grade change. We considered half-grade change or greater in any subregion as change between 60–84 months. BMLs and meniscal tears and extrusions at the 60-month visit were included as covariates in sensitivity analyses. BML was graded in volume based on the WORMS score (grade 0–3)(22, 23).

Assessment of daily walking—Daily walking steps and amount of minutes walking at MVPA were measured using a StepWatch Activity Monitor (Orthocare Innovations, Mountlake Terrace, WA) at the 60-month visit. The StepWatch is a small waterproof accelerometer attached to the ankle. MOST participants were given written and verbal instructions for attaching the monitor each morning and removing it at bedtime for 7 consecutive days. No feedback was given to the user on amount of steps they had walked during the day. The StepWatch had to be worn for 3 valid days, as it has been shown that 3 days provides an acceptable estimate of weekly activity using this monitor(24, 25). A valid day included wearing the StepWatch for 10 hours per day, but not necessarily continuously(26). The 10 hour requirement has been shown to represent more than 66% of walking hours(26). Time with no steps for 180 consecutive minutes did not count towards

the 10-hour requirement(26–28). Since the StepWatch captures steps from only one leg, we doubled the number of steps so that we report the number of steps taken by the person. The step data were converted as described by White et al(27): the number of steps were reduced by 25% for all the participants so that this number could be comparable to pedometer data that are reported in other studies(29). The data are reported as average steps/day and tertiles. Furthermore, we reported number of minutes per day with walking at MVPA(8). High test-retest reliability has been reported for the measurement of daily walking using the monitor(24).

Other variables—Data on weight and height, age, alignment, and history of knee injuries were collected at the 60-month visit. Calculation of body mass index (BMI) was based on weight/(height*height) in meter. Knee pain was examined using the Western Ontario McMaster Universities Osteoarthritis Index (WOMAC) pain subscale(30). The subscale ranges from 0–20 where 0 represents no pain and 20 represents extreme pain.

Statistical analyses

To examine the association of daily walking with structural change, we first classified the participants' daily steps into tertiles, which we named low, moderate, and high levels of daily walking. Next, we calculated the association of low, moderate (referent), and high levels of daily walking with radiographic worsening (coded as no change vs. any change) using logistic regression models, reporting odds ratios (ORs) and 95% confidence intervals (CIs). In addition, we analyzed the association between daily walking as a continuous variable and radiographic worsening. In separate models, we carried out similar analyses for medial and lateral radiographic worsening. We repeated the analyses with MVPA as the independent variable to assess for an association with radiographic worsening. Potential confounders included age, sex, BMI, WOMAC knee pain, KL grade, alignment, and history of knee injury at the 60-month visit. Generalized estimation equations (GEEs) were used to account for correlation between two knees within an individual(31).

Logistic regression models were used to estimate the risk of cartilage loss in medial and lateral tibiofemoral compartments, respectively, with low or high compared to moderate levels of daily walking, and with daily walking as a continuous variable. Our analyses of MRI data were specific to the 5 subregions in each compartment (e.g. for medial, one of them is central medial femur) and we used GEE to account for the correlations of change in these 5 subregions. We also carried out analyses using MVPA as a continuous variable, and with a cut off between those walking no minutes on MVPA vs. those walking any minutes of MVPA. The analysis on cartilage loss was not adjusted for cartilage morphology at 60 months, as this may introduce biased results(32).

Sensitivity analyses were conducted, including the primary analysis for radiographic worsening and cartilage loss, but with an additional interaction term between the 60-month cartilage morphology status and daily walking incorporated. We also adjusted for 60-month BML and meniscus status (tear and extrusion). Furthermore, we conducted analyses between cartilage loss and daily walking divided into < or 10000 steps/day. Sensitivity analyses were additionally performed on those without a history of knee injury at the 60-month visit,

due to possible residual confounding by knee injuries(33). Analyses were conducted to assess differences in sex, age, knee injury, BMI, WOMAC pain between those included in the radiographic and MRI models (with KL grade 0–2) and the remainder of the MOST cohort (with KL grade 0–2). Statistical significance was set for p-values < 0.05. Analyses were conducted using SAS and STATA intercooled 13.

RESULTS

Of 2898 knees (1687 people) with x-rays at the 60-month visit that were not end stage or already replaced, and 2011 knees (1233 people) with MRI's, 890 knees (510 people) were missing for JSN analyses and 1232 knees (1007 people) were missing for MRI analyses (Figure 1). The 1179 study participants included (2008 knees: 1019 left and 989 right knees) in the JSN analyses were on average ($\pm$ SD) 67.0 ($\pm$ 7.6) years (range: 55–84) and had a mean BMI of 29.8 ($\pm$ 5.3)(range: 18–57). There were 59% females. The mean WOMAC pain score was 1.9($\pm$ 2.5)(range: 0–15). Mean steps/day were 6981 ($\pm$ 2630)(min-max 774–19450), and the average minutes per day of MVPA were 8.8 ($\pm$ 11.2)(range: 0–80). Corresponding numbers for those 779 included in the analyses for cartilage loss were: mean age 67.0 ($\pm$ 7.6) years, mean BMI 29.5 ($\pm$ 4.7), 61% were women, with a mean steps/day of 7185 ($\pm$ 2565). Table 1 shows characteristics of study participants across knees divided into tertiles of steps/day. Sixteen percent (calculated from 2008 knees) had no minute of MVPA per day. At 60 months 51.6%, 20.5%, and 27.9% of knees had KL grade 0, 1, and 2, respectively. Distribution of cartilage loss between 60–84 months across tertiles of daily walking is given in Table 2.

Analyses showed no significant differences for those with KL grade 0–2 who were included in the regression model (n= 1179) vs. those with KL grades 0–2 who were not included (n=510) for BMI, sex, age, or knee injury. Those included in our analyses had significantly less WOMAC pain than those not included (1.9 vs. 2.9, $p<0.001$). Correspondingly, no differences were found for BMI, sex, or knee injury between those 779 who were included in the analyses on cartilage loss compared to the 454 who were not included. However, those not included were 1 year older ($p=0.03$), and had 0.6 points higher WOMAC pain score ($p=0.002$).

Associations between structural changes and daily walking

No significant associations were found between radiographic worsening and levels of daily walking (Table 3). Furthermore, no significant associations were found between medial radiographic worsening and low (adjusted OR 0.86, 95% CI 0.59, 1.27) and high (adjusted OR 1.21, 95% CI 0.82, 1.79), vs. moderate levels of daily walking, or lateral radiographic worsening and low (adjusted OR 1.10, 95% CI 0.66, 1.83) and high (adjusted OR 0.62, 95% CI 0.33, 1.16) vs. moderate level of daily walking. Furthermore, no significant associations were found between walking with MVPA and radiographic worsening (Table 3).

Significantly increased odds for cartilage loss in the medial compartment was seen for the lowest level of daily walking (<6078 steps/day) vs. those in the moderate level (6078–7938 steps/day) for unadjusted analysis (OR 1.70 95% OR 1.06, 2.73), but not for adjusted analyses. Figure 2 shows the adjusted association between tertiles of daily walking and

proportion of medial cartilage loss. Analyzing daily walking as a continuous variable did not reveal a significant relationship with cartilage loss, and likewise, comparing those walking 10000 vs. <10000 steps/day did not reveal a statistically association in the adjusted model. No significant interactions were detected for daily walking and cartilage morphology status at the 60-month visit for either lateral ($p=0.106$) or medial ($p=0.274$) compartments. No significant association was found between walking at MVPA and medial cartilage loss (Figure 3) or lateral cartilage loss. Furthermore, omitting those with a history of knee injury at the 60-month visit did not change the relationship between daily walking and medial or lateral cartilage loss.

DISCUSSION

The results of this study did not confirm our hypotheses that those in the lowest and highest levels of daily walking had higher risk of structural changes compared to those with a moderate level of walking. Furthermore, we did not find an association of walking with MVPA and structural changes over two years.

Even though not directly comparable, our results are in contrast to the literature suggesting associations between physical activity and structural changes in the knee joint(2, 34, 35). Animal studies(36–38), as well as studies on knee injured patients(39, 40), and population-based cohort studies have suggested that high activity level may increase the risk of development and progression of knee OA(1), although conflicting results have been reported(41, 42). No universal definitions of high, moderate, and low activity level exist for walking, even though suggestions exist(43). The MOST cohort does not represent a population with high activity level, such as those in which an association of activity and structural worsening have been found (>10000 steps/day)(2), and our tertiles may not represent the low, medium and high activity levels for other populations. Tudor-Locke et al. (43) have recommended that <5000 steps/day reflect a “sedentary lifestyle index”, 5000–7499 steps/day reflects “low activity”, 7500–9999 reflects “somewhat active”, and $>10\ 000$ reflects a “highly active” level. The MOST data showed that the participants walked on average 6981 (± 2630)(radiographic group) and 7185(± 2565)(MRI group) steps/day, with tertiles lower than the suggested classifications, but with mean values similar to the recommended 7000 steps in the American College of Sports Medicine Position Stand(44). Our third tertile walked >7950 steps/day, reflecting “somewhat active” according to Tudor-Locke et al’s suggestion. Furthermore, 16% of the study participants walked less than 1 minute with MVPA, indicating a general low activity level in the MOST cohort(45).

Dore et al(2) followed community-dwelling adults over 60 years of age for 2.7 years with measurements of daily walking at baseline using a pedometer and MRI findings both at baseline and after 2.7 years. They found that walking more than 10 000 steps/day was associated with increase in BMLs, meniscal pathology, and cartilage defect score. Furthermore, they found an interaction effect between daily walking and cartilage volume, indicating that activity level was protective against volume loss in individuals with more baseline cartilage volume, but led to increased cartilage loss in those with less baseline cartilage volume. The authors concluded that individuals with knee abnormalities should not walk more than 10 000 steps/day. Dore et al. did not exclude those with moderate and severe

radiographic OA and reported a prevalence of OA of 57% as compared to about 28% mild radiographic OA in our cohort. This may be the main reason for the different results across these two studies, in addition to the number of people walking >10000 steps/day. In another study, Lin et al.(1) investigated physical activity in relation to change in T2 values in cartilage over 4 years using data from the Osteoarthritis Initiative. They found that both high and very low activity scores as measured with the Physical Activity for the Elderly (PASE) Scale were associated with greater progression of T2 relaxation time in asymptomatic people without radiographic OA. They did not control for knee injury in their analyses, but according to their results and like ours, knee injury did not differ between tertiles of activity level. Currently, it is not known if progression in T2 values leads to cartilage morphology changes. Our analyses showed that a history of knee injury measured at the 60-month visit was significantly associated to medial cartilage loss (results not shown). Thus, knee injury seems to be more important for medial structural worsening than activity level, at least when average activity level is relatively low as seen in the MOST cohort.

Our data showed no relation between structural change and minutes per day of walking with MVPA, but this is limited by a narrow range of step frequency in our subjects. Future studies should examine whether prolonged bouts of activity intensity as would be encouraged in exercise programs might influence joint structures.

Our study had weaknesses: Only one time point of physical activity measurement may not be optimal to assess daily walking levels as predictors of structural change. However, activity and step monitors have shown to be valid and reliable in measuring peoples' physical activity level(24). StepWatch recorded steps/day has high concurrent validity in comparison with several reference standard measures of step frequency(46, 47), and is 96% accurate with counting steps in older adults(48). Furthermore, following a prospective cohort of people over many years, high dropout rates may be expected. After selection of study participants according to our inclusion and exclusion criteria, we still included as many as 1179 people (2008 knees) for the radiographic worsening analyses and 779 people for the MRI analyses. Due to the challenge with studying risk factors for progression of osteoarthritis in knees with advanced disease, we narrowed our sample to people at risk of developing or with mild OA. Studying cartilage loss in this population with no or mild OA using MRI with 1.0T magnet may not be optimal compared to newer MRI techniques(49). Finally, even though our data included a substantial number of knees with worsening as seen in Table 3, a longer follow-up would have been more ideal to detect radiographic structural changes in the knee joint.

Results from the MOST study indicated no association between daily walking and structural changes over two years in people at risk of or with mild knee OA.

Acknowledgments

Sources of support in the form of grants or industrial support

MOST data are funded by NIH and NIA grants (AG18820, AG18832, AG18947, and AG19069). The Research Council of Norway funds a postdoctoral position for the corresponding author.

Reference List

1. Lin W, Alizai H, Joseph GB, Srikkum W, Nevitt MC, Lynch JA, et al. Physical activity in relation to knee cartilage T2 progression measured with 3 T MRI over a period of 4 years: data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage*. 2013; 21:1558–1566. [PubMed: 23831632]
2. Dore DA, Winzenberg TM, Ding C, Otahal P, Pelletier JP, Martel-Pelletier J, et al. The association between objectively measured physical activity and knee structural change using MRI. *Ann Rheum Dis*. 2013; 72:1170–1175. [PubMed: 22896739]
3. Jones G, Schultz MG, Dore D. Physical activity and osteoarthritis of the knee: can MRI scans shed more light on this issue? *Phys Sportsmed*. 2011; 39:55–61. [PubMed: 22030941]
4. Martin KR, Kuh D, Harris TB, Guralnik JM, Coggon D, Wills AK. Body mass index, occupational activity, and leisure-time physical activity: an exploration of risk factors and modifiers for knee osteoarthritis in the 1946 British birth cohort. *BMC Musculoskelet Disord*. 2013; 14:219. [PubMed: 23883324]
5. Richmond SA, Fukuchi RK, Ezzat A, Schneider K, Schneider G, Emery CA. Are joint injury, sport activity, physical activity, obesity, or occupational activities predictors for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther*. 2013; 43:515-B19. [PubMed: 23756344]
6. Otterness IG, Eskra JD, Bliven ML, Shay AK, Pelletier JP, Milici AJ. Exercise protects against articular cartilage degeneration in the hamster. *Arthritis Rheum*. 1998; 41:2068–2076. [PubMed: 9811063]
7. Sun HB. Mechanical loading, cartilage degradation, and arthritis. *Ann N Y Acad Sci*. 2010; 1211:37–50. [PubMed: 21062294]
8. Marshall SJ, Levy SS, Tudor-Locke CE, Kolkhorst FW, Wooten KM, Ji M, et al. Translating physical activity recommendations into a pedometer-based step goal: 3000 steps in 30 minutes. *Am J Prev Med*. 2009; 36:410–415. [PubMed: 19362695]
9. Tudor-Locke C, Sisson SB, Collova T, Lee SM, Swan PD. Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population. *Can J Appl Physiol*. 2005; 30:666–676. [PubMed: 16485518]
10. Miller RH, Edwards WB, Brandon SC, Morton AM, Deluzio KJ. Why don't most runners get knee osteoarthritis? A case for per-unit-distance loads. *Med Sci Sports Exerc*. 2014; 46:572–579. [PubMed: 24042311]
11. Yusuf HR, Croft JB, Giles WH, Anda RF, Casper ML, Caspersen CJ, et al. Leisure-time physical activity among older adults. United States, 1990. *Arch Intern Med*. 1996; 24:1321–1326. [PubMed: 8651841]
12. Felson DT, Niu J, Guermazi A, Roemer F, Aliabadi P, Clancy M, et al. Correlation of the development of knee pain with enlarging bone marrow lesions on magnetic resonance imaging. *Arthritis Rheum*. 2007; 56:2986–2992. [PubMed: 17763427]
13. Segal NA, Nevitt MC, Gross KD, Hietpas J, Glass NA, Lewis CE, et al. The Multicenter Osteoarthritis Study: opportunities for rehabilitation research. *PM R*. 2013; 5:647–654. [PubMed: 23953013]
14. Zhang Y, Neogi T, Hunter D, Roemer F, Niu J. What effect is really being measured? An alternative explanation of paradoxical phenomena in studies of osteoarthritis progression. *Arthritis Care Res (Hoboken)*. 2014; 66:658–661. [PubMed: 24877249]
15. Zhang Y, Niu J, Felson DT, Choi HK, Nevitt M, Neogi T. Methodologic challenges in studying risk factors for progression of knee osteoarthritis. *Arthritis Care Res (Hoboken)*. 2010; 62:1527–1532. [PubMed: 20617531]
16. Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis*. 1957; 16:494–502. [PubMed: 13498604]
17. Altman RD, Gold GE. Atlas of individual radiographic features in osteoarthritis, revised. *Osteoarthritis Cartilage*. 2007; 15(Suppl A):1–56. [PubMed: 16891130]
18. Englund M, Felson DT, Guermazi A, Roemer FW, Wang K, Crema MD, et al. Risk factors for medial meniscal pathology on knee MRI in older US adults: a multicentre prospective cohort study. *Ann Rheum Dis*. 2011; 70:1733–1739. [PubMed: 21646417]

19. Felson DT, Nevitt MC, Yang M, Clancy M, Niu J, Torner JC, et al. A new approach yields high rates of radiographic progression in knee osteoarthritis. *J Rheumatol.* 2008; 35:2047–2054. [PubMed: 18793000]
20. Englund M, Guermazi A, Roemer FW, Aliabadi P, Yang M, Lewis CE, et al. Meniscal tear in knees without surgery and the development of radiographic osteoarthritis among middle-aged and elderly persons: The Multicenter Osteoarthritis Study. *Arthritis Rheum.* 2009; 60:831–839. [PubMed: 19248082]
21. Roemer FW, Guermazi A, Lynch JA, Peterfy CG, Nevitt MC, Webb N, et al. Short tau inversion recovery and proton density-weighted fat suppressed sequences for the evaluation of osteoarthritis of the knee with a 1.0 T dedicated extremity MRI: development of a time-efficient sequence protocol. *Eur Radiol.* 2005; 15:978–987. [PubMed: 15633060]
22. Peterfy CG, Guermazi A, Zaim S, Tirman PF, Miaux Y, White D, et al. Whole-Organ Magnetic Resonance Imaging Score (WORMS) of the knee in osteoarthritis. *Osteoarthritis Cartilage.* 2004; 12:177–190. [PubMed: 14972335]
23. Roemer FW, Guermazi A, Javaid MK, Lynch JA, Niu J, Zhang Y, et al. Change in MRI-detected subchondral bone marrow lesions is associated with cartilage loss: the MOST Study. A longitudinal multicentre study of knee osteoarthritis. *Ann Rheum Dis.* 2009; 68:1461–1465. [PubMed: 18829615]
24. Mudge S, Taylor D, Chang O, Wong R. Test-retest reliability of the StepWatch Activity Monitor outputs in healthy adults. *J Phys Act Health.* 2010; 7:671–676. [PubMed: 20864764]
25. Mudge S, Stott NS. Test–retest reliability of the StepWatch Activity Monitor outputs in individuals with chronic stroke. *Clin Rehabil.* 2008; 22:871–877. [PubMed: 18955419]
26. Song J, Semanik P, Sharma L, Chang RW, Hochberg MC, Mysiw WJ, et al. Assessing physical activity in persons with knee osteoarthritis using accelerometers: data from the osteoarthritis initiative. *Arthritis Care Res (Hoboken).* 2010; 62:1724–1732. [PubMed: 20806273]
27. White DK, Tudor-Locke C, Felson DT, Gross KD, Niu J, Nevitt M, et al. Walking to meet physical activity guidelines in knee osteoarthritis: is 10,000 steps enough? *Arch Phys Med Rehabil.* 2013; 94:711–717. [PubMed: 23228625]
28. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc.* 2008; 40:181–188. [PubMed: 18091006]
29. Feito Y, Bassett DR, Thompson DL, Tjo BM. Effects of body mass index on step count accuracy of physical activity monitors. *J Phys Act Health.* 2012; 9:594–600. [PubMed: 21946229]
30. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol.* 1988; 15:1833–1840. [PubMed: 3068365]
31. Zhang Y, Zhang B, Wise B, Niu J, Zhu Y. Statistical approaches to evaluating the effect of risk factors on the pain of knee osteoarthritis in longitudinal studies. *Curr Opin Rheumatol.* 2009; 21:513–519. [PubMed: 19584728]
32. Glymour MM, Weuve J, Berkman LF, Kawachi I, Robins JM. When is baseline adjustment useful in analyses of change? An example with education and cognitive change. *Am J Epidemiol.* 2005; 1:267–278. [PubMed: 15987729]
33. Urquhart DM, Wluka AE, Teichtahl AJ, Cicuttini FM. The effect of physical activity on the knee joint: is it good or bad? *Br J Sports Med.* 2007; 41:546–547. [PubMed: 17726198]
34. Stehling C, Lane NE, Nevitt MC, Lynch J, McCulloch CE, Link TM. Subjects with higher physical activity levels have more severe focal knee lesions diagnosed with 3T MRI: analysis of a non-symptomatic cohort of the osteoarthritis initiative. *Osteoarthritis Cartilage.* 2010; 18:776–786. [PubMed: 20202488]
35. Wang Y, Simpson JA, Wluka AE, Teichtahl AJ, English DR, Giles GG, et al. Is physical activity a risk factor for primary knee or hip replacement due to osteoarthritis? A prospective cohort study. *J Rheumatol.* 2011; 38:350–357. [PubMed: 20952471]

36. Julkunen P, Halmesmaki EP, Iivarinen J, Rieppo L, Narhi T, Marjanen J, et al. Effects of growth and exercise on composition, structural maturation and appearance of osteoarthritis in articular cartilage of hamsters. *J Anat.* 2010; 217:262–274. [PubMed: 20646109]
37. Beckett J, Jin W, Schultz M, Chen A, Tolbert D, Moed BR, et al. Excessive running induces cartilage degeneration in knee joints and alters gait of rats. *J Orthop Res.* 2012; 30:1604–1610. [PubMed: 22508407]
38. Franciozi CE, Tarini VA, Reginato RD, Goncalves PR, Medeiros VP, Ferretti M, et al. Gradual strenuous running regimen predisposes to osteoarthritis due to cartilage cell death and altered levels of glycosaminoglycans. *Osteoarthritis Cartilage.* 2013; 21:965–972. [PubMed: 23602983]
39. Øiestad BE, Engebretsen L, Storheim K, Risberg MA. Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. *Am J Sports Med.* 2009; 37:1434–1443. [PubMed: 19567666]
40. Lohmander LS, Englund PM, Dahl LL, Roos EM. The Long-term Consequence of Anterior Cruciate Ligament and Meniscus Injuries: Osteoarthritis. *Am J Sports Med.* 2007; 29:1756–1769. [PubMed: 17761605]
41. Hansen P, English M, Willick SE. Does running cause osteoarthritis in the hip or knee? *PM R.* 2012; 4:S117–S121. [PubMed: 22632690]
42. Barbour KE, Hootman JM, Helmick CG, Murphy LB, Theis KA, Schwartz TA, et al. Meeting physical activity guidelines and the risk of incident knee osteoarthritis: a population-based prospective cohort study. *Arthritis Care Res (Hoboken).* 2014; 66:139–146. [PubMed: 23983187]
43. Tudor-Locke C, Bassett DR, Swartz AM, Strath SJ, Parr BB, Reis JP, et al. A preliminary study of one year of pedometer self-monitoring. *Ann Behav Med.* 2004; 28:158–162. [PubMed: 15576253]
44. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011; 43:1334–1359. [PubMed: 21694556]
45. White DK, Tudor-Locke C, Felson DT, Gross KD, Niu J, Nevitt M, et al. Do radiographic disease and pain account for why people with or at high risk of knee osteoarthritis do not meet physical activity guidelines? *Arthritis Rheum.* 2013; 65:139–147. [PubMed: 23124774]
46. Foster RC, Lanningham-Foster LM, Manohar C, McCrady SK, Nysse LJ, Kaufman KR, et al. Precision and accuracy of an ankle-worn accelerometer-based pedometer in step counting and energy expenditure. *Prev Med.* 2005; 41:778–783. [PubMed: 16125760]
47. Karabulut M, Crouter SE, Bassett DR Jr. Comparison of two waist-mounted and two ankle-mounted electronic pedometers. *Eur J Appl Physiol.* 2005; 95:335–343. [PubMed: 16132120]
48. Resnick B, Nahm ES, Orwig D, Zimmerman SS, Magaziner J. Measurement of activity in older adults: reliability and validity of the Step Activity Monitor. *J Nurs Meas.* 2001; 9:275–290. [PubMed: 11881269]
49. Wang Y, Wluka AE, Jones G, Ding C, Cicuttini FM. Use magnetic resonance imaging to assess articular cartilage. *Ther Adv Musculoskelet Dis.* 2012; 4:77–97. [PubMed: 22870497]

Figure 1.
Flow chart of the study participants.

Figure 2.

The binscatter shows proportion of medial cartilage loss for tertiles of steps/day (dashed lines) adjusted for 60-month age, sex, BMI, knee injury, pain, radiographic grade, and alignment. Medial cartilage loss includes any change in the five medial subregions femur central and posterior, and tibia anterior, central, and posterior.

Figure 3.
Proportion of medial cartilage loss given for moderate or greater intensity.

Table 1

Characteristics of study participants at the 60-month visit (n=2008 knees)

	Tertile 1 (<5860 steps) N=670	Tertile 2 (5860–7846 steps) N=669	Tertile 3 (>7846 steps) N=669	p-value
Age (mean±sd)	69.7±7.8	66.5±7.3	64.0±6.4	<0.001
Body mass index (mean±sd)	31.6±5.7	29.6±4.8	27.8±4.2	<0.001
WOMAC pain (mean±sd)	2.5±3.1	1.7±2.5	1.5±2.2	<0.001
Mechanical alignment (mean±sd)	−1.04±2.9	−1.11±2.8	−1.37±2.7	n.s
MVPA (mean±sd)	2.6±3.3	7.3±7.3	17.2±14.8	<0.001
KL grade				
0	297 (44.3)	360 (53.8)	379 (56.7)	
1	149 (22.3)	122 (18.2)	141 (21.0)	
2	224 (33.4)	187 (28.0)	149 (22.3)	<0.001
Knee injury				
No	514 (76.7)	506 (75.6)	507 (75.8)	
Yes	156 (23.3)	163 (24.4)	162 (24.2)	n.s
OARSI grade change*				
No	540 (80.6)	560 (83.7)	579 (86.5)	
Yes	130 (19.4)	109 (16.3)	90 (13.5)	<0.05

Sd, standard deviation; WOMAC, Western Ontario McMaster Universities Arthritis Index; MVPA, minutes of moderate to vigorous intensity; KL, Kellgren and Lawrence OARSI, Osteoarthritis Research International; n.s, non-significant;

* Change between 60 to 84 months visits. Analyses for continuous variables are performed using Analysis of variance (ANOVA). Analyses for categorical variables are conducted using Chi square test.

Table 2

Number of knees with any cartilage loss (half a grade change) between 60 to 84 months by tertiles of daily walking (n=779).

	<6078 steps N=260 knees	6078–7938 steps N=260 knees	>7938 steps N=259 knees
LATERAL*			
Femur lateral central	13 (5.0)	15 (5.8)	12 (4.6)
Femur lateral posterior	6 (2.3)	9 (3.5)	6 (2.3)
Tibia lateral anterior	0 (0)	6 (2.3)	0 (0)
Tibia lateral central	29 (11.1)	19 (7.3)	14 (5.4)
Tibia lateral posterior	21 (8.1)	15 (5.8)	17 (6.6)
MEDIAL*			
Femur medial central	32 (12.3)	26 (10.0)	28 (10.8)
Femur medial posterior	20 (7.7)	15 (5.8)	9 (3.5)
Tibia medial anterior	4 (1.5)	1 (0.4)	5 (1.9)
Tibia medial central	35 (13.5)	12 (4.6)	25 (9.7)
Tibia medial posterior	13 (5.0)	5(1.9)	10(3.8)

* Lateral and medial femoral anterior compartments are omitted due to lack of loading during walking. Numbers indicate numbers of knees followed by % of 260 knees.

Table 3

The association between activity levels and structural worsening

TERTILES	N (%)** of knees with worsening	Crude OR (95% CI)	Adjusted OR (95% CI)
Radiographic worsening			
<5859 steps	130 (19.4)	1.22 (0.91, 1.65)	0.91 (0.64, 1.27)
5859–7846 steps	109 (16.3)	1.0 (referent)	1.0 (referent)
>7846 steps	90 (13.5)	0.79 (0.57, 1.09)	0.99 (0.69, 1.42)
Intensity (minutes)		0.99 (0.98, 1.00)	1.01 (0.99, 1.02)
Lateral cartilage loss*			
<6078 steps	69 (26.5)	0.94 (0.54, 1.63)	0.82 (0.45, 1.51)
6078–7938 steps	64 (24.6)	1.0 (referent)	1.0 (referent)
>7938 steps	49 (18.9)	0.70 (0.39, 1.28)	0.74 (0.43, 1.28)
MVPA (minutes)		0.99 (0.97, 1.01)	1.00 (0.98, 1.02)
Medial cartilage loss*			
<6078 steps	104 (40.0)	1.70 (1.06, 2.73)	1.37 (0.81, 2.32)
6078–7938 steps	59 (22.7)	1.0 (referent)	1.0 (referent)
>7938 steps	77 (29.7)	1.23 (0.75, 2.03)	1.37 (0.80, 2.33)
MVPA (minutes)		0.98 (0.97, 1.00)	0.99 (0.97, 1.01)

OR, odds ratio; CI, confidence intervals; MVPA, moderate to vigorous intensity. Bold numbers indicate $p < 0.05$.

* Increase in WOMS score between 60–84 months.

** Percentage of number in the tertiles ($n=669$ knees for JSN, and $n=260$ people for cartilage loss). The analyses are adjusted for the following confounders measured at the 60-month visit: age, sex, BMI, WOMAC knee pain, KL grade, alignment, and history of knee injury.