CORRESPONDENCE

Prevalence Trends in Lifestyle-Related Risk Factors: Two Cross-Sectional Analyses With a Total of 8728 Participants From the Study of Health in Pomerania From 1997 to 2001 and 2008 to 2012

by Prof. Dr. med. Henry Völzke et al. in issue 11/2015

Validated Questionnaire Should Be Used

The decrease by almost 40% in physical inactivity in their leisure time in 70–81 year olds in Mecklenburg-Western Pomerania in the 10-year comparison is to be welcomed (1). On reading the report of this very interesting population-based study, I was not quite clear whether the data on the very complex and multi-dimensional behavior of habitual physical activity had been collected by using a single item question or the Baecke questionnaire (2)—as an earlier publication by the authors would lead me to believe (3). For this reason I would ask the authors to clarify this briefly. I would like to add that modified versions of Baecke's original questionnaire have shown greater test-retest reliability in older study populations. In future it would be desirable, depending on the situation and the cases, if the operationalizing instruments were to consider subjective expectations of benefit and needs in relation to physical activity more strongly.

DOI: 10.3238/arztebl.2015.0563a

REFERENCES

1. Völzke H, Ittermann T, Schmidt CO, et al.: Prevalence trends in lifestyle-related risk factors—two cross-sectional analyses with a total of 8728 participants from the Study of Health in Pomerania from 1997 to 2001 and 2008 to 2012. *Dtsch Arztebl Int* 2015; 112: 185–92.
2. Baecke JA, Burema J, Frijters JE: A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982; 36: 936–42.
3. Völzke H, Alte D, Schmidt CO, et al.: Cohort profile: the study of health in Pomerania. *Int J Epidemiol* 2011; 40: 294–307.

Dr. oec. troph. Martin Hofmeister
Verbraucherzentrale Bayern e.V.,
Referat Lebensmittel und Ernährung, München
hofmeister@vzbayern.de

Lack of Physical Activity

The article describes extremely interesting prevalence data from one of the most important national studies on cardiovascular risk factors. Most risk factors were defined in a sensible and differentiated manner (for example, diabetes), but questions remain with regard to the topic of physical inactivity.

Participants who exercise less than one hour per week are described as inactive. Persons who exercise for one hour per week therefore do not have this risk factor. This leaves out the remaining 167 hours of the week. More recent data have shown unequivocally that the extent of actual inactivity (time spent sitting down) is a risk factor that is independent of the extent of physical exercise (2). One hour of exercise is in no way sufficient

to balance this potential risk and does not conform to international recommendations.

For completeness's sake we wish to add that, in actual fact, the most important predictor for health risk is physical performance—this includes even subjective fitness, as experienced by individual subjects (less fit/same level of fitness/fitter than people of the same age and sex) and reported in the Copenhagen City Heart Study (3).

The classification relating to the risk factor physical inactivity in the present study is therefore not up to date, in spite of the undisputed merits of the program.

DOI: 10.3238/arztebl.2015.0563b

REFERENCES

1. Völzke H, Ittermann T, Schmidt CO, et al.: Prevalence trends in lifestyle-related risk factors—two cross-sectional analyses with a total of 8728 participants from the Study of Health in Pomerania from 1997 to 2001 and 2008 to 2012. *Dtsch Arztebl Int* 2015; 112: 185–92.
2. Biswas A, Oh PI, Faulkner GE, et al.: Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med* 2015; 162: 123–32.
3. Holtermann A, Marott JL, Gyntelberg F, et al.: Self-reported cardiorespiratory fitness: prediction and classification of risk of cardiovascular disease mortality and longevity—a prospective investigation in the Copenhagen City Heart Study. *J Am Heart Assoc* 2015; 4: e001495.

Prof. Dr. med. Burkhard Weisser
Christian-Albrechts-Universität zu Kiel
Sportmedizin
bweisser@email.uni-kiel.de

In Reply:

Hofmeister and Weisser comment on the results relating to the prevalence trend of physical inactivity and criticize the crude categorization, while using a simple question on exercise of at least one hour per week (1). As they rightly emphasize, Baecke's validated questionnaire has become the standard instrument for epidemiological studies (2). One limitation of the SHIP-0 study is the fact that this questionnaire was incompletely included in SHIP-0, and only the variables used in our article allow comparability with SHIP-Trend-0. Both authors' comments are therefore completely justified. In the studies subsequent to SHIP-0, Baecke's questionnaire was completely integrated.

DOI: 10.3238/arztebl.2015.0563c

REFERENCES

1. Völzke H, Ittermann T, Schmidt CO, et al.: Prevalence trends in lifestyle-related risk factors—two cross-sectional analyses with a total of 8728 participants from the Study of Health in Pomerania from 1997 to 2001 and 2008 to 2012. *Dtsch Arztebl Int* 2015; 112: 185–92.
2. Baecke JA, Burema J, Frijters JE: A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982; 36: 936–42.

Prof. Dr. med. Henry Völzke
Institut für Community Medicine, SHIP/ Klinisch-Epidemiologische Forschung
Universitätsmedizin Greifswald
voelzke@uni-greifswald.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.