Technical Section [TECHNICAL NOTES AND TIPS

A simple technique to improve venting of the femur

S McHale, R Yarlagadda

Plymouth Hospitals NHS Trust, UK

CORRESPONDENCE TO

Stephen McHale, E: stevemchale@hotmail.com

BACKGROUND

Prophylactic cephalomedullary nailing is a recommended technique for prophylactic fixation in femurs at risk of fracture due to metastatic bone disease.¹ During the procedure, raised intramedullary pressure can cause fat and neoplastic embolus, risking cardiovascular instability and death.² We describe a simple technique to improve venting of the femur using easily available equipment.

TECHNIQUE

At the beginning of the procedure, a stab incision is made and blunt dissection is performed to the lateral femur. The femur is opened with a 4.5mm drill bit under fluoroscopic guidance at least three cortical diameters proximal to the anticipated position of the tip of the nail. A 14Fr suction catheter (ConvaTec, Uxbridge, UK) is cut to a length of approximately 15cm and slid over a rigid guidewire as included on the nailing set. This is used to guide the tip of the suction catheter to the venting hole. The guidewire is then removed and suction connected.

Figure 1 The suction catheter is cut to an appropriate length and then threaded with the rigid guidewire.

DISCUSSION

This technique is simple to perform with easily available equipment and it may reduce intramedullary pressure from the start of the nailing procedure.

References

1. British Orthopaedic Association, British Orthopaedic Oncology Society. *Metastatic Bone Disease: A Guide to Good Practice*. London: BOA; 2001.
2. Kröpfl A, Davies J, Berger U *et al*. Intramedullary pressure and bone marrow fat extravasation in reamed and unreamed femoral nailing. *J Orthop Res* 1999; **17**: 261–268.

Wound dressing following debridement for Fournier's gangrene

JM Henderson, NNS Patel

Oxford University Hospitals NHS Trust, UK

CORRESPONDENCE TO

John Henderson, E: john_henderson@doctors.net.uk

The aggressive debridement of necrotic tissue required in Fournier's gangrene often leaves a substantial skin defect around the scrotum and perineal area (Fig 1). These patients may return to theatre multi-

Figure 1 Skin defect following debridement

ple times for 'relook' surgery and further debridement. Temporarily covering the open wound in the perineal area pending definitive reconstruction presents a challenge and faecal contamination is a risk. We demonstrate a technique using lengths of povidone-iodine soaked 90cm × 5m gauze roll (Synergy Health, Swindon, UK). Each testis is wrapped individually before a length of gauze is folded across the site 3–4 times and secured to the wound edges with skin staples (Figs 2–4).

Figure 2 Each testis is wrapped individually.

Figure 4 The dressing is secured with staples.

A consistent and reliable technique for tensioning a suture button syndesmosis fixation system

T Koç¹, N Cullen², HP Taylor¹

¹Poole Hospital NHS Foundation Trust, UK

²Royal National Orthopaedic Hospital Trust, UK

CORRESPONDENCE TO

Togay Koç, E: togay.koc@doctors.org.uk

The authors declare they have no conflicts of interest.

BACKGROUND

The TightRope® syndesmosis fixation system (Arthrex, Naples, FL, US) is a suture-button device for the dynamic stabilisation of the distal tibiofibular syndesmosis.¹ This implant avoids the need for routine removal.² It is important to tighten the suture sufficiently in order to maintain reduction of the syndesmosis. We describe the use of a traditional technique for tightening sutures applied to this novel fixation system.

TECHNIQUE

The distal fibula is exposed and the distal tibiofibular syndesmosis is reduced according to the surgeon's preference. The cortices are drilled and the TightRope® is passed as per the implant's surgical technique instructions.³ With the buttons flush to the cortices and/or plate and the foot in a natural plantarflexed position, a standard knot is prepared (Fig 1). This is tightened over a closed pair of Mayo scissors held by an assistant (Fig 2). As the surgeon maintains tension on the knot, the assistant opens and closes the scissors (Fig 3) 3–4 times to tighten the suture before slipping the scissors out, thereby allowing the

Figure 3 A length of gauze roll is folded across the wound 3–4 times.