Published in final edited form as:

Child Care Health Dev. 2014 September ; 40(5): 706–714.

Paternal ADHD Symptoms and Child Conduct Problems: Is Father Involvement Always Beneficial?

Abigail Mintz Romirowsky, Ph.D. and

graduate of the Clinical Psychology Program at the University of Maryland, College Park and is a postdoctoral fellow at Children's National Medical Center. amintz@cnmc.org, 401-374-8349, 1402 Swann St NW #3, Washington DC 20009, United States

Andrea Chronis-Tuscano, Ph.D.

Associate Professor of Psychology at the University of Maryland, College Park. Dr. Chronis-Tuscano directs the Maryland Attention-Deficit/Hyperactivity Disorder (ADHD) Program, and she is also an Adjunct Professor of Pediatrics at the George Washington University School of Medicine, Children's National Medical Center. achronis@umd.edu, 301-405-9640, 1123K Biology/Psychology Building College Park, Maryland 20742, United States

Abstract

Background—Maternal psychopathology robustly predicts poor developmental and treatment outcomes for children with attention-deficit/hyperactivity disorder (ADHD). Despite the high heritability of ADHD, few studies have examined associations between paternal ADHD symptoms and child adjustment, and none have also considered degree of paternal involvement in childrearing. Identification of modifiable risk factors for child conduct problems is particularly important in this population given the serious adverse outcomes resulting from this comorbidity.

Methods—This cross-sectional study examined the extent to which paternal involvement in childrearing moderated the association between paternal ADHD symptoms and child conduct problems among 37 children with ADHD and their biological fathers.

Results—Neither paternal ADHD symptoms nor involvement was independently associated with child conduct problems. However, the interaction between paternal ADHD symptoms and involvement was significant, such that paternal ADHD symptoms were positively associated with child conduct problems only when fathers were highly involved in childrearing.

Conclusions—The presence of adult ADHD symptoms may determine whether father involvement in childrearing has a positive or detrimental influence on comorbid child conduct problems.

Keywords

paternal; ADHD; conduct problems; involvement; parenting

Introduction

In the United States, 3-7% of school-age children are diagnosed with attention-deficit/hyperactivity disorder (ADHD; American Psychiatric Association, 2000). Children with ADHD display developmentally-aberrant symptoms of hyperactivity, impulsivity, and/or

inattention with associated functional impairment across important life areas (e.g., family conflict, academic difficulty, poor social relationships; Coie & Dodge, 1998; Mikami & Pfiffner, 2008). ADHD is also highly comorbid with conduct disorder (CD; 2.6-17.1%; Bird, Gould, & Staghezza-Jaramillo, 1994; Romano, Tremblay, Vitarro, Zoccolillo, Pagani, 2005) and oppositional defiant disorder (ODD; 36.5-65%; Biederman et al., 1996b; Yang, Wang, Qian, Biederman, & Faraone, 2004). Indeed, the presence of comorbid CD/ODD symptoms (collectively referred to as “conduct problems”) substantially increases the risk for very serious outcomes including substance abuse and criminal behavior (Elkins, McGue, & Iacono, 2007; Harty, Miller, Newcorn, & Halperin, 2009; Henry, Caspi, Moffitt, & Silva, 1996; Lahey et al., 1998; Lynam, 1998). Thus, identifying risk factors for conduct problems among children with ADHD is a research priority.

Maternal psychopathology is a robust risk factor for child conduct problems among children with ADHD (Chronis et al., 2007), yet the impact of paternal psychopathology remains relatively understudied. Because ADHD is highly heritable (Faraone et al., 2005), parental ADHD symptoms has emerged as a focus of intervention and research. Adults with ADHD experience the same core symptoms of inattention and hyperactivity/impulsivity as children (Faraone et al., 2000b), and these symptoms profoundly impact parenting ability (Johnston, Mash, Miller, & Ninowski, 2012). Specifically, parents with ADHD often display impulsive decision-making, emotional reactivity, reduced vigilance, and poor planning. Indeed, among families of children with ADHD, mothers and fathers with elevated ADHD symptoms themselves display more laxness, inconsistent discipline, poor parental monitoring, over reactivity, and coerciveness (Arnold, O’Leary, & Edwards, 1997; Chronis-Tuscano et al., 2008; Harvey, Danforth, McKee, Ulaszek, & Friedman, 2003; Murray & Johnston, 2006; Rogers, Wiener, Marton, & Tannock, 2009). Paternal childhood ADHD is also associated with child conduct problems in this population (Chronis et al., 2003); however, no studies have examined the relationship between paternal ADHD and child conduct problems.

Developmental research has identified paternal involvement as an important predictor of child psychosocial outcomes. For example, early involvement by both residential and non-residential fathers predicts lower levels of adolescent conduct problems and police contact (Coley & Medeiros, 2007; Flouri & Buchanan, 2002). Within families of children with ADHD, having a biological father present in the home is related to lower levels of child conduct problems (Pfiffner, McBurnett, & Rathouz, 2001).

What remains unclear, however, is the impact of paternal involvement on child behavior when fathers have psychopathology (e.g., ADHD). Considering that fathers of children with ADHD often report maladaptive involvement with their children (Arnold et al., 1997; Harvey et al., 2003; Rogers et al., 2009) and are 4-5 times more likely to have ADHD themselves (Chronis et al., 2003), research is needed to understand the relations between paternal ADHD symptoms, paternal involvement, and child conduct problems. To our knowledge, the current study is the first to examine whether paternal involvement moderates the relationship between paternal ADHD symptoms and concurrent child conduct problems.

Method

Participants

The current study sample (N=37) was recruited from a larger sample of mothers who participated in three prior studies conducted in our laboratory (N=195; Chronis-Tuscano et al., 2013; Chronis-Tuscano et al., 2011; Raggi, Chronis-Tuscano, Fishbein, & Groomes, 2009). Mothers were originally recruited via mailings to local ADHD groups, schools, and health providers in the Washington, DC metropolitan area. Across these original studies, children met Diagnostic and Statistical Manual of Mental Disorders (4th ed.; *DSM-IV*; American Psychiatric Association, 1994) ADHD criteria, were between the ages of 6–12 years old, and had an estimated IQ of at least 70. Mothers who agreed to be informed of future research opportunities were recontacted to obtain permission to contact the child's father to elicit his participation in the current study. Because ADHD is highly heritable (Faraone et al., 2005; Thapar, Langley, Owen, & O'Donovan, 2007), we included only biological fathers. Also, because parents of children with ADHD are more likely to be divorced or unmarried (Wymbs et al., 2008), we made considerable attempts to recruit non-residential biological fathers. See Figure 1 for a complete consort diagram.

Families who participated in the current study had a higher mean family income ($t(202) = -2.19, p < .05$) and fewer non-residential fathers ($\chi^2(1, N = 202) = 10.26, p < .01$) than the larger sample from which we recruited. However, no differences between the full sample and this subsample emerged for maternal education, child conduct problems, and percentage of fathers with a history of psychiatric diagnosis or treatment. Considerable variability was observed in child conduct problems (i.e., 48.6 % of children with ODD and 8.1% with CD) and paternal ADHD (i.e., 24% of fathers with childhood ADHD and 13% with adult ADHD). Additionally, our sample shows considerable racial/ethnic diversity. Sample characteristics are presented in Table 1.

Procedure

Measures of child psychopathology and impairment were completed by mothers and teachers at the time of their participation in the original studies. During the current study, measures of paternal psychopathology and involvement were completed by fathers via mailed questionnaires and a structured telephone interview (Figure 1). Considering the difficulty of including fathers in clinical research (Phares, 1992), a very brief battery of measures was selected, and no travel to the university was required (Mitchell et al., 2007). Fathers were compensated \$25 for completing questionnaires and \$25 for completing the telephone interview.

Materials

Child ADHD Diagnosis and Conduct Problems—A comprehensive child ADHD assessment was conducted (Pelham, Fabiano, & Massetti, 2005), and the presence of an ADHD diagnosis was required for inclusion in the original studies. ADHD symptoms were considered present if endorsed as clinically significant by the mother on the Schedule for Affective Disorders for School-Aged Children-Present and Lifetime Version (K-SADS-PL; Kaufman et al., 1997) or Disruptive Behavior Disorders checklist (DBD; Pelham, Gnagy,

Greenslade, & Milich, 1992), or the teacher on the DBD. The parent-reported DBD demonstrated good internal consistency ($\alpha=.89$). Diagnostic reliability on the KSADS has been reported elsewhere for these samples (.85-.86; Chronis-Tuscano et al., 2013; Chronis-Tuscano et al., 2011; Raggi et al., 2009). The Children's Impairment Rating Scale (CIRS; Fabiano et al., 2006) was also completed by both mothers and teachers in order to establish the presence of cross-situational impairment, a requirement for a DSM-IV diagnosis and for inclusion in these studies (APA, 2000).

To measure conduct problems, symptoms of ODD and CD were considered present if endorsed as clinically significant by the mother on the K-SADS or DBD, or the teacher on the DBD. Conduct problems were examined continuously in the current study.

Paternal ADHD Symptoms—Fathers were administered the Mini International Neuropsychiatric Interview (M.I.N.I Plus; Sheehan et al., 1998) by telephone to assess for symptoms of ADHD (childhood and adulthood), antisocial personality disorder, alcohol dependence, and non-alcohol substance dependence (all measured continuously). The MINI was selected for its strong psychometric properties, including concordance with the Structured Clinical Interview for DSM-IV (SCID; First, Gibbon, Spitzer, & Williams, 1996; Sheehan et al., 1998) and brief administration time. Telephone administration of diagnostic interviews is supported by prior studies (Rohde, Lewinsohn, & Seeley, 1997; Sobin et al., 1993) and was preferable in the current study in order to minimize participant burden.

Paternal Involvement—During the current study, fathers also completed the engagement/activities section of the Early Head Start Father Interview (Cabrera et al., 2004), to measure the frequency with which they performed specific activities with the target child in the past month (e.g., "Played outside in the yard, a park, or a playground with him/her?"). We modified the original measure by removing items that were not relevant to children ages 6-12 and adding items addressing essential activities for children with ADHD (e.g., "Attended a parent-teacher conference at his/her school?", "Took him/her to the doctor?"). Good internal consistency has been observed for both the original measure ($\alpha = .77-.86$; Cabrera et al., 2004) and our modified version ($\alpha=.93$), the latter of which also shows convergent validity with the Alabama Parenting Questionnaire Involvement subscale ($r=.62$; Shelton, Frick, & Wootton, 1996). Scores were entered as a frequency count (i.e., number of times fathers participated in each activity, summed across items) and examined continuously.

Results

Prior to the main analyses, relationships between all study variables were examined (Table 2). Significant correlations suggested that fathers of minority children and those with higher levels of past alcohol dependence were less involved in caregiving, whereas those with higher income levels were more involved. Interestingly, no significant correlations emerged between fathers' level of involvement and continuous ADHD symptoms, substance dependence, or antisocial problems. Next, child and parent demographic variables were entered into two separate preliminary regression analyses predicting child conduct problems

(see Table 1). No demographic variables were significantly associated with child conduct problems.

A series of linear regression analyses were conducted to examine the extent to which father involvement moderated the relationship between paternal ADHD symptoms and child conduct problems. On the first step of these analyses, father involvement and paternal ADHD symptoms (both examined continuously) were entered simultaneously. In the second step, we examined whether the interaction between paternal ADHD symptoms and involvement explained variance in child conduct problems above and beyond the first step. Analyses were completed separately for paternal childhood ADHD symptoms and paternal adulthood ADHD symptoms.

Neither paternal ADHD symptoms nor paternal involvement were independently associated with child conduct problems. However, paternal involvement moderated the relationship between paternal ADHD symptoms and child conduct problems (Table 3). Specifically, the interaction terms accounted for 11.9% and 20.1% of the variance in child conduct problems for childhood and adulthood ADHD symptoms, respectively. No other form of paternal psychopathology or their interactions with father involvement were significant.

To examine the two-way interactions, the effect of paternal ADHD symptoms on child conduct problems was examined when father involvement was low (-1 SD) and high ($+1$ SD) (Aiken & West, 1991). When father involvement was low, paternal ADHD symptoms were *not* significantly related to child conduct problems ($p=0.21$ for childhood ADHD and $p=0.41$ for adult ADHD). However, in the presence of high father involvement, paternal ADHD symptoms were significantly positively related to child conduct problems ($p=0.04$ and $p=0.004$ for childhood and adulthood ADHD symptoms, respectively; Figure 2).

Discussion

This cross-sectional study extends the growing literature on associations between paternal ADHD symptoms and child conduct problems by considering level of paternal involvement as a potential moderator. Indeed, among fathers with higher levels of involvement, paternal ADHD symptoms were positively associated with child conduct problems; yet, among fathers with lower levels of involvement, there was no significant relationship. This finding represents an interesting exception to the general finding within the developmental psychology literature that higher levels of paternal involvement relate to *better* psychosocial outcomes in children (Coley & Medeiros, 2007; Flouri & Buchanan, 2002). It is likely that the relation between paternal involvement and child behavior depends on paternal mental health, in this case paternal ADHD symptoms.

The results of this study should be considered in light of its limitations. First, fathers are extraordinarily difficult to recruit for participation in research (Addis & Mahalik, 2003; Mitchell et al., 2007), and this is compounded in clinical samples (Cassano, Adrian, Veits, & Zeman, 2006; Pfiffner et al, 2001). As a result, this study was limited by a small and potentially select sample. Non-residential fathers comprised only 10.8% of the sample, which was lower than both that of the larger sample from which we recruited (38%) and

what would be expected within families of children with ADHD (Wymbs et al., 2008). Additionally, our subsample had a significantly higher mean income level than the larger sample from which we recruited. These sample characteristics may limit the generalizability of our findings to the broader population of families of children with ADHD. Future studies should recruit larger samples that are more representative of the ADHD population, perhaps through community-based establishments or using internet-based data collection.

A second important limitation concerns the timing of variable measurement. This study was cross-sectional in nature. Additionally, because this was an add-on study, child conduct problems were assessed at the time families participated in the original studies, prior to the measurement of paternal psychopathology and involvement. As a result, we cannot assume that paternal ADHD symptoms and involvement pre-dated child conduct problems (Mash & Dozois, 2003). Future longitudinal studies may clarify whether high levels of both paternal ADHD symptoms and involvement is a potential risk factor for the development of later conduct problems among children with ADHD.

Third, the manner in which fathers were recruited was limited in our study. We required mothers' permission to contact fathers, and prior research has found that mothers often function as "gatekeepers" to protect fathers' time (Mitchell et al., 2007). Phone calls to elicit participation and complete phone interviews were also made by a female investigator. Future studies which employ face-to-face visits and involve male research assistants in contacting fathers may be more successful, as these procedures have been found to be considerably more effective for recruiting fathers, regardless of income-level and residential-status (Mitchell et al., 2007). These limitations aside, this study was unique in that we collected direct reports from fathers about their ADHD symptoms, which is recommended but rarely obtained in this literature (Pfiffner et al., 2001).

Finally, future studies should consider the *quality* of father involvement, rather than simply the *frequency* of father involvement in various childrearing activities. It is likely that maladaptive parenting is the mechanism by which involvement by fathers with elevated ADHD symptoms have children with higher levels of conduct problems (Arnold et al., 1997). For instance, paternal inattention may impact fathers' ability to monitor and maintain consistency, whereas impulsivity may promote harsh and emotionally-reactive discipline, two aspects of parenting which have been associated with child conduct problems (Pfiffner, McBurnett, Rathouz, & Judice, 2005). Future research should therefore seek to more fully understand the extent to which paternal parenting underlies the current study finding.

Despite these limitations, this study is among the very first to consider both paternal ADHD and paternal involvement in relation to child conduct problems, and its results preliminarily suggest that paternal psychological health may determine whether father involvement in childrearing has a positive or detrimental influence on child conduct problems. As such, this study addresses a general lack of research related to fathers and specifically emphasizes the importance of examining modifiable risk factors, especially paternal involvement, among families of children with ADHD in order to prevent the escalation of problems in this population. We hope that our provocative findings will stimulate larger studies on this topic.

Within a clinical context, these findings suggest the importance of routinely assessing paternal ADHD and involvement as one component of child ADHD assessment. Moreover, it may be especially important to engage highly-involved fathers with elevated ADHD symptoms in treatment (Fabiano, 2007). Indeed, maternal ADHD symptoms have been shown to be associated with diminished treatment effects for children with ADHD (Chronis-Tuscano et al., 2011; Sonuga-Barke, Daley, & Thompson, 2002). Thus, researchers and clinicians should consider integrating treatment for maternal and paternal ADHD symptoms, when present, as one component of comprehensive treatment for their children (Chronis-Tuscano & Stein, 2012; Johnston et al., 2012).

Funding & Acknowledgments

This research was supported by the University of Maryland College Park (Summer Research Award to Dr. Andrea Chronis-Tuscano), the National Institute of Mental Health (Grants R34 MH073567-01 and R03 MH070666-1 to Dr. Andrea Chronis-Tuscano), APA Division 53 Dissertation Award (to Dr. Veronica Raggi), and the Milton Dean Havron Social Sciences Award for Academic Excellence (to Dr. Veronica Raggi). A debt of gratitude is owed to the many doctoral students, postdoctoral fellows, and undergraduates who contributed to this research, and especially to Dr. Natasha Cabrera for her guidance regarding study design.

References

- Addis ME, Mahalik JR. Men, masculinity, and the contexts of help seeking. *American Psychologist*. 2003; 58:5–14. [PubMed: 12674814]
- Aiken, LS.; West, SG. Multiple regression: Testing and interpreting interactions. Sage; Newbury Parge, CA: 1991.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed. The American Psychiatric Association; Washington, DC: 1994.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed. - Text Revision. The American Psychiatric Association; Washington, DC: 2000.
- Arnold EH, O'Leary SG, Edwards GH. Father involvement and self-reported parenting of children with attention deficit-hyperactivity disorder. *Journal of Consulting and Clinical Psychology*. 1997; 65:337–342. [PubMed: 9086700]
- Biederman J, Faraone SV, Milberger S, Garcia-Jetton J, Chen L, Mick E, et al. Is childhood oppositional defiant disorder a precursor to adolescent conduct disorder? Findings from a four-year follow-up study of children with ADHD. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1996b; 35:1193–204. [PubMed: 8824063]
- Bird HR, Gould MS, Staghezza-Jaramillo BM. The comorbidity of ADHD in a community sample of children aged 6 through 16 years. *Journal of Child and Family Studies*. 1994; 3:365–378.
- Cassano M, Adrian M, Veits G, Zeman J. The inclusion of fathers in the empirical investigation of child psychopathology: An update. *Journal of Clinical Child and Adolescent Psychology*. 2006; 35:583–589. [PubMed: 17007604]
- Caspi A, Taylor A, Smart M, Jackson J, Tagami S, Moffitt TE. Can women provide reliable information about their children's fathers? Cross-informant agreement about men's lifetime antisocial behaviour. *Journal of Child Psychology and Psychiatry*. 2001; 42:915–920. [PubMed: 11693586]
- Chronis-Tuscano A, Clarke TL, O'Brien KA, Raggi VL, Diaz Y, Lewinsohn PM. Development & Preliminary Evaluation of an Integrated Treatment for Depressed Mothers of Children with Attention-Deficit/Hyperactivity Disorder. *Journal of Consulting & Clinical Psychology*. 2013 Advance online publication. doi: 10.1037/a0032112.
- Chronis AM, Lahey BB, Pelham WE, Kipp H, Baumann B, Lee SS. Psychopathology and substance abuse in parents of young children with attention - deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2003; 42:1424–1432. [PubMed: 14627877]

- Chronis AM, Lahey BB, Pelham WE, Williams SH, Baumann BL, Kipp H. Maternal depression and early positive parenting predict future conduct problems in young children with attention-deficit/hyperactivity disorder. *Developmental Psychology*. 2007; 43:70–82. [PubMed: 17201509]
- Chronis-Tuscano A, O'Brien KA, Jones HA, Clarke TL, Raggi VL, Seymour KE. The relation between maternal ADHD symptoms & improvement in child behavior following brief behavioral parent training is mediated by change in negative parenting. *Journal of Abnormal Child Psychology*. 2011; 39(7):1047–1057. [PubMed: 21537894]
- Chronis-Tuscano A, Raggi VL, Clarke TL, Rooney ME, Diaz Y, Pian J. Associations between maternal attention-deficit/hyperactivity disorder symptoms and parenting. *Journal of Abnormal Child Psychology*. 2008; 36:1237–1250. [PubMed: 18553132]
- Chronis-Tuscano A, Stein MA. Pharmacotherapy for parents with Attention-Deficit Hyperactivity Disorder (ADHD): Impact on maternal ADHD and parenting. *CNS Drugs*. 2012; 26:725–732. [PubMed: 22647070]
- Coie, JD.; Dodge, KA. Aggression and antisocial behavior. In: Damon, W.; Eisenberg, N., editors. *Handbook of child psychology*. 5th ed. Wiley; New York: 1998. p. 779-862.
- Coley RL, Medeiros BL. Reciprocal longitudinal relations between nonresident father involvement and adolescent delinquency. *Child Development*. 2007; 78:132–147. [PubMed: 17328697]
- Elkins IJ, McGue M, Iacono WG. Prospective effects of attention-deficit/hyperactivity disorder, conduct disorder, and sex on adolescent substance use and abuse. *Archives of General Psychiatry*. 2007; 64:1145–1152. [PubMed: 17909126]
- Fabiano GA, Pelham WE, Waschbusch DA, Gnagy EM, Lahey BB, Chronis AM. A practical measure of impairment: psychometric properties of the impairment rating scale in samples of children with attention deficit hyperactivity disorder and two school-based samples. *Journal of Clinical Child and Adolescent Psychology*. 2006; 35:369–385. [PubMed: 16836475]
- Fabiano GA. Father Participation in Behavioral Parent Training for ADHD: Review and Recommendations for Increasing Inclusion and Engagement. *Journal of Family Psychology*. 2007; 21:638–693.
- Faraone SV, Biederman J, Spencer T, Wilens T, Seidman LJ, Mickac E, Doyle AE. Attention-deficit/hyperactivity disorder in adults: An overview. *Biological Psychiatry*. 2000; 48:9–20. [PubMed: 10913503]
- Faraone SV, Perlis RH, Doyle AE, Smoller JW, Goralnick JJ, Holmgren MA, Sklar P. Molecular Genetics of Attention-Deficit/Hyperactivity Disorder. *Biological Psychiatry*. 2005; 57:1313–1323. [PubMed: 15950004]
- First, MB.; Gibbon, M.; Spitzer, RL.; Williams, JB. *Structured Clinical Interview for DSM-IV Axis I Disorders*. Biometrics Research; New York, NY: 1996.
- Flouri E, Buchanan A. Father involvement in childhood and trouble with the police in adolescence: Findings from the 1958 British cohort. *Journal of Interpersonal Violence*. 2002; 17:689–701.
- Harty SC, Miller CJ, Newcorn JH, Halperin JM. Adolescents with childhood ADHD and comorbid disruptive behavior disorders: Aggression, anger, and hostility. *Child Psychiatry and Human Development*. 2009; 40:85–97. [PubMed: 18597170]
- Harvey E, Danforth JS, McKee TE, Ulaszek WR, Friedman JL. Parenting of children with Attention-Deficit/Hyperactivity Disorder (ADHD): The role of parental ADHD symptomatology. *Journal of Attention Disorders*. 2003; 7:31–42. [PubMed: 14738179]
- Henry B, Caspi A, Moffitt TE, Silva PA. Temperamental and familial predictors of violent and nonviolent criminal convictions: Age 3 to age 18. *Developmental Psychology*. 1996; 32:614–623.
- Johnston C, Mash EJ, Miller N, Ninowski JE. Parenting in adults with attention-deficit/hyperactivity disorder (ADHD). *Clinical Psychology Review*. 2012; 32:215–228. [PubMed: 22459785]
- Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, Williamson D. Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): Initial reliability and validity data. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1997; 36:980–988. [PubMed: 9204677]
- Lahey BB, Loeber R, Quay HC, Applegate B, Shaffer D, Waldman I, Bird HR. Validity of DSM-IV subtypes of conduct disorder based on age of onset. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1998; 37:435–442. [PubMed: 9549965]

- Lynam DR. Early identification of the fledgling psychopath: Locating the psychopathic child in the current nomenclature. *Journal of Abnormal Psychology*. 1998; 107:566–575. [PubMed: 9830244]
- Mash, EJ.; Dozois, JJA. Child psychopathology: A developmental-systems perspective. In: Mash, EJ.; Barkley, RA., editors. *Child Psychopathology*. 2nd edition. The Guilford Press; New York: 2003. p. 1-71.
- Mikami AY, Pfiffner LJ. Siblings relationships among children with ADHD. *Journal of Attention Disorders*. 2008; 11:482–492. [PubMed: 17494830]
- Mitchell SJ, See HM, Tarkow AKH, Cabrera N, McFadden KE, Shannon JD. Conducting studies with fathers: Challenges and opportunities. *Applied Development Science*. 2007; 11:239–244.
- Murray C, Johnston C. Parenting in mothers with and without attention-deficit/ hyperactivity disorder. *Journal of Abnormal Psychology*. 2006; 115:52–61. [PubMed: 16492095]
- Pelham WE, Gnagy EM, Greenslade KE, Milich R. Teacher ratings of DSM-III—R symptoms for the disruptive behavior disorders. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1992; 31:210–218. [PubMed: 1564021]
- Pelham WE, Fabiano GA, Massetti GM. Evidence-based assessment of attention deficit hyperactivity disorder in children and adolescents. *Journal of Clinical Child and Adolescent Psychology*. 2005; 34:449–76. [PubMed: 16026214]
- Pfiffner LJ, McBurnett K, Rathouz PJ. Father absence and familial antisocial characteristics. *Journal of Abnormal Child Psychology*. 2001; 29:357–367. [PubMed: 11695539]
- Pfiffner LJ, McBurnett K, Rathouz PJ, Judice S. Family correlates of oppositional and conduct disorders in children with attention deficit/hyperactivity disorder. *Journal of Abnormal Child Psychology*. 2005; 33:551–563. [PubMed: 16195950]
- Phares V. Where's poppa? The relative lack of attention to the role of fathers in child and adolescent psychopathology. *American Psychologist*. 1992; 47:656–664. [PubMed: 1642375]
- Piacentini J, Cohen P, Cohen J. Combining discrepant diagnostic information from multiple sources: are complex algorithms better than simple ones? *Journal of Abnormal Child Psychology*. 1992; 20:51–63. [PubMed: 1548394]
- Raggi VL, Chronis-Tuscano A, Fishbein H, Groomes A. Development of a brief, behavioral homework intervention for middle school students with attention-deficit/hyperactivity disorder. *School Mental Health*. 2009; 1:61–77.
- Rohde P, Lewinsohn PM, Seeley JR. Comparability of telephone and face-to-face interviews in assessing Axis I and II disorders. *American Journal of Psychiatry*. 1997; 154:1593–1598. [PubMed: 9356570]
- Rogers MA, Wiener J, Marton I, Tannock R. Parental involvement in children's learning: Comparing parents of children with and without Attention-Deficit/Hyperactivity Disorder (ADHD). *Journal of School Psychology*. 2009; 47:167–185. [PubMed: 19328285]
- Romano E, Tremblay RE, Vitaro F, Zoccolillo M, Pagani L. Sex and informant effects on diagnostic comorbidity in an adolescent community sample. *Canadian Journal of Psychiatry/Revue Canadienne de Psychiatrie*. 2005; 50:479–489.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E. The Mini International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*. 1998; 59(Supplement 20):22–33. [PubMed: 9881538]
- Shelton KK, Frick PJ, Wootton J. Assessment of parenting practices in families of elementary school-age children. *Journal of Clinical Child Psychology*. 1996; 25:317–329.
- Singh I. Boys will be boys: Fathers' perspectives on ADHD symptoms, diagnosis, and drug treatment. *Harvard Review of Psychiatry*. 2003; 11:308–316. [PubMed: 14713567]
- Sobin C, Weissman MM, Goldstein RB, Adams P, Wickramaratne P, Warner V, Lish JD. Diagnostic interviewing for family studies: Comparing telephone and face-to-face methods for the diagnosis of lifetime psychiatric disorders. *Psychiatric Genetics*. 1993; 3:227–233.
- Sonuga-Barke EJS, Daley D, Thompson M. Does maternal ADHD reduce the effectiveness of parent training for preschool children's ADHD? *Journal of the American Academy of Child and Adolescent Psychiatry*. 2002; 41:696–702. [PubMed: 12049444]

- Thapar A, Langley K, Owen MJ, O'Donovan MC. Advances in genetic findings on attention deficit hyperactivity disorder. *Psychological Medicine*. 2007; 37:1681–1692. [PubMed: 17506925]
- Wymbs BT, Pelham WE, Molina BSG, Gnagy EM, Wilson TK, Greenhouse JB. Rate and predictors of divorce among parents of youths with ADHD. *Journal of Consulting and Clinical Psychology*. 2008; 76:735–744. [PubMed: 18837591]
- Yang L, Wang YF, Qian QJ, Biederman J, Faraone SV. DSM-IV subtypes of ADHD in a Chinese outpatient sample. *Journal of American Academy of Child and Adolescent Psychiatry*. 2004; 43:248–250.

Key Messages

1. The impact of father involvement on child development may be complicated by the presence of paternal psychopathology, specifically adult ADHD symptoms.
2. As with mothers, high levels of ADHD symptoms in fathers may be associated with related impairments in the quality of parenting.
3. Though mothers are traditionally the primary point of contact for child ADHD interventions, clinicians should attempt to engage fathers of children with ADHD.

Figure 1.

Consort Diagram

Note. CIRS=Children's Impairment Rating Scale; DBD=Disruptive Behavior Disorders Scale; KSADS=Schedule for Affective Disorders for School-Aged Children; MINI=Mini International Neuropsychiatric Interview.

Figure 2.
Interaction of Paternal ADHD Symptoms and Paternal Involvement Predicting Child Conduct Problems (Based on Median Split)

Note. ADHD=attention-deficit/hyperactivity disorder.

* $p < .05$, ** $p < .01$

Table 1

Sample Demographic and Diagnostic Characteristics (N = 37)

<i>Child Demographic Variables</i>		
Age		8.29 (2.00)
Gender		
Male	26 (70.3%)	
Female	11 (29.7%)	
Ethnic Characteristics		
White	23 (62.2%)	
African American	5 (13.5%)	
Hispanic or Latino	1 (2.7%)	
Biracial	7 (19.4%)	
<i>Child Clinical Characteristics (KSADS/DBD)</i>		
ODD Diagnosis	18 (48.6%)	
CD Diagnosis	3 (8.1%)	
Conduct Problems		4.59 (2.95)
<i>Maternal Demographic Variables</i>		
Age		41.03 (5.83)
Total Family Income		\$126,897.87 (\$67,159.07) Range=\$22,000-\$300,000
Highest Level of Education (years post middle school; proxy for SES)		7.70 (3.09)
<i>Paternal Demographic Variables</i>		
Age		42.94 (7.19)
Residential Status		
Residential	33 (89.2)	
Non-residential	4 (10.8)	
<i>Paternal Clinical Characteristics (MINI)</i>		
ADHD (Childhood Diagnosis)	9 (24.3)	
ADHD (Adulthood Diagnosis)	5 (13.5)	
Depression (Lifetime Diagnosis)	5 (13.5)	
Alcohol Dependence (Lifetime Diagnosis)	12 (32.4)	
Substance Dependence (Lifetime Diagnosis)	5 (13.5)	
ASPD (Lifetime Diagnosis)	1 (2.7)	
Any Psychiatric Disorder (Lifetime Diagnosis)	19 (51.4)	

Note. Results presented as M (SD) or as percent (n). ADHD=attention-deficit/hyperactivity disorder, CD=conduct disorder, DBD=Disruptive Behavior Disorders Scale; DSM-IV=Diagnostic and Statistical Manual, Fourth Edition; KSADS=Schedule for Affective Disorders for School-Aged Children; MINI=Mini International Neuropsychiatric Interview; ODD=oppositional defiant disorder; SES=socioeconomic status.

Table 2
Correlation Matrix of Independent, Dependent, Moderator and Demographic Variables

Variables	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1. Child Age	-	-0.05	0.17	-0.02	-0.28	0.00	0.21	0.04	-0.01	-0.11	-0.20	0.10	-0.13	0.17	-0.07	-0.17	0.11
2. Child Gender		-	0.14	-0.02	-0.08	-0.00	0.15	0.25	-0.06	-0.02	-0.06	-0.23	-0.15	-0.26	-0.14	0.19	0.05
3. Child Minority Status			-	-0.14	0.12	-0.15	0.11	-0.22	0.03	-0.02	0.06	0.05	-0.16	0.04	0.08	-0.43**	-0.12
4. Child ADHD Medication Status				-	0.05	0.08	-0.28	0.09	-0.09	-0.21	-0.06	-0.28	0.05	-0.18	-0.14	0.27	-0.07
5. Child Conduct Problems					-	-0.09	0.12	0.00	0.04	0.18	-0.01	-0.12	-0.03	-0.05	-0.12	0.03	-0.16
6. Father Age						-	-0.17	0.42*	-0.05	0.03	-0.17	-0.05	-0.12	-0.22	-0.23	0.15	0.79**
7. Father Residential Status							-	0.00	0.28	-0.35*	0.21	0.26	0.25	-0.37*	0.06	-0.02	-0.23
8. Total Household Income								-	-0.13	-0.10	0.03	-0.11	-0.03	-0.15	-0.21	0.52**	0.55**
9. Father ADHD (Childhood)									-	0.87**	0.45**	0.24	0.32	0.31*	0.53**	-0.19	-0.20
10. Father ADHD (Adulthood)										-	0.47**	0.37*	0.44**	0.31*	0.47**	-0.12	-0.19
11. Father Alcohol Dependence (Past)											-	0.47**	0.79**	0.07	0.52**	-0.13	-0.24
12. Father Alcohol Dependence (Current)												-	0.56**	-0.01	0.31	-0.45**	-0.15
13. Father Substance Dependence (Past)													-	0.23	0.48**	-0.19	-0.16
14. Father Depression (Lifetime Diagnosis)														-	0.13	-0.06	-0.18
15. Father ASPD (Lifetime)															-	-0.26	-0.25
16. Father Involvement																-	0.19
17. Maternal Age																	-

Note. Results reported as Pearson product-moment correlations or Kendall's tau as appropriate. Psychopathology variables refer to continuous symptoms unless otherwise noted. ADHD=attention-deficit/hyperactivity disorder, ASPD=antisocial personality disorder.

* $p < .05$.

10>d
**

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Paternal ADHD Symptoms and Paternal Involvement Predicting Child Conduct Problems

	<i>df</i>	<i>F</i>	<i>R</i> ²	<i>R</i> ²	SE	β
Step 1	2, 33	.022	.001	.001		
Childhood ADHD					.185	.027
Involvement					.025	.030
Step 2	1, 32	2.581	.195	.194**		
Childhood ADHD $\times$ Involvement					.007	1.554**
	<i>df</i>	<i>F</i>	<i>R</i> ²	<i>R</i> ²	SE	β
Step 1	2,33	.503	.030	.030		
Adulthood ADHD					.164	.171
Involvement					.024	.046
Step 2	1,32	3.935	.269	.240**		
Adulthood ADHD $\times$ Involvement					.006	1.744**

Note. ADHD=attention-deficit/hyperactivity disorder.

*
p<.05,

**
p<.01