CORRESPONDENCE

Diving Medicine in Clinical Practice

by Dr. med. Lars Eichhorn, Prof. Dr. med. Dr. Sportwiss. Dieter Leyk
in issue 9/2015

Dr. med. Ralf Cüppers

Arzt für Psychotherapeutische Medizin, Flensburg
ralf@psychotherapeutische-medizin.net

Conflict of interest statement

The author declares that no conflict of interest exists.

Psychological Contraindications

On the one hand, any licensed physician is allowed to attest a prospective diver's fitness to dive, even without a relevant further qualification. On the other hand, s/he should certainly not do so after merely reading this short CME article. Not for nothing does the fitness to dive manual comprise more than 200 printed pages, which a doctor should know before attesting anything.

There is no such thing as "unfitness to dive." Aviation physicians have the right to bar a would-be pilot from flying a plane; the reason is that such an undertaking may put others at risk in case of a crash. Diving medicine specialists cannot bar persons from diving because recreational diving puts only the divers themselves at risk. Diving is always the diver's own responsibility. A medical certificate attesting fitness to dive mainly serves to protect diving schools and tour operators. Making such a certificate a requirement will have a single positive effect in that recreational divers are obliged to submit to a diving medical consultation, in which they will receive extensive and individual information on what their personal risk factors are. Such advice is the main task of a diving medicine specialist; the paper certificate is an optional extra.

What the CME article is unforgivably lacking is any mention of psychological contraindications. Most life-threatening diving incidents are caused by psychological slips. Diving medicine specialists should elicit information on how someone deals with stress, how willing they are to experience fear, and how they handle high-risk behavior. A diagnosis of a "phobic patient" is a relative contraindication, as is that of a "carefree/careless person." Hyperventilation in a scuba diver having a panic attack under water is not a good scenario. Personality disorders require critical evaluation. Further contraindications include depression, suicidality, manic hubris, psychotic symptoms with misjudgment of reality, and addictive behaviors including and excluding substances. It goes without saying that recreational divers should be sober and not currently be taking psychotropic drugs.

In my view, the psychological situation of recreational divers is more important than their lung function or cardiac output when the objective is to prevent diving accidents by means of giving high-quality advice.

DOI: 10.3238/arztebl.2015.0614a

REFERENCES

1. Wendling J, Ehm O, Ehrsam R, Kness P, Nussberger P (eds.): Tauchtauglichkeit Manual: Richtlinien für die Untersuchung von Sporttauchern der GTÜM (Deutschland), SGUHM (Schweiz) und ÖGTH (Österreich); 2nd edition. Hyperbaric Editions c/o Dr. Jürg Wendling 2001.
2. Eichhorn L, Leyk D: Diving medicine in clinical practice. Dtsch Arztebl Int 2015; 112: 147–58.

Emergency Equipment on Diving Boats

Diving sites used during holidays almost never have rapid access to a decompression chamber. Boats rarely carry oxygen on board. For this reason I have the following advice for diving stations, which is put into practice in a few cases: keep a full, compressed air cylinder in the boat. Attach a breathing regulator on a 10-meter high-pressure hose (available from retailers of diving accessories) and lower down. In the first signs of decompression sickness, take a dive down to the regulator, preferably accompanied (your own cylinder is most probably empty by this stage) and decompress once more, slowly. A fast and effective measure.

DOI: 10.3238/arztebl.2015.0614b

REFERENCES

1. Eichhorn L, Leyk D: Diving medicine in clinical practice. Dtsch Arztebl Int 2015; 112: 147–58.

Dr. med. Rolf Stockhausen

Meerbusch
info@schnarchen.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank Dr Cüppers and agree completely, that studying a CME article is absolutely no substitute for a specialist training qualification in diving medicine. This is obvious from our article, in which we explicitly recommend (for example, in the Overview section) attending the courses of the German Society for Diving and Hyperbaric Medicine (GTÜM) and obtain GTÜM certification by passing the two relevant courses that the association offers, "Medical Evaluation for Diving" (Tauchmedizinische Untersuchungen, GTÜM Course I) and "The Diving Physician Course" (Taucherarzt, GTÜM Course IIa). We also clearly pointed out the GTÜM's pertinent examination standards and recommendations, seeking expert advice, making available advice over the telephone from diving medicine specialists, and relevant literature. Our article aimed to inform the wide readership of *Deutsches Ärzteblatt* about the physiological-physical basics of diving, about fitness to dive examinations and medical diving advice, as well as about diving emergencies and their treatment.

Attentive readers could not fail to notice that the psychological state of divers is a subject we dealt with in our article. When conducting the fitness to dive examination, doctors should pay attention to the potential diver's psychological condition, not only in the context of a thorough medical history, but they should also conduct a critical analysis of a person's medication (see the

Box in our article). The GTÜM explicitly provides for considering the psychological state of divers with its questionnaire (www.gtuem.org).

With regard to diving incidents, Cüppers thinks that examining the psychological situation of recreational divers is more important. We do not have any robust data to contribute to this evaluation. It is therefore important during a careful diving medical examination to identify all potential risk factors, if possible. This obviously includes the psychological sector.

Dr Stockhausen rightly mentions the often lacking emergency equipment on dive boats. However, we counsel against the suggested “in-water decompression” (a repeated dive on noticing the first symptoms of decompression sickness), for a multitude of reasons: the initially mild symptoms of decompression sickness are often progressive and can deteriorate in an unforeseeable manner. Options for intervention in deep water are extremely limited, and adequate monitoring of a diver is not possible. A depth of 10 meters (=2 bar) is mostly not sufficient to eliminate the bubbles responsible for the symptoms. Although additional administration of 100% O₂ would mean that the higher concentration gradient in the lung would favor elimination of nitrogen, the induced high partial O₂ pressures are dangerous with regard to the potential of oxygen to trigger seizures. Furthermore, the gas reserves on board are not likely to be sufficient for recompression of adequate duration (the US Navy Table 6, for

example, which is often used for the purposes of treatment, provides for a treatment duration of >280 min). Furthermore, the boat cannot return to base with a diver dangling from a rope, so the transfer into a decompression chamber is delayed further.

To conclude, divers should be clearly instructed to ask dive base operators for adequate emergency equipment and to find out about the emergency management on site.

DOI: 10.3238/arztebl.2015.0614c

REFERENCES

1. Eichhorn L, Leyk D: Diving medicine in clinical practice. Dtsch Arztebl Int 2015; 112: 147–58.

Dr. med. Lars Eichhorn

Klinik und Poliklinik für Anästhesiologie und Operative Intensivmedizin,
Universitätsklinikum Bonn
lars.eichhorn@ukb.uni-bonn.de

Prof. Dr. med. Dr. Sportwiss. Dieter Leyk

Zentrales Institut des Sanitätsdienstes der Bundeswehr Koblenz, Laborabteilung IV – Wehr-
medizinische Ergonomie und Leistungsphysiologie;
Deutsche Sporthochschule Köln, Forschungsgruppe Leistungsepidemiologie, Institut für
Physiologie und Anatomie

Conflict of interest statement

Prof. Dr. med. Dr. Sportwiss. Dieter Leyk and Dr. med. Lars Eichhorn declare that no conflict of interest exists. Dr. Eichhorn is the recipient of a scholarship from the Else-Kröner-Fresenius-Stiftung.