Published in final edited form as:

Crit Care Clin. 2015 October ; 31(4): 705–723. doi:10.1016/j.ccc.2015.06.007.

Acute Kidney Injury in The Surgical Patient

Charles Hobson, MD, MHA^{1,2}, Girish Singhanian, MD³, and Azra Bihorac, MD, MS^{3,4}

¹Department of Surgery, Malcom Randall VA Medical Center, Gainesville, FL

²Department of Health Services Research, Management, and Policy, University of Florida, Gainesville, FL

³Department of Medicine, University of Florida, Gainesville, FL

⁴Department of Anesthesiology, University of Florida, Gainesville, FL

Synopsis

Perioperative acute kidney injury (AKI) is a common, morbid and costly surgical complication. Current efforts to understand and manage AKI in surgical patients focus on prevention, mitigation of further injury when AKI has occurred, treatment of associated conditions and facilitation of renal recovery. Lesser severity AKI is now understood to be much more common, and more morbid, than was previously believed. The ability to detect AKI within hours of onset would be helpful in protecting the kidney and in preserving renal function, and several imaging and biomarker modalities are currently being evaluated. Automated, rapid and noninvasive collection and analysis of preoperative and operating room data raises the prospect of real time risk prediction for perioperative AKI. Prevention of AKI is of paramount importance, and goal-directed fluid management, remote ischemic preconditioning and early involvement of a nephrologist have all been shown to be effective.

Introduction

Perioperative acute kidney injury (AKI), characterized by persistent oliguria and/or an increase in serum creatinine levels, is a common perioperative complication and is associated with up to a tenfold increase in hospital cost and mortality, decreased long-term survival and an increased risk for chronic kidney disease (CKD) and hemodialysis after discharge [1-11]. Depending upon the type of surgical procedure that a patient undergoes, AKI complicates the perioperative hospital stay for up to 50% of surgical patients [1, 2, 12-16]. Yet AKI remains among the most underdiagnosed postoperative complications despite increasing understanding of its epidemiology and outcomes. Considering the high

Corresponding Author: Azra Bihorac MD MS, Department of Anesthesiology, PO Box 100254, Gainesville, FL 32610-0254, Telephone: (352) 273-9009, Fax: (352) 392-7029, abihorac@anest.ufl.edu.

Co-author: Girish Singhanian MD, Department of Medicine, PO Box 100254, Gainesville, FL 32610-0254, Telephone: (352) 273-9009, Fax: (352) 392-7029, girish.singhanian@medicine.ufl.edu, Co-author: Charles Hobson MD MHA, Department of Surgery, NF/SG VAMC, Gainesville, FL 32608 Telephone: 352-514-2854, chobson@ufl.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

prevalence of AKI and the deleterious effect when it occurs, our effort must focus on AKI prevention, mitigation of further injury when AKI has already occurred, treatment of negative effects on other organs, and facilitation of renal recovery in patients with established AKI. Although the understanding of the mechanisms of AKI has grown substantially, and the emergence of new biomarkers and imaging techniques has provided new tools for early risk stratification and diagnosis, the translation of these discoveries into clinical practice has been slow. The well-defined timing of surgical physiologic stress on the kidney in the perioperative period provides a unique opportunity for early risk stratification to guide perioperative assessment and preventive therapies to achieve these goals.

Definitions, Epidemiology and Outcomes Associated with Perioperative AKI

Prior to 2004 the reported incidence of hospital-acquired AKI varied significantly from 1 to 31% due to the incoherent criteria used to define AKI and the focus placed on the most severe AKI defined either by a large increase in serum creatinine or the need for renal replacement therapy (RRT) [17, 18]. In 2004 the Acute Dialysis Quality Initiative published Risk, Injury, Failure, Loss, and End-stage Kidney (RIFLE), a consensus definition for AKI that for the first time included less severe AKI stages and provided a structured classification for severity and recovery[19]. The recent Kidney Disease: Improving Global Outcomes (KDIGO) guidelines have expanded the AKI criteria to include changes as small as 0.3 mg/dL [20]. The reported epidemiology and outcomes of AKI have been under rapid evolution since the publication of these new definitions.

Surgical societies and registries have been slow in adopting these new definitions. The American College of Surgeons' Committee on Trauma defines acute renal failure as a serum creatinine increase greater or equal to 3.5 mg/dL, but in a large multicenter trauma study only 15% of all patients with AKI defined by RIFLE criteria had a peak creatinine value greater than 3 mg/dL [2]. The American College of Surgeons' National Surgical Quality Improvement Project (NSQIP) is the largest existing prospective surgical database that quantifies 30-day risk-adjusted surgical outcomes for patients undergoing major surgery, and it defines AKI as a rise in serum creatinine greater than 2 mg/dL from the patient's baseline or as the acute need for RRT [21]. Not surprisingly, studies using the ACS NSQIP database have demonstrated a substantial 30-day mortality associated with AKI and an incidence as low as 1%, creating the perception that postoperative AKI is a rare and often fatal complication after surgery[22]. In a recent single-center cohort study of > 20,000 postoperative patients, the ACS NSQIP definition for AKI severely underestimated the incidence of AKI defined by consensus criteria[12].

The incidence of AKI in recent studies using the current consensus criteria ranges from 25% for trauma patients[2] to as high as 50% for patients undergoing aortic surgery or liver transplantation [23, 24]. AKI has been demonstrated to be a common and serious postoperative complication associated with increased risk for short and long term mortality, increased incidence of CKD, increased incidence of other postoperative complications, and much higher resource utilization compared to patients with no postoperative AKI[3, 12, 25-34]. The risk-adjusted association between postoperative change in serum creatinine and

adverse clinical outcomes is continuous and observed at even lower cut-offs than in the original RIFLE definition [12, 35], and it has been shown that the adverse effects of AKI, as defined by consensus criteria, persistent for years even for those patients considered to have partial or even full recovery by the time of hospital discharge[3, 4]. The risk-adjusted average cost of care for these patients was \$42,600 for patients with any AKI compared to \$26,700 for patients without AKI[1]. Thus prevention of postoperative AKI should be seen as an important target in standardized surgical practice and in studies focusing on quality measures that could translate into improved care for the surgical patient.

Risk Factors for AKI

A number of preoperative and intraoperative factors have emerged as important and common predictors for AKI across different surgical populations (Table 1). Scoring systems have been used for years in an attempt to measure risk factors for adverse outcomes after surgery. Several existing scoring systems that measure the risk for AKI after cardiac surgery rely mostly on preoperative variables (reviewed in [36]). More novel preoperative factors such as total lymphocyte count <1,500 cells/ μ L and elevated C-reactive protein were identified to be associated with AKI after cardiac surgery [37, 38]. Genetic polymorphisms for selected inflammatory and vasoconstrictor genes (alleles angiotensinogen 842C and interleukin 6–572C in Caucasians, and endothelial nitric oxide synthase 894T and angiotensin-converting enzyme deletion and insertion in African Americans) provided two- to four-fold improvement over clinical factors alone in explaining post-cardiac surgery AKI [39]. For patients undergoing non-cardiac surgery, preoperative factors unique for the surgery type need to be considered [40]. Women undergoing surgery for any type of cancer, and especially those with metastatic cancer, had significantly higher odds of developing AKI [14]. Among patients undergoing orthotopic liver transplantation the Model for End-Stage Liver Disease score, but not pre-transplantation creatinine values, were predictive of AKI [41]. A low Norton scale score (a measure of a patient's risk for developing a pressure ulcer) and preoperative use of diuretics and NSAIDs were all associated with AKI following total hip arthroplasty [42, 43]. Among patients undergoing endovascular abdominal aortic aneurysm repair the use of fenestrated grafts and increasing contrast dose carried a higher risk for AKI [44]. In a large prospective study of severe trauma patients any increase in serum creatinine on admission above that expected (based on a patient's age, gender and race), an increase in lactic acid, low body temperature and any transfusion of packed red blood cells and cryoprecipitate within the first 24 hours of trauma were associated with the increased risk for subsequent AKI [2]. Attempting to create a meaningful risk calculation for AKI from these disparate factors has been challenging.

While some of these factors are unavoidable others are both preventable and largely ignored in routine clinical practice. Preoperative assessment of kidney health using readily available clinical tests - urinary albuminuria and estimated glomerular filtration rate using serum creatinine - is one of the least utilized yet most valuable clinical resources not only for the assessment of the risk for AKI but also the overall risk for postoperative morbidity and mortality. Chronic kidney disease (CKD) affects 5% of the US population[45] and is an independent predictor of mortality and cardiovascular events [46]. A systematic review of 31 cohort studies of patients undergoing elective surgery demonstrated a graded relationship

between CKD severity and postoperative death, comparable to that seen with diabetes, stroke and coronary disease [47]. In a recent analysis of the ACS NSQIP database the adjusted hazard ratio for 30-day mortality ranged from 2.30 for stage 3 CKD to 3.05 for stage 5 CKD compared with no CKD [48]. Furthermore, preoperative proteinuria, independent of preoperative eGFR and other comorbidities, was not only associated with the risk of AKI but was also a powerful independent risk factor for long-term all-cause mortality and ESRD after cardiac surgery [49, 50]. However the importance of CKD as a perioperative risk factor is still not widely appreciated among physicians involved in perioperative decision making.

Some perioperative risk indicators consider CKD an important prognostic factor in postoperative risk assessment [51-53] whereas others do not [54, 55]. One difficulty in increasing the awareness of CKD as an important perioperative risk factor is the complicated relationship between serum creatinine and eGFR calculated with commonly used estimation equations such as the Chronic Kidney Disease Epidemiology Collaboration equation [56]. Especially among women and the elderly the serum creatinine alone may not give an accurate picture of CKD as creatinine values within normal limits may correspond to a low eGFR. Thus in the perioperative setting use of estimation equations to assess eGFR, rather than relying on serum creatinine values alone, is a strategy to assure that CKD is appropriately assessed as a risk factor for not only AKI but overall postoperative mortality.

One factor that has received intense interest recently has been the possible effect of preoperative medications on preoperative complications. Statins (or HMG-CoA reductase inhibitors) have received the most attention due to their important pleiotropic effects including reduced decreased vascular inflammation and improved endothelial function [57]. The data on preoperative statin use and risk for AKI is evolving and contradictory. The subgroup analysis of a systematic review of preoperative use of statins among cardiac surgery patients (including four studies with a total of 367 patients, including 19 patients with renal failure) demonstrated no benefit from preoperative statin use [58]. A study from The Cleveland Clinic reporting no difference in AKI or hospital mortality between 4,683 statin users propensity matched with 22,000 non-statin users was limited to elective surgery cases and reported a very low incidence of AKI (6%), hospital mortality (0.6%), and need for dialysis (0.05%) rendering it difficult to compare to other studies. In a small cohort of 151 vascular surgery patients, Kor et al found no difference with preoperative statin use using moderate to severe AKI (incidence 7%), need for RRT (3%) and mortality (5%) as endpoints [58, 59]. In contrast a population-based Canadian retrospective cohort study including 213,347 older patients who underwent major elective surgery demonstrated a 16% lower odds of severe AKI, 17% lower odds of acute dialysis and 21% lower odds of mortality for patients on a preoperative statin [60]. In a retrospective cohort of 98,939 patients undergoing major open abdominal, cardiac, thoracic, or vascular procedure preoperative statin use was associated with a 20%-26% reduction in the incidence of postoperative acute kidney injury defined by consensus criteria [61]. These retrospective results, and the intriguing pleiotropic actions of statins, have prompted several prospective trials of the effect of statins on perioperative complications.

A number of modifiable intraoperative factors have been identified as risk factors for AKI for both cardiac and non-cardiac surgery, including use of cardiopulmonary bypass (CPB), hemodilution, intraoperative transfusion and hemoglobin levels, hypotension, oxygen delivery and any use of diuretics, vasopressors and inotropes [62-65]. Rewarming after CPB and hyperthermic perfusion during CPB are novel risk factors for AKI after cardiac surgery [66, 67]. A recent systematic review of perioperative interventions aimed to optimize global blood flow showed no difference in mortality, but the rates of renal failure (relative risk (RR) 0.71, 95% confidence interval (CI) 0.57 to 0.90) were reduced [68]. Goal-directed intraoperative management to reduce the risk of postoperative AKI through optimizing renal perfusion is both feasible and underutilized.

[Tags: Preoperative risk factors, surgical populations, scoring systems, cardiac surgery, trauma surgery, transplant surgery, Model for End-Stage Liver Disease score, MELD, Norton scale, albuminuria, estimated glomerular filtration rate, eGFR, preoperative medications, statins, intraoperative factors, goal-directed management]

Risk Stratification for AKI

Given that AKI is common after surgery, and associated with significant morbidity and cost, the ability to detect AKI within hours of onset would likely be helpful in implementing measures to protect the kidney from further injury and to preserve renal function. In the presence of common clinical risk factors for perioperative AKI, such as hypotension or hypoxemia or certain comorbidities, it is difficult to discriminate those who will imminently develop AKI from those who will not before change in creatinine or urine output is measurable. Widespread adoption of effective preventive interventions is only likely with the assistance of a test or tests that can be quickly done at the bedside and that reliably discern those patients at increased risk for AKI from those at low risk.

Use of imaging techniques

Ultrasound and Doppler imaging of the kidney has been used for years in the assessment of CKD in the transplanted kidney and in renal artery stenosis. Doppler imaging detects macroscopic vascular abnormalities as well as microvascular changes in blood flow in the kidney. Renal resistive index (RRI), determined by Doppler ultrasonography, quantifies changes in renal vascular resistance and recent studies have shown that an elevated RRI is associated with an increased risk for AKI [69]. In patients with septic shock an increased RRI has been shown to be associated with AKI [70-72]. The RRI, when used in the immediate postoperative period after cardiac surgery with cardiopulmonary bypass, predicts delayed AKI and its severity [73]. When measured using intraoperative trans-esophageal echocardiography, for patients undergoing cardiac surgery, the predictive results for AKI are comparable to RRI obtained via translumbar ultrasound [74]. The association between elevated RRI and AKI holds for non-cardiac surgeries like orthopedic surgeries and in critically ill patients in the medical ICU [75, 76]. Renal resistive index has also been shown to be helpful in predicting the progression of postoperative AKI after cardiac surgery [77]. While RRI is closely related to renal vascular resistance, it has become clear in recent studies that there are other factors that affect RRI especially in the unstable patient, and the indications and utility of the test for predicting AKI are still unresolved [78, 79]. Another

newly developed technique for assessing renal microvasculature is contrast enhanced ultrasonography (CEUS) to assess renal perfusion [80]. A recent study of patients undergoing elective cardiac surgery, who were considered to be at risk of AKI and were studied with CEUS, showed that renal perfusion decreased within 24 hours after surgery. The technique of CEUS has been validated in the quantification of microcirculatory flow in the liver and the heart, and shows early promise in assessing the risk of AKI [81].

Magnetic resonance imaging (MRI) is another emerging modality in the early diagnosis of AKI. Newer and less toxic contrast agents, including ultra-small particles of iron oxide (USPIO), are being studied for imaging renal blood flow and volume [82]. Blood oxygenation level dependent (BOLD) MRI, which uses deoxyhemoglobin as an endogenous contrast agent for the noninvasive assessment of tissue oxygen bioavailability, has been used to evaluate intrarenal oxygenation [83]. Changes in medullary blood flow related to the use of nephrotoxic agents including nonsteroidal anti-inflammatory drugs, intravenous contrast agents and calcineurin inhibitors are effectively demonstrated with BOLD MRI [84]. It has recently been used to study changes in medullary blood flow associated with hypertension and chronic kidney disease [85-87]. BOLD MRI has been used to study renal oxygenation and function in animal models of AKI [88, 89], however one recent study using BOLD MRI in patients with AKI found no correlation between MRI findings and GFR [90]. As with the Doppler ultrasound derived RRI, the indications and utility of BOLD MRI for predicting AKI in surgical patients are still unresolved.

Use of urine and plasma biomarkers

Another approach for rapid diagnosis of organ injury is the analysis of serum and/or urine biomarkers that reveal early evidence of cellular stress or injury. Biomarkers for cardiac injury, such as serum troponin, are useful in the evaluation and treatment of patients with chest pain because they help clinicians identify those patients who have undergone recent myocardial injury. Many serum and urine biomarkers have been studied for their ability to predict AKI (reviewed in [91-95]). Difficulties in finding biomarkers with good predictive power include the variable time from insult to the development of AKI, the association of biomarkers with both CKD and AKI, and their association with the diverse underlying disease processes that can cause AKI. Furthermore, surgical and critically ill patients are exposed to systemic inflammation with cellular stress and injuries in several organs, repetitive exposure to invasive procedures and hemodynamic perturbations requiring fluid therapy and vasopressor support, blood transfusion and nephrotoxic drugs. The abundance of these potential mediators and confounders of AKI may cause non-specific increases in biomarkers reflecting overall illness severity rather than specific organ damage.

Among surgical patients, the use of biomarkers for AKI prediction was most studied after cardiac procedures with the most promising results demonstrated for the use of plasma and urine neutrophil gelatinase-associated lipocalin (NGAL) (reviewed in [95, 96]). Systemic inflammation induces NGAL synthesis by extrarenal tissues and the release of NGAL from neutrophils mainly in the dimeric form. A recent systematic review summarized studies that measured NGAL in more than 7000 patients after cardiac surgery, and showed moderate overall discriminatory ability with area under the receiver operating characteristic curve

(ROC-AUC) between 0.82 and 0.83. Studies including more than 8500 critically ill patients, mostly recruited from mixed medical/surgical ICU populations, demonstrated similar overall predictive performance with ROC-AUC between 0.79 and 0.80 [97, 98]. The inability to distinguish systemic inflammatory effects from organ-specific increases in NGAL, and the lack of a diagnostic platform to differentiate specific biological forms of NGAL has hampered widespread clinical use of this biomarker.

Cystatin C (CyC) is the most studied novel functional kidney biomarker [93]. This cysteine protease inhibitor is produced by all nucleated cells of the body, released into the bloodstream at a constant rate, excreted through glomerular filtration into primary urine and subsequently completely reabsorbed and catabolized in proximal renal tubules. As a consequence, CyC is not normally found in urine in significant amounts and urinary CyC may reflect tubular damage. Due to the constant rate of its production, plasma CyC concentrations may be a better marker of GFR than creatinine in surgical patients. However, its ability to provide early risk stratification for postoperative AKI or RRT requirement remains uncertain among surgical patients [93].

Recently two urinary biomarkers, tissue inhibitor of metalloproteinases-2 (TIMP-2) and insulin-like growth factor binding protein 7 (IGFBP7), have been validated as markers of risk for AKI[99, 100]. These markers were recently approved for use by the US Food and Drug Administration, becoming the first AKI biomarkers to do so. A multicenter study involving 420 patients found that urinary [TIMP-2]·[IGFBP7] was predictive of moderate to severe AKI in critically ill patients within 12 hours. Unlike prior studies evaluating these markers or others for AKI[99], the endpoint was adjudicated by a committee of three independent experts who were blinded to the results of the test. The ROC-AUC for the urinary [TIMP-2]·[IGFBP7] test was 0.82 (95% CI 0.76-0.88) and was superior to simultaneously measured serum creatinine and other existing biomarkers for predicting the risk of imminent AKI. Patients with a urinary test result above the pre-specified high sensitivity cutoff value of 0.3 (ng/ml)²/1000 had seven times the risk for AKI (95% CI 4-22). Meersch et al examined the sensitivity and specificity of the [TIMP-2]·[IGFBP7] test for any AKI stage among patients undergoing cardiac surgery and found a sensitivity of 0.92 and specificity of 0.81 for a cutoff value of 0.5 (ROC-AUC of 0.84) using the highest urinary [TIMP-2]·[IGFBP7] achieved in the first 24 hours following surgery [101]. Interestingly TIMP-2 and IGFBP7 are both associated with G1 cell-cycle arrest, an epithelial cellular protective mechanism, rather than with cellular necrosis or apoptosis. Epithelial cells, by virtue of their anatomy and function, are susceptible to multiple environmental stressors: toxin exposure, oxidative stress and inflammation among others. When DNA may be damaged, or when bioenergetic resources are scarce, epithelial cells may enter cell-cycle arrest to protect themselves and TIMP-2 and IGFBP7 become elevated. Thus these urinary biomarkers may indicate risk for injury before any actual AKI takes place.

Use of clinical prediction scores

One very different approach towards early diagnosis and prediction of organ injury is automated analysis of the large amounts of clinical data obtained during routine care to detect critical incidents or trends that might be predictive of injury. Accurate risk

stratification of patients in real time could enable the selection of optimal therapy in a timely fashion to prevent AKI altogether, or to mitigate the effects of the complication even before signs and symptoms arise, and could be tailored to a patients' personal clinical profile. Despite the acquisition of multiple continuously recorded physiological signals during modern perioperative management, the use of this data for the development of risk and prediction models has been limited to the prediction of broad outcomes such as postoperative mortality rather than to specific morbidity events including AKI [102-104]. A majority of the predictive scores and algorithms have been limited either to a specific type of surgery or preoperative risk factors only, or have used the occurrence of the most severe AKI as an end-point while excluding the more prevalent mild and moderate AKI.

Most of the studies that developed and validated predictive models or clinical scores for AKI were performed among cardiac surgery patients. A recent systematic review [36] evaluated the available risk models for AKI after cardiac surgery and reported four clinical risk scores for AKI requiring dialysis [105-108], and three scores to predict a broader definition of AKI [109-111]. These scores predicted a probability of severe AKI between <1% and >20%, with the ROC-AUC varying between 0.77 and 0.84. For patients undergoing non-cardiac surgery a few predictive models, developed in small cohorts utilizing limited intraoperative data or in larger cohorts but using only severe AKI as an end-point, are further limited by the lack of validation studies and provide only modest predictive accuracy [22, 112-114]. Recently UK AKI in Cardiac Surgery Collaborators group have developed and validated a new risk prediction score for any stage AKI after cardiac surgery with ROC-AUC of 0.74, providing better discrimination compared to previously published scores [115].

The volume of physiologic data routinely acquired during intraoperative hemodynamic monitoring is rarely used in published risk scores and when used, it is usually summarized by some reductionist approach (mean, lowest value, etc) rather than applied in their continuity and complexity and almost never in automatized fashion [13, 22, 106, 112, 113, 116-121]. Lack of sophistication in both data collection and analysis of real time physiologic data has limited this approach. Automated risk scores, developed using machine learning to automatically analyze physiological time series data, have already been shown to predict neonatal clinical outcomes more accurately than can be achieved with any pre-existing scoring system [122]. Machine learning applied to the automated, rapid, noninvasive measurements obtained in the operating room and ICU raises the prospect of real time risk prediction for perioperative AKI and studies utilizing them are starting to emerge [35, 123-125].

[Tags: Imaging, ultrasound, Doppler, renal resistive index, RRI, contrast enhanced ultrasound, CEUS, blood oxygenation level dependent MRI, BOLD MRI, urine and plasma biomarkers, clinical prediction scores, automated analysis, predictive models, real time physiologic data, automated risk scores, machine learning]

Prevention and Treatment of Perioperative AKI

One of the challenges in managing AKI is the paucity of interventions to treat it once it occurs. Given that reality, prevention of AKI is of paramount importance. Many interventions have been studied in an attempt to prevent or ameliorate perioperative AKI [13, 15, 16]. Although patients with CKD have a higher risk for adverse perioperative events, preoperative optimization of renal function using pharmacological therapy such as angiotensin converting enzyme inhibitors (ACE-I), diuretic therapy, or regular visits to a nephrologist to prevent a decline in kidney function around the time of surgery were not proven to mitigate the risk of AKI [113, 126]. Some writers advocate avoidance of ACE-I or angiotensin receptor blocker therapies around the time of surgery especially when hypotension is anticipated [126, 127]. Multiple pharmacologic therapies have been unsuccessfully tested for the prevention of perioperative AKI, including scavengers of oxygen free radicals such as mannitol and N-acetylcysteine, dopamine, fenoldapam, loop diuretics and atrial natriuretic peptide (reviewed in [13, 15, 16, 128]). A recent systematic review and meta-analysis including 1079 patients in five randomized control trials demonstrated no benefit for the prophylactic perioperative use of sodium bicarbonate for prevention of AKI after cardiac surgery. In contrast the use of sodium bicarbonate prolonged the duration of mechanical ventilation and ICU length of stay, and increased the risk of alkalemia [129]. A recent large randomized clinical trial of patients undergoing noncardiac surgery found that neither aspirin nor clonidine administered perioperatively reduced the risk of AKI (13.4% for aspirin vs 12.3% for placebo; 13.0% for clonidine vs 12.7% for placebo), whereas both aspirin and clonidine were associated with clinically important adverse effects [130].

Some aspects of perioperative management do appear to have an important effect on kidney function. The use of off-pump technique in cardiac surgery has demonstrated benefit in randomized control trials and meta-analysis [116]. Two large meta-analyses and a systematic review have showed that intraoperative interventions associated with goal-directed fluid management were associated with a significant reduction in the incidence of all severity stages of AKI [68, 131]. Initiating appropriate hemodynamic monitoring to allow the anesthesiologist to optimize intravascular volume, cardiac output or oxygen delivery in high risk patients resulted in a decreased risk of perioperative AKI both if started preoperatively (odds ratio 0.70, 95% CI 0.53 to 0.94; $P=0.02$) or intraoperatively (OR 0.47, 95% CI 0.27 to 0.81; $P=0.006$) [131]. One contentious and unresolved issue in goal directed fluid management and resuscitation is the optimal endpoint and how to measure it. An optimal mean arterial pressure (MAP) for the kidney at risk for AKI is unknown, and may be different from that obtained peripherally [81]. The ability to assess renal perfusion at the bedside using Doppler ultrasound may provide better fluid and vasoactive medication management for the unstable patient at risk for AKI. One study in patients with septic shock showed that RRI decreased significantly when MAP was increased using norepinephrine from 65 to 75 mmHg [70]. Another prospective study of patients who had experienced sustained hypotension showed that an average MAP between 72 and 82 mmHg during the first three days after hypotension was associated with a lower incidence of RIFLE-AKI compared to patients with an average MAP below 72mmHg [132]. Further studies exploring

the optimal resuscitation endpoints for the kidney to prevent AKI, and how to measure those endpoints, are needed. Another interesting meta-analysis involving 1600 patients in ten trials demonstrated that volatile anesthetics may provide renal protection in patients undergoing cardiac surgery and supports the notion that further research of high methodologic quality is needed to define optimal intraoperative management of patients at high risk of AKI [133].

Another evolving concept is the use of remote ischemic preconditioning to prevent AKI in certain patient populations. Ischemia reperfusion injury occurs whenever a tissue bed becomes temporarily ischemic and is then restored to normal perfusion. The kidneys are particularly sensitive to Ischemia reperfusion injury due to their high metabolic and oxygen demands and complex microvasculature [69]. However, in addition to causing injury, programmed brief and intermittent ischemia is known to have cytoprotective effects. Local ischemic preconditioning has been shown to be effective against various types of AKI [134, 135]. Remote ischemic preconditioning, involving programmed brief and repeated ischemia of a remote tissue such as limb skeletal muscle, has been shown to be as effective as local ischemic preconditioning in preventing cellular damage. It is thought to attenuate injury through up regulation of a variety of intracellular kinases, resulting in modification of mitochondrial function, metabolic down-regulation and temporary cell-cycle arrest [136]. Recently in a randomized controlled trial remote ischemic preconditioning was found to be protective against contrast medium induced AKI for patients undergoing elective coronary angiography and who were judged to be at high risk for AKI [134]. Similar results were found in a randomized controlled trial of 120 patients undergoing elective cardiac surgery, in which the patients randomized to RIPC had significantly less risk of postoperative AKI [130]. While these early trials provide the promise of a novel, noninvasive and virtually morbidity-free therapy to prevent AKI, further investigation is needed to define the indications and utility of this approach.

One important potential method of limiting the consequences of AKI in surgical patients is early and continued involvement of a nephrologist. A study from two hospitals in the United Kingdom with no on-site nephrology services showed that, compared to hospitals with nephrology consultation available, there were significant shortcomings in AKI recognition and management which were associated with poor survival and increased rates of CKD [137]. In prospective observational studies of the patient admitted to the ICU it has been shown that delayed nephrology consultation was associated with higher mortality [138, 139] and increased dialysis dependence rates at hospital discharge [139]. In a recent prospective controlled nonrandomized study patients in an early nephrology consultation group (seen within 18 hours of onset of AKI) had significantly lower risk of further decrease in kidney function [140].

Nephrology referral has also been shown to be important in follow up care after an episode of AKI. A study of patients who sustained AKI between 2003 and 2008 in a Veterans Administration hospital, and were considered to be at risk for subsequent worsening of renal function, showed that the cumulative incidence of outpatient nephrology referrals for these patients was only 8.5% [141]. Another more recent study of Veterans Administration patients admitted to the hospital with AKI showed that measurement of serum creatinine during outpatient follow-up was common, but measurement of proteinuria, parathyroid

hormone or serum phosphorus was rare [142]. The importance of this low referral rate is emphasized in a cohort study of hospitalized adults with AKI who received temporary inpatient dialysis and survived for 90 days following discharge, in which patients with early nephrology follow-up had significantly lower all-cause mortality [143].

Given the association between AKI and the later development of CKD and other complications, follow up care for patients who sustain AKI in the hospital could have important public health and socioeconomic impact [144, 145]. In 2012 the KDIGO AKI workgroup released guidelines recommending that patients be evaluated 3 months after AKI for the new onset of CKD or worsening of any pre-existing CKD [146]. Patients with CKD are to be managed according to the Kidney Disease Outcome Quality Initiative (KDOQI) CKD guidelines, and patients with no CKD are to be managed according to the KDOQI guidelines for patients “at risk” for CKD. Finally, in addition to nephrology consultation for the patient who has sustained AKI, it is imperative that primary care practitioners understand the risks associated with even mild degrees of AKI suffered by their patients, both to initiate timely nephrology involvement and to optimally manage patients at risk for the development of CKD [147-149]. The current attempts at better coordinating care for the patient with chronic disease, including the patient-centered medical home and accountable-care organizations with robust electronic medical recordkeeping, may help improve the care of patients how have sustained AKI [150-152].

[Tags: prevention, preoperative optimization, pharmacological therapy, off-pump technique, hemodynamic monitoring, renal perfusion, optimal resuscitation endpoints, remote ischemic preconditioning, nephrologist, nephrology referral, follow up care, care coordination]

Summary

Acute and chronic kidney injury and dysfunction play important roles in affecting perioperative outcomes. AKI is a common complication after surgery and mild to moderate AKI is more common than severe AKI. All stages of AKI severity are associated with increased short and long term morbidity and mortality. Clinical risk factors for AKI are similar but not identical in different surgical populations. There appears to be no single therapy that will prevent perioperative AKI. Considering the high prevalence of AKI and the deleterious effect when it occurs, our effort must focus on prevention of AKI, mitigation of further injury when AKI has already occurred, treatment of negative effects on other organs, and facilitation of renal recovery in patients with established AKI. This clinical pathway requires a medical team of experts, including all primary healthcare providers who manage surgical and critically ill patients, backed up by bedside nurses, pharmacists and nephrologists. Every patient admitted needs a comprehensive and systematic assessment of kidney health following a new paradigm. Current strategies should focus on better management of the preoperative risks and susceptibilities for AKI by more accurate assessment of the patient's renal reserve and susceptibility to new injury. Standardized approach for intraoperative management for patients at high risk for AKI need to focus on avoidance of hemodynamic derangements that have been shown to impact renal function. In the early postoperative period the magnitude of exposures to insult and the extent of the sustained renal distress or damage need to be evaluated using a combination of clinical

parameters, novel biomarkers and evolving imaging techniques. The determination of potential causes of AKI, the initiation of treatment and then continued reassessment in response to that therapy should follow promptly afterwards with the early involvement of nephrology teams.

Acknowledgments

Conflicts of Interest and Source of Funding: AB is supported by Center for Sepsis and Critical Illness Award P50 GM-111152 from the National Institute of General Medical Sciences and has received research grants from Society of Critical Care Medicine and Astute Medical, Inc.

References

1. Hobson C, Ozrazgat-Baslanti T, Kuxhausen A, et al. Cost and Morality Associated With Postoperative Acute Kidney Injury. *Ann Surg.* 2014
2. Bihorac A, Delano MJ, Schold JD, et al. Incidence, clinical predictors, genomics, and outcome of acute kidney injury among trauma patients. *Ann Surg.* 2010; 252:158–65. [PubMed: 20562612]
3. Hobson CE, Yavas S, Segal MS, et al. Acute kidney injury is associated with increased long-term mortality after cardiothoracic surgery. *Circulation.* 2009; 119:2444–53. [PubMed: 19398670]
4. Bihorac A, Yavas S, Subbiah S, et al. Long-term risk of morality and acute kidney injury during hospitalization after major surgery. *Ann Surg.* 2009; 249:851–8. [PubMed: 19387314]
5. Wald R, Quin R, Luo J. Chronic dialysis and death among survivors of acute kidney injury requiring dialysis. *JAMA.* 2009; 302:1179–85. [PubMed: 19755696]
6. van Kuijk JP, Flu WJ, Chonchol M, et al. Temporary perioperative decline of renal function is an independent predictor for chronic kidney disease. *Clin J Am Soc Nephrol.* 2010; 5:1198–204. [PubMed: 20430939]
7. Ishani A, Nelson D, Clothier B, et al. The magnitude of acute serum creatinine increase after cardiac surgery and the risk of chronic kidney disease, progression of kidney disease, and death. *Arch Intern Med.* 2011; 171:226–33. [PubMed: 21325112]
8. James MT, Ghali WA, Knudtson ML, et al. Associations between acute kidney injury and cardiovascular and renal outcomes after coronary angiography. *Circulation.* 2011; 123:409–16. [PubMed: 21242477]
9. Thakar CV, Christianson A, Himmelfarb J, et al. Acute kidney injury episodes and chronic kidney disease risk in diabetes mellitus. *Clin J Am Soc Nephrol.* 2011; 6:2567–72. [PubMed: 21903988]
10. Coca SG, Singanamala S, Parikh CR. Chronic kidney disease after acute kidney injury: a systematic review and meta-analysis. *Kidney Int.* 2012; 81:442–8. [PubMed: 22113526]
11. Chawla LS, Amdur RL, Shaw AD, et al. Association between AKI and long-term renal and cardiovascular outcomes in United States veterans. *Clin J Am Soc Nephrol.* 2014; 9:448–56. [PubMed: 24311708]
12. Bihorac A, Brennan M, Ozrazgat Baslanti T, et al. National Surgical Quality Improvement Program Underestimates the Risk Associated with Mild and Moderate Postoperative Acute Kidney Injury. *Crit Care Med.* 2013; 41:2570–83. [PubMed: 23928835]
13. Borthwick E, Ferguson A. Perioperative acute kidney injury: risk factors, recognition, management, and outcomes. *BMJ.* 2010; 341:c3365. [PubMed: 20603317]
14. Vaught A, Ozrazgat-Baslanti T, Javed A, et al. Acute kidney injury in major gynaecological surgery: an observational study. *BJOG.* 2014
15. Calvert S, Shaw A. Perioperative acute kidney injury. *Perioper Med (Lond).* 2012; 1:6. [PubMed: 24764522]
16. Thakar CV. Perioperative Acute Kidney Injury. *Adv Chronic Kidney Dis.* 2013; 20:67–75. [PubMed: 23265598]
17. Hoste EA, Kellum JA. Incidence, classification, and outcomes of acute kidney injury. *Contrib Nephrol.* 2007; 156:32–8. [PubMed: 17464113]

18. Ricci Z, Cruz DN, Ronco C. Classification and staging of acute kidney injury: beyond the RIFLE and AKIN criteria. *Nat Rev Nephrol.* 2011; 7:201–8. [PubMed: 21364520]
19. Bellomo R, Ronco C, Kellum JA, et al. Acute renal failure - definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care.* 2004; 8:R204–12. [PubMed: 15312219]
20. KDOQI. Clinical Practice Guideline for Acute Kidney Injury: AKI Definition. *Kidney Int.* 2012; 2:19–36.
21. American College of Surgeons National Surgical Quality Improvement Program. User Guide for the 2010 Participant Use Data File. Chicago, IL 60611-3211: American College of Surgeons; 2010.
22. Kheterpal S, Tremper K, Heung M, et al. Development and validation of an acute kidney injury risk index for patients undergoing general surgery: results from a national data set. *Anesthesiology.* 2009; 110:505–15. [PubMed: 19212261]
23. Arnaoutakis GJ, Bihorac A, Martin TD, et al. RIFLE criteria for acute kidney injury in aortic arch surgery. *J Thorac Cardiovasc Surg.* 2007; 134:1554–61. [PubMed: 18023682]
24. Kundakci A, Pirat A, Komurcu O, et al. Rife criteria for acute kidney dysfunction following live transplantation: incidence and risk factors. *Transplant Proc.* 2010; 42:4171–4. [PubMed: 21168655]
25. Bihorac A, Schold JD, Hobson CE. Long-term morality associated with acute kidney injury requiring dialysis. *JAMA.* 2010; 303:229. author reply -30. [PubMed: 20085947]
26. Ozrazgat Baslanti T, Kuxhausen A, Hobson CE, et al. Effect of Postoperative Acute Kidney Injury on Hospital Cost. *Crit Care Med.* 2012; 40
27. Dimick JB, Pronovost PJ, Cowan JA, et al. Complications and costs after high-risk surgery: where should we focus quality improvement initiatives? *J Am Col Surg.* 2003; 196:671–8.
28. Dimick JB, Chen SL, Taheri PA, et al. Hospital costs associated with surgical complications: a report from the private-sector National Surgical Quality Improvement Program. *J Am Col Surg.* 2004; 199:531–7.
29. Thakar CV, Christianson A, Freyberg R, et al. Incidence and outcomes of acute kidney injury intensive care units: A Veterans Administration study*. *Crit Care Med.* 2009; 37:2552–8. [PubMed: 19602973]
30. Duran PA, Concepcion LA. Survival after acute kidney injury requiring dialysis: Long-term flow up. *Hemodial Int.* 2014; 18:S1–S6. [PubMed: 25330825]
31. Coca SG, Yusuf B, Shlipak MG, et al. Long-term risk of mortality and other adverse outcomes after acute kidney injury: a systematic review and meta-analysis. *Am J Kidney Dis.* 2009; 53:961–73. [PubMed: 19346042]
32. Lafrance JP, Miller DR. Acute kidney injury associates with increased long-term morality. *J Am Soc Nephrol.* 2010; 21:345–52. [PubMed: 20019168]
33. Amdur RL, Chawla LS, Amodeo S, et al. Outcomes following diagnosis of acute renal failure in U.S. veterans: focus on acute tubular necrosis. *Kidney Int.* 2009; 76:1089–97. [PubMed: 19741590]
34. Wald R, Quinn RR, Luo J, et al. Chronic dialysis and death among survivors of acute kidney injury requiring dialysis. *JAMA.* 2009; 302:1179–85. [PubMed: 19755696]
35. Ozrazgat Baslanti T, Korenkevych D, Momcilovic P, et al. Mathematical modeling of the association between the pattern of change in postoperative serum creatinine and hospital morality. *Crit Care Med.* 2012; 40
36. Huen SC, Parikh CR. Predicting acute kidney injury after cardiac surgery: a systematic review. *Ann Thorac Surg.* 2012; 93:337–47. [PubMed: 22186469]
37. Lomivorotov V, Efremov SM, Boboshko VA, et al. Preoperative total lymphocyte count in peripheral blood as a predictor of poor outcome in adult cardiac surgery. *J Cardiothorac Vasc Anesth.* 2011; 25:975–80. [PubMed: 21354824]
38. Kim DH, Shim JK, Hong SW, et al. Predictive value of C-reactive protein for major postoperative complications following off-pump coronary artery bypass surgery: prospective and observational trial. *Circ J.* 2009; 73:872–7. [PubMed: 19282606]

39. Stafford-Smith M, Podgoreanu M, Swaminathan M, et al. Association of genetic polymorphisms with risk of renal injury after coronary bypass graft surgery. *Am J Kidney Dis.* 2005; 45:519–30. [PubMed: 15754274]
40. van Kuijk JP, Flu WJ, Valentijn TM, et al. Preoperative left ventricular dysfunction predispose to postoperative acute kidney injury and long-term morality. *J Nephrol.* 2011; 24:764–70. [PubMed: 21360471]
41. Romano TG, Schmidtbauer I, Silva FM, et al. Role of MELD score and serum creatinine as prognostic tools for the development of acute kidney injury after liver transplantation. *PLoS One.* 2013; 8:e64089. [PubMed: 23717537]
42. Asleh K, Sever R, Hilu S, et al. Association between low admission Norton scale scores and postoperative complications after elective THA in elderly patients. *Orthopedics.* 2012; 35:e1302–6. [PubMed: 22955393]
43. Aveline C, Leroux A, Vautier P, et al. Risk factors for renal dysfunction after total hip arthroplasty. *Ann Fr Anesth Reanim.* 2009; 28:728–34. [PubMed: 19709848]
44. Brooks CE, Middleton A, Dhillon R, et al. Predictors of creatinine rise post-endovascular abdominal aortic aneurysm repair. *ANZ J Surg.* 2011; 81:827–30. [PubMed: 22295426]
45. Lamb EJ, Levey AS, Stevens PE. The Kidney Disease Improving Global Outcomes (KDIGO) guideline update for chronic kidney disease: evolution not revolution. *Clin Chem.* 2013; 59:462–5. [PubMed: 23449698]
46. Matsushita K, van der Velde M, Astor BC, et al. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular morality in general population cohorts: a collaborative meta-analysis. *The Lancet.* 2010; 375:2073–81.
47. Mathew A, Devereaux PJ, O'Hare A, et al. Chronic kidney disease and postoperative morality: a systematic review and meta-analysis. *Kidney Int.* 2008; 73:1069–81. [PubMed: 18288098]
48. Gaber AO, Moore LW, Aloia TA, et al. Cross-sectional and case-control analyses of the association of kidney function staging with adverse postoperative outcomes in general and vascular surgery. *Ann Surg.* 2013; 258:169–77. [PubMed: 23478526]
49. Wu VC, Huang TM, Wu PC, et al. Preoperative proteinuria is associated with long-term progression to chronic dialysis and morality after coronary artery bypass grafting surgery. *PLoS One.* 2012; 7:e27687. [PubMed: 22276092]
50. Huang TM, Wu VC, Young GH, et al. Preoperative Proteinuria Predicts Adverse Renal Outcomes after Coronary Artery Bypass Grafting. *J Am Soc Nephrol.* 2011; 22:156–63. [PubMed: 21115618]
51. Fleisher LA, Eagle KA. Clinical practice. Lowering cardiac risk in noncardiac surgery. *N Engl J Med.* 2001; 345:1677–82. [PubMed: 11759647]
52. Kertai MD, Boersma E, Klein J, et al. Optimizing the prediction of perioperative mortality in vascular surgery by using a customized probability model. *Arch Intern Med.* 2005; 165:898–904. [PubMed: 15851641]
53. Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation.* 1999; 100:1043–9. [PubMed: 10477528]
54. Detsky AS, Abrams HB, McLaughlin JR, et al. Predicting cardiac complications in patients undergoing non-cardiac surgery. *J Gen Intern Med.* 1986; 1:211–9. [PubMed: 3772593]
55. Goldman L, Caldera DL, Nussbaum SR, et al. Multifactorial index of cardiac risk in noncardiac surgical procedures. *N Engl J Med.* 1977; 297:845–50. [PubMed: 904659]
56. Levey AS, Stevens LA, Schmid CH, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med.* 2009; 150:604–12. [PubMed: 19414839]
57. Davignon J. Beneficial cardiovascular pleiotropic effects of statins. *Circulation.* 2004; 109:III39–43. [PubMed: 15198965]
58. Liakopoulos OJ, Kuhn EW, Slottosch I, et al. Preoperative statin therapy for patients undergoing cardiac surgery. *Cochrane Database Syst Rev.* 2012; 4:CD008493.
59. Kor DJ, Brown MJ, Iscimen R, et al. Perioperative statin therapy and renal outcomes after major vascular surgery: a propensity-based analysis. *J Cardiothorac Vasc Anesth.* 2008; 22:210–6. [PubMed: 18375322]

60. Molnar AO, Coca SG, Devereaux PJ, et al. Statin Use Associates with a Lower Incidence of Acute Kidney Injury after Major Elective Surgery. *J Am Soc Nephrol.* 2011; 22:939–46. [PubMed: 21493769]
61. Brunelli SM, Waikar S, Bateman BT, et al. Preoperative Statin Use and Postoperative Acute Kidney Injury. *Am J Med.* 2012; 125:1195–+. [PubMed: 23062398]
62. Karkouti K, Wijeyesundera DN, Yau TM, et al. Acute kidney injury after cardiac surgery: focus on modifiable risk factors. *Circulation.* 2009; 119:495–502. [PubMed: 19153273]
63. Haase M, Bellomo R, Story D, et al. Effect of mean arterial pressure, haemoglobin and blood transfusion during cardiopulmonary bypass on post-operative acute kidney injury. *Nephrology Dialysis Transplantation.* 2012; 27:153–60.
64. de Somer F, Mulholland J, Bryan M, et al. O₂ delivery and CO₂ production during cardiopulmonary bypass as determinants of acute kidney injury: time for a goal-directed perfusion management? *Critical Care.* 2011; 15:R192. [PubMed: 21831302]
65. Parolari A, Pesce LL, Pacini D, et al. Risk factors for perioperative acute kidney injury after adult cardiac surgery: role of perioperative management. *Ann Thorac Surg.* 2012; 93:584–91. [PubMed: 22269725]
66. Newland RF, Tully PJ, Baker RA. Hyperthermic perfusion during cardiopulmonary bypass and postoperative temperature are independent predictors of acute kidney injury following cardiac surgery. *Perfusion-Uk.* 2013; 28:223–31.
67. Boodhwani M, Rubens FD, Wozny D, et al. Effects of mild hypothermia and rewarming on renal function after coronary artery bypass grafting. *Ann Thorac Surg.* 2009; 87:489–95. [PubMed: 19161766]
68. Grocott Michael, PW.; Dushianthan, A.; Hamilton Mark, A., et al. *Cochrane Database Syst Rev.* John Wiley & Sons, Ltd; 2012. Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery.
69. Ninet S, Schnell D, Dewitte A, et al. Doppler-based renal resistive index for prediction of renal dysfunction reversibility: A systematic review and meta-analysis. *J Crit Care.* 2015
70. Derudder S, Cheisson G, Mazoit JX, et al. Renal arterial resistance in septic shock: effects of increasing mean arterial pressure with norepinephrine on the renal resistive index assessed with Doppler ultrasonography. *Intensive Care Med.* 2007; 33:1557–62. [PubMed: 17486316]
71. Lerolle N, Guerot E, Faisy C, et al. Renal failure in septic shock: predictive value of Doppler-based renal arterial resistive index. *Intensive Care Med.* 2006; 32:1553–9. [PubMed: 16941165]
72. Gornik I, Godan A, Gašparovi V. Renal resistive index at ICU admission and its change after 24 hours predict acute kidney injury in sepsis. *Critical Care.* 2014; 18:P366.
73. Bossard G, Bourgoin P, Corbeau JJ, et al. Early detection of postoperative acute kidney injury by Doppler renal resistive index in cardiac surgery with cardiopulmonary bypass. *Br J Anaesth.* 2011; 107:891–8. [PubMed: 21940396]
74. Kararmaz A, Kemal Arslantas M, Cinel I. Renal Resistive Index Measurement by Transesophageal Echocardiography: Comparison With Translumbar Ultrasonography and Relation to Acute Kidney Injury. *J Cardiothorac Vasc Anesth.*
75. Darmon M, Schortgen F, Vargas F, et al. Diagnostic accuracy of Doppler renal resistive index for reversibility of acute kidney injury in critically ill patients. *Intensive Care Med.* 2011; 37:68–76. [PubMed: 20862450]
76. Marty P, Szatjnic S, Ferre F, et al. Doppler renal resistive index for early detection of acute kidney injury after major orthopaedic surgery: a prospective observational study. *Eur J Anaesthesiol.* 2015; 32:37–43. [PubMed: 25014511]
77. Guinot PG, Bernard E, Abou Arab O, et al. Doppler-based renal restive index can assess progression of acute kidney injury in patients undergoing cardiac surgery. *J Cardiothorac Vasc Anesth.* 2013; 27:890–6. [PubMed: 23731713]
78. Viazi F, Leocini G, Derchi LE, et al. Ultrasound Doppler renal restive index: a useful tool for the management of the hypertensive patient. *J Hypertens.* 2014; 32:149–53. [PubMed: 24172238]
79. Dewitte A, Coquin J, Meyssignac B, et al. Doppler resistive index to reflect regulation of renal vascular tone during sepsis and acute kidney injury. *Crit Care.* 2012; 16:R165. [PubMed: 22971333]

80. Mahoney M, Sorace A, Waram J, et al. Volumetric contrast-enhanced ultrasound imaging of renal perfusion. *J Ultrasound Med*. 2014; 33:1427–37. [PubMed: 25063408]
81. Harrois A, Duranteau J. Contrast-enhanced ultrasound: a new vision of microcirculation in the intensive care unit. *Critical Care*. 2013; 17:449. [PubMed: 24103639]
82. Choyke P, Kobayashi H. Functional magnetic resonance imaging of the kidney using macromolecular contrast agents. *Abdom Imaging*. 2006; 31:224–31. [PubMed: 16283582]
83. Prasad PV, Edelman RR, Epstein FH. Noninvasive evaluation of intrarenal oxygenation with BOLD MRI. *Circulation*. 1996; 94:3271–5. [PubMed: 8989140]
84. Hofmann L, Simon-Zoula S, Nowak A, et al. BOLD-MRI for the assessment of renal oxygenation in humans: acute effect of nephrotoxic xenobiotics. *Kidney Int*. 2006; 70:144–50. [PubMed: 16641929]
85. Vink E, Boer A, Verloop W, et al. The effect of renal denervation on kidney oxygenation as determined by BOLD MRI in patients with hypertension. *Eur Radiol*. 2015:1–9. [PubMed: 25182628]
86. Vink EE, de Boer A, Hoogduin HJ, et al. Renal BOLD-MRI relates to kidney function and activity of the renin-angiotensin-aldosterone system in hypertensive patients. *J Hypertens*. 2015
87. Pruijm M, Hofmann L, Piskunowicz M, et al. Determinants of renal tissue oxygenation as measured with BOLD-MRI in chronic kidney disease and hypertension in humans. *PLoS One*. 2014; 9:e95895. [PubMed: 24760031]
88. Oostendorp M, de Vries EE, Slenter JMGM, et al. MRI of renal oxygenation and function after normothermic ischemia-reperfusion injury. *NMR Biomed*. 2011; 24:194–200. [PubMed: 20954164]
89. Li LP, Lu J, Zhou Y, et al. Evaluation of Intrarenal Oxygenation in Iodinated Contrast-Induced Acute Kidney Injury -Susceptible Rats by Blood Oxygen Level -Dependent Magnetic Resonance Imaging. *Invest Radiol*. 2014; 49:403–10. [PubMed: 24566288]
90. Inoue T, Kozawa E, Okada H, et al. Noninvasive evaluation of kidney hypoxia and fibrosis using magnetic resonance imaging. *J Am Soc Nephrol*. 2011; 22:1429–34. [PubMed: 21757771]
91. Vanmassenhove J, Vanholder R, Nagler E, et al. Urinary and serum biomarkers for the diagnosis of acute kidney injury: a in-depth review of the literature. *Nephrol Dial Transplant*. 2013; 28:254–73. [PubMed: 23115326]
92. Osterman M, Philips BJ, Forni LG. Clinical review: Biomarkers of acute kidney injury: where are we now? *Crit Care*. 2012; 16:233. [PubMed: 23014769]
93. Wasung ME, Chawla LS, Madero M. Biomarkers of renal function, which and when? *Clin Chim Acta*. 2015; 438:350–7. [PubMed: 25195004]
94. Charlton JR, Portilla D, Okusa MD. A basic science view of acute kidney injury biomarkers. *Nephrology Dialysis Transplantation*. 2014; 29:1301–11.
95. Koyner JL, Parikh CR. Clinical Utility of Biomarkers of AKI in Cardiac Surgery and Critical Illness. *Clin J Am Soc Nephro*. 2013; 8:1034–42.
96. Martensson J, Bellomo R. The rise and fall of NGAL in acute kidney injury. *Blood Purif*. 2014; 37:304–10. [PubMed: 25170751]
97. Haase M, Bellomo R, Devarajan P, et al. Accuracy of neutrophil gelatinase-associated lipocalin (NGAL) in diagnosis and prognosis in acute kidney injury: a systematic review and meta-analysis. *Am J Kidney Dis*. 2009; 54:1012–24. [PubMed: 19850388]
98. Haase-Fielitz A, Haase M, Devarajan P. Neutrophil gelatinase-associated lipocalin as a biomarker of acute kidney injury: a critical evaluation of current status. *Ann Clin Biochem*. 2014; 51:335–51. [PubMed: 24518531]
99. Kashani K, Al-Khafaji A, Ardiles T, et al. Discovery and validation of cell cycle arrest biomarkers in human acute kidney injury. *Crit Care*. 2013; 17:R25. [PubMed: 23388612]
100. Bihorac A, Chawla LS, Shaw AD, et al. Validation of Cell-Cycle Arrest Biomarkers for Acute Kidney Injury Using Clinical Adjudication. *Am J Respir Crit Care Med*. 2014
101. Meersch M, Schmidt C, Van Aken H, et al. Urinary TIMP-2 and IGFBP7 as early biomarkers of acute kidney injury and renal recovery following cardiac surgery. *PLoS One*. 2014; 9:e93460. [PubMed: 24675717]

102. Lake AP, Williams EG. ASA classification and perioperative variables: graded anaesthesia score? *Br J Anaesth.* 1997; 78:228–9. [PubMed: 9068348]
103. Copeland GP, Jones D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg.* 1991; 78:355–60. [PubMed: 2021856]
104. Gawande A, Kwaan MR, Regenbogen SE, et al. An Apgar score for surgery. *J Am Col Surg.* 2007; 204:201–8.
105. Chertow GM, Lazarus JM, Christiansen CL, et al. Preoperative Renal Risk Stratification. *Circulation.* 1997; 95:878–84. [PubMed: 9054745]
106. Thakar CV, Arrigain S, Worley S, et al. A clinical score to predict acute renal failure after cardiac surgery. *J Am Soc Nephrol.* 2005; 16:162–8. [PubMed: 15563569]
107. Mehta RH, Grab JD, O'Brien SM, et al. Beside tool for predicting the risk of postoperative dialysis in patients undergoing cardiac surgery. *Circulation.* 2006; 114:2208–16. quiz. [PubMed: 17088458]
108. Wijeyesundera DN, Karkouti K, Dupuis JY, et al. Derivation and validation of a simplified predictive index for renal replacement therapy after cardiac surgery. *JAMA.* 2007; 297:1801–9. [PubMed: 17456822]
109. Aronson S, Fontes ML, Miao Y, et al. Risk index for perioperative renal dysfunction/failure: critical dependence on pulse pressure hypertension. *Circulation.* 2007; 115:733–42. [PubMed: 17283267]
110. Palomba H, de Castro I, Neto AL, et al. Acute kidney injury prediction following elective cardiac surgery: AKICS Score. *Kidney Int.* 2007; 72:624–31. [PubMed: 17622275]
111. Brown JR, Cochran RP, Leavitt BJ, et al. Multivariable prediction of renal insufficiency developing after cardiac surgery. *Circulation.* 2007; 116:1139–43. [PubMed: 17846294]
112. Rueggberg A, Boehm S, Napieralski F, et al. Development of a risk stratification model for predicting acute renal failure in orthotopic liver transplantation recipients. *Anaesthesia.* 2008; 63:1174–80. [PubMed: 18803627]
113. Kheterpal S, Tremper KK, Englesbe MJ, et al. Predictors of postoperative acute renal failure after noncardiac surgery in patients with previously normal renal function. *Anesthesiology.* 2007; 107:892–902.
114. Abelha FJ, Botelho M, Fernandes V, et al. Determinants of postoperative acute kidney injury. *Crit Care.* 2009; 13:R79. [PubMed: 19463152]
115. Birnie K, Verheyden V, Pagano D, et al. Predictive models for kidney disease: improving global outcomes (KDIGO) defined acute kidney injury in UK cardiac surgery. *Crit Care.* 2014; 18:606. [PubMed: 25673427]
116. Nigwekar SU, Kandula P, Hix JK, et al. Off-pump coronary artery bypass surgery and acute kidney injury: a meta-analysis of randomized and observational studies. *Am J Kidney Dis.* 2009; 54:413–23. [PubMed: 19406542]
117. Seabra VF, Alobaidi S, Balk EM, et al. Off-pump coronary artery bypass surgery and acute kidney injury: a meta-analysis of randomized controlled trials. *Clin J Am Soc Nephrol.* 2010; 5:1734–44. [PubMed: 20671222]
118. Kolhe NV, Stevens PE, Crowe AV, et al. Case mix, outcome and activity for patients with severe acute kidney injury during the first 24 hours after admission to an adult, general critical care unit: application of predictive models from a secondary analysis of the ICNAR Case Mix Programme database. *Crit Care.* 2008; 12(Suppl 1):S2. [PubMed: 19105800]
119. Thakar CV, Liangos O, Yared JP, et al. Predicting acute renal failure after cardiac surgery: validation and re-definition of a risk-stratification algorithm. *Hemodial Int.* 2003; 7:143–7. [PubMed: 19379354]
120. Candela-Toha A, Elias-Martin E, Abaira V, et al. Predicting Acute Renal Failure after Cardiac Surgery: External Validation of Two New Clinical Scores. *Clin J Am Soc Nephrol.* 2008; 3:1260–5. [PubMed: 18463173]
121. Josephs SA, Thakar CV. Perioperative risk assessment, prevention, and treatment of acute kidney injury. *Int Anesthesiol Clin.* 2009; 47:89–105. [PubMed: 19820480]
122. Saria S, Rajani AK, Gould J, et al. Integration of early physiological responses predicts later illness severity in preterm infants. *Sci Transl Med.* 2010; 2:48ra65.

123. Ng SY, Sanagou M, Wolfe R, et al. Prediction of acute kidney injury within 30 days of cardiac surgery. *The Journal of Thoracic and Cardiovascular Surgery*. 2014; 147:1875–83.e1. [PubMed: 23993316]
124. Celi LA, Tang RJ, Villarroel MC, et al. A Clinical Database-Driven Approach to Decision Support: Predicting Morality Among Patients with Acute Kidney Injury. *J Healthc Eng*. 2011; 2:97–110. [PubMed: 22844575]
125. Celi LA, Galvin S, Davidzon G, et al. A Database-driven Decision Support System: Customized Morality Prediction. *J Pers Med*. 2012; 2:138–48. [PubMed: 23766893]
126. Thakar CV, Kharat V, Blanck S, et al. Acute kidney injury after gastric bypass surgery. *Clin J Am Soc Nephrol*. 2007; 2:426–30. [PubMed: 17699447]
127. Kheterpal S, Khodaparast O, Shanks A, et al. Chronic angiotensin-converting enzyme inhibitor or angiotensin receptor blocker therapy combined with diuretic therapy is associated with increased episodes of hypotension in noncardiac surgery. *J Cardiothorac Vasc Anesth*. 2008; 22:180–6. [PubMed: 18375317]
128. Patel NN, Rogers CA, Angelini GD, et al. Pharmacological therapies for the prevention of acute kidney injury following cardiac surgery: a systematic review. *Heart Fail Rev*. 2011; 16:553–67. [PubMed: 21400231]
129. Tie HT, Luo MZ, Luo MJ, et al. Sodium bicarbonate in the prevention of cardiac surgery-associated acute kidney injury: a systematic review and meta-analysis. *Crit Care*. 2014; 18:517. [PubMed: 25212718]
130. Garg AX, Kurz A, Sessler DI, et al. Perioperative aspirin and clonidine and risk of acute kidney injury: A randomized clinical trial. *JAMA*. 2014; 312:2254–64. [PubMed: 25399007]
131. Brienza N, Giglio MT, Marucci M, et al. Does perioperative hemodynamic optimization protect renal function in surgical patients? A meat-analytic study. *Crit Care Med*. 2009; 37:2079–90. [PubMed: 19384211]
132. Badin J, Boulain T, Ehrmann S, et al. Relation between mean arterial pressure and renal function in the early phase of shock: a prospective, explorative cohort study. *Crit Care*. 2011; 15:R135. [PubMed: 21645384]
133. Cai J, Xu R, Yu X, et al. Volatile anesthetics in preventing acute kidney injury after cardiac surgery: A systematic review and meta-analysis. *The Journal of Thoracic and Cardiovascular Surgery*. 2014; 148:3127–36. [PubMed: 25218542]
134. Lee HT, Emala CW. Protective effects of renal ischemic preconditioning and adenosine pretreatment: role of A1 and A3 receptors. *Am J Physiol-Renal Physiol*. 2000; 278:F380–F7. [PubMed: 10710542]
135. Turman MA, Bates CM. Susceptibility of human proximal tubular cells to hypoxia: effect of hypoxic preconditioning and comparison to glomerular cells. *Ren Fail*. 1997; 19:47–60. [PubMed: 9044451]
136. Kharbanda RK, Nielsen TT, Redington AN. Translation of remote ischaemic preconditioning into clinical practice. *The Lancet*. 2009; 374:1557–65.
137. Meran S, Wonnacott A, Amphlett B, et al. How good are we at managing acute kidney injury in hospital? *Clinical Kidney Journal*. 2014; 7:14–50.
138. Ponce D, Zorzeonn CdPF, dos Santos NY, et al. Early nephrology consultation can have an impact on outcome of acute kidney injury patients. *Nephrology Dialysis Transplantation*. 2011; 26:3202–6.
139. e Silva VTC, Liaño F, Muriel A, et al. Nephrology referral and outcomes in critically ill acute kidney injury patients. *PLoS One*. 2013; 8:e70482. [PubMed: 23936440]
140. Balasubramanian G, Al-Aly Z, Moiz A, et al. Early nephrologist involvement in hospital-acquired acute kidney injury: a pilot study. *Am J Kidney Dis*. 2011; 57:228–34. [PubMed: 21195518]
141. Siew ED, Peterson JF, Eden SK, et al. Outpatient nephrology referral rates after acute kidney injury. *J Am Soc Nephrol*. 2012; 23:305–12. [PubMed: 22158435]
142. Matheny ME, Peterson JF, Eden SK, et al. Laboratory Test Surveillance following Acute Kidney Injury. *PLoS One*. 2014; 9:e103746. [PubMed: 25117447]
143. Harel Z, Wald R, Bargman JM, et al. Nephrologist follow-up improves all-cause mortality of severe acute kidney injury survivors. *Kidney Int*. 2013; 83:901–8. [PubMed: 23325077]

144. Lameire NH, Bagga A, Cruz D, et al. Acute kidney injury: a increasing global concern. *The Lancet*. 2013; 382:170–9.
145. Kirwan, C.; Prowle, J. *Annual Update in Intensive Care and Emergency Medicine*. Vol. 2013. Springer; 2013. Acute Kidney Injury Is a Chronic Disease that Requires Long-Term Follow-up; p. 723–37.
146. Kellum JA, Lameire N. Diagnosis, evaluation, and management of acute kidney injury: a KDIGO summary (Part 1). *Crit Care*. 2013; 17:204. [PubMed: 23394211]
147. Bowman BT, Kleiner A, Bolton WK. Comanagement of diabetic kidney disease by the primary care provider and nephrologist. *Med Clin North Am*. 2013; 97:157–73. [PubMed: 23290736]
148. Diamantidis, CJ.; Powe, NR.; Jar, BG., et al. *Clin J Am Soc Nephro*. CJN; 2011. Primary care-specialist collaboration in the care of patients with chronic kidney disease; p. 06240710
149. Richards N, Harris K, Whitfield M, et al. Primary care-based disease management of chronic kidney disease (CKD), based on estimated glomerular filtration rate (eGFR) reporting, improves patient outcomes. *Nephrology Dialysis Transplantation*. 2008; 23:549–55.
150. Stoves J, Conolly J, Cheung CK, et al. Electronic consultation as an alternative to hospital referral for patients with chronic kidney disease: a novel application for networked electronic health records to improve the accessibility and efficiency of healthcare. *Qual Saf Health Care*. 2010; 19:e54–e. [PubMed: 20554576]
151. DuBose TD, Behrens MT, Berns A, et al. The patient-centered medical home and nephrology. *J Am Soc Nephrol*. 2009; 20:681–2. [PubMed: 19297553]
152. Chang J, Ronco C, Rosner MH. Computerized decision support systems: improving patient safety in nephrology. *Nat Rev Nephrol*. 2011; 7:348–55. [PubMed: 21502973]

Key Points

- Acute and chronic kidney injury and dysfunction play important roles in affecting perioperative outcomes.
- AKI is a common complication after surgery and mild to moderate AKI is more common than severe AKI.
- All stages of AKI severity are associated with increased short and long term morbidity and mortality.
- Clinical risk factors for AKI are similar but not identical in different surgical populations.

Table 1
Risk Factors for AKI

Type of Surgery	Preoperative Risk Factors	Intraoperative Risk Factors
Cardiac Surgery	Advanced age Female sex Baseline renal function Diabetes mellitus Poor glycemic control Congestive heart failure Low ejection fraction Peripheral vascular disease Intra-aortic balloon pump use Chronic obstructive pulmonary disease Peripheral vascular disease Hypertension Lower preoperative hemoglobin Atrial fibrillation Preoperative total lymphocyte count <1,500 cells/ μ L Gene polymorphisms (Alleles angiotensinogen 842C and interleukin 6 -572C in Caucasians Alleles endothelial nitric oxide synthase 894T and angiotensin-converting enzyme deletion and insertion in African Americans)Elevated preoperative C-reactive protein	Emergent reoperation Valve replacement surgery Surgery on the thoracic aorta Deep hypothermic circulatory arrest Low-output syndrome Vasopressors need prior to cardiopulmonary bypass Use of Cardiopulmonary bypass Volume of blood transfusion Intraoperative nadir hamoglobin level Intraoperative hypotension (<50 mmHg) Urine output Surgery requiring cardiopulmonary bypass Cardiopulmonary bypass duration Low pump flow Low perfusion pressure Severe haemodilution Low oxygen delivery (DO ₂) and low DO ₂ /VCO ₂ ratios Hyperthermia (the arterial outlet temperature >37 degrees C) Intraoperative inotropes Furosemide administration ICU admission temperature after CPB Aprotinin use
Non-Cardiac Surgery	Advanced age Male sex Baseline renal function Diabetes mellitus Liver disease Peripheral vascular disease Chronic obstructive pulmonary disease Left ventricular dysfunction High body mass index American Society of Anesthesiologists physical status Preoperative albumin <3.2 Preoperative Anemia Use of hydroxyethyl starch fluids Model for End-Stage Liver Disease (MELD) score, Hepatorenal syndrome type II and Hepatitis C (orthotopic liver transplantation) Low Norton scores (performed by nurses) and preoperative diuretics and NSAIDS (total hip arthroplasty)	Emergent surgery High risk surgery Reoperation Open vascular surgery procedure Surgery for malignant gynecological tumors Prolonged surgical times (>4 h) Aortic cross-clamp time Fenestrated grafts and contrast dose for endovascular procedures Intraoperative hypotension Intraoperative vasopressor use Number of transfused packed red blood cells Prolonged dopamine use Lactic acidosis Lateral decubitus positioning in laparoscopic surgery Administration of furosemide or mannitol Duration of anhepatic phase (orthotopic liver transplantation)