Fibroblast growth factor 9 is a novel modulator of negative affect

Elyse L. Aurbach^{a,b,1}, Edny Gula Inui^{a,1}, Cortney A. Turner^a, Megan H. Hagenauer^a, Katherine E. Prater^{a,b}, Jun Z. Li^a, Devin Absher^c, Najmul Shah^a, Peter Blandino Jr.^a, William E. Bunney^c, Richard M. Myers^d, Jack D. Barchas^e, Alan F. Schatzberg^f, Stanley J. Watson Jr.^a, and Huda Akil^{a,2}

^aMolecular and Behavioral Neuroscience Institute, University of Michigan, Ann Arbor, MI 48109; ^bNeuroscience Graduate Program, University of Michigan, Ann Arbor, MI 48109; ^cDepartment of Psychiatry & Human Behavior, University of California, Irvine, CA 92697; ^dHudsonAlpha Institute for Biotechnology, Huntsville, AL 35806; ^eDepartment of Psychiatry, Cornell University, New York, NY 10065; and ^fDepartment of Psychiatry, Stanford University, Stanford, CA 94305

Contributed by Huda Akil, August 6, 2015 (sent for review July 21, 2015; reviewed by Victoria Arango and James P. Herman)

Both gene expression profiling in postmortem human brain and studies using animal models have implicated the fibroblast growth factor (FGF) family in affect regulation and suggest a potential role in the pathophysiology of major depressive disorder (MDD). FGF2, the most widely characterized family member, is down-regulated in the depressed brain and plays a protective role in rodent models of affective disorders. By contrast, using three microarray analyses followed by quantitative RT-PCR confirmation, we show that FGF9 expression is up-regulated in the hippocampus of individuals with MDD, and that FGF9 expression is inversely related to the expression of FGF2. Because little is known about FGF9's function in emotion regulation, we used animal models to shed light on its potential role in affective function. We found that chronic social defeat stress, an animal model recapitulating some aspects of MDD, leads to a significant increase in hippocampal FGF9 expression, paralleling the elevations seen in postmortem human brain tissue. Chronic intracerebroventricular administration of FGF9 increased both anxiety- and depression-like behaviors. In contrast, knocking down FGF9 expression in the dentate gyrus of the hippocampus using a lentiviral vector produced a decrease in FGF9 expression and ameliorated anxiety-like behavior. Collectively, these results suggest that high levels of hippocampal FGF9 play an important role in the development or expression of mood and anxiety disorders. We propose that the relative levels of FGF9 in relation to other members of the FGF family may prove key to understanding vulnerability or resilience in affective disorders.

fibroblast growth factor 9 | major depression | anxiety | hippocampus | affect

The neurotrophic hypothesis of major depressive disorder (MDD) posits that the neurobiological basis for mood disorders may be due to the dysregulation of growth factors and their effects on brain circuitry (1, 2). This hypothesis is supported by gene expression profiling experiments in postmortem human brains that implicate the fibroblast growth factor (FGF) family and other neurotrophins in MDD (3, 4). To date, only a few growth factors involved in mood disorders, such as brain-derived neurotrophic factor (BDNF) and FGF2, have been studied extensively.

Our laboratory and others have demonstrated that FGF2 is down-regulated in the frontal cortices (3), hippocampus (5), and locus coeruleus (4) in depressed individuals. This consistent decrease in FGF2 expression across regions is striking and underscores the likely importance of FGF2 in mediating affect.

Follow-up studies from our laboratory and others have focused on the FGF family and have demonstrated a key role of FGF2 in the control of emotional behavior. Rats exposed to chronic social defeat stress, an animal model recapitulating some aspects of MDD, showed decreased hippocampal FGF2 expression relative to unstressed controls (6), whereas subchronic and chronic administration of FGF2 had antidepressant properties (7, 8). Moreover, administration of antidepressants and anxiolytics induced FGF2 expression (9–11), and recent studies suggested that this

induction is necessary for the effectiveness of the antidepressants (8). Paralleling the human depression studies, hippocampal FGF2 was basally decreased in high-anxiety animals, but chronic administration in adulthood (12) or a one-time administration of FGF2 during early development (13) alleviated their anxious phenotype. Additionally, anxiety-like behavior in outbred rats was increased by knocking down FGF2 in the dentate gyrus (DG) of the hippocampus (14). Collectively, these studies show that FGF2 is an endogenous anxiolytic and antidepressant that plays both an organizational role during development and an ongoing role during adulthood to modulate emotional reactivity. The clinical relevance of FGF2 is further supported because its role in MDD was first discovered in the human brain (3).

Other members of the FGF family showed different patterns of dysregulation in postmortem tissue from individuals with MDD. Most notably, FGF9 had a pattern of dysregulation opposite to

Significance

Molecular mechanisms mediating negative emotion and contributing to major depression remain elusive: here, we present evidence implicating fibroblast growth factor 9 (FGF9) as a key mediator. We use whole-transcriptome studies of postmortem human tissue to demonstrate that FGF9 is elevated in depression. Reverse translation animal studies demonstrate that both endogenous and exogenous FGF9 promotes anxiety- and depression-like behavior. Conversely, localized blockade of endogenous FGF9 expression decreases anxiety behavior. To our knowledge, this paper is the first description of hippocampal FGF9 function and the first evidence implicating FGF9 in negative affect. Thus, FGF9 represents a novel target for treating affective disorders. Moreover, our findings suggest that FGF2 and FGF9 work in functional opposition; we hypothesize that the balance between FGF factors may prove critical for optimal regulation of mood.

Author contributions: E.L.A., E.G.I., C.A.T., M.H.H., J.Z.L., W.E.B., R.M.M., J.D.B., A.F.S., S.J.W., and H.A. designed research; E.L.A., E.G.I., M.H.H., K.E.P., J.Z.L., D.A., N.S., R.M.M., and S.J.W. performed research; E.L.A., M.H.H., J.Z.L., D.A., P.B., W.E.B., R.M.M., J.D.B., A.F.S., and S.J.W. contributed new reagents/analytic tools; E.L.A., E.G.I., M.H.H., and K.E.P. analyzed data; E.L.A., E.G.I., C.A.T., M.H.H., K.E.P., and H.A. wrote the paper; C.A.T. provided coordination of the Pritzker Database; W.E.B. provided coordination, administration, and oversight of the Brain Bank; and R.M.M., J.D.B., A.F.S., and S.J.W. created the Consortium infrastructure for the human studies.

Reviewers: V.A., Columbia University; and J.P.H., University of Cincinnati.

Conflict of interest: E.L.A., C.A.T., M.H.H., J.Z.L., D.A., P.B., W.E.B., R.M.M., J.D.B., A.F.S., S.J.W., and H.A. are members of the Pritzker Neuropsychiatric Research Consortium, which is supported by the Pritzker Neuropsychiatric Disorders Research Fund, LLC (Fund). There exists a shared intellectual property agreement between the academic and philanthropic entities of the Consortium. The Fund has no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

¹E.L.A. and E.G.I. contributed equally to this work.

²To whom correspondence should be addressed. Email: akil@umich.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1510456112/-DCSupplemental.

FGF2: FGF9 expression was increased in the frontal cortices (3) and locus coeruleus (4) in patients with MDD. This relationship is intriguing because additional evidence suggests that FGF2 and FGF9 may be functionally opposed. For example, FGF2 expression was decreased and FGF9 expression was increased in an in vitro cellular model of chronic stress (15).

Despite the evidence that FGF9 is altered in MDD, little is known about the function of this molecule in the brain. FGF9 is primarily expressed by neurons in the cortex, hippocampus, thalamus, cerebellum, and spinal cord (16–20), although it is also expressed by glia in the hindbrain and spinal cord (21). FGF9 interacts with several of the tyrosine kinase FGF receptors (FGFR) (22), binding preferentially to FGFR3 (23). Functionally, the literature suggests that FGF9 may promote cell survival. For example, FGF9 acts as a survival factor for nigrostriatal and mesencephalic dopamine neurons (24), and FGF9 treatment can increase the survival and soma size of cultured basal forebrain cholinergic neurons (25). FGF9 also weakly promoted proliferation and survival of cultured adult subventricular neural progenitor cells, inhibiting astrocyte differentiation (26). However, these are all studies of exogenous FGF9, which may be mimicking the actions of other FGFs. Moreover, none of the studies to date have examined the role of FGF9 in the hippocampus or its role in the regulation of emotionality.

Here, we specifically sought to elucidate the potential role of hippocampal FGF9 in the regulation of emotions and mood, given our early observations that FGF9 expression was altered in the postmortem hippocampus of individuals with MDD. We have carried out follow-up analyses in the human brain using multiple data sets to ascertain the relationship between FGF9 and other members of the FGF family. In addition, because it is difficult to determine whether gene dysregulation in postmortem human brain tissue contributes to the development of MDD or is a result of the disease process, we used a series of animal experiments to examine the role of FGF9 in affective behavior. To this end, we examined the effects of psychosocial stress, an animal model recapitulating some aspects of depression, on hippocampal FGF9 expression. We also characterized the effects of FGF9 administration and knockdown on anxiety-like and depression-like behavior and hippocampal gene expression. The combination of these various strands of evidence provides strong support for FGF9 as an endogenous molecule that promotes negative affect and may play a role in vulnerability to major depression.

Results

Human Studies.

FGF9 expression was increased in the hippocampus of depressed individuals. Based on published data indicating that specific growth factor transcripts are altered in depressed brains (3–5), we selectively examined their differential expression in major depression using three human hippocampal microarray datasets. We examined diagnosis-related gene expression using a linear model that controlled for a variety of confounding pre- and postmortem variables and found that hippocampal FGF9 expression was consistently increased in subjects with MDD relative to controls in all three datasets, across platforms (Table 1). Using quantitative real-

time polymerase chain reaction (qRT-PCR), we confirmed that FGF9 expression was increased by 32% in individuals with MDD relative to controls, with a Cohen's *d* effect size of 0.57, a medium-sized effect (Table 1). In the microarray data, FGFR1 and FGFR2 appeared to be down-regulated in individuals with MDD, but these effects were borderline significant and platform-specific (Table S1). Expression of BDNF, glial-derived neurotrophic factor (GDNF), FGF2, and FGFR3 did not differ in MDD subjects relative to controls (Table S1). Thus, these analyses pointed to FGF9 as one of the most clearly and consistently altered growth factors in MDD: we always observed it to be up-regulated in the depressed hippocampus. **FGF9 expression is correlated with expression of other members of the FGF family.** We further explored whether the relationship between the expression of hippocampal FGF9 and other candidate members of the FGF family differed by diagnosis (Fig. 1 *A–D*). We found that the expression of FGF9 was significantly negatively related to FGF2, FGFR1, FGFR2, and FGFR3. A negative correlation between FGF9 and FGFR1 was present in MDD but not in control tissue (Fig. S14). These correlations suggest a general dysregulation in the FGF system in MDD, with FGF9 appearing to be one of the key players.

Animal Studies.

Repeated social stress increased both social avoidance behavior and FGF9 gene expression. We subjected rats to repeated social stress over 10 days. Rats subjected to the stress paradigm showed significantly less weight gain during the experiment (Fig. S2 *A* and *B*) compared with handled controls. These stressed rats also showed increased social avoidance behavior in the social interaction test (Fig. 2*A*). Hippocampal FGF9 mRNA expression was significantly increased following repeated social stress compared with handled controls (Fig. 2 *B* and *C*); we observed this increase by tracing hippocampal subregions CA1, CA2, CA3, and dentate gyrus from mRNA in situ hybridization (ISH) autoradiograms (*Materials and Methods*). Thus, this model replicated the observations in the human MDD brain.

Chronic FGF9 microinjections increased both anxiety-like and depression-like behavior. We performed a dose-finding study by acutely administering FGF9 or vehicle (intracerebroventricular, i.c.v.) to examine the effect of exogenous FGF9 on spontaneous affective behavior. We administered a range of doses over five orders of magnitude (Fig. S3 *A–C*): of these, we found that 20 ng FGF9 altered both anxiety- and depression-like behavior acutely, so we chose to use this dose to determine whether long-term exposure to FGF9 was sufficient to alter spontaneous affective behavior. Chronic FGF9 administration increased anxiety-like behavior on the elevated plus maze (EPM). FGF9-microinjected animals spent less percent time in the open arms of the EPM compared with vehicle controls (Fig. 3*A*). We observed a decrease in the percent time spent climbing and an increase in the percent time spent immobile in animals injected with FGF9 compared with vehicle controls on the forced swim test (FST) (Fig. 3*B*). Chronic FGF9 microinjections did not alter locomotor activity (Fig. S3*D*), indicating that the effects were specific to affective-like

Table 1. FGF9 was increased in the postmortem hippocampus from individuals with MDD relative to nonpsychiatric controls

Candidate	Microarray: Affymetrix HG-U133A	Microarray: Illumina HT-12	Microarray: Illumina Ref-8_v2	qPCR	qPCR: effect size (Cohen's <i>D</i>)
Control (<i>n</i>)	56	24	45	22	
MDD (<i>n</i>)	36	21	33	23	
FGF9 (raw <i>P</i> value)	↑ (0.0029)	↑ (0.0295)	↑ (0.0027)	↑ 32% (0.0272)	0.57 (medium)

Significant effects are bolded. The most conservative *P* value is reported when there were multiple probes.

Fig. 1. Postmortem hippocampal FGF9 expression correlates with expression of other FGFs. FGF9 expression in postmortem hippocampal tissue is negatively correlated with expression of (A) FGF2 ($R^2 = 0.15$, $P < 0.001$), (B) FGFR1 ($R^2 = 0.09$, $P < 0.001$), (C) FGFR2 ($R^2 = 0.17$, $P < 0.001$), and (D) FGFR3 ($R^2 = 0.19$, $P < 0.001$).

behavior. Chronic FGF9 administration also decreased FGFR1 expression in the dentate gyrus (Fig. S1 B and C).

Animals administered LVshFGF9 had significantly less FGF9 expression in the dentate gyrus relative to LVshNS. To determine whether endogenous FGF9 expression was necessary for the expression of affective behavior, we used a lentiviral vector that used RNA interference to knock down expression of FGF9 (Fig. S4). Four weeks after microinjecting the knockdown (LVshFGF9) or control (LVshNS) viruses bilaterally into the dentate gyrus, we assessed effects on behavior and euthanized all animals (Fig. S5A). We confirmed the impact of the localized viral injection by examining enhanced green fluorescent protein (eGFP) expression, a marker for successful viral transduction, with concurrent changes in FGF9 expression in the dentate gyrus. Animals who received the LVshFGF9 knockdown virus showed significantly less FGF9 expression relative to LVshNS control animals, indicating that LVshFGF9 was successful in reducing FGF9 expression by ~30% (Fig. 4 A and B). In contrast, when we examined expression of other members of the FGF family, we found no effect of FGF9 knockdown (Fig. S5 B–E). Therefore, the LVshFGF9 virus was both effective and selective in reducing FGF9 expression in the dentate gyrus.

The LVshFGF9 and LVshNS viruses primarily infected dentate granule neurons. To examine the cell types impacted by lentiviral infection, we performed triple-label immunohistochemistry, using antibodies against eGFP to mark transduced cells, NeuN to label neurons, and GFAP to identify astrocytes. We observed that the vast majority of colocalization occurred between eGFP and NeuN, with very few GFAP-labeled cells showing concurrent eGFP expression (Fig. 4C). The pattern of transduction and colocalization did not differ between LVshNS and LVshFGF9 animals. Therefore, the virus preferentially transduced dentate granule neurons, suggesting that the behavioral effects of FGF9 knockdown are likely mediated through these cells.

FGF9 knockdown decreased spontaneous anxiety-like behavior on the EPM. Only animals who expressed eGFP in the dentate gyrus bilaterally in at least four serial sections were included, leaving 8 animals per group. LVshFGF9 animals spent significantly less time in the closed arms of the EPM than did LVshNS control animals (Fig. 4D), suggesting an anxiolytic effect. Similarly, LVshFGF9 animals spent more time in the open arms than did LVshNS animals, although this trend did not reach statistical significance (Fig. 4D). The impact of FGF9 knockdown was selective to anxiety-like behavior: we observed no differences between groups in the total distance traveled (Fig. S5F). There were no differences in depression-like behavior on the FST (Fig. 4E).

Discussion

This series of studies is the first, to our knowledge, to link the dysregulation of hippocampal FGF9 to affective disorders. Using microarray data from several analyses with qRT-PCR validation, we showed that FGF9 expression is increased in the postmortem hippocampus of individuals diagnosed with MDD relative to nonpsychiatric controls. We demonstrated that FGF9 expression is negatively correlated with FGF2 expression, as well as with

three FGF receptors (FGFR1, FGFR2, and FGFR3). To address the functional significance of these observations in human brain, we performed a complementary series of experiments in a rodent model to better characterize the role of FGF9 in affective processes. Chronic social defeat stress decreased social interaction and body weight in our animals and was associated with increased hippocampal FGF9 expression. Chronic FGF9 administration increased both anxiety- and depression-like behavior. Conversely, knocking down FGF9 expression in the dentate gyrus decreased anxiety-like behavior. These animal studies provide converging, complementary evidence for the role of FGF9 as an anxiogenic and prodepressant agent in the rodent brain. Together, this body of work suggests that FGF9 is an endogenous factor that enhances vulnerability to negative affect. Indeed, we believe that FGF9 is the first growth factor to be described as having a sustained anxiogenic and prodepressant role in the hippocampus. Thus, an agent that blocks the effects of FGF9 may be useful in treating human affective disorders.

FGF9 Expression Is Dysregulated in the Postmortem Hippocampus from Individuals Diagnosed with MDD. Using microarray and qRT-PCR, we observed a significant increase in hippocampal FGF9 expression in individuals with MDD. We did not observe a decrease in BDNF, GDNF, or FGF2 expression. In previous studies, other groups have observed a decrease in hippocampal FGF2 expression in depressed patients using mRNA ISH (5). The reason behind the discrepancy in our results is not clear, but may be due to differences in technique (microarray vs. mRNA ISH) or because tissue pH was higher in our study, and low pH can impact gene expression (27). Moreover, factors such as comorbid anxiety or a treatment history that includes antidepressant drugs would likely influence these transcripts, and we could not fully account for these factors.

We uncovered a significant negative correlation between hippocampal FGF9 expression and FGF2 expression in postmortem hippocampal tissue, and this relationship may indicate that these transcripts work in functional opposition. In support of this idea, we have previously observed contrasting patterns of FGF2 and FGF9 expression in other brain regions from individuals diagnosed with MDD. Specifically, FGF2 expression was reduced and FGF9 expression was elevated in the frontal cortices (3) and locus coeruleus (4) of the depressed brain relative to controls. The negative correlations between FGF9 expression and other FGFs also underscore the biological significance of FGF9 dysregulation in affective disorders: these relationships indicate that FGF9 may be acting in a complex network of molecules, and the alterations in

Fig. 2. Social defeat stress increases FGF9 expression in the rat hippocampus. (A) Rats subjected to repeated social stress showed increased social avoidance [$t(17) = 2.5$, $P < 0.03$; $n = 8$ –10 animals per group]. (B) Representative pseudocolored autoradiograms from mRNA in situ hybridization against FGF9 from handled controls (Left) and repeated social stress animals (Right). (C) Repeated social stress significantly increased FGF9 gene expression in hippocampal subregions: CA1, CA2, CA3, and dentate gyrus [$t(8) = 2.46$, $P < 0.04$, $t(8) = 3.41$, $P < 0.01$, $t(8) = 3.12$, $P < 0.02$, $t(8) = 3.15$, $P < 0.02$, respectively; $n = 5$ animals per group]. All values are mean $\pm$ SEM.

Fig. 3. FGF9 administration altered anxiety- and depression-like behavior in rats. (A) Chronic FGF9 microinjections increased anxiety-like behavior as evidenced by a decrease in the percent time spent on the open arm compared with vehicle controls [$F_{(1,16)} = 8.1$, $P < 0.05$; $n = 8$ –10 animals per group]. (B) Chronic FGF9 microinjections increased depression-like behavior as shown by decreased percent time spent climbing and increased percent time spent immobile in the FST compared with vehicle controls [$F_{(1,14)} = 7.3$, $P < 0.05$, $F_{(1,14)} = 6.6$, $P < 0.05$, respectively; $n = 8$ –10 animals per group]. Chronic FGF9 microinjections did not alter locomotor activity (Fig. S3), indicating that the effects were specific to affective-like behavior. All values are mean $\pm$ SEM.

FGF9 expression are concurrent with changes in other FGF family members that we have previously observed to impact affective behavior. However, these human studies are, by nature, correlational, and do not address whether these changes are a by-product of an underlying pathology associated with MDD or if they are part of the disease process. Moreover, many of the patients had a treatment history including antidepressants and/or other psychoactive drugs. Both the clinical and the toxicology data revealed a great deal of heterogeneity in drug exposure, making it difficult to analyze the impact of these drugs on the expression of FGF9. Thus, it will be important for future animal studies to ascertain whether psychoactive drugs can modify the expression of FGF9. As noted above, such studies on FGF2 indicate that the dysregulation we observed in MDD is not secondary to these treatments, but occurs despite them: antidepressants induce FGF2 and mediate their actions in part through that induction. Whether antidepressants inhibit FGF9 expression remains to be ascertained.

FGF9 Expression Mediates Affect in an Animal Model. We previously demonstrated that hippocampal FGF2 expression is decreased in rats subjected to 4 days of social defeat stress (6). We also found that hippocampal FGF2 expression correlates with variability in anxiety- and depression-like behavior. For example, endogenous hippocampal FGF2 expression correlated with spontaneous open arm time on the EPM in outbred rats (14). Moreover, animals that naturally exhibit high anxiety-like behavior have low levels of hippocampal FGF2 (12, 13). In contrast, we have demonstrated here that ten days of social defeat stress up-regulates hippocampal FGF9 expression while decreasing body weight and social interactions, correlates of increased depression-like behavior. These results are congruent with our observations from human studies and indicate that hippocampal FGF9 may play an important role in stress responsiveness.

Chronic administration of FGFs produced coordinate effects on anxiety- and depression-like behavior: FGF2 reduced anxiety- (12) and depression-like (7) behavior. In contrast, we demonstrate here that chronic FGF9 administration increases anxiety-like and depression-like behavior, demonstrating again that the two FGFs have opposing effects. Given that antidepressant and anxiolytic medications induce growth factor expression (9–11), it may be that administering growth factors directly activates the same physiological mechanisms as classical anxiolytics and antidepressants. Indeed, work from others indicates FGF2 is necessary for the positive effect of antidepressants (8). However, one limitation of our FGF9 administration studies is the lack of

anatomical specificity: that is, that administration into the lateral ventricle likely affects many brain regions in addition to the hippocampus. Other regions, including the prefrontal cortices, are also likely to be involved in the behavioral expression of these changes in negative affect, and future studies can better elucidate the function of FGF9 in these regions.

Altering endogenous gene expression using transgenic mice or viral-mediated knockdown can be used to demonstrate that hippocampal growth factor expression is necessary for appropriate spontaneous regulation of affective behavior. We previously used a lentiviral vector to knock down FGF2 expression in the dentate gyrus of outbred rats, increasing anxiety-like behavior on the EPM (14). Similarly, knocking down hippocampal FGF2 eliminated basal behavioral differences in selectively bred rats that typically exhibit differences in spontaneous anxiety-like behavior (28). In contrast, we demonstrate here that using a lentiviral vector to knock down FGF9 expression bilaterally in the dentate gyrus decreases anxiety-like behavior. Interestingly, unlike the results of the administration experiments, our knockdown experiments have produced effects on anxiety-like, but not depression-like, behavior.

Fig. 4. Lentiviral-mediated FGF9 knockdown reduced FGF9 expression in dentate granule neurons and decreased anxiety-like behavior. (A) Representative pseudocolored autoradiograms from mRNA in situ hybridization against FGF9 for animals transduced with a control, nonsilencing virus (LVshNS; Left) and with the FGF9 knockdown virus (LVshFGF9; Right). Dentate gyrus is enlarged for ease of viewing. (B) FGF9 expression is significantly reduced in the dentate gyrus of rats transduced with the LVshFGF9 virus relative to LVshNS controls [$t_{(19)} = 3.53$, $P < 0.01$; $n = 8$ animals per group]. (C) Triple-label immunohistochemistry demonstrating high cell-type specificity of viral transduction; eGFP = green, NeuN = red, GFAP = blue. The virus preferentially infects dentate granule neurons. (Scale bar, 100 μ m.) (D) Knocking down FGF9 expression in the dentate gyrus decreased closed arm time [$t_{(14)} = 2.51$, $P = 0.025$] and increased open arm time [$t_{(14)} = 2.02$, $P = 0.063$] in the elevated plus maze ($n = 8$ animals per group). (E) FGF9 knockdown did not impact climbing, swimming, or immobility behavior in the forced swim test [$t_{(14)} = 1.91$, $P = 0.07$; $t_{(14)} = 1.13$, $P = 0.28$; $t_{(14)} = 1.68$, $P = 0.11$, respectively; $n = 8$ animals per group]. All values are mean $\pm$ SEM.

adding a loop sequence (UUCAAGAGA) and restriction enzyme sites. The shRNA constructs were synthesized (Invitrogen) and subcloned into the pLentiLox3.7 lentiviral vector, which contains an eGFP reporter tag. A non-silencing shRNA (shNS) sequence was used as a control (14). shRNA3 was selected for use *in vivo* because it was most effective at reducing FGF9 expression *in vitro* (Fig. S4) and showed the least toxic effects on cultured cells (<10% cell death). The vector containing shRNA3 (hereafter referred to as shFGF9) and the shNS vector were submitted to the University of Michigan Viral Vector core for synthesis. The returned lentiviruses (100 $\times$ concentration; LVshFGF9 and LVshNS) were used in FGF9 knockdown experiments *in vivo*.

All animals underwent microinjection surgery (groups: LVshFGF9, LVshNS). Under isoflurane anesthesia, a 33-gauge microinjector was lowered bilaterally just above the dentate gyrus of the hippocampus (coordinates from bregma: A/P -5.0 , M/L ± 3.5 , D/V -3.6 – -3.8). One microliter of LVshNS or LVshFGF9 was infused over 4 min (14), and 2 min were allowed for diffusion. We allowed 4 wk for recovery, and then animals were subjected to behavioral testing in the EPM and FST (SI Materials and Methods and Fig. S5A).

Behavioral testing. We monitored locomotor activity for 1 h (SI Materials and Methods). We used the FST to determine depressive-like behavior (SI Materials and Methods). We used the EPM to examine anxiety-like behavior (SI Materials and Methods).

mRNA ISH. All rats were euthanized by rapid decapitation; brains were removed, snap-frozen, and stored at -80°C . Ten-micrometer sections were taken every 200 μm through the hippocampus, and ISH methodology was performed as previously described (34) (SI Materials and Methods). All *in situ* probes were synthesized in-house, and exposure times were experimentally determined for optimal signal (SI Materials and Methods). mRNA expression signals from autoradiographic films were quantified using computer-assisted optical densitometry software, Scion Image (Scio Corporation) or ImageJ (National Institutes of Health). For all experiments except FGF9 knockdown, integrated optical densities were determined by outlining a hippocampal subfield (CA1, CA2, CA3, and dentate gyrus) on each hemisphere, correcting for background. For quantification of sections from animals in the FGF9 knockdown study, we used a modified approach that enabled us to

quantify expression of a gene target only with evidence of successful viral transduction (SI Materials and Methods). eGFP autoradiograms were also analyzed animal by animal: animals in the knockdown study were only included in the final behavioral analysis if they showed robust expression of eGFP in the dentate gyrus of both hemispheres in at least four adjacent sections ($\geq 800\ \mu\text{m}$), without labeling of other structures. After exclusion, eight animals were included in each group.

Immunohistochemistry. Animals ($n = 4$ animals/group; LVshNS, LVshFGF9) underwent the same surgical procedures as above. Three weeks after surgery, animals were transcardially perfused using 4% (wt/vol) paraformaldehyde (PFA). Free-floating sections (45 μm) were blocked for 1 h at room temperature and incubated overnight at room temperature with chicken anti-GFP, 1:2,000 (Abcam); rabbit anti-NeuN, 1:300 (Abcam); and mouse anti-GFAP, 1:500 (Millipore). Sections were washed and transferred to secondary solution (AlexaFluor488 goat anti-chicken IgG, 1:200; AlexaFluor568 goat anti-rabbit IgG, 1:200; AlexaFluor 647 goat anti-mouse, 1:200; Life Technologies) for 2 h at room temperature. Sections were washed, mounted, and coverslipped using ProLong Gold Antifade Reagent with DAPI (Fisher Scientific). Slides were imaged on an Olympus Fluoview FV1000 confocal microscope (SI Materials and Methods).

Statistical analyses. Analysis of human data are described above. All animal behavioral tests were analyzed by one-way ANOVA or Student *t* test.

ACKNOWLEDGMENTS. We thank all the members of the Pritzker Consortium (especially the University of California, Irvine Brain Bank staff), and Robert Thompson, Juan Lopez, Adriana Medina, and David Krowleski for brain dissections. We also thank Audrey Seasholtz, Gwen Krowleski, Jack Parent, Hui Li, Jennifer Fitzpatrick, Jim Stewart, Fei Li, Amy Tang, Mary Hoversten, and Sharon Burke. This work was supported by the Pritzker Neuropsychiatric Research Consortium, Office of Naval Research Grants ONR N00014-15-1-2224 and ONR N00014-12-1-0366, National Institutes of Health Grant R01MH104261, and the Hope for Depression Research Foundation. E.L.A. was supported by National Science Foundation Graduate Research Fellowship Program and NIH Grants 5T32DA007281-17 and 5T32EY017878-10.

- Turner CA, Watson SJ, Akil H (2012) The fibroblast growth factor family: Neuro-modulation of affective behavior. *Neuron* 76(1):160–174.
- Duman RS, Monteggia LM (2006) A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 59(12):1116–1127.
- Evans SJ, et al. (2004) Dysregulation of the fibroblast growth factor system in major depression. *Proc Natl Acad Sci USA* 101(43):15506–15511.
- Bernard R, et al. (2011) Altered expression of glutamate signaling, growth factor, and glia genes in the locus coeruleus of patients with major depression. *Mol Psychiatry* 16(6):634–646.
- Gaughran F, Payne J, Sedgwick PM, Cotter D, Berry M (2006) Hippocampal FGF-2 and FGF1 mRNA expression in major depression, schizophrenia and bipolar disorder. *Brain Res Bull* 70(3):221–227.
- Turner CA, Calvo N, Frost DO, Akil H, Watson SJ (2008) The fibroblast growth factor system is downregulated following social defeat. *Neurosci Lett* 430(2):147–150.
- Turner CA, Gula EL, Taylor LP, Watson SJ, Akil H (2008) Antidepressant-like effects of intracerebroventricular FGF2 in rats. *Brain Res* 1224:63–68.
- Elsayed M, et al. (2012) Antidepressant effects of fibroblast growth factor-2 in behavioral and cellular models of depression. *Biol Psychiatry* 72(4):258–265.
- Mallei A, Shi B, Mochetti I (2002) Antidepressant treatments induce the expression of basic fibroblast growth factor in cortical and hippocampal neurons. *Mol Pharmacol* 61(5):1017–1024.
- Bachis A, Mallei A, Cruz MI, Wellstein A, Mochetti I (2008) Chronic antidepressant treatments increase basic fibroblast growth factor and fibroblast growth factor-binding protein in neurons. *Neuropharmacology* 55(7):1114–1120.
- Gómez-Pinilla F, Dao L, Choi J, Ryba EA (2000) Diazepam induces FGF-2 mRNA in the hippocampus and striatum. *Brain Res Bull* 53(3):283–289.
- Perez JA, Clinton SM, Turner CA, Watson SJ, Akil H (2009) A new role for FGF2 as an endogenous inhibitor of anxiety. *J Neurosci* 29(19):6379–6387.
- Turner CA, Clinton SM, Thompson RC, Watson SJ, Jr, Akil H (2011) Fibroblast growth factor-2 (FGF2) augmentation early in life alters hippocampal development and rescues the anxiety phenotype in vulnerable animals. *Proc Natl Acad Sci USA* 108(19):8021–8025.
- Eren-Kocak E, Turner CA, Watson SJ, Akil H (2011) Short-hairpin RNA silencing of endogenous fibroblast growth factor 2 in rat hippocampus increases anxiety behavior. *Biol Psychiatry* 69(6):534–540.
- Salaria S, et al. (2006) Microarray analysis of cultured human brain aggregates following cortisol exposure: Implications for cellular functions relevant to mood disorders. *Neurobiol Dis* 23(3):630–636.
- Tagashira S, Ozaki K, Ohta M, Itoh N (1995) Localization of fibroblast growth factor-9 mRNA in the rat brain. *Brain Res Mol Brain Res* 30(2):233–241.
- Nakamura S, et al. (1997) Motor neurons in human and rat spinal cord synthesize fibroblast growth factor-9. *Neurosci Lett* 221(2-3):181–184.
- Todo T, et al. (1998) Neuronal localization of fibroblast growth factor-9 immunoreactivity in human and rat brain. *Brain Res* 783(2):179–187.
- Garcés A, Nishimune H, Philippe J-M, Pettmann B, deLapeyrière O (2000) FGF9: A motoneuron survival factor expressed by medial thoracic and sacral motoneurons. *J Neurosci Res* 60(1):1–9.
- Lin Y, et al. (2009) Neuron-derived FGF9 is essential for scaffold formation of Bergmann radial fibers and migration of granule neurons in the cerebellum. *Dev Biol* 329(1):44–54.
- Nakamura S, et al. (1999) Glial expression of fibroblast growth factor-9 in rat central nervous system. *Glia* 28(1):53–65.
- Goetz R, Mohammadi M (2013) Exploring mechanisms of FGF signalling through the lens of structural biology. *Nat Rev Mol Cell Biol* 14(3):166–180.
- Ornitz DM, et al. (1996) Receptor specificity of the fibroblast growth factor family. *J Biol Chem* 271(25):15292–15297.
- Huang J-Y, Hong Y-T, Chuang J-I (2009) Fibroblast growth factor 9 prevents MPP⁺-induced death of dopaminergic neurons and is involved in melatonin neuroprotection *in vivo* and *in vitro*. *J Neurochem* 109(5):1400–1412.
- Kanda T, et al. (2000) Self-secretion of fibroblast growth factor-9 supports basal forebrain cholinergic neurons in an autocrine/paracrine manner. *Brain Res* 876(1-2):22–30.
- Lum M, Turbic A, Mitrovic B, Turnley AD (2009) Fibroblast growth factor-9 inhibits astrocyte differentiation of adult mouse neural progenitor cells. *J Neurosci Res* 87(10):2201–2210.
- Li JZ, et al. (2004) Systematic changes in gene expression in postmortem human brains associated with tissue pH and terminal medical conditions. *Hum Mol Genet* 13(6):609–616.
- Chaudhury S, et al. (2014) FGF2 is a target and a trigger of epigenetic mechanisms associated with differences in emotionality: Partnership with H3K9me3. *Proc Natl Acad Sci USA* 111(32):11834–11839.
- Lamers F, et al. (2011) Comorbidity patterns of anxiety and depressive disorders in a large cohort study: The Netherlands Study of Depression and Anxiety (NESDA). *J Clin Psychiatry* 72(3):341–348.
- American Psychiatric Association (2000) *Diagnostic and Statistical Manual of Mental Disorders* (American Psychiatric Association, Washington, DC), 4th Ed.
- Li JZ, et al. (2013) Circadian patterns of gene expression in the human brain and disruption in major depressive disorder. *Proc Natl Acad Sci USA* 110(24):9950–9955.
- Tomita H, et al. (2004) Effect of agonal and postmortem factors on gene expression profile: Quality control in microarray analyses of postmortem human brain. *Biol Psychiatry* 55(4):346–352.
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C(T)}$ Method. *Methods* 25(4):402–408.
- Kabbaj M, Devine DP, Savage VR, Akil H (2000) Neurobiological correlates of individual differences in novelty-seeking behavior in the rat: Differential expression of stress-related molecules. *J Neurosci* 20(18):6983–6988.