Published in final edited form as:

Compr Physiol. 2014 October ; 4(4): 1737–1774. doi:10.1002/cphy.c140021.

Molecular Mechanisms and Regulation of Urinary Acidification

Ira Kurtz^{1,2}

¹Division of Nephrology, David Geffen School of Medicine, Los Angeles, CA, USA

²Brain Research Institute UCLA, Los Angeles, CA, USA

Abstract

The H^+ concentration in human blood is kept within very narrow limits, ~ 40 nM, despite the fact that dietary metabolism generates acid and base loads that are added to the systemic circulation throughout the life of mammals. One of the primary functions of the kidney is to maintain the constancy of systemic acid-base chemistry. The kidney has evolved the capacity to regulate blood acidity by performing three key functions: 1) reabsorb HCO_3^- that is filtered through the glomeruli to prevent its excretion in the urine; 2) generate a sufficient quantity of new HCO_3^- to compensate for the loss of HCO_3^- resulting from dietary metabolic H^+ loads and loss of HCO_3^- in the urea cycle; and 3) excrete HCO_3^- (or metabolizable organic anions) following a systemic base load. The ability of the kidney to perform these functions requires that various cell types throughout the nephron respond to changes in acid-base chemistry by modulating specific ion transport and/or metabolic processes in a coordinated fashion such that the urine and renal vein chemistry is altered appropriately. The purpose of the article is to provide the interested reader with a broad review of a field that began historically ~ 60 years ago with whole animal studies, and has evolved to where we are currently addressing questions related to kidney acid-base regulation at the single protein structure/function level.

Introduction

Systemic acid-base chemistry is very tightly controlled in normal humans, in spite of the fact that metabolism of typical diets generates a net H^+ load (405). Moreover, there is an inverse association between the blood HCO_3^- concentration and the dietary H^+ load that is greater among middle-aged and elderly persons than younger adults (36). Even small changes in systemic acid-base chemistry may affect bone, skeletal muscle and insulin resistance (1, 213, 469). In postmenopausal women administration of base orally has been shown to improve bone density and decrease renal nitrogen excretion (222, 337, 640).

Diets rich in arginine, lysine, phosphoproteins, and sulfur containing amino acids when metabolized generate net H^+ that are delivered into the venous system (430). The H^+ combine with HCO_3^- (and to some extent other body buffers) extracellularly and intracellularly according to the following reaction: $H^+ + HCO_3^- \rightarrow CO_2 + H_2O$. CO_2 generated in the reaction is excreted by the lungs thereby preventing the systemic CO_2

concentration from increasing. It is the function of the kidneys to regenerate and delivery to the renal veins the required quantity of HCO_3^- lost in this process (288, 404). This function of the kidney is distinguished from the additional requirement performed predominantly by the proximal tubule (and to a lesser extent the thick ascending limb) to reabsorb HCO_3^- from the glomerular filtrate such that typically less than 0.1% of the filtered HCO_3^- is excreted in the urine (589).

Whole Body Acid-base Balance

The generation of HCO_3^- is an essential function of the kidney ensuring that whole body HCO_3^- content remains relatively constant despite the loss of HCO_3^- due to dietary H^+ loads. The kidney generates new HCO_3^- from metabolizable organic anions including α -ketoglutarate (derived from glutamine), lactate, citrate, and fatty acids (183, 287, 288). These biochemical processes occur predominantly in the proximal tubule and to a lesser extent other nephron segments. In the process of generating α -ketoglutarate that is subsequently converted into 2 HCO_3^- in the Krebs cycle and in gluconeogenesis, 2 NH_4^+ are produced according to the following mitochondrial reactions (256, 622, 795, 796):

- 1) glutamine + H_2O $\rightarrow$ glutamate + NH_4^+ (catalyzed by phosphate dependent glutaminase)
- 2) glutamate + H_2O + NAD $\rightarrow$ α -ketoglutarate + NADH + NH_4^+ (catalyzed by glutamate dehydrogenase)

The partitioning of the generated NH_4^+ generated by the kidney between the renal vein and urine is an important factor at the whole organ level that has additional significant effects on systemic acid-base balance. Although new HCO_3^- generated in the proximal tubule is delivered to the renal veins, NH_4^+ produced in the kidney is partitioned between the urine and the renal veins. Of the approximately 75 mEq of NH_4^+ per day produced on an average North American diet, approximately 50% is transferred to the renal veins and an equal amount is excreted (534, 535, 686). Importantly, the NH_4^+ delivered to the renal veins represents a substance that is HCO_3^- consuming, resulting in a decrease the systemic pH. Specifically, NH_4^+ delivered to the circulation upon reaching the liver consumes HCO_3^- in the urea cycle according to the following reaction $2 \text{NH}_4^+ + 2 \text{HCO}_3^- \rightarrow \text{urea} + 3 \text{H}_2\text{O} + \text{CO}_2$ (298). This has led to the concept of “effective new HCO_3^- production” defined as HCO_3^- produced by the kidney, which has not been consumed in the urea cycle and thus is available as a buffer for modulating systemic acid-base balance.

The excretion of ~50% of total NH_4^+ production requires the coordinated intertubular transport of $\text{NH}_3/\text{NH}_4^+$ from its production site predominantly in proximal tubule cells to the collecting duct lumen. In metabolic acidosis, both the quantity of new HCO_3^- produced in the proximal tubule is increased and the intertubular transport of $\text{NH}_3/\text{NH}_4^+$ is altered such that a greater percentage of total NH_4^+ produced is excreted in the urine (less being delivered to the renal veins) thereby significantly increasing the effective new HCO_3^- delivery to the systemic circulation (101, 535, 686, 687). By altering the total new HCO_3^- production rate and renal vein/urine NH_4^+ partitioning, the kidney possesses powerful mechanisms for compensating for the loss of HCO_3^- due to dietary metabolic H^+ generation and in the urea cycle.

A second source of new HCO_3^- generation is derived from tubular luminal protonation of HPO_4^{2-} and subsequent excretion of H_2PO_4^- (titratable acid) (431, 615). Approximately 75% of titratable acid (TA) is derived from H_2PO_4^- excretion whereas the remainder is composed of protonated creatinine and other less defined organic anions. The site of protonation depends on the luminal pH and the pK of the protonated anion. H_2PO_4^- generation takes place primarily in the proximal tubule as the luminal pH decreases from 7.4 to ~ 6.7 (475), whereas the protonation of other organic anions (average pK ~ 4.4) occurs in the collecting duct (325).

Human urine contains not only NH_4^+ , H_2PO_4^- , H^+ -creatinine, but also ~90 organic anions totaling ~40 mEq per day (147, 172, 432). Approximately 50% of these organic anions have metabolizable carboxyl groups that can be converted into HCO_3^- at the pH of normal blood (123, 354). Normal urine also contains ~10 mEq per day of metabolizable organic cations (172). Under control conditions and following exogenous acid-or alkali loading, the net renal excretion of H^+ or base is traditionally calculated according to the following equation: $\text{NAE (meq/time)} = \text{volume/time} \times [\text{NH}_4^+ + \text{TA} - \text{HCO}_3^-]$. The urinary excretion of free H^+ is omitted in the calculation because its contribution is minimal i.e. at a urine pH of 4.5, the concentration of free H^+ is 32 μmol per liter. A similar calculation can be used to quantitate the magnitude of renal vein acid or base delivery to the systemic circulation. Because of the magnitude of metabolizable organic anion excretion in urine (representing loss of HCO_3^-), several authors have indicated that NAE should be calculated as: $\text{NAE (meq/time)} = \text{volume/time} \times [\text{NH}_4^+ + \text{TA} - \text{metabolizable organic anions} - \text{HCO}_3^-]$ (172, 588). The excretion of various metabolizable organic anions is altered predictably during exogenous acid and alkali loading and during normal variations in protein intake (123, 246, 290, 292, 537, 656).

Acid/alkali loading induces changes in NAE associated with alterations in the expression and function of H^+ /base transporters throughout the nephron involving complex signaling pathways that will be detailed below. The response of the kidney to exogenous or endogenous acid/alkali loads results in a change in both new effective HCO_3^- generation and HCO_3^- transport (absorption and secretion) that ultimately contributes to achieving a new steady state.

Proximal Tubule

General Properties of Proximal Tubule HCO_3^- Transport

The proximal tubule is the site in the nephron where the majority (~ 75%) of the filtered HCO_3^- load is reabsorbed into the systemic circulation. The glomerular filtrate has a HCO_3^- concentration of ~ 25 meq/l and assuming that the glomerular filtration rate (GFR) is ~ 144 liters/day, the amount of HCO_3^- entering the proximal tubule is ~ 3.6 moles/day. The urinary excretion of HCO_3^- is typically less than 5 mmoles/day. The proximal tubule is divided longitudinally morphologically into a convoluted and straight segment and each portion of the proximal tubule has a different capacity to absorb HCO_3^- (76, 454, 455). HCO_3^- absorption across the apical membrane is indirect and requires that H^+ are secreted from the cells into the lumen thereby decreasing the luminal bicarbonate concentration from 25 to ~ 10 mM, with an associated decrease in luminal pH from 7.4 to ~ 6.8 according to the

following reaction: $\text{H}^+ + \text{HCO}_3^- \rightarrow \text{H}_2\text{CO}_3 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$. Less HCO_3^- is absorbed quantitatively in the proximal straight tubule. The proximal convoluted tubule is further subdivided into an early S1 segment, and an S2 segment that consists of the end of the convoluted portion followed by a straight segment located in the cortex. The S3 straight segment representing the terminal portion of the proximal tubule is located in the outer stripe of the outer medulla. There are additional differences in the magnitude of HCO_3^- transport depending on whether the segment is from S1, S2, and S3 superficial or juxtamedullary nephrons (3, 58, 387).

The route of HCO_3^- absorption in the proximal tubule is transcellular and active since it occurs against a chemical (HCO_3^-) gradient, and the fact that there is a minimal transepithelial voltage (V_{TE}) of -2 mV (early proximal tubule) and $+2$ mV (late proximal tubule) to drive transport paracellularly (53, 227). Given that the proximal tubule cell apical membrane voltage is ~ -50 to -70 mV relative to the lumen, with an intracellular pH (pHi) of ~ 7.2 , luminal H^+ secretion is active occurring against an electrochemical gradient (78, 610, 813). Luminal acidification could theoretically be mediated by direct luminal H^+ secretion, HCO_3^- absorption, CO_3^{2-} absorption, or CO_2 secretion. To address this question several groups examined whether in the $\text{H}^+ + \text{HCO}_3^- \leftrightarrow \text{H}_2\text{CO}_3 \leftrightarrow \text{CO}_2 + \text{H}_2\text{O}$ reaction, H_2CO_3 and CO_2 are at equilibrium and whether slowing of the interconversion of H_2CO_3 to $\text{CO}_2 + \text{H}_2\text{O}$ with carbonic anhydrase inhibitors resulted in a “disequilibrium pH” (199, 585, 719). In the absence of carbonic anhydrase inhibition, the measured and predicted pH values were found to be essentially equal. However, carbonic anhydrase inhibition generated an acid disequilibrium pH (measured luminal pH lower than predicted) demonstrating that net luminal H^+ secretion occurred. Direct HCO_3^- absorption would have resulted in an alkaline disequilibrium pH. In addition, these experiments showed that the luminal conversion of H_2CO_3 into CO_2 is normally catalyzed by carbonic anhydrase.

Carbonic anhydrase enzymes play a key role in proximal tubule transepithelial HCO_3^- absorption. Each proximal tubule compartment (cell, lumen, peritubular space) has specific CA enzymes that accelerate the interconversion of H_2CO_3 and $\text{CO}_2 + \text{H}_2\text{O}$. In the cytoplasm CAII (29 kDa monomeric protein) is the predominant isoform accounting for most of the carbonic anhydrase activity in the renal cortex (575). Carbonic anhydrase inhibitors affect carbonic anhydrase activity in all three compartments (lumen, cell, and peritubular fluid) and because of their high permeability essentially block all proximal tubule HCO_3^- absorption (171). Several studies have examined the role of the carbonic anhydrase activity in the luminal or basolateral compartments specifically. Luminal perfusion with impermeant CA inhibitors block HCO_3^- absorption almost completely (464). Two CA isoforms have been localized to the brush border of the proximal tubule. CAIV is a glycosylphosphatidylinositol (GPI) anchored membrane bound enzyme with S2 expression $> \text{S1} \gg \text{S3}$ (122, 459, 637). In addition to CAIV, CAXIV has been localized to the brush border membrane of S1 and S2 segments in the mouse proximal tubule (355, 509). The lack of luminal CA activity in S3 proximal tubules was corroborated in the isolated perfused rabbit proximal tubule using a luminal fluorescent pH dye that demonstrated the presence of an acid disequilibrium pH (408). On the basolateral membrane immunofluorescence staining has localized the transmembrane enzyme CA XII to the S1 and S2 proximal tubule segments (411, 541). In addition to CAXII, CAIV is also expressed on the basolateral membrane of

the proximal tubule (122, 459, 637). There appears to be species differences in the relative activity/abundance of these enzymes. In mouse kidney, CA isoforms other than CAIV (e.g. CAXII) account for the bulk of membrane-associated CA activity whereas in rabbit, CAIV accounts for most of the membrane-associated CA activity. The role of basolateral membrane bound carbonic anhydrase activity is highlighted in rabbit isolated perfused proximal tubule experiments where basolateral addition of an impermeant carbonic anhydrase inhibitor blocks ~ 60% of HCO_3^- absorption (698).

Apical Membrane Transporters: H^+ -ATPase; NHE3; NHE8; Organic Anion Transporters; CFEX

H^+ -ATPase— HCO_3^- -stimulated oligomycin-insensitive H^+ -ATPase activity and ATP-dependent H^+ translocation in brush border membrane vesicles was first demonstrated by Kinne-Saffrin and Kinne (367-369). Additional experiments in rat proximal tubules also provided evidence for a Na^+ -independent apical H^+ transport process. Yoshitomi et al reported that in the rat perfused tubule *in vivo*, following perfusion in Na^+ -free solutions, pHi decreased and then recovered (812). Steady state pHi under similar conditions was more alkaline than in the presence of Na^+ suggesting the presence of a Na^+ -independent H^+ /base transport process (19). In the S3 proximal tubule perfused *in vitro*, following acute intracellular acidification, pHi recovered in the absence of Na^+ (400). The pHi recovery was inhibited by the H^+ -ATPase inhibitors *N,N'*-dicyclohexylcarbodiimide (DCCD) and *N*-ethylmaleimide (NEM), and by glycolytic ATP inhibition. In similar studies, Nakhoul et al did not report predictable inhibition by DCCD (521). Additional evidence for an apical H^+ -ATPase was derived from experiments by Frömter et al using *in vivo* rat proximal tubules perfused in the absence of organics. In the early proximal tubule the V_{TE} was + 1 mV and was 0.2 mV in the late proximal tubule) (225). The positive V_{TE} was insensitive to ouabain and acetazolamide and therefore compatible with electrogenic H^+ secretion (226). These findings were ultimately confirmed by labeling of the proximal tubule with H^+ -ATPase specific antibodies demonstrating that the largest expression was at the base of the microvilli (118). Studies by Bank et al used luminal DCCD to block ~ 21% of HCO_3^- absorption but whether complete inhibition occurred is unknown (49) and there are no experiments in mice lacking apical H^+ -ATPase activity (congenital or conditional). It is currently believed that the apical H^+ -ATPase is responsible for ~ 35% of transepithelial bicarbonate absorption. An additional role for H^+ -ATPase activity in the endosomal pathway was suggested from studies of $\alpha 4$ subunit-deficient mice that have proximal tubule transport abnormalities including proteinuria, phosphaturia, accumulation of material in lysosomes, and abnormal endocytic trafficking (304). Apical H^+ -ATPase expression appears to be upregulated in proximal tubule cells by PKA, and AMP-activated protein kinase (AMPK) decreases the PKA-induced increase in H^+ -ATPase expression (10). Additional regulatory pathways involved in modulating H^+ -ATPase activity are highlighted below in the context of the collecting duct.

Several studies have also reported a role for Cl^- in modulating proximal tubule H^+ -ATPase activity. In rabbit S3 proximal tubules perfused *in vitro*, Na^+ -independent pHi recovery following an acid load is dependent on intracellular Cl^- (400) and the shuttling of cytoplasmic H^+ -ATPase containing vesicles to the apical membrane is also Cl^- -dependent

(477). In rat proximal tubules (superficial S1 and S2 segments), H^+ -ATPase activity was shown to be Cl^- -dependent (830) and evidence was provided that CFTR and ClC-5 may play a role (141).

Na^+/H^+ exchangers: NHE3 and NHE8—Electroneutral Na^+/H^+ exchange activity was first demonstrated in rat proximal tubule apical membrane vesicles by Murer et al (513). Additional studies in vesicles demonstrated that external protons compete with Na^+ for a single site with a K_{Na^+} of ~ 10 meq/l (pH 7.5) and a K_{H^+} of 35 nM (43, 373, 755). Lithium and amiloride were shown to compete with Na^+ at an external binding site by noncompetitive and competitive mechanisms (331, 371, 372). In the *in vitro* perfused salamander proximal tubule Boron and Boulpaep demonstrated a luminal Na^+ -dependent pHi recovery process that was amiloride-inhibitable (104). Sasaki et al. reported a similar transport process in the *in vitro* perfused rabbit proximal tubule as did Alpern and Chambers in the *in vivo* perfused rat proximal convoluted tubule (21). Transepithelial HCO_3^- absorption was almost completely inhibited by the removal of Na^+ (lumen, bath), by Na^+ - K^+ -ATPase inhibition, and by luminal amiloride (21). In total, these data from various preparations and experimental conditions provided convincing functional evidence for an apical membrane electroneutral Na^+/H^+ exchanger that plays a key role in both luminal acidification and transepithelial HCO_3^- transport.

Na^+/H^+ exchangers (NHE proteins) are now known to be encoded by separate 9 separate genes (191). Two Na^+/H^+ exchangers, NHE3 and NHE8, have been localized to the apical membrane of the proximal tubule (96, 99). Several lines of evidence indicate that in the adult, NHE3 is responsible for mediating apical the Na^+/H^+ exchange that was first detected physiologically (28, 30, 32, 62, 65, 87, 133, 134, 532, 533, 555, 566, 624, 691, 692, 751, 798, 817). In general NHEs are ~ 100 kDa proteins with an intracellular N-terminal domain, an ~ 450 aa transmembrane domain composed of 11 or 12 transmembrane regions that mediate Na^+/H^+ exchange, and a C-terminal cytoplasmic tail involved in regulation NHE activity (191). Transmembrane region IV is thought to be involved in ion transport, and transmembrane region IX is involved in amiloride inhibition (12).

The transporter has been localized by immunocytochemistry to the proximal tubule brush border (30, 87, 533, 691). In addition, the inhibitory profile of expressed NHE3 to amiloride, amiloride analogues, and HOE694 is comparable to the native apical membrane Na^+/H^+ exchanger (532, 692, 798). Regulation of expressed NHE3 by acidosis, thyroid hormone and glucocorticoids, angiotensin II, endothelin, parathyroid hormone, and protein kinase A parallels the regulation of native apical membrane Na^+/H^+ exchange by these factors (28, 32, 62, 65, 133, 134, 164, 566, 817).

The contribution of apical NHE3 mediated H^+ secretion to proximal tubule HCO_3^- absorption has been estimated by several approaches. Howlin et al and Chan and Giebisch found that luminal amiloride inhibited HCO_3^- absorption in the rat proximal convoluted tubule but the completeness of Na^+/H^+ antiporter inhibition remained unclear (152, 317). In separate experiments using high concentrations of luminal amiloride and a more potent analogue, t-butyl amiloride, it was concluded that $\sim 65\%$ of transepithelial HCO_3^- absorption was Na^+/H^+ antiporter dependent (573). In NHE3 $^{-/-}$ mice proximal tubule HCO_3^-

absorption was decreased to ~40% of control (624, 751) and the H⁺ pump inhibitor bafilomycin decreased HCO₃⁻ transport by 60% providing evidence for a component of H⁺-ATPase mediated transport (751). Disruption of the NHE3 *Slc9a3* gene may lead to compensatory changes in apical membrane H⁺-ATPase activity and potentially other NHE isoforms, making it difficult to assess the contribution of NHE3 to transepithelial bicarbonate absorption from these studies (162, 752). Moreover attempts to assess the contribution NHE3 to HCO₃⁻ absorption in tubules perfused in Na⁺-free solutions, or by inhibition of the Na⁺-K⁺-ATPase are also difficult to interpret because of their potential effect on altering the activity of basolateral NBCe1-A (see below). Until conditional knockout studies of NHE3 are done, the best assessment of the contribution of NHE3 relies on studies using inhibitors.

NHE8 may also contribute to bafilomycin-insensitive HCO₃⁻ absorption and has been localized to the proximal tubule by immunocytochemistry, Western blotting, and in situ hybridization (269, 270). It is currently thought that in neonates, NHE8 likely accounts for the majority of apical membrane NHE activity since NHE3 is poorly expressed and ethylisopropylamiloride (EIPA) doses that block NHE8 but not NHE3 suppress NHE activity (68, 98, 701). In adults, although NHE8 may be predominantly an intracellular organelle exchanger (98), studies in senile rats suggest that age-related decrease in proximal tubule acidification is due to a decrease in NHE8 expression rather than NHE3 or H⁺-ATPase activity (220). The exact mechanism for the age-related switch in apical NHE3 and NHE8 expression is unknown but appears to involve both thyroid hormone and glucocorticoids (63, 98, 234).

Organic Anion Transport—In the *Ambystoma* proximal tubule perfused *in vitro*, Siebens and Boron demonstrated the functional presence of an apical membrane Na⁺-lactate cotransporter in parallel with a basolateral membrane H⁺-lactate cotransporter (or equivalently an OH⁻/lactate exchanger) (652). In the rabbit S3 proximal tubule, Nakhoul and Boron reported apical membrane Na⁺-acetate cotransport activity, and both apical and basolateral H⁺-acetate cotransport (or equivalently OH⁻/acetate exchange) (519, 521). The low intracellular Na⁺ concentration would generate a chemical driving force for apical Na⁺-acetate uptake and subsequent equivalent H⁺ efflux (with acetate) across either the apical and/or basolateral membranes. Geibel examined the importance of this system to transepithelial H⁺ secretion and in the presence of physiologic HCO₃⁻-buffered solutions, concluded it does not play an important role (238).

Cl⁻/Base Exchange: CFEX (PAT1)—Lucci and Warnock examined the effect of anion transport inhibitors on the perfused rat proximal tubule and provided the first evidence that apical Na⁺/H⁺ exchange is coupled to Cl⁻/OH⁻ (or Cl⁻/HCO₃⁻ exchange) (465). Baum et al used luminal stilbene and furosemide inhibitors in the *in vitro* perfused rabbit proximal convoluted tubule to inhibit Cl⁻ absorption (60). Several labs showed that pH gradients can drive ³⁶Cl transport in voltage clamped apical membrane brush border vesicles and that Cl⁻ gradients could drive pH changes transport measured with acridine orange (131, 650, 756). Chen et al reported similar results using a Cl⁻ fluorescent probe (156). In contrast, Seifter et al failed to document pH gradient driven Cl⁻ transport under voltage clamped conditions

(641) and other groups were also unable to measure Cl^- -induced pH changes (142, 329). In the rabbit proximal straight tubule, Ishibashi et al were also unable to detect Na^+ -dependent apical Cl^- transport using Cl^- electrodes (326). Similarly, Schwartz was unable in the rabbit proximal tubule to demonstrate an effect of luminal Cl^- on luminal HCO_3^- appearance (631). Several studies have failed to document a significant effect of Cl^- removal on HCO_3^- transport and luminal DIDS (an inhibitor of anion exchange) has even been reported to stimulate HCO_3^- absorption (60, 129, 486, 607). Specifically, following removal of luminal and basolateral Cl^- , the steady state luminal HCO_3^- concentration in rabbit proximal convoluted and straight tubules perfused at slow flow rates was Cl^- independent (129, 486). These findings have been criticized in that apical and basolateral Cl^- was replaced with nitrate that can potentially be transported by anion exchangers (449). Additional studies however in proximal convoluted tubules showed that Cl^- substitution with isethionate also did not alter transepithelial HCO_3^- absorption (607).

Karniski and Aronson advanced the understanding of the underlying transport mechanisms in rabbit brush border vesicles, by first showing the presence of formate-driven Cl^- transport and Cl^- -driven formate transport whose activity was significantly greater than Cl^-/OH^- exchange possibly accounting for the previous discrepant results (351). Additional studies also revealed the presence of a Cl^- /oxalate exchange mechanism (350). In the *in vivo* perfused rat proximal tubule, Alpern and subsequently Baum in the rabbit proximal tubule confirmed the presence of apical Cl^- /formate exchange by measuring Cl^- -induced pH changes in the presence of formate (17, 59). Formate stimulated proximal tubule volume absorption in tubules perfused with high Cl^- /low HCO_3^- (low pH) solutions mimicking late proximal tubule composition (617, 618). The finding that there was associated cell volume changes and that basolateral DPC (Cl^- channel blocker) inhibited the formate stimulation of transport provided further evidence that the effect on NaCl transport was transcellular rather than paracellular (616, 745, 747). Wang et al reported additional evidence for such a transport process by showing that formate and oxalate stimulate volume and Cl^- absorption in the rat proximal tubule (745, 747). These studies led to the acceptance of a transport model with parallel apical Cl^- /formate and Na^+/H^+ (NHE3) exchangers and apical formic acid recycling. Saleh et al subsequently identified an apical formate/ OH^- exchange process (distinct from the Cl^- /formate exchange) that also could contribute formate recycling (605). As predicted, amiloride blocked Cl^- transport in formate/oxalate-free conditions, and EIPA inhibited the formate-stimulated Cl^- transport (60, 574, 745). Unlike formate, oxalate-stimulated Cl^- transport was not EIPA inhibitable and ultimately modeled to be mediated by sulfate/oxalate exchange coupled to Na^+ -sulfate cotransport (40, 41, 745).

An initial difficulty with the formic acid recycling model resulted from the fact that the measured formic acid permeability was too low to account for the observed transport rates (571). Unstirred layer effects resulting in formic acid accumulation above the bulk concentration was raised as a possible explanation, but further analysis suggested that brush border microvilli were too short to result in appreciable differences in the formic acid concentration (389, 390). Furthermore, the activity of apical Cl^- /formate and Cl^-/OH^- exchange was subsequently shown to vary among proximal tubule segments potentially accounting for the discrepancy in earlier vesicle studies. Accordingly, Kurtz et al were

unable to detect apical Cl^- /formate exchange in isolated perfused superficial rabbit proximal straight tubule monitoring Cl^- -induced changes in pHi however apical Cl^- / OH^- exchange activity was present (406). Furthermore, Sheu et al. detected apical Cl^- /formate exchange in the superficial proximal convoluted tubule but not in the juxtamedullary segment where Cl^- / OH^- exchange activity was found (648). These findings suggested that apical Cl^- / OH^- exchange may play a more important role in late proximal tubule segments where the generation of a lumen to peritubular Cl^- gradient favors transcellular electroneutral passive Cl^- transport.

Identifying the transporter(s) mediating proximal tubule apical anion exchange subsequently focused on the SLC26 gene family (15). Although initially SLC26A4 (pendrin) was initially thought to be a candidate, immunocytochemistry studies were negative (375) and in pendrin knockout mice brush border Cl^- /formate exchange was unaffected (352). A second protein, CFEX (or PAT1) encoded by SLC26A6 that mediated Cl^- /oxalate, Cl^- /formate, Cl^- / OH^- , and Cl^- / HCO_3^- exchange in expression systems, was shown to be localized to the apical membrane of mouse proximal tubules (160, 375, 799). In CFEX^{-/-} mice, oxalate stimulation of proximal tubule volume absorption was absent however formate stimulation of proximal tubule transport was partially impaired (754). Currently it is thought that CFEX (PAT1) mediates apical membrane Cl^- /oxalate exchange *in vivo*. Interestingly, in CFEX^{-/-} mice, baseline volume (NaCl) reabsorption (in the absence of added formate or oxalate) was normal suggesting that CFEX does not mediate significant Cl^- absorption via Cl^- / OH^- and Cl^- / HCO_3^- exchange. Although intracellular pH studies of CFEX^{-/-} proximal tubules demonstrated that CFEX mediates Cl^- / HCO_3^- exchange (754), under physiologic conditions, the transporter preferentially mediates Cl^- absorption via a Cl^- /oxalate exchange process. The reason other CFEX modes of anion transport are not apparent might result from the concentration and affinity of the various transported substrates *in vivo*. In addition, CFEX may interact with other proteins *in vivo* such as CAII, which can modulate its transport properties, and furthermore, expression studies may not simulate *in vivo* proximal tubule protein-protein interactions (26).

Mathematical models of proximal tubule transport showed that the number of Cl^- /formate exchangers has no significant effect on NaCl transport whereas changing the density of luminal NHE3 does have a significant effect (783). Subsequent studies in NHE3 deficient mice demonstrated that formate unlike oxalate failed to stimulate rat proximal tubule volume absorption (752). Petrovic et al provided evidence that formate may directly stimulate NHE3 in the mouse proximal tubule (560). Currently, it is unclear whether formate can directly interact with NHE3 or an accessory protein to modulate apical Na^+/H^+ exchange. In addition, it is unclear why deletion of *Slc26a6* in mice leads to decreased proximal tubule NHE3 activity.

Basolateral Membrane: NBCe1-A; NHE1; $\text{Cl}^-/\text{HCO}_3^-$ Exchange (Na^+ -Coupled and Na^+ -Independent; TASK2; H^+ -Coupled Organic Anion Transport (see above)

Electrogenic Na^+ -Base Cotransport: NBCe1-A—In the *in vivo* perfused rat proximal convoluted tubule, Burckhardt et al first suggested the presence of a basolateral membrane HCO_3^- conductance that was sensitive to SITS (128). Biagi and Sohtell reported similar

findings in rabbit proximal tubules (80) and Alpern demonstrated barium-sensitive K^+ induced pH_i changes that suggested the transport process carried a net negative charge (19). Definitive experiments by Boron and Boulpaep showed that basolateral transport was coupled to Na^+ (103). Studying the *in vitro* perfused salamander proximal tubule, lowering peritubular HCO_3^- led to the depolarization of the basolateral membrane, and decreased pH_i and intracellular Na^+ . Reduction of peritubular Na^+ also caused a similar basolateral membrane depolarization and decrease in pH_i. These changes were Cl^- -independent and stilbene inhibitable and were compatible with an electrogenic $Na^+-HCO_3^-$ cotransport process. Biagi and Sohtell (79, 80), Alpern (19, 20), Yoshitomi et al (812), and Sasaki et al (611), subsequently confirmed these findings in mammalian proximal tubules, and Lopes et al reported a similar transport process in *Necturus* proximal tubules (460). The $Na^+-HCO_3^-$ cotransport process was also studied in basolateral membrane vesicles by Akiba et al (6), Grassl and Aronson (271, 272), and Soleimani et al (661, 662). $^{22}Na^+$ flux was found to be stimulated by HCO_3^- gradients, Na^+ gradients could drive HCO_3^- flux (measured by pH changes with acridine orange), and the transport process was modulated by changes in vesicle potential and was stilbene-inhibitable. Soleimani et al concluded that the ion transport stoichiometry was 1:3 (positive charge: negative charge) and that the species transported were 1 Na^+ : 1 HCO_3^- : 1 CO_3^{2-} (662). Using BCECF to characterize the quantitate the activity of basolateral electrogenic $Na^+-HCO_3^-$ cotransport in S1-S3 superficial and juxtamedullary proximal tubule segments, Abuladze et al reported significant differences axially along the proximal tubule with the activity in S1 proximal tubules ~ 3-5 fold greater than S2 segments, and the activity in S2 segments ~ 8 times greater than the S3 segment (3).

Electrogenic $Na^+-HCO_3^-$ cotransport is now known to underlie the majority of base efflux in the mammalian proximal tubule. First, depolarizing the basolateral membrane blocks transepithelial HCO_3^- absorption (607). Second, Na^+ removal almost completely blocks transepithelial HCO_3^- absorption whereas Cl^- removal is without effect (129, 152, 155, 486, 607). Third, Cl^- removal has no effect on basolateral membrane H^+ /base permeability whereas Na^+ removal has a significant effect (19, 570, 613, 812). Fourth, basolateral stilbene inhibitors block HCO_3^- absorption (151). These studies argue against a significant role for basolateral Na^+ -coupled Cl^-/HCO_3^- exchange in modulating net transepithelial HCO_3^- transport (that would be expected to function as a Na^+ -base influx and Cl^- efflux process).

The molecular identity of the electrogenic $Na^+-HCO_3^-$ cotransporter was first determined by Romero et al who cloned the protein from *Ambystoma* kidney (595). The transporter (NBCe1-A) has also been identified in mammals (encoded by the *SLC4A4* gene) (4) and is expressed predominantly in S1 and S2 proximal tubules (480, 620). NBCe1-A is a homodimer with each monomer functioning independently (345). Each NBCe1-A monomer has a large N-terminal cytoplasmic region, a transmembrane region with 14 transmembrane regions, and a short C-terminal cytoplasmic tail (827, 828). Three NBCe1 variants (encoded by the *SLC4A4* gene) have been identified at the protein level in mammals (NBCe1-A, -B, -C) and two variants at the transcript level in mice (NBCe1-D and -E) that are expressed in various tissues (2, 4, 456, 482). These NBCe1 variants differ in their N-terminal regions and/or C-terminal tails, but share the same predicted topological structure and

transmembrane (transporting) region as NBCe1-A. Mutations in these NBCe1 variants might account for some of the extrarenal manifestations of patients with autosomal recessive proximal RTA (pRTA) due to mutations in the transporter (see below).

The direction of ion flux mediated by NBCe1-A in the proximal tubule is determined by the electrochemical driving force (μ) through the transporter, which is a function of the basolateral membrane potential, the substrate ion activities, and the NBCe1-A charge transport stoichiometry (407, 829). It has been implicitly assumed (in lieu of *in vivo* measurements of the native human transporter) that the NBCe1-A has charge transport stoichiometry of 1:3 ($\text{Na}^+\text{-CO}_3^{2-}\text{-HCO}_3^-$ cotransport mode) in the human proximal tubule largely based on *in vivo* measurements in other species (512, 564, 812). When expressed in a mammalian expression system (829) or oocytes (425), human NBCe1-A has a charge transport stoichiometry of 1:2. Various factors including phosphorylation state, (276), cell-type (275) and cell Ca^{2+} (511) have been reported to modulate its value and yet the relevance of these findings to the proximal tubule *in vivo* is unclear.

Recent electrophysiologic data using NO_3^- as a surrogate for CO_3^{2-} transport indicated that human NBCe1-A functioning with a charge transport stoichiometry of 1:2 mediates $\text{Na}^+\text{-CO}_3^{2-}$ cotransport (829). Preliminary data in the *Xenopus* oocyte expression system using surface pH electrodes has also suggested that rat NBCe1-A transports $\text{Na}^+\text{-CO}_3^{2-}$ (424) recent analysis using published rat proximal tubule data showed that NBCe1-A functioning as a $\text{Na}^+\text{-CO}_3^{2-}$ cotransporter (1:2 charge transport stoichiometry) would also be capable of mediating cell to peritubular base flux while being more sensitive than in the 1:3 mode of changes in the electrochemical gradient across the transporter (829).

Na^+/H^+ Exchange: NHE1—Boron and Boulpaep first reported apical and basolateral Na^+/H^+ exchange processes in the salamander proximal tubule (104). In renal cortical basolateral membrane fractions, initial studies could not detect Na^+/H^+ exchange activity (330, 602), whereas Grassl and Aronson subsequently reported pH gradient driven Na^+ uptake that was not amiloride inhibitable (271). Experiments in rat and rabbit proximal convoluted tubules, showed that peritubular Na^+ changes altered pH_i but the effect was not amiloride sensitive (19, 392). In the rabbit S3 proximal straight tubule, Kurtz provided evidence for an amiloride sensitive basolateral membrane Na^+/H^+ exchanger (401) and Giebel et al found a similar process in juxtamedullary S1 and S2 proximal tubules (237). In the rabbit, immunocytochemistry studies have detected basolateral NHE1 in all proximal tubule segments whereas rat proximal tubules are unstained (88). Cytoplasmic CAII can interact with NHE1 and the association between these proteins increases through an ERK-dependent pathway (446). Holthouser et al reported that in human kidney proximal tubule cell lines, ouabain-mediated regulation of $\text{Na}^+\text{-K-ATPase}$ activity is dependent on its interaction with NHE1 (310).

$\text{Na}^+\text{-Coupled and Na}^+\text{-Independent Cl}^-/\text{HCO}_3^-$ Exchange—The presence of basolateral $\text{Cl}^-/\text{HCO}_3^-$ exchange was first detected in *Necturus* proximal tubules (202). The Na^+ -dependence of the transport process was subsequently shown by Guggino (280). These and other studies concluded the transporter was a $\text{Na}^+\text{-coupled Cl}^-/\text{HCO}_3^-$ exchange process (279, 649) whereas Alpern and Chambers reported the presence of both $\text{Na}^+\text{-}$

coupled and Na^+ -independent $\text{Cl}^-/\text{HCO}_3^-$ exchange (20). Because the data could also have been mediated by electrogenic Na^+ -base transport in parallel with a basolateral Cl^- conductive pathway, additional studies were done which concluded that in the rat proximal tubule greater than 25% of basolateral H^+ /base flux is mediated by a Na^+ -coupled $\text{Cl}^-/\text{HCO}_3^-$ exchange process (570) and similar findings were obtained in the rabbit proximal straight tubule (613). The rabbit S3 proximal tubule was reported to have a Na^+ -independent $\text{Cl}^-/\text{HCO}_3^-$ exchange process only, in addition to electrogenic Na^+ -base transport (401, 520). Grassl et al studying rabbit basolateral membrane vesicles provided evidence for electroneutral $\text{Cl}^-/\text{HCO}_3^-$ exchange that was not Na^+ -coupled whereas in rat vesicles, Na^+ -dependence could be demonstrated (272). Similarly Chen and Verkman in rabbit basolateral membrane vesicles found evidence for a Na^+ -coupled $\text{Cl}^-/\text{HCO}_3^-$ exchange process (157). In summary, it appears that the basolateral membrane of the proximal tubule mediates $\text{Cl}^-/\text{HCO}_3^-$ exchange however its Na^+ -dependence still remains controversial. Moreover, there is currently no data that sheds light on the molecular nature of the specific protein(s) involved.

TASK2—TASK2 pH-sensitive K^+ channels are expressed on the basolateral membrane of proximal tubule cells and TASK2^{-/-} mice have metabolic acidosis and elevated urine pH (757). During HCO_3^- administration, the fractional excretion of Na^+ and Cl^- is increased. The acid-base abnormalities in these mice are thought to be potentially due to a depolarization of the basolateral membrane potential with impaired NBCe1-A base efflux leading to a pRTA phenotype. Thus far, mutations in TASK2 have not been reported in humans.

$\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ Permeability—Net luminal H^+ secretion in the proximal tubule is potentially modulated by its $\text{H}^+/\text{OH}^-/\text{HCO}_3^-$ permeability properties. The apical H^+/OH^- permeability using different preparations is ~ 0.5 cm/sec when corrected for surface area (328, 586). Passive H^+/OH^- flux is predicted to be quantitatively unimportant in the early proximal tubule where the luminal pH is initially 7.4 and cell pH is ~ 7.3 (571) whereas in the later portion of proximal tubule, where the luminal pH decreases to ~ 6.8 , it is estimated that $\sim 30\%$ of secreted H^+ would diffuse back into the cells. The basolateral membrane H^+/OH^- permeability of 0.67 cm/s approximates the apical membrane permeability and plays a very minor role in basolateral base efflux. The estimated transepithelial H^+/OH^- permeability (0.3 cm/sec) is identical with the transcellular H^+/OH^- permeability (calculated from the apical and basolateral values measured separately) suggesting that paracellular H^+/OH^- flux is insignificant (291, 571, 632). Estimates of the transepithelial HCO_3^- permeability of $1.7 - 3.5 \times 10^{-5}$ cm/sec in rat proximal convoluted tubule and 2.0×10^{-5} cm/sec in rabbit proximal convoluted tubules, also indicate that passive HCO_3^- flux in the early proximal tubule is insignificant (22, 153, 309). However, because of a lower luminal pH (and HCO_3^- concentration) in the late proximal tubule the calculated peritubular to luminal HCO_3^- backleak becomes more quantitatively significant (22).

Regulation of Proximal Tubule H⁺/Base Transporters and Whole Tubule Transport

NHE3

In the current topological model of NHE3, the cytoplasmic C-terminus functions as a regulatory domain that dynamically interacts with accessory proteins that modulate NHE3 function and membrane expression. Several NHE3 binding partners and regulatory proteins have been reported including megalin, PDZK1, DPPIV, and Shank2, however their functional role in most instances is unknown (86, 242-245, 295). Of the known regulatory factors, one of the most important is inhibition by cAMP-dependent phosphorylation via PKA (342, 503, 775). In experiments designed to stimulate cAMP acutely, early studies showed that the appearance of luminal cAMP as an index of cytosolic cAMP in S1 tubules correlated inversely with the proximal tubular HCO₃⁻ transport rate (452). NHE3 is also phosphorylated via PKC (349, 492, 774, 789), and regulated through G-proteins (11), and tyrosine kinase (244). Rapid regulation of NHE3 is mediated by changes in brush border insertion/retrieval NHE3 (94, 212, 305, 821). Early micropuncture and tubule perfusion studies had shown that PTH inhibits proximal tubule HCO₃⁻ absorption due to inhibition of transcellular H⁺ secretion (48, 322, 490, 491, 577). Luminal Na⁺/H⁺ exchange activity was decreased in response to PTH in perfused tubules (184, 289), which mirrored the effect of cAMP in tubule suspensions (190) and brush border vesicles (342). PTH decreased the sensitivity of NHE3 to changes in intracellular pH (342, 499, 566) and Na⁺/H⁺ exchange activity increased in parathyroidectomized animals (174). Vesicles studies showed that the effect of PTH on Na⁺/H⁺ exchange activity and NHE3 phosphorylation occurred within minutes (212). The scaffolding protein NHERF-1, which contains a C-terminal PDZ domain was shown to play a key role in cAMP mediated inhibition of NHE3 (778, 781, 831). Specifically, NHE3 and NHERF-1 are linked to the actin cytoskeleton with the PKA anchoring protein/actin binding protein ezrin (an ERM family actin binding protein) and the interactions between these proteins is required for PKA-mediated inhibition of NHE3 (181, 723, 776, 777, 779, 780, 782, 819, 831).

The signaling pathways involved in NHE3 modulation have been for the most part performed in the OKP cell line that expresses NHERF-1, and has Na⁺-H⁺ activity kinetics similar to NHE3 (32, 416). These cells were used as a model system for analyzing the interaction between NHERF-1 and NHE3, and NHERF-1 and ezrin (773, 777). Ezrin, NHERF-1 and NHE3 have been immunolocalized to the brush border of the rat proximal tubule where they form a protein complex (724). In mice lacking NHERF-1, NHE3 inhibition by cAMP is absent and the function of NHE3 in proximal tubule cells cultured from these mice is not regulated by PTH (182, 647, 779). cAMP inhibition of NHE3 also involves EPAC (exchange protein directly activated by cAMP) (311). Both EPAC and PKA each contribute ~50% of the cAMP inhibitory effect on NHE3 (514). NHERF-1 is required for cAMP inhibition of NHE3, and occurs via both PKA- and EPAC-dependent mechanisms (514). Finally PP1, a calyculin A sensitive phosphatase stimulates NHE3 by dephosphorylating the transporter (201).

The NHE3 residue(s) that are phosphorylated via PKA/cAMP of NHE3 remains controversial (503, 822). Kurashima reported the phosphorylation of NHE3-Ser⁶³⁴ and that Ser⁶⁰⁵ is also required for the effect but remains unphosphorylated (398). Zhao reported that PKA induces the phosphorylation of NHE3-Ser⁵⁵² and Ser⁶⁰⁵ (822). Further studies by Kocinsky showed that in normal rats, NHE3-Ser⁵⁵² is significantly more phosphorylated than Ser⁶⁰⁵ and is localized to the coated pit region of the brush border that has inactive NHE3 (378, 379). In addition, phosphorylation at both sites does not alter NHE3 function.

PTH may also modulate NHE3 activity via PKC through phospholipase C activation (318, 492, 774). Although the data is somewhat conflicting, acute PKC activation appears to increase transcellular HCO₃⁻ absorption whereas chronic PKC activation decreases transport (64, 744). Changes in cytosolic Ca²⁺ do not affect the intrinsic function of NHE3 unlike NHE1, which has a cytosolic calmodulin binding domain (residues 636-656) (729, 731).

Casein kinase 2 stimulates the basal activity of NHE3 by interacting with the C-terminal tail and phosphorylating Ser⁷¹⁹, which increases plasma membrane delivery of newly synthesized NHE3 (606). PTH decreases the membrane expression of the transporter over a period of hours (212). This effect has been observed in brush border vesicles and OKP cells (175). Disruption of the actin cytoskeleton leads to an increase in NHE3 expression (148). The exact role of actin in the retrieval/insertion of NHE3 in the brush border is unclear however its role in PKA mediated NHE3 phosphorylation has been shown (397, 681). NHE3 appears to transfer to the base of the microvilli following mediated PKA phosphorylation (806). Retrieval of NHE3 with megalin to a vesicle compartment is also involved (85, 86). PTH has also been shown to decrease NHE3 expression due to altered NHE3 mRNA stability with an associated inhibitory effect on the *SLC9A3* promoter (77).

Indirect binding of ezrin to NHE3 via NHERF-1 (NHE3 amino acids 585 to 689) also regulates the transporter signaling through cAMP (776, 818, 819), cGMP (144), Ca²⁺ (363, 427), and lysophosphatidic acid (161, 426). Ezrin can also bind directly to NHE3 (amino acids 475 to 589) affecting NHE3 trafficking (delivery from the synthetic pathway, basal exocytosis) and NHE3 movement within the brush border that may alter NHE3 endocytosis (145). Ca²⁺ induced NHE3 inhibition requires PKC α , which binds to E3KARP decreasing surface NHE3 expression likely by inducing endocytosis (427). CHP1 (calcineurin homologous protein) increases NHE3 membrane abundance and ezrin phosphorylation (Thr-567) (187). NHE3 basal activity is currently thought to be regulated by sequential effects of Akt and GSK3 acting on the same NHE3 C-terminal ezrin binding domain (658).

NBCe1-A

NBCe1-A has a unique N-terminal autostimulatory domain (ASD) that enhances the function of the transporter via an unknown mechanism (482). Mg²⁺ causes NBCe1-A rundown in *Xenopus* oocytes, that may involve a Mg²⁺-dependent phosphatase (5'-lipid phosphatase) which dephosphorylates PIP₂ (797). PIP₂ *per se* activates NBCe1-A, however the mechanism has not been determined (797). Various systemic and hormonal factors alter the expression and/or function of NBCe1-A. Angiotensin II has a biphasic effect on NBCe1 mediated transport (176, 177, 313, 558, 823) and NBCe1-A inhibition via the AT_{1B} receptor

leads to decreased NBCe1-A surface expression mediated by Ca^{2+} -insensitive PKC ϵ (558, 559). Endocytosis of NBCe1-A is modulated by PKCs (PKC $\alpha\beta\gamma$) and a novel PKC δ (54, 557, 815). ATP increases the NBCe1-A function via an unidentified protein kinase (511).

The function and/or expression of NBCe1-A is also modulated by PTH, norepinephrine, dopamine, and changes in blood pressure. PTH decreases the function NBCe1-A in the rat (possibly via a cAMP) whereas in the rabbit, PTH is without effect (609). Infusion of norepinephrine in rats increases the expression of the transporter (663) whereas dopamine decreases the activity of NBCe1-A (396). NBCe1-A expression in the SHR rat was increased ~2 fold above control rats via an unknown mechanism (664).

NBCe1-A expression is altered in various models of renal tubular acidosis (RTA). The transporter is upregulated in lithium induced distal RTA (dRTA) (364), however in hyperkalemic dRTA induced by ureteral obstruction its expression in the proximal tubule is decreased (741). In humans, hypothyroidism is associated with incomplete dRTA, and NBCe1-A expression is decreased in a rat hypothyroid model (505). Concomitant NH_4Cl loading increases the expression of NBCe1 (505). The calcineurin inhibitor FK506 which has been reported to cause both pRTA and hyperkalemic distal RTA, induces a decrease in NBCe1-A expression (506)

Carbonic anhydrase II (CAII) binds to NBCe1-A at a C-terminal D⁹⁸⁶NDD⁹⁸⁹ motif and Gross et al proposed that CAII and NBCe1 form an intermolecular transport metabolon (277). Pushkin et al. and Becker et al. provided further evidence for functional interaction between NBCe1 and CAII (69, 578). CAIV and CAIX were subsequently shown to bind extracellular loop 4 in NBCe1-B (and presumably NBCe1-A) (25, 531). Not all groups however have been able to demonstrate functional interaction with CAII (461, 563, 803). Interestingly, the loss of CAII function results in a more mild phenotype compared to patients with NBCe1 mutations (444, 659). Moreover, mutations in CAIV cause retinitis pigmentosa (RP17) without a pRTA phenotype having been reported (584). In *Xenopus* oocytes, CAI, CAII and CAIII stimulated NBCe1-A transport is due to carbonic anhydrase enzymatic activity rather than intramolecular proton shuttling (623).

Proximal Tubule HCO_3^- Transport

Acute and Chronic Changes in Acid-Base Chemistry and K^+ —The first studies by Malnic and Mello Aires addressing the dependence of luminal HCO_3^- concentration on HCO_3^- absorption using the split droplet technique showed that the kinetics were first order hence the rate of luminal H^+ secretion was proportional to the luminal HCO_3^- concentration (476). These findings were confirmed by other studies using a similar preparation, and in separate tubule perfusion experiments (22, 703). *In vivo* microperfusion studies that accounted for passive HCO_3^- flux demonstrated that there is a dependent on H^+ secretion below a luminal HCO_3^- concentration of ~ 45 meq/l (22). Since HCO_3^- cannot be transported by NHE3, these findings were likely dependent the effect of changes in the luminal HCO_3^- concentration on luminal pH and/or cell pH, since both factors can potentially alter the function of NHE3 and NBCe1-A independently.

Transepithelial HCO_3^- absorption varies inversely with the peritubular HCO_3^- concentration because of its effect on transcellular H^+ secretion with minimal effect on paracellular HCO_3^- diffusion (23, 151, 239, 493, 608, 703, 826). The effect had been attributed to a change in peritubular pH (with secondary changes in cell pH) modulating the function of basolateral Na^+ -base cotransport (NBCe1-A). An elevation of peritubular pCO_2 increases proximal tubule HCO_3^- absorption while a decrease has the opposite effect (169, 333, 438, 493, 608, 826). The exact mechanism is complex to address because of the rapid diffusion of CO_2 across all proximal tubular compartments with accompanying changes in pH and HCO_3^- . Moreover, apical and basolateral membrane H^+ /base transport processes have different sensitivities to changes in H^+ activity in each compartment. For example, apical NHE3 is allosterically regulated by an acidic cytoplasmic pH thereby increasing its Na^+ - H^+ exchange function greater than would be predicted from the concentration of the transported ions (42). A potential explanation for these findings is derived from studies of NHE1 that have identified residues 567-635 in the cytoplasmic C-terminus as playing an important role in cytoplasmic pH “sensing” (730).

NHE3 stimulation by a decrease in pH requires the activation of c-Src that complexes with Pyk2 (445). A second pathway involves ERK1,2/c-fos and both pathways converge to activate the endothelin ET_B receptor ultimately increasing NHE3 membrane expression (568). In both rat and rabbit proximal tubules, ET-1 stimulates apical Na^+ / H^+ exchange (203, 282). In OKP cells transfected with the ET_B receptor, ET-1 stimulates NHE3 activity and is associated with phosphorylation of the transporter and increased plasma membrane expression (164, 554, 555). Metabolic acidosis is associated with increased endothelin-1 (ET-1) mRNA expression (415, 448). Increased ET-1 expression appears to involve the AP-1 transcription factor (312). The metabolic acidosis stimulation of NHE3 activity is blocked in mice with targeted disruption of the ET_B receptor (415, 448). Tyrosine kinase, CaM kinase, and an increase in cell Ca^{2+} are involved (164, 165). Cortisol levels increase in metabolic acidosis and adrenalectomized rats fail to manifest a metabolic acidosis induced stimulation of Na^+ / H^+ activity (370, 481). Glucocorticoids act concomitantly with metabolic acidosis to increase NHE3 trafficking and plasma membrane expression whereas aldosterone has no effect (29).

Using “out of equilibrium” solutions, Boron et al was able to analyze the effect of pH, HCO_3^- , and CO_2 individually and found that pH per se was not a factor in modulating proximal tubule HCO_3^- transport (825, 826). The authors proposed that the basolateral membrane possesses HCO_3^- and CO_2 sensors that modulate transepithelial HCO_3^- transport. Genistein (tyrosine kinase inhibitor) eliminated the CO_2 sensitivity of HCO_3^- transport (825). Basolateral CO_2 sensing signaled through the apical membrane AT2 receptor via secreted angiotensin II peptide (824).

Chronic changes in systemic acid-base balance appear to have profound effect on proximal tubule H^+ /base transport processes and transepithelial HCO_3^- transport. These studies are complicated by the concomitant changes in renal hemodynamics and hormonal milieu that can independently modulate HCO_3^- transport. Kunau et al showed that chronic metabolic acidosis stimulates proximal tubule HCO_3^- absorption to a greater extent than more severe acute metabolic acidosis (394). Cogan showed that chronic hypercapnia is more potent than

acute hypercapnia (168). Cohn et al demonstrated in apical vesicles that the effect of chronic metabolic acidosis on Na^+/H^+ exchange activity was retained *in vitro* at pH 7.4 (memory effect) suggesting a change in membrane expression or structure (174). Further studies showed that the effect was indeed due to an increase in V_{max} (370, 690). Chronic metabolic acidosis also stimulated basolateral electrogenic Na^+ -base transport (NBCe1-A) that was retained *in vitro* at pH 7.4 (7, 570) whereas chronic metabolic alkalosis had the converse effect. NaCl and NaHCO_3 administration could modulate proximal tubule HCO_3^- absorption by decreasing the expression of NBCe1-A thereby contributing to the recovery from metabolic alkalosis and/or volume overload (33). Unlike the effect of metabolic acid-base disorders, mixed results have been reported in chronic respiratory acidosis (7).

Acid-base disorders are often accompanied by changes in K^+ balance. K^+ depletion increases bicarbonate absorption by ~ 25% increase whereas a lowering peritubular K^+ has no effect (151). These results differ from studies in OKP cells bathed in media with low K^+ where Na^+/H^+ exchange activity was stimulated that has been attributed to a decrease in pHi (31). In K^+ depletion, the expression of NBCe1-A is increased providing a potential mechanism for enhancing proximal tubule HCO_3^- transport during hypokalemic metabolic alkalosis (34).

Luminal Flow Rate, Extracellular Volume, Renin-Angiotensin System, Aldosterone and Catecholamines—The phenomenon of glomerulotubular balance where increased GFR leads to enhanced proximal tubule HCO_3^- absorption prevents significant renal excretion of HCO_3^- . Experiments in the perfused proximal tubule both *in vivo* and *in vitro* have shown that tubular HCO_3^- absorption is flow-dependent (24, 151, 196). Possible mechanisms include: 1) an alteration in the axial HCO_3^- concentration profile; 2) an effect on paracellular permeability; 3) effect on an unstirred layer in the brush border; 4) flow is transduced into a torque by its effect on brush border microvilli. Each of these potential factors has been studied. Chan et al (151) and Alpern et al (24) showed that changes in the axial HCO_3^- concentration profile can only account for a small portion of the effect. Du et al (196) and Chan et al (153) also showed that changes in paracellular permeability are not involved. Preisig et al showed that increased flow results in an increase in pHi due to enhanced apical Na^+/H^+ exchange (569). Involvement of apical Na^+/H^+ exchange was further shown in mice with targeted disruption of NHE3 where glomerulotubular balance is impaired (135). Du et al demonstrated flow-dependence in the mouse proximal tubule and concluded that the apical H^+ -ATPase activity was also involved (196). Other studies have documented an increase NHE3 or NBCe1-A activity in models of chronically increased luminal flow that are also associated with renal hypertrophy (uninephrectomy, increased protein intake) (173, 297, 524, 572).

A brush border acidic unstirred layer would be predicted to impede H^+ secretion wherein as luminal flow increases, the effect of the unstirred layer would diminish thereby enhancing luminal H^+ secretion. However, experiments in the rat proximal tubule indicated that there is not a significant brush border diffusion barrier (24, 569). An alternate mechanism was proposed whereby luminal flow varies linearly with the torque on the brush border microvilli (195, 196, 742). The flow-dependence of both volume and HCO_3^- absorption in the mouse proximal tubule was modeled where the magnitude of the torque was the “sensor”

that signaled enhanced H^+ secretion and HCO_3^- absorption. Over a 5-fold increase in luminal flow, microvillus torque increased 2-fold, and correlated with HCO_3^- transport changes. The actin cytoskeleton played an important role in affecting both NHE3 and H^+ -ATPase function. Du et al re-analyzed the rabbit tubule perfusion data of Burg and Orloff and found that the reported changes in flow would increase microvillus torque by ~43% that could account for the observed increase in volume absorption (130, 195). To account for these species differences, it was suggested that the rabbit tubule is more distensible so that the changes in luminal diameter prevent significant changes in microvillus torque that occur in the mouse.

Proximal tubule HCO_3^- absorption is also modulated by the extracellular volume status and pressure independent of altered luminal flow rate. Volume expansion has been shown decrease HCO_3^- absorption in the rat proximal tubule although the magnitude of the effect differs among studies (84, 170). The underlying mechanism involves several factors including an increase in paracellular backleak (due changes in paracellular HCO_3^- permeability), and modulation of net transcellular H^+ secretion resulting possibly from alteration in peritubular pH and/or HCO_3^- (18, 23). Interestingly, parathyroidectomized rats have a significantly decreased response without the underlying mechanism(s) having been defined (496). A decrease in dietary Na^+ was also shown to induce an increase in brush border Na^+/H^+ exchange *in vitro* due to an increase in V_{max} (504). Conversely, chronic dietary Na^+ loading induced the phosphorylation of NHE3-Ser⁵⁵², and redistribution of NHE3 to the base of the microvilli driven by myosin VI, and redistribution of the NHE3 regulator DPPIV (807).

Because the proximal tubule synthesizes and secretes angiotensinogen, the luminal concentration of angiotensin II is 1 – 10 nM while the systemic concentration is significantly less (0.01-0.1 nM) (108, 565, 642). In the *in vivo* perfused rat proximal tubule, angiotensin II was shown to enhance HCO_3^- absorption (452, 453). In the *in vitro* perfused rabbit proximal tubule, the mechanism of the stimulation involved both apical Na^+/H^+ exchange and basolateral Na^+ -base transport (236). Angiotensin II increased the V_{max} of apical Na^+/H^+ exchange and at higher concentrations also increased H^+ -ATPase activity in a proximal tubule OKP cell line by enhancing plasma membrane expression involving signaling through tyrosine kinase, PI3K, and P38 (140, 603). Angiotensin II has also been reported to increase apical H^+ -ATPase membrane vesicle insertion in rat proximal tubule fragments (727). Peritubular angiotensin II has a biphasic effect on transepithelial Na^+ transport in rat and rabbit proximal tubules where lower concentrations are stimulatory and higher concentrations that are likely unphysiologic are inhibitory; a similar phenomenon was shown in OKP cells examining NHE3 activity (134, 316, 629). The inhibitory effect was reported to involve angiotensin III interacting with AT2 receptor (284, 538, 539). Studies in rat tubules and cortical homogenates have concluded that angiotensin II signals through a decrease in adenylyl cyclase activity, however in OKP cells, NHE3 activity was stimulated by angiotensin II in the absence of changes in cAMP (134, 452, 794). In both OKP cells and rats treated chronically with angiotensin II, NHE3 activity was increased likely due to increased functional protein content (189, 802). Other investigators found no effect on NHE3 content but detected increased NBCe1-A protein that was blocked using an AT1

receptor blocker (700). An additional effect involves angiotensin II mediated translocation of NHE3, NHERF-1, ezrin and the H⁺-ATPase from the tip to the base of the brush border microvilli (433, 593). Others have shown that PKC, CaM kinase, c-Src, and PLA2 are involved in angiotensin II signaling and that a functional intracellular angiotensin system is active in the proximal tubule (205). Acid-base disturbances such as metabolic acidosis can also increase AT1 receptor expression and thereby modulate angiotensin II signaling (518).

Unlike angiotensin II, the plasma peptide ANG-(1–7) concentration increases following extracellular volume expansion (341, 551, 552). In vivo stationary microperfusion studies in rat proximal tubules have shown that luminal ANG-(1–7) has a biphasic dose-dependent effect on HCO₃[−] reabsorption mediated by the Mas receptor via NHE3 (143). Unlike angiotensin II high concentrations of ANG-(1–7) stimulate HCO₃[−] reabsorption whereas low concentrations inhibit transport. In the rat proximal straight tubule, basolateral ANG-(1–7) has a biphasic mediated by AT₁ receptors such that at physiologic levels, it increases HCO₃[−] reabsorption, whereas at high concentrations, it decreases transport (231).

Basolateral membrane transporters are also affected by renin-angiotensin signaling. Specifically, angiotensin II (see above) increases NBCe1-A activity in part via PKC and by increasing transporter density (594, 599). Additional effects include a stimulation of the basolateral Na⁺-K⁺-ATPase via phosphorylation (810), and an increase basolateral membrane K⁺ conductance (176). Apical Na⁺ uptake pathways appear to predominate since angiotensin II increases intracellular Na⁺ (587).

Aldosterone in rats increased proximal tubule NHE3 brush border protein content and stimulates transepithelial water reabsorption (393). In microperfusion studies of rat S2 proximal tubules, Pergher found that aldosterone (luminal or peritubular) stimulated HCO₃[−] absorption via glucocorticoid receptors through a nongenomic mechanism (556). In cultured human proximal tubule cells, aldosterone signaling was reported to be mediated through the EGF receptor (194). Basolateral NHE1 is regulated by aldosterone through genomic (aldosterone receptor) and nongenomic mechanisms. At lower concentrations, aldosterone increases intracellular Ca²⁺ and stimulates basolateral NHE1, whereas higher concentrations (10^{−6} M) decrease intracellular Ca²⁺ and inhibit the transporter (428). The glucocorticoid receptor appears to be involved in the rapid nongenomic effects of aldosterone on NHE1 activity that may play a role during changes in volume status.

Proximal tubule transport is reduced ~40% following renal denervation (72). Conversely, an increase in renal nerve activity can stimulate transport by ~30% (70). In the rabbit perfused proximal tubule, norepinephrine increased both HCO₃[−] and Cl[−] transport (67). In rabbit proximal tubule suspensions and rat proximal tubules perfused *in vivo*, catecholamines stimulated Na⁺/H⁺ exchange activity (150, 525). NHE3 stimulation is mediated by α₂ receptor binding associated with decreased adenylyl cyclase signaling through Gα_i, and α₁ receptor binding signaling through MAPK (451). β₂ receptor stimulation increases the association of NHERF-1 with the receptor resulting in increased Na⁺/H⁺ exchange by preventing cAMP driven downregulation of NHE3 (285). Sympathetic nerve activity may also stimulate NBCe1-A through PKC (598). Renal nerves also modulate local angiotensin synthesis and potentially AT1 receptor density (386, 580, 581). Proximal tubule Na⁺

transport is modulated by NO that is generated from renal nerve activity or interstitial endothelial cells or (447). NO release in pericapillary perfusion experiments increased apical Na^+/H^+ exchange (37). Mice lacking neuronal NOS have decreased proximal tubule HCO_3^- transport (749). Wang et al shed light on these findings by showing that tubule perfusion with sodium nitroprusside was dose-dependent where a low dose increased transport, a high dose decreased transport, and NOS inhibition decreased HCO_3^- absorption (743). NO signaling appears to be mediated through cGMP (338). NO stimulates cultured human proximal tubule cells expressing soluble guanylate cyclase to secrete cGMP resulting in inhibition of apical NHE3 through Src (612).

Dopamine, Glucocorticoids, Insulin, and Uroguanylin—Dopamine receptors are present on the luminal and basolateral membranes of the proximal tubule (527). In the proximal straight tubule, and tubule suspensions, dopamine inhibits Na^+ transport (73, 417). Other studies suggest that prior exposure to norepinephrine is required for the dopamine inhibitory effect (66). The inhibitory effect of dopamine is mediated through NHE3 (cAMP/PKA) and the Na^+/K^+ -ATPase (PKC) (241). In the rat *in vivo*, dopamine mediated phosphorylated NHE3 shifts from the brush border to a subapical region (379). Dopamine also decreases the activity of basolateral NBCe1-A (396).

Glucocorticoids stimulate NHE3 activity through several mechanisms. Acutely, dexamethasone administration increases NHE3 trafficking to the apical membrane following phosphorylation of NHE3-Ser⁶⁶³ via glucocorticoid-dependent kinase 1 (SGK1) (97, 739). CIC-5 appears to play a key role in basal and dexamethasone-stimulated exocytosis of NHE3 (450). As discussed in the context of metabolic acidosis, glucocorticoids synergistically increase NHE3 trafficking to the plasma membrane (29). Systemic administration of glucocorticoids or *in vitro* exposure of proximal tubule cell cultures or OKP cells stimulate NHE3 activity via the glucocorticoid receptor by modulating NHE3 mRNA abundance (62, 65, 740).

Insulin stimulates fluid absorption in the proximal tubule perfused *in vitro* (61) and alters NHE3 activity both acutely and chronically via separate mechanisms. Although the mechanism of acute activation is unknown, several potential pathways have been ruled out including phosphatidylinositol 3-kinase-serum- and glucocorticoid-dependent kinase 1 (PI3K-SGK1), cbl/CAP/TC10, NHE3-megalin interaction, NHE3 phosphorylation, and increased apical membrane trafficking (228). Chronically, insulin stimulates NHE3 activity via PI3K-SGK1 (228). Glucocorticoids also augment the effect of insulin on NHE3 potentially via the induction of SGK1 (374).

Uroguanylin is a member of the guanylin family of peptides that are involved in pH and volume regulation (434). Renoguanylin was first isolated from eels and is expressed in kidney and intestine (816). In microperfused rat proximal tubules renoguanylin decreased HCO_3^- absorption via an inhibition of apical Na^+/H^+ exchange and H^+ -ATPase via a protein kinase dependent pathway (434). In additional studies in rat proximal tubules and LLC-PK cells, NHE3 inhibition involved the activation of both cGMP/PKG and cAMP/PKA, NHE3 phosphorylation, and decreased NHE3 surface expression (435).

Isolated Familial Proximal Renal Tubular Acidosis: Mutations in NBCe1—Of the proteins responsible for H^+ /base transport in the proximal tubule, only mutations in NBCe1 cause a severe form of isolated pRTA as no mutations in NHE3 have been described and CAII mutations often cause a mild form of combined proximal and distal RTA. pRTA is the end result of both acquired and genetic abnormalities in multiple pathways that mediate proximal tubular HCO_3^- absorption (reviewed in detail (296)). Defective proximal tubule HCO_3^- absorption can occur in isolation or accompanied by other proximal tubule transport defects (Fanconi's syndrome). Mutations in NBCe1 are the only known cause of isolated familial pRTA (321). While other proximal tubular transport functions are normal, patients have extrarenal abnormalities including short stature, neurological findings (mental retardation, basal ganglia calcifications, migraine headaches), ocular abnormalities (cataracts, band keratopathy, glaucoma), tooth (enamel) defects, and elevated amylase and lipase (402). The extrarenal manifestations are likely due to local abnormalities in H^+ /base transport due to defective NBCe1 variant function. The additional role that systemic acidemia may play remains to be determined.

2 nonsense mutations (Q29X, W516X), 2 frameshift deletions (2311 delA, and a C-terminal tail 65bp-del), and 8 missense mutations (R298S, S427L, T485S, G486R, R510H, L522P, A799V, and R881C), have been reported. All cases reported thus far are inherited in an autosomal recessive fashion. Whether individuals heterozygous for NBCe1 mutations have subtle defects in proximal tubule bicarbonate transport and/or mild ocular, brain, growth, and pancreatic enzyme abnormalities is unknown. Mice with targeted disruption of NBCe1 have a shortened lifespan and neurologic and ocular manifestations do not occur suggesting that these abnormalities require a longer time period to develop (235, 413). Headaches have been reported in patients with R510H, L522P, R881C, 2311 delA, and 65bp-del mutations, and also in heterozygous family members of a patient with a 65 base-pair C-terminal deletion and the L522P mutation (680). It has been hypothesized that headaches are due to ER retained misfolded NBCe1-B in brain astrocytes associated with abnormal NMDA-mediated neuronal hyperactivity (804).

Autosomal dominant isolated pRTA has also been reported in patients with decreased bone density and short stature (109, 353, 429). Various proximal tubule proteins involved in H^+ /base transport including CA II, IV, and XIV; NBCe1; NHE3; NHE8; NHERF-1 and -2, and PAT1 (CFEX) were not found to have mutations in their coding regions. It remains to be determined whether any of these genes are mutated in their intron and promoter regions.

Loop of Henle

Approximately 10-20% of the filtered HCO_3^- load entering the nephron is absorbed in the loop of Henle (139) and the major site of active HCO_3^- absorption is thought to be the thick ascending limb (TAL; (126, 198, 255, 267). The initial segment of the loop of Henle, the thin descending limb, has apical and basolateral Na^+/H^+ exchangers that likely function to regulate pH_i (399); with little else known about its H^+ /base transport properties. Passive processes are potentially involved that promote HCO_3^- absorption in the thin descending limb. Osmotic equilibration due to water absorption in the thin descending limb (384, 385) would be predicted result in an increase in the concentration of HCO_3^- and an elevation of

the luminal $p\text{CO}_2$. The subsequent flux of CO_2 down its concentration (depending on the CO_2 permeability of the thin descending limb) would increase the luminal pH resulting in the back-titration of HCO_3^- from other luminal buffers (NH_4^+ and H_2PO_4^-). This process would effectively mediate the efflux of HCO_3^- in the form of CO_2 . The elevated luminal pH also plays an important role in the transport of ammonia from the thin limbs to the collecting ducts (257). Unlike the descending limb of Henle, the ascending limb is water impermeable with a high Na^+ and Cl^- permeability (323, 324). Na^+ transport is mediated by passive diffusion whereas Cl^- transport occurs via passive diffusion and an undefined carrier-mediated process. Whether HCO_3^- transport occurs in this segment is unknown.

The main site of HCO_3^- absorption is the medullary and cortical TAL (mTAL and cTAL respectively) (255, 267). In the TAL, in addition to apical NHE3 which plays the same role as in the proximal tubule, NHE2 and the vacuolar H^+ -ATPase have also been localized to the apical membrane however their role in HCO_3^- absorption is unclear (30, 89, 118, 414, 601, 678, 748, 766). As in the proximal tubule, luminal HCO_3^- is absorbed across the apical membrane indirectly via H^+ secretion mediated by NHE3 that combines with luminal HCO_3^- generating CO_2 and H_2O that are transferred across the apical membrane (414, 748). In addition, an apical K^+ -dependent HCO_3^- transport process that opposes transepithelial HCO_3^- absorption has also been described (763). K^+ -dependent ATP-ase activity has also been reported (814). TAL cells express cytoplasmic CAII and XV, apical and basolateral CAIV, basolateral CAXII, and apical CAXIV however species differences exist (411, 575, 600, 633). Loop diuretics such as furosemide that block cellular Na^+ uptake via NKCC2 stimulate HCO_3^- transport potentially via enhanced NHE3 activity or changes in membrane potential (255).

A basolateral stilbene-sensitive Na^+ -base transport process was initially described in the rat and mouse TAL however its molecular identity is unknown (357, 391). Both electrogenic Na^+ -base transport and the AE2 anion exchanger (16, 204, 224, 436, 583) that is also expressed on the basolateral membrane may potentially contribute to cell HCO_3^- efflux and transepithelial HCO_3^- absorption. HCO_3^- uptake across the basolateral membrane of the mTAL is currently thought to be mediated in part by NBCn1, an electroneutral stilbene-insensitive sodium bicarbonate cotransporter that could also play a role in cellular $\text{NH}_3 + \text{H}^+$ efflux and transepithelial tubular ammonia absorption (106, 410, 721). A basolateral K^+ -dependent HCO_3^- transport process has been reported in the rat mTAL (93). These transporters could play important roles in transepithelial HCO_3^- absorption, intracellular pH regulation, cell volume regulation, and $\text{NH}_3/\text{NH}_4^+$ transport although their exact roles in each of these processes is currently unresolved.

Basolateral membrane ClC-K1 and ClC-K2 chloride channels (or their human orthologs ClC-Ka and ClC-Kb, respectively), play an important role in transcellular Cl^- absorption (209, 211, 377, 706). Bartter's syndrome (a cause of Cl^- -resistant metabolic alkalosis) types 3 and 4 is caused by mutations in genes coding for ClC-Kb and the regulatory subunit, Barttin (90, 388, 655). Barttin promotes ClC-K insertion into the plasma membrane (209, 412, 732). Rarely Bartter's syndrome can be caused by compound mutations in ClC-Kb and ClC-Ka (619). The membrane expression and physiologic role differ among species and in

that $\text{ClC-K2}^{-/-}$ mice die at a young age and $\text{ClC-K1}^{-/-}$ mice have nephrogenic diabetes insipidus (478).

The TAL is also an important sight for transporting $\text{NH}_3/\text{NH}_4^+$ from the lumen to the peritubular interstitium for subsequent transport into the collecting duct (233, 248, 267, 358, 359, 764, 771). NH_4^+ is transported across the apical membrane via NKCC2 and potentially via the ROMK potassium channel. The very low apical membrane NH_3 permeability prevents the backflux of NH_3 from cell to lumen following apical NH_4^+ uptake (358). The basolateral Na^+/H^+ exchanger NHE4 mediates the cell peritubular efflux of NH_4^+ (NH_3 may also exit the TAL passively) (107, 315). In metabolic acidosis, NHE4 mediated NH_4^+ efflux is stimulated (107).

Regulation of Thick Ascending Limb Bicarbonate Transport

Acute and Chronic Changes in Acid-base Chemistry

As in the proximal tubule changes in acid-base status alter HCO_3^- transport in the TAL. Acute and chronic metabolic acidosis stimulate HCO_3^- transport in both rat microperfusion and isolated perfused mTAL studies (138, 249). Chronic metabolic acidosis also stimulates mTAL ammonia absorption (249). These effects are associated with the upregulation of NHE3 and AE2 (HCO_3^- transport); and NBCn1 (ammonia transport) (360, 410, 414, 583). In the cortex, sodium bicarbonate loading had no effect on AE2 expression unlike potassium bicarbonate loading that decreased AE2 expression, whereas in the outer medulla, sodium bicarbonate loading increased AE2 expression (583).

Basolateral Na^+/H^+ exchange mediated by NHE1 and NHE4 play an important role in modulating the function of apical NHE3, $\text{NH}_3/\text{NH}_4^+$ transport, and cell volume regulation (107, 268). Changes in NHE1 mediated transport are coupled to apical NHE3 via the actin cytoskeleton (759). In $\text{NHE1}^{-/-}$ mice, HCO_3^- transport in mTALs is significantly reduced (268). Furthermore, the inhibition of HCO_3^- transport by NGF is mediated through mPI3K-mTOR and ERK signaling pathways that result in NHE1 inhibition (253, 263, 765). NHE1 is also involved in the inhibition of mTAL HCO_3^- absorption by LPS (760) (see below).

Aldosterone, Glucocorticoids, Peptide Hormones, PGE2, and Sepsis

Although less studied than the proximal tubule, various hormones and peptides modulate HCO_3^- transport in the loop of Henle and TAL (139). *In vivo* microperfusion studies in rats showed that infusion of angiotensin II stimulates loop segment HCO_3^- absorption thought to involve the TAL although the exact tubule segment involved cannot be discerned from these studies (138). In similar microperfusion studies, the *in vivo* administration of high dose aldosterone and glucocorticoids restored HCO_3^- absorption to basal levels in adrenalectomized rats (704). Aldosterone inhibits TAL HCO_3^- transport via a nongenomic ERK-dependent inhibition of apical NHE3 (261, 264, 758) and angiotensin II inhibits HCO_3^- absorption through a cytochrome P-450 dependent pathway (260). Furthermore, glucagon and arginine vasopressin (AVP) inhibit transport via a cAMP dependent process (82, 252, 495). In the mTAL nerve growth factor inhibits HCO_3^- transport but stimulates absorption in the presence of AVP through distinct signaling pathways (253, 765). The effect of AVP on transport is distinct from the effect of osmolality (250, 251). Specifically,

an increase in osmolality inhibits luminal NHE3 and transport via tyrosine kinase signaling whereas a decrease in osmolality increases HCO_3^- transport through PI-3 kinase signaling (250, 251, 258, 766). The effect of AVP but not hyperosmolality can be reversed by PGE_2 through PKC and G protein (pertussis toxin-sensitive) pathways (254, 259). The stimulatory effect of hyposmolality on HCO_3^- absorption is blocked by AVP and cAMP (258). PTH directly inhibits transport in the isolated perfused rat mTAL (252). *In vivo* PTH stimulates HCO_3^- transport however this effect is complicated by potential changes in the luminal HCO_3^- concentration profile (83).

HCO_3^- absorption in the mTAL is inhibited by gram-negative lipopolysaccharide (LPS) by activation of TLR4 (toll-like receptor 4) and by gram-positive lipoteichoic acid and peptidoglycan through TLR2 (toll-like receptor 2) activation (262, 265, 266). Basolateral LPS inhibits blocks HCO_3^- absorption via through ERK-dependent inhibition of NHE3 (262, 761). Apical LPS inhibits HCO_3^- absorption via TLR4/MyD88-dependent activation of the PI3K-Akt-mTOR pathway coupled to inhibition of NHE1 (760, 762).

Distal Convoluted Tubule/Connecting Tubule

The distal convoluted tubule (DCT) is the next segment of the nephron that is capable of HCO_3^- transport. HCO_3^- absorption has been documented in micropuncture studies and in microperfusion experiments although the magnitude varies among studies (136, 137, 154, 395, 439, 440, 463). In the early DCT, apical H^+ secretion is mediated by Na^+/H^+ exchange (NHE2) and H^+ -ATPase activity (46, 149, 750). In the latter portion of the DCT, luminal H^+ secretion is mediated by H^+ -ATPase and H^+ - K^+ -ATPase transport (216, 717, 750, 785). It is difficult from micropuncture and *in vivo* microperfusion studies to clearly determine whether the next portion of the nephron, the connecting tubule (CT), and even the early cortical collecting duct (CCD) is accounting for the reported experimental results. The colonic form of the H^+ - K^+ -ATPase has been localized to the connecting and early cortical collecting duct (216, 717). In addition to HCO_3^- absorption, studies by Levine have also provided evidence for luminal HCO_3^- secretion (443, 750). This transport process is likely mediated by luminal pendrin, which will be discussed below.

Collecting Duct: Cortical, Outer Medullary and Inner Medullary Collecting Ducts

The collecting duct is subdivided anatomically into the CCD, and outer medullary (OMCD) and inner medullary collecting ducts (IMCD), with each tubule segment having unique H^+ /base transport properties. The CCD has a heterogenous cell population with ~ 60% of the cells being principal cells (PCs) that are responsible for Na^+ absorption, K^+ secretion, and water absorption in response to AVP (528). PCs also possess H^+ /base transport pathways that are involved in pH_i regulation rather than transcellular H^+ /base transport (146, 753, 768). Approximately 40% of the cells in this tubule segment are intercalated cells (ICs) that are further subdivided into Type A and Type B subtypes (sometimes referred to as alpha and beta IC cells respectively). Morphologically Type A ICs have apical microplicae and microvilli and intramembranous rod-shaped particles whereas Type B ICs have few apical microvilli and have basolateral rod-shaped particles (473, 714, 715). Type A ICs mediate

HCO_3^- absorption whereas Type B ICs secrete HCO_3^- . Type A cells have an apical H^+ -ATPase and basolateral anion exchange mediated predominantly by AE1, whereas Type B ICs have apical $\text{Cl}^-/\text{HCO}_3^-$ exchange mediated by pendrin (SLC26A4) and basolateral or diffuse H^+ -ATPase expression (14, 117, 118, 365, 500, 501, 712). These properties are best distinguished and characterized in the rat.

Among mammalian species, heterogeneity exists in the expression of these transporters and the overall HCO_3^- transport properties of ICs. For example, at the whole tubule level, rabbit CCDs secrete HCO_3^- whereas rat CCDs absorb HCO_3^- (44, 240, 457, 485, 626, 628, 669). Intramembranous rod-shaped particles that are thought to be associated with the H^+ -ATPase are present in the apical membrane of rat Type A ICs and absent in rat Type B ICs (714). In the majority of rabbit ICs, these particles are present apically and basolaterally to varying degrees (592). In addition, rat Type B ICs have basolateral H^+ -ATPase staining whereas in the rabbit staining is typically diffuse (628). Moreover, the majority of ICs in the rabbit unlike the rat have H^+ -ATPase restricted to cytoplasmic vesicles (713). In the mouse CCD, Type B ICs are less common than in rat or rabbit and lack a dark cytoplasm with numerous mitochondria typical of other species (685). In addition, in mouse Type B ICs, small vesicles without studs are localized beneath the apical gray zone (685). AE1 in the rat is localized to the basolateral membrane of Type A ICs (711, 712) whereas in the rabbit, it is primarily located in multivesicular bodies and cytoplasmic vesicles and occasionally on a portion of the basolateral membrane (471). Differences in IC CA isoform expression among various species also exists (see below).

Cells that don't fit the classic model of Type A or Type B ICs have also been described. Intercalated cells in the rabbit CCD termed γ or G cells have been identified functionally as having apical and basolateral Na^+ -independent $\text{Cl}^-/\text{HCO}_3^-$ exchange, however the percentage of ICs with this property is controversial (207, 772). While the apical transporter is thought to be pendrin, the transporter responsible for basolateral $\text{Cl}^-/\text{HCO}_3^-$ exchange is not known. Based on their specific ultrastructural properties, Kim et al (362) and Madsen et al (474) reported ICs in the CNT that differed from classic Type A and Type B ICs that are called non-A non-B ICs. Subsequent studies in mice and rats showed that these cells express apical H^+ -ATPase and pendrin, but not basolateral AE1 (362, 365, 685). Breton and Brown have suggested that they are a modified Type B IC (115). The categorization of ICs into various subtypes becomes more complex given that the expression of the H^+ /base transporters (pendrin, H^+ -ATPase, AE1) in these cells may be modulated by systemic acid-base status and other factors (55, 639).

In addition to the aforementioned transport proteins, Type A and Type B ICs possess other key transporters whose molecular identity is unclear. In Type A ICs, basolateral Cl^- channels have been demonstrated in the CCD and in other collecting duct segments (380-382, 516). In Type B ICs, a basolateral Cl^- conductive pathway contributes to transcellular Cl^- transport that is stimulated by cAMP, modulated by the intracellular HCO_3^- concentration, and is inhibited by DPC and anthracene-carboxylate (479, 625, 682, 683). ClC-3 mRNA is expressed in Type B ICs (530) however it has not been localized in these cells at the protein level. Type A and B cells also possess an unidentified basolateral

Na^+/H^+ exchange process likely plays a role in pH_i regulation rather than transcellular transport (500, 768).

Unlike the CCD, which can secrete or absorb HCO_3^- , the OMCD only absorbs HCO_3^- (44, 457, 487). In the outer stripe portion (OMCD_{os}) approximately 2/3 of the cells are PCs and 1/3 are Type A ICs (715). Type A ICs are distinguished from PCs in that they have a basolateral Cl^- conductive pathway, a lower basolateral membrane voltage, and essential no apical conductive pathways (381). In the OMCD Type A ICs generate the lumen positive transepithelial voltage due to electrogenic apical H^+ secretion; with basolateral HCO_3^- efflux mediated by the AE1 $\text{Cl}^-/\text{HCO}_3^-$ exchanger coupled to Cl^- recycling across the basolateral membrane (628). Given the lack of Cl^- chloride channels on the apical membrane, Cl^- is thought to enter the lumen paracellularly driven by the positive luminal transtubular potential. Type A ICs in the OMCD also express the basolateral SLC26A7 anion exchanger and endosomes containing the transporter are targeted to the basolateral membrane during hypertonicity and K^+ depletion (51, 561, 801). These cells also have intracellular H^+/K^+ -ATPase activity (see below) (790).

In the rat and rabbit in the OMCD_{os} and inner stripe portion of the OMCD (OMCD_{is}), ~33% of the cells are Type A ICs (715). In the rabbit OMCD_{is} , the outer portion of the OMCD_{is} has Type A ICs, however the inner portion has cells which cannot be classified as PCs or ICs morphologically and have been termed inner stripe cells (592). Furthermore in the OMCD_{is} there is cell-cell heterogeneity with regards to both H^+ -ATPase and AE1 expression (118, 627, 628). The cells on electron microscopy (in the rabbit) have apical intramembranous particles that are thought to represent H^+ -ATPase transporters (592) and there is functional evidence for both apical Na^+ -independent H^+ transport and basolateral $\text{Cl}^-/\text{HCO}_3^-$ exchange (112, 301, 302). An unidentified basolateral Na^+/H^+ exchange process is also present (112, 301). HCO_3^- absorption appears to be mediated in part by an apical H^+/K^+ -ATPase and a Na^+ -dependent process that has not been characterized (38, 39, 697, 811). Finally unlike the CCD, there is physiologic evidence for apical membrane carbonic anhydrase activity (670). The basolateral membrane of ICs in the OMCD_{is} is predominantly Cl^- -conductive (380, 516) that can potentially be modulated by HCO_3^- (302).

The IMCD is divided anatomically into either the initial (IMCD_i) and terminal segment (IMCD_t), or categorized into thirds (IMCD_1 , IMCD_2 , and IMCD_3) (167, 470). As in the OMCD, species differences are apparent. In the rabbit, the IMCD_1 cells are of a single type and resemble inner stripe cells with respect to their staining positive for carbonic anhydrase and Na^+/K^+ -ATPase (592), whereas in the rat ~ 10% of the cells are Type A ICs (167, 470). Like the OMCD_{is} , in the IMCD_i there is functional evidence for apical carbonic anhydrase activity (lack of an acid luminal disequilibrium pH) (734). Unlike Type A ICs, IMCD cells lack staining for the H^+ -ATPase, AE1, and H^+/K^+ -ATPase. In the rat IMCD_i and IMCD_t segments, there is evidence for H^+ secretion and HCO_3^- absorption via a Na^+ -independent mechanism (736, 737). Although there is conflicting evidence for an H^+ -ATPase, the data in rat IMCD_t suggests there is H^+/K^+ -ATPase activity (737). In the rabbit, no evidence for luminal H^+ secretion has been found (327). Basolateral base transport is mediated by a $\text{Cl}^-/\text{HCO}_3^-$ exchanger that is possibly AE2 (16, 217, 668, 676). In the basolateral membrane

there is functional evidence for a Na^+/H^+ exchange process (307) and a HCO_3^- conductance whose molecular identities are unknown (667).

H^+ -ATPase (V-ATPase), Pendrin, OXGR1, NDCBE, AE4, KCC4, H^+ - K^+ -ATPase, Carbonic Anhydrase

H^+ -ATPase—The H^+ -ATPase is assembled into two domains: A V_0 transmembrane domain and a V_1 cytoplasmic domain (ATP6V1) (111, 166, 221, 725). The V_1 domain is composed of eight separate subunits with a specific stoichiometry whereas the V_0 domain is composed of four subunits. ATP hydrolysis in the V_1 domain occurs at the B/A subunit interface. H^+ in the V_0 domain are translocated between the a- and c-ring. In addition to transcellular and organelle H^+ transport, additional functions include the regulation of GTPase activity (314), modulation of Wnt (180), and notch signaling (805).

Transcellular H^+ flux in the CCD depends on the cell membrane voltage, which is influenced by Cl^- (possibly directly) and by other electrogenic processes such as Na^+ transport in PCs. Although the ClC-5 Cl^- channel colocalizes with the apical H^+ -ATPase in Type A ICs, Cl^- channel activity has not been demonstrated in the native tubule (380, 381, 604). The ClC-5 chloride channel that co-localizes with the apical H^+ -ATPase in Type A ICs may play a role in endocytosis rather than transepithelial transport (604). Regulation of H^+ secretion in Type A ICs is also mediated by several factors including recycling of subapical H^+ -ATPase containing vesicles, modulation of the interaction between the two H^+ -ATPase domains which has been studied in lower organisms (188, 343, 344, 510, 722), and potentially changes in the coupling efficiency between the enzymatic and H^+ translocating machinery.

Recycling of vesicles represents an important mechanism for modifying the number of H^+ -ATPase molecules expressed on the cell surface (120, 472). The actin cytoskeleton plays an important role where subunits B1, B2, and C bind to actin (158, 308, 720). Inhibition of RhoA depolymerizes actin and increases H^+ -ATPase membrane expression (651). Changes in systemic acid-base status also modulate the recycling machinery. An increase in the pCO_2 stimulates exocytosis of vesicles that is inhibited by colchicine (634). During exocytosis the H^+ -ATPase interacts with the SNARE complex (47). Although the exact mechanism is unclear, vesicles coated on their surface with H^+ -ATPase enzyme direct the recycling process (116) or help to recruit coat proteins to endosomes (498).

Alterations in pH are potentially sensed by soluble adenylyl cyclase (sAC), which is expressed in ICs and has been found to co-immunoprecipitate with the H^+ -ATPase (545). It is hypothesized that an increase in cytoplasmic HCO_3^- concentration increases cAMP leading to PKA activation and increased plasma membrane H^+ -ATPase expression (542, 543, 545). The PDZ motif in C-terminal tail of the electroneutral sodium bicarbonate cotransporter NBCn1 also interacts with H^+ -ATPase B1 subunit in keeping with the presence of a HCO_3^- -sensing signaling pathway (579). The specific H^+ -ATPase subunit that is phosphorylated is unclear but has been shown to involve the A subunit (27, 286). pH sensing may signal through GPR4 as a mechanism for increasing cAMP (247, 546). Studies in other cell types suggest that purinergic receptor and Ca^{2+} signaling may also be involved (71, 159).

In Type A ICs, angiotensin II increases H^+ secretion likely via stimulation of translocation from a cytoplasmic pool to the apical membrane (550, 596, 728). PKC signaling is involved following binding of angiotensin II to the AT1 receptor (596). Aldosterone stimulates proton secretion in the collecting duct acutely via a nongenomic effect mediated by G α_q , PKC, and the ERK1/2 MAPK kinase pathway (792, 793). Chronic exposure to aldosterone increases translocation to the plasma membrane (792). Loss of aldosterone signaling leads to hyperkalemic dRTA partially due to loss of aldosterone-dependent signaling of H^+ -ATPase mediated Type A IC apical H^+ secretion (409).

Patients with mutations in the B1 subunit have autosomal recessive hypokalemic dRTA (347). Since B1 $^{-/-}$ mice don't have a hyperchloremic metabolic acidosis and it has been suggested that B1 subunit mutations in humans prevents the adaptive formation and expression of B2 subunit containing H^+ -ATPase transporters. (219, 547). Patients with mutations in the $\alpha 4$ subunit have autosomal recessive hypokalemic dRTA (660). Studies in $\alpha 4^{-/-}$ mice have led to the suggestion that patients with ATP6V0A4 gene mutations have both proximal and distal acidification defects accounting for their severe metabolic acidosis (304, 526). Loss of the $\alpha 4$ subunit also leads to downregulation of other H^+ -ATPase subunits and may affect the assembly of the transporter. The transcription factor Fox1 directly regulates the expression of the $\alpha 4$ subunit and Fox1 $^{-/-}$ mice have loss of the $\alpha 4$ subunit in addition to pendrin and AE1 (95, 718).

Pendrin—Pendrin, the apical Cl^-/HCO_3^- exchanger in Type B ICs and non-A, non B ICs (365, 597) was first shown to be an iodine transporter that was mutated in Pendred syndrome (goiter and deafness) (192, 210). Whether patients with Pendred syndrome have a defect in Type B IC HCO_3^- secretion is unknown. Mice with targeted disruption of pendrin either lack CCD HCO_3^- secretion (597) or have reduced apical Cl^-/HCO_3^- exchange and CCD HCO_3^- secretion (35) suggesting that at least in mouse, additional apical Cl^-/HCO_3^- exchange process(es) may be present. Loss of pendrin in mice also modulates the expression of several Na^+ , Ca^{2+} , H^+ /base and NH_3 transporting proteins (52, 366, 549). Pendrin overexpression in mice results in Cl^- -sensitive hypertension (335). The transporter in CCD Type B cells has the following halide affinity, $Cl^- \sim Br^- > I^- > F^-$ (206), with an apparent affinity for luminal Cl^- of ~ 10 mM in the rabbit (625). Pendrin mediated transport is modulated by changes in luminal Cl^- concentration and likely plays an important role in the recovery phase of Cl^- -sensitive metabolic alkalosis following administration of Cl^- -containing salts (232, 443, 669). Pendrin mediated HCO_3^- secretion via Type B ICs is acetazolamide inhibitable (696) and stimulated by cAMP (625, 626). cAMP/PKA signaling appears to mediate the isoproterenol (β -adrenergic agonist) induced increase in pendrin membrane expression/activity (45). Changes in whole body Cl^- balance, and water balance also modulate pendrin expression (507, 582, 710, 735). Pendrin expression is also affected by acid and base loading, and mineralocorticoids (223, 507, 562, 709, 726) but is not involved in Type B IC pHi regulation (767).

Recent studies have shown that luminal α -ketoglutarate (α -KG) via the OXGR1 α -KG receptor that is localized to the apical membrane of Type B and non-A, non-B ICs in the CNT and CCD modulates pendrin and Type B IC function (688). In isolated perfused CCDs, α -KG stimulates apical Cl^-/HCO_3^- exchange, and transtubular Na^+ and Cl^- transport that is

thiazide inhibitable. Base loading significantly increased urinary α -KG levels providing a luminal signal to enhance CCD Type B IC cell HCO_3^- secretion. $\text{OXGR1}^{-/-}$ mice do not respond to luminal α -KG with enhanced $\text{Cl}^-/\text{HCO}_3^-$ exchange and have reduced ability to excrete a base load.

ENaC, NDCBE, Pendrin and Type B IC Na^+ Transport—There is recent evidence suggesting that the function of Type B ICs modulates Na^+ transport in PCs, and that type B ICs per se transport Na^+ . $\text{Pendrin}^{-/-}$ mice have decreased expression of ENaC in PCs and changes in luminal HCO_3^-/pH (pendrin function) can alter ENaC Na^+ transport (549, 738). Secondly, studies in mice suggest that Type B ICs in addition to apical pendrin, have a thiazide sensitive Na^+ -coupled $\text{Cl}^-/\text{HCO}_3^-$ exchange process that is thought to be mediated by NDCBE (*Slc4a8*) providing an additional mechanism for coupling the transport of Na^+ and Cl^- (278, 437). The coupling of pendrin ($2\text{Cl}^-/2\text{HCO}_3^-$ exchange) and NDCBE (Na^+ -coupled $\text{Cl}^-/2\text{HCO}_3^-$) transport would be predicted to mediate the net cell influx of NaCl . Thirdly, Type B ICs $\text{B1}^{-/-}$ mice have renal loss of Na^+ , Cl^- , K^+ , polyuria, decreased ENaC expression in PCs, and decreased pendrin expression in Type B ICs (278). The expression of ENaC, the large conductance calcium-activated potassium channel, and aquaporin 2 were normalized when Type B IC ATP-triggered PGE2 paracrine signaling was blocked.

In addition to purported NDCBE, another member of the SLC4 base transporting family AE4 has been localized to CCD IC cells. AE4 was initially thought to mediate $\text{Cl}^-/\text{HCO}_3^-$ exchange (693) however its amino acid sequence more closely resembles members of the family that mediate Na^+ -coupled base transport (403). The expression pattern varies among studies and between species. AE4 has been immunolocalized to the apical and lateral membranes of rabbit CCD Type A ICs (376). A separate study in rabbit showed apical co-localization with peanut lectin (a Type B IC marker) (800). In the rat, AE4 is expressed on the basal and lateral membrane of CCD Type A and B ICs (376). In the mouse AE4 is localized to the basolateral membrane of CCD Type B ICs (95, 306). Mice with targeted disruption of AE4 do not have an overt phenotype however CCD Type B IC function was not reported (657).

KCC4—Mice with loss of the KCl cotransporter KCC4 develop dRTA and deafness suggesting a role for the transporter in basolateral Cl^- efflux in Type A ICs (100). In the rabbit, KCC4 has been immunolocalized to the basolateral membrane of cells in the DCT, CNT, that dissipates gradually with the transition to the CCT (708).

$\text{H}^+-\text{K}^+-\text{ATPase}$ — $\text{H}^+-\text{K}^+-\text{ATPases}$ are ATPases (E1, E2 or P-type) secrete H^+ in exchange for K^+ electroneutrally (179, 274, 281). The transporter consists of an α and β subunit, and 4 α subunit isoforms and 2 β subunit variants have been characterized. The known variants differ in their inhibitory profiles (127). The gastric or $\text{HK}\alpha 1$ variant is inhibited by SCH28080 and omeprazole and not ouabain. The colonic form, $\text{HK}\alpha 2$, is inhibited by ouabain but not SCH 28080. The $\text{HK}\alpha 4$ form found in skin is sensitive to ouabain and SCH 28080. There is evidence that $\text{HK}\alpha 1$ and $\text{HK}\alpha 2$ mediate the functional activity in the mouse collecting duct (132, 186, 274, 281). The inhibitory profile in the CCD and OMCD of K^+ -replete rats is the same as the gastric isoform (type I profile) whereas in K^+ -depleted animals activity was partially SCH 28080- and ouabain-sensitive and K^+ could be replaced by Na^+

(type III profile) (127). Type I activity was not detectable in mice with disruption of HK α 1 (*Atp4a* gene) and the type III activity was preserved, whereas in mice with loss of HK α 2 (*Atp12a*), type I activity was present and type III activity was lost (186). These findings and subsequent studies by Shao et al (646) suggest that either the kidney also expresses an HK α 2 different from the colonic variant, or technical experimental reasons account for the findings. Mice with loss of both HK α 1 and HK α 2 have normal acid-base parameters possibly due to upregulation of compensatory mechanisms (494, 666). Lynch et al examined acid secretion in ICs of CCDs from HK α 1^{-/-} and HK α 2^{-/-} or combined HK α 1^{-/-}/HK α 2^{-/-} mice and demonstrated a decrease in H⁺ transport in both Type A and Type B ICs (467, 468). Interestingly, 35-70% of HCO₃⁻ transport in the OMCD_{is} is thought to be H⁺-K⁺-ATPase dependent (283, 697). In the CCD, acute respiratory acidosis appears to stimulate H⁺-K⁺-ATPase transport (654). Calcitonin and isoproterenol also activate H⁺-K⁺-ATPase activity via ERK and cAMP (218, 418). HK α 2^{-/-} mice have reduced colonic ENaC activity, which is of interest since Na⁺-transporting PCs in the rabbit collecting duct (in addition to IC and OMCD cells) express HK α 2c (717). Whether the collecting duct H⁺-K⁺-ATPase plays a role in mediating Na⁺ transport *per se* is unclear, however a low Na⁺ diet has been reported to stimulate its activity (178, 653).

Carbonic Anhydrase—In general, HCO₃⁻ transport in the collecting duct is sensitive to carbonic anhydrase inhibition (457, 483, 484, 487, 591, 702). ICs in the collecting duct in all species stain for cytoplasmic CAII (119, 121, 458, 665) however differences exist among various species in the expression of membrane anchored CAIV (122, 459, 694). The expression of both CAII and CAIV is increased by metabolic acidosis (694, 791). In addition to playing a role in H⁺/base transport, studies in mice suggest that CAII may be important for the development of IC cells and for the expression of other transporters. In CAII^{-/-} mice there is a loss of Type A and Type B ICs (110), and pendrin expression is significantly decreased (679). Cells in the CCD lack luminal carbonic anhydrase and because of this, a spontaneous luminal disequilibrium pH is present due to net luminal H⁺ secretion. In addition to the CCD, the superficial DCT, OMCD_{os}, and IMCD lack luminal carbonic anhydrase. The basolateral membrane of PCs stains for CAXII whose significance is unclear but may play a role in pHi regulation (411, 638).

NH₃-NH₄⁺ Transport: Rhesus Proteins RhBG and RhCG and Sulfatides—The acidification of the collecting duct lumen is associated with the secretion of NH₃ from the interstitium and its protonation in the lumen to NH₄⁺ (770, 771). NH₃ permeates the collecting duct plasma membrane via specific rhesus (Rh) membrane proteins. RhBG and RhCG are expressed in murine and human collecting ducts in essentially the same pattern except that in humans, PCs don't express either protein. RhBG is expressed basolaterally in the CNT, Type A ICs, and non-A, non-B cells (294, 716), whereas RhCG is expressed on both the apical and basolateral cell membranes of the DCT, CNT, Type A ICs, and non-A, non-B cells (293, 361). Chronic metabolic acidosis increases RhCG expression in both OMCD and IMCD ICs (643) and in the OMCD_{is} increases apical RhCG expression in ICs and PCs (644). In mice lacking IC RhBG basal urinary NH₄⁺ excretion is essentially normal, however following acid-loading or K⁺-depletion, NH₄⁺ excretion is significantly decreased (91, 92). Mice with targeted loss of RhCG in the collecting duct or specifically in ICs have

normal acid–base parameters with only mildly reduced urinary NH_4^+ excretion, however following acid-loading, a more severe metabolic acidosis develops with decreased urinary NH_4^+ excretion (421, 422). In mice with deletion of RhCG in the collecting duct with intact RhCG in the CNT, a less severe phenotype exists suggesting that the CNT and possibly the late DCT contribute to RhCG mediated NH_3 permeation (422). Complete loss of RhCG leads to an 80% decrease in CCD NH_3 permeability (105). Mice with RhCG haploinsufficiency have a 40% decrease in CCD NH_3 permeability and following chronic acid loading have a blood HCO_3^- that is less than controls (105). Combined collecting duct $\text{RhBG}^{-/-}/\text{RhCG}^{-/-}$ mice have normal acid-base parameters but following acid-loading have a more severe metabolic acidosis than controls (423). These studies in mice establish the role of Rhbg and Rhcg in the renal response to extrarenal metabolic acidosis. Interestingly, Rh gas channels also mediate CO_2 permeation however the relevance to collecting duct H^+ /base transport is currently unknown (515). Recently, highly charged anionic glycosphingolipids (sulfatides) have been reported to play an important role in maintaining the high concentration of NH_4^+ in the papillary interstitium required for collecting duct NH_3 secretion (673).

Regulation of Collecting Duct Acidification

Acid-Base and Electrolyte (Na^+ , Cl^- , K^+ , and Ca^{2+}) Chemistry

In the CCD and OMCD, an acute decrease in basolateral pH and HCO_3^- stimulates bicarbonate absorption in perfused tubules *in vitro* (113, 327, 334). Raising the basolateral pCO_2 in the CCD has led to conflicting results that may be due to differences unidirectional HCO_3^- absorption versus secretion at baseline (113, 489). In the OMCD, raising the basolateral pCO_2 increases HCO_3^- absorption (489). Although not yet determined, the stimulation of transport could result changes in pHi and or the intracellular HCO_3^- concentration that kinetically alter the rate of apical H^+ -ATPase and basolateral $\text{Cl}^-/\text{HCO}_3^-$ exchange in Type A ICs. In addition, in the CCD, Type B and non-A, non-B IC HCO_3^- secretion could decrease. As discussed above other mechanisms including H^+ -ATPase pH sensing and recycling of H^+ -ATPase containing vesicles into the plasma membrane likely play an important role in modulating collecting duct acidification in response to acute acidosis (metabolic and respiratory). The insulin receptor-related receptor (InsR-RR) is expressed on the basolateral membrane of Type B ICs and non-A and non-B ICs that may act as an alkaline pH sensor signaling through ERK1/2, the actin cytoskeleton, and modulation of pendrin expression (56, 185, 536).

Chronic changes in systemic acid-base balance induce changes in collecting duct transport that can be demonstrated *in vitro* in dissected tubules (memory effect). This is best illustrated by studies showing the direction of net HCO_3^- transport (absorption and secretion) can be predictably altered in CCDs dissected from animals with metabolic acidosis or alkalosis (44, 484, 485, 671). Similar effects have been described in rat DCTs perfused *in vivo* (320, 442). HCO_3^- transport by the IMCD is also regulated in metabolic acidosis and alkalosis (74, 75, 702, 736). In contrast the OMCD does not appear to be regulated by systemic acid-base balance (420, 457, 487).

In the CCD, early studies suggested that Type A and Type B ICs interconvert in response to *in vivo* acid-base changes (636). Subsequent studies reported modulation of transport and anion exchanger expression in CCDs exposed to acidic peritubular fluid *in vitro* (576, 614, 639, 695). Al-Awqati and colleagues reported that the changes in the transport properties of ICs are due to hensen/DMBT1 that interacts with galectin-3 and cypA (9, 630, 635, 639, 684, 705) and cyclophilins are also reportedly involved (8, 230, 553). Other studies have suggested that there is remodeling of individual Type A and Type B cells and intermediary hybrid cells rather than a strict change in polarity (55, 117, 628, 714). A reversal of polarity is now considered unlikely given that IC expression of apical AE1 and basolateral pendrin have never been detected.

In the mouse, Type A ICs proliferate in response to metabolic acidosis that involves signaling through GDF-15 (200). More recent studies in rat have shown that chronic metabolic acidosis leads to Type A IC proliferation in the CCD and OMCD (784). PCs also proliferate but Type B ICs under the same conditions do not. Because acetazolamide induced the same response as NH_4Cl loading, systemic rather than urinary pH likely mediated the proliferative response (784).

Unlike peritubular pH, luminal pH changes are unlikely to alter intracellular pH in the DCT, CCD, and OMCD_{is} because of the low $\text{H}^+/\text{HCO}_3^-$ apical permeability (113, 154, 302, 668, 767). The lack of change in pH_i also suggests that the H^+ -ATPase in Type A and IMCD cells is rather insensitive to changes in luminal pH. In the DCT, luminal flow rate and the axial profile of the luminal HCO_3^- concentration modulate HCO_3^- absorption as in the proximal tubule (136, 154, 320). In Type B ICs that secrete HCO_3^- via pendrin, changes in the luminal HCO_3^- concentration also alter the rate of apical $\text{Cl}^-/\text{HCO}_3^-$ exchange (767).

Increasing the Na^+ concentration in the lumen of the CCD increases PC Na^+ transport and in the absence of transportable anions (particularly in the presence of aldosterone), leads to an increased transepithelial voltage (383, 419). In this setting although Type A ICs don't absorb Na^+ , it would be predicted that their apical membrane would depolarize due to circular intraepithelial current loops leading to enhanced H^+ secretion. Accordingly, diseases such as Liddle's syndrome where PC ENaC mediated Na^+ transport is increased (567) secondarily enhances Type A IC H^+ secretion. Conversely blocking ENaC in PCs with amiloride would in the CCD be predicted to hyperpolarize the apical membrane of ICs as has been shown in the OMCD_{os} (382) leading to decreased electrogenic H^+ secretion and HCO_3^- absorption (529).

The luminal Cl^- concentration is a factor determining the rate of Type B cell apical $\text{Cl}^-/\text{HCO}_3^-$ exchange, and the magnitude paracellular Cl^- transport (232, 443, 625, 669). In Cl^- -sensitive metabolic alkalosis, increasing the Cl^- concentration in the lumen of the CCD likely enhances pendrin (K_m for luminal $\text{Cl}^- \sim 5\text{-}10\text{ mM}$) mediated HCO_3^- secretion associated with an increase in basolateral H^+ -ATPase transport (711) contributing to the correction of the acid-base disorder. The apparent K_m of basolateral $\text{Cl}^-/\text{HCO}_3^-$ exchange for basolateral Cl^- in the OMCD_{is} is $\sim 115\text{ mM}$ (302) however whether clinical variations in the peritubular Cl^- concentration are physiologically relevant remains to be determined.

Determining the direct role of K^+ on collecting duct H^+ /base transport is complex given that its effect(s) *in vivo* are potentially mediated by changes in extracellular and/or intracellular K^+ . Moreover, in clinical disorders affecting K^+ balance, there often are accompanying changes in mineralocorticoid levels and systemic acid-base balance that can independently modulate tubule transport. In the DCT, hypokalemia enhances HCO_3^- absorption (136, 137). CCDs dissected from K^+ depleted animals (with hypoaldosteronism and metabolic acidosis) have increased HCO_3^- absorption *in vitro* whereas OMCD HCO_3^- absorption is decreased (488). K^+ depletion is also associated with increased collecting duct H^+ - K^+ -ATPase activity as discussed (193), and increased expression of apical H^+ -ATPase, basolateral AE1, and basolateral *Slc26a7* (50, 672).

Ca^{2+} (5.0 mM) and the CaSR agonist neomycin significantly enhance H^+ -ATPase activity in the OMCD (590). In TRPV5^{-/-} mice, activation of apical CaSR (Ca^{2+} -sensing receptor) increased luminal Ca^{2+} concentration in the DCT and collecting duct could potentially increase H^+ -ATPase activity lowering urinary pH and downregulate AQP2 (aquaporin 2) thereby preventing stone formation (590). In addition, disruption of the *Atpv1b1* gene in TRPV5^{-/-} mice results in normalization of urinary pH and tubular precipitation of Ca^{2+} -phosphate in the medullary collecting duct. These observations indicate that increased H^+ -ATPase-mediated urinary acidification in TRPV5^{-/-} mice protects against renal Ca^{2+} -phosphate stone formation.

Peptide Hormones, Renin-Angiotensin System, Mineralocorticoids, and Prostaglandins

In the rat DCT and CCD, ADH converts HCO_3^- secretion to net HCO_3^- absorption however the mechanism is unknown (82, 441, 689). Glucagon stimulates HCO_3^- secretion and/or decreases HCO_3^- absorption in the DCT and collecting duct potentially via signaling increase through cAMP (208, 495).

Isoproterenol or VIP increases HCO_3^- absorption in the rat DCT, (441). In the CCD, isoproterenol increases cAMP and stimulates HCO_3^- secretion likely via an increase in apical pendrin activity and the basolateral Cl^- conductance (299, 625, 626, 682). H^+ -ATPase expression is modulated by cAMP/PKA and sAC as discussed whereas signaling through PKC doesn't appear to play a role (247, 303, 542, 543, 545).

PTH increases adenylate cyclase in the DCT and CNT (508, 540). In metabolic acidosis PTH is increased and the increase in collecting duct acidification induced by PTH may be due to enhanced phosphate delivery where phosphate acts as a buffer or a non-reabsorbable anion (81, 497). Metabolic acidosis may also downregulate the PTH receptor (340).

In the DCT, ET-1 administration *in vivo* increases acidification by stimulating H^+ secretion and by decreasing and decreasing HCO_3^- secretion (following HCO_3^- loading) (787). Metabolic acidosis stimulates endothelial ET-1 secretion (788). In the DCT, inhibition of the ET_B receptor following protein loading (mild metabolic acidosis) *in vivo* blocks the aldosterone induced stimulation of H^+ secretion (356). In the CCD, binding of endothelin to the ET_B receptor decreases HCO_3^- secretion signaling through NO-guanylate cyclase (699).

Prorenin binds to the (pro)renin receptor ((P)RR) inducing renin-angiotensin system activity. The signal transduction pathway is independent of angiotensin II generation (522, 523). (P)RR has been identified as an accessory protein of the H^+ -ATPase (466) and colocalizes with the H^+ -ATPase in the collecting duct (5). Evidence in Madin-Darby canine kidney (MDCK) cells suggests that (P)RR and H^+ -ATPase activity is required for prorenin-induced activation of ERK1/2 (5). Activation of the H^+ -ATPase by aldosterone and angiotensin II is also dependent on ERK1/2 (596). Knockdown of (P)RR expression in MDK cells blocked prorenin and AVP-induced H^+ -ATPase activation perhaps via changes in membrane insertion, suggesting that (P)RR is needed for prorenin-dependent and -independent activation of the H^+ -ATPase (462). The importance of the prorenin induced stimulation *in vivo* is unclear given the unphysiologic concentration of prorenin required.

The effect of angiotensin II on collecting duct H^+ /base transport is complex. Angiotensin II has been reported to increase luminal HCO_3^- secretion in the rabbit CCD (769) and Type B IC H^+ -ATPase activity in the mouse CNT and CCD (728); whereas in rat, HCO_3^- absorption is increased (746). In the mouse, Pech et al (548, 550) have shown that angiotensin II directly stimulates the Type A IC H^+ -ATPase with increased plasma membrane expression and that Type B IC HCO_3^- secretion is secondarily increased potentially via a lowering of the luminal HCO_3^- concentration (increased driving force) and generation of luminal CO_2 (501). In the mouse although angiotensin II increases Type B IC H^+ -ATPase membrane expression, pendrin expression is unchanged (728) HCO_3^- absorption is decreased in the OMCD in response to angiotensin II (733).

There are several mechanisms by which mineralocorticoids stimulate luminal H^+ secretion in the collecting duct. In the CCD, mineralocorticoids stimulate PC ENaC mediated Na^+ transport that indirectly depolarizes the apical membrane of Type A ICs resulting in an increase in H^+ secretion (544). Furthermore there is a direct effect on Type IC H^+ -ATPase transport in both the CCD and OMCD_{is} (383, 674). H^+ secretion in the rat IMCD is also stimulated by mineralocorticoids (197). A nongenomic mechanism causes rapid translocation of H^+ -ATPase to the apical membrane signaling through $G(\alpha_q)$ protein-coupled receptors, and intracellular Ca^{2+} signaling, PKC, and ERK1/2 are involved (792). In addition, cAMP/PKA has a modulatory role. In mice lacking the V1a receptor that develop type 4 RTA fludrocortisone ameliorated the acidosis by restoring excretion of urinary ammonium via increased expression of H^+ - K^+ -ATPase and RhCG and decreased H^+ -ATPase expression (332). The activity and expression of the H^+ - K^+ -ATPase regulated by mineralocorticoids involves increased α_2 subunit mRNA (273). HCO_3^- secretion by the Type B IC is stimulated by mineralocorticoids (383, 674). This effect is blocked by ameliorating the accompanying metabolic alkalosis with acid-loading (232).

As discussed in Type B ICs, proton pump inactivation induces release of PGE₂ via calcium-coupled purinergic receptor activation (278). In the OMCD_{is} PGE₂ inhibits HCO_3^- absorption (300). Base loading increases the urinary excretion of PGI₂ associated with increased HCO_3^- secretion in the rat DCT via changes in cAMP signaling (786).

Isolated Familial Distal RTA: Mutations in CAII, AE1, H⁺-ATPase Subunits

Patients with mutations (23 mutations reported thus far) in CAII have autosomal recessive combined pRTA and dRTA because of the presence of CAII in proximal tubule cells, Type A intercalated cells, and medullary collecting duct cells. (57). CAII deficiency tends to have a higher incidence in the Arabian Peninsula possibly due to consanguinity with more than 70% of the cases described from this region (214, 319). A splice site mutation at the junction of exon 2–intron 2 (c.232+1 G > A) is commonly detected in these patients (215). Extrarenal manifestations include osteopetrosis, intracerebral calcification and developmental delay (645). Mild hearing loss due to a conductive defect has also been reported (517).

AE1 is a 12-14 transmembrane spanning protein whose topologic structure has been recently modeled after prokaryotic CIC channels (102). Two variants, kAE1 (kidney AE1) and eAE1 (erythrocyte AE1) are transcribed by the *SLC4A1* gene that differ in their N-terminal sequence; kAE1 is shortened by 65 residues in its N-terminus because of alternative promoter usage. eAE1 is expressed in the erythrocyte and kAE1 is localized to the basolateral membrane of Type A ICs (114, 621, 820). AE1 mutations causing dRTA were first reported by Bruce et al and subsequently by Karet *et al* (124, 348). Mutations in AE1 also cause hereditary spherocytosis (HS) without dRTA (336). Only in very rare instances do patients have both dRTA and red cell hemolysis (125). Thus far, 16 separate mutations in AE1 have been reported to cause dRTA with associated hypokalemia, nephrocalcinosis, hypercalciuria, and nephrolithiasis; both autosomal recessive and dominant inheritance has been described. Interestingly, the mutated proteins have been found to function essentially normally *in vitro* suggesting other mechanisms such as misfolding and ER or Golgi retention, or mistargeting to the plasma membrane play an important role. Importantly, there are no patient biopsy results that have addressed this issue; rather, interferences are made on mutated AE1 ER/golgi retention, or mistargeting using *in vitro* cell model systems. Loss of polarized expression has been reported in the context of the C-terminal R901X truncation mutant, AE1-G609R, and M909T, whereas intracellular retention in polarized cells characterizes the C479W, R589H, S613F, and G701D mutants (summarized in (13, 57)). AE1 is a dimer and it has been shown that in the context of autosomal dominant mutations, mixed mutant/wild-type dimers are retained intracellularly and cannot be rescued by the wild-type monomer (809). In contrast, mutations inherited autosomal recessively can be expressed on the plasma membrane normally as mutant/wild-type dimers (809). Recent experiments have shown that trafficking of kAE1 mutants can be partially restored *in vitro* by various treatments (163).

Karet et al described patients with autosomal recessive dRTA and sensorineural hearing loss that have mutations in the *ATP6V1B1* gene encoding the H⁺-ATPase B1 subunit (347). The hearing loss is progressive however occasionally a conductive component is present and bilaterally enlarged vestibular aqueducts have been reported (339). In addition, the patients are hypokalemic with nephrocalcinosis, and can have hypercalciuria and rickets. Karet et al also reported mutations in the *ATP6V0A1* gene encoding the $\alpha 4$ subunit in patients with dRTA and normal hearing (346). The patients also had hypokalemia with nephrocalcinosis, and several had hypercalciuria. Auditory brain stem response tests were normal. Subsequent studies of patients with mutations in the $\alpha 4$ subunit did have hearing loss, such that hearing

loss *per se* can no longer be considered diagnostic for a specific H⁺-ATPase subunit abnormality (502, 675, 707). Underlying mechanisms include abnormal pump assembly, impaired function, abnormal targeting, and loss of interaction with phosphofructokinase-1 (PFK-1) (229, 677, 808).

Acknowledgement

The preparation of this manuscript was supported in part by funds from the NIH (R01-DK077162), the Allan Smidt Charitable Fund, the Factor Family Foundation, and the Arvey Foundation.

Reference

1. Abramowitz MK, Hostetter TH, Melamed ML. Lower serum bicarbonate and a higher anion gap are associated with lower cardiorespiratory fitness in young adults. *Kidney Int.* 81:1033–1042. [PubMed: 22297677]
2. Abuladze N, Lee I, Newman D, Hwang J, Boorer K, Pushkin A, Kurtz I. Molecular cloning, chromosomal localization, tissue distribution, and functional expression of the human pancreatic sodium bicarbonate cotransporter. *J Biol Chem.* 1998; 273:17689–17695. [PubMed: 9651366]
3. Abuladze N, Lee I, Newman D, Hwang J, Pushkin A, Kurtz I. Axial heterogeneity of sodium-bicarbonate cotransporter expression in the rabbit proximal tubule. *Am J Physiol.* 1998; 274:F628–633. [PubMed: 9530281]
4. Abuladze N, Song M, Pushkin A, Newman D, Lee I, Nicholas S, Kurtz I. Structural organization of the human NBC1 gene: kNBC1 is transcribed from an alternative promoter in intron 3. *Gene.* 2000; 251:109–122. [PubMed: 10876088]
5. Advani A, Kelly DJ, Cox AJ, White KE, Advani SL, Thai K, Connelly KA, Yuen D, Trogadis J, Herzenberg AM, Kuliszewski MA, Leong-Poi H, Gilbert RE. The (Pro)renin receptor: site-specific and functional linkage to the vacuolar H⁺-ATPase in the kidney. *Hypertension.* 2009; 54:261–269. [PubMed: 19546380]
6. Akiba T, Alpern RJ, Eveloff J, Calamina J, Warnock DG. Electrogenic sodium/bicarbonate cotransport in rabbit renal cortical basolateral membrane vesicles. *J Clin Invest.* 1986; 78:1472–1478. [PubMed: 3782468]
7. Akiba T, Rocco VK, Warnock DG. Parallel adaptation of the rabbit renal cortical sodium/proton antiporter and sodium/bicarbonate cotransporter in metabolic acidosis and alkalosis. *J Clin Invest.* 1987; 80:308–315. [PubMed: 3038953]
8. Al-Awqati Q, Gao XB. Differentiation of intercalated cells in the kidney. *Physiology (Bethesda).* 26:266–272. [PubMed: 21841074]
9. Al-Awqati Q, Vijayakumar S, Hikita C, Chen J, Takito J. Phenotypic plasticity in the intercalated cell: the hensin pathway. *Am J Physiol.* 1998; 275:F183–190. [PubMed: 9691006]
10. Al-Bataineh MM, Gong F, Marciszyn A, Myerburg MM, Pastor-Soler NM. Regulation of the Proximal Tubule Vacuolar H⁺-ATPase by PKA and AMP-Activated Protein Kinase. *Am J Physiol Renal Physiol.*
11. Albrecht FE, Xu J, Moe OW, Hopfer U, Simonds WF, Orlowski J, Jose PA. Regulation of NHE3 activity by G protein subunits in renal brush-border membranes. *Am J Physiol Regul Integr Comp Physiol.* 2000; 278:R1064–1073. [PubMed: 10749796]
12. Alexander RT, Grinstein S. Tethering, recycling and activation of the epithelial sodium-proton exchanger, NHE3. *J Exp Biol.* 2009; 212:1630–1637. [PubMed: 19448073]
13. Alper SL. Familial renal tubular acidosis. *J Nephrol.* 23(Suppl 16):S57–76. [PubMed: 21170890]
14. Alper SL, Natale J, Gluck S, Lodish HF, Brown D. Subtypes of intercalated cells in rat kidney collecting duct defined by antibodies against erythroid band 3 and renal vacuolar H⁺-ATPase. *Proc Natl Acad Sci U S A.* 1989; 86:5429–5433. [PubMed: 2526338]
15. Alper SL, Sharma AK. The SLC26 gene family of anion transporters and channels. *Mol Aspects Med.* 34:494–515. [PubMed: 23506885]

16. Alper SL, Stuart-Tilley AK, Biemesderfer D, Shmukler BE, Brown D. Immunolocalization of AE2 anion exchanger in rat kidney. *Am J Physiol.* 1997; 273:F601–614. [PubMed: 9362338]
17. Alpern RJ. Apical membrane chloride/base exchange in the rat proximal convoluted tubule. *J Clin Invest.* 1987; 79:1026–1030. [PubMed: 3558814]
18. Alpern RJ. Bicarbonate-water interactions in the rat proximal convoluted tubule. An effect of volume flux on active proton secretion. *J Gen Physiol.* 1984; 84:753–770. [PubMed: 6096481]
19. Alpern RJ. Mechanism of basolateral membrane H⁺/OH⁻/HCO₃⁻ transport in the rat proximal convoluted tubule. A sodium-coupled electrogenic process. *J Gen Physiol.* 1985; 86:613–636. [PubMed: 2999293]
20. Alpern RJ, Chambers M. Basolateral membrane Cl⁻/HCO₃⁻ exchange in the rat proximal convoluted tubule. Na-dependent and -independent modes. *J Gen Physiol.* 1987; 89:581–598. [PubMed: 2953859]
21. Alpern RJ, Chambers M. Cell pH in the rat proximal convoluted tubule. Regulation by luminal and peritubular pH and sodium concentration. *J Clin Invest.* 1986; 78:502–510. [PubMed: 3016029]
22. Alpern RJ, Cogan MG, Rector FC Jr. Effect of luminal bicarbonate concentration on proximal acidification in the rat. *Am J Physiol.* 1982; 243:F53–59. [PubMed: 7091367]
23. Alpern RJ, Cogan MG, Rector FC Jr. Effects of extracellular fluid volume and plasma bicarbonate concentration on proximal acidification in the rat. *J Clin Invest.* 1983; 71:736–746. [PubMed: 6826733]
24. Alpern RJ, Cogan MG, Rector FC Jr. Flow dependence of proximal tubular bicarbonate absorption. *Am J Physiol.* 1983; 245:F478–484. [PubMed: 6414312]
25. Alvarez BV, Loiselle FB, Supuran CT, Schwartz GJ, Casey JR. Direct extracellular interaction between carbonic anhydrase IV and the human NBC1 sodium/bicarbonate co-transporter. *Biochemistry.* 2003; 42:12321–12329. [PubMed: 14567693]
26. Alvarez BV, Vilas GL, Casey JR. Metabolon disruption: a mechanism that regulates bicarbonate transport. *The EMBO journal.* 2005; 24:2499–2511. [PubMed: 15990874]
27. Alzamora R, Thali RF, Gong F, Smolak C, Li H, Baty CJ, Bertrand CA, Auchli Y, Brunisholz RA, Neumann D, Hallows KR, Pastor-Soler NM. PKA regulates vacuolar H⁺-ATPase localization and activity via direct phosphorylation of the a subunit in kidney cells. *J Biol Chem.* 285:24676–24685. [PubMed: 20525692]
28. Ambuhl PM, Amemiya M, Danczkay M, Lotscher M, Kaissling B, Moe OW, Preisig PA, Alpern RJ. Chronic metabolic acidosis increases NHE3 protein abundance in rat kidney. *Am J Physiol.* 1996; 271:F917–925. [PubMed: 8898023]
29. Ambuhl PM, Yang X, Peng Y, Preisig PA, Moe OW, Alpern RJ. Glucocorticoids enhance acid activation of the Na⁺/H⁺ exchanger 3 (NHE3). *J Clin Invest.* 1999; 103:429–435. [PubMed: 9927505]
30. Amemiya M, Loffing J, Lotscher M, Kaissling B, Alpern RJ, Moe OW. Expression of NHE-3 in the apical membrane of rat renal proximal tubule and thick ascending limb. *Kidney Int.* 1995; 48:1206–1215. [PubMed: 8569082]
31. Amemiya M, Tabei K, Kusano E, Asano Y, Alpern RJ. Incubation of OKP cells in low-K⁺ media increases NHE3 activity after early decrease in intracellular pH. *Am J Physiol.* 1999; 276:C711–716. [PubMed: 10069999]
32. Amemiya M, Yamaji Y, Cano A, Moe OW, Alpern RJ. Acid incubation increases NHE-3 mRNA abundance in OKP cells. *Am J Physiol.* 1995; 269:C126–133. [PubMed: 7631739]
33. Amlal H, Chen Q, Greeley T, Pavelic L, Soleimani M. Coordinated down-regulation of NBC-1 and NHE-3 in sodium and bicarbonate loading. *Kidney Int.* 2001; 60:1824–1836. [PubMed: 11703600]
34. Amlal H, Habo K, Soleimani M. Potassium deprivation upregulates expression of renal basolateral Na⁺-HCO₃⁻ cotransporter (NBC-1). *Am J Physiol Renal Physiol.* 2000; 279:F532–543. [PubMed: 10966933]
35. Amlal H, Petrovic S, Xu J, Wang Z, Sun X, Barone S, Soleimani M. Deletion of the anion exchanger Slc26a4 (pendrin) decreases apical Cl⁻/HCO₃⁻ exchanger activity and impairs bicarbonate secretion in kidney collecting duct. *Am J Physiol Cell Physiol.* 299:C33–41. [PubMed: 20375274]

36. Amodu A, Abramowitz MK. Dietary acid, age, and serum bicarbonate levels among adults in the United States. *Clin J Am Soc Nephrol*. 8:2034–2042. [PubMed: 24052219]
37. Amorena C, Castro AF. Control of proximal tubule acidification by the endothelium of the peritubular capillaries. *Am J Physiol*. 1997; 272:R691–694. [PubMed: 9124496]
38. Armitage FE, Wingo CS. Luminal acidification in K-replete OMCDi: contributions of H-K-ATPase and bafilomycin-A1-sensitive H-ATPase. *Am J Physiol*. 1994; 267:F450–458. [PubMed: 8092259]
39. Armitage FE, Wingo CS. Luminal acidification in K-replete OMCDi: inhibition of bicarbonate absorption by K removal and luminal Ba. *Am J Physiol*. 1995; 269:F116–124. [PubMed: 7631825]
40. Aronson PS. Role of SLC26A6-mediated Cl(–)-oxalate exchange in renal physiology and pathophysiology. *J Nephrol*. 23(Suppl 16):S158–164. [PubMed: 21170874]
41. Aronson PS, Giebisch G. Mechanisms of chloride transport in the proximal tubule. *Am J Physiol*. 1997; 273:F179–192. [PubMed: 9277578]
42. Aronson PS, Nee J, Suhm MA. Modifier role of internal H⁺ in activating the Na⁺-H⁺ exchanger in renal microvillus membrane vesicles. *Nature*. 1982; 299:161–163. [PubMed: 7110335]
43. Aronson PS, Suhm MA, Nee J. Interaction of external H⁺ with the Na⁺-H⁺ exchanger in renal microvillus membrane vesicles. *J Biol Chem*. 1983; 258:6767–6771. [PubMed: 6304054]
44. Atkins JL, Burg MB. Bicarbonate transport by isolated perfused rat collecting ducts. *The American journal of physiology*. 1985; 249:F485–489. [PubMed: 4051002]
45. Azroyan A, Morla L, Crambert G, Laghmani K, Ramakrishnan S, Edwards A, Doucet A. Regulation of pendrin by cAMP: possible involvement in beta-adrenergic-dependent NaCl retention. *Am J Physiol Renal Physiol*. 302:F1180–1187. [PubMed: 22262479]
46. Bailey MA, Giebisch G, Abbiati T, Aronson PS, Gawenis LR, Shull GE, Wang T. NHE2-mediated bicarbonate reabsorption in the distal tubule of NHE3 null mice. *J Physiol*. 2004; 561:765–775. [PubMed: 15604231]
47. Banerjee A, Shih T, Alexander EA, Schwartz JH. SNARE proteins regulate H(+)–ATPase redistribution to the apical membrane in rat renal inner medullary collecting duct cells. *J Biol Chem*. 1999; 274:26518–26522. [PubMed: 10473613]
48. Bank N, Aynedjian HS. A micropuncture study of the effect of parathyroid hormone on renal bicarbonate reabsorption. *J Clin Invest*. 1976; 58:336–344. [PubMed: 956369]
49. Bank N, Aynedjian HS, Mutz BF. Proximal bicarbonate absorption independent of Na⁺-H⁺ exchange: effect of bicarbonate load. *Am J Physiol*. 1989; 256:F577–582. [PubMed: 2539746]
50. Barone S, Amlal H, Kujala M, Xu J, Karet F, Blanchard A, Kere J, Soleimani M. Regulation of the basolateral chloride/base exchangers AE1 and SLC26A7 in the kidney collecting duct in potassium depletion. *Nephrol Dial Transplant*. 2007; 22:3462–3470. [PubMed: 17804457]
51. Barone S, Amlal H, Xu J, Kujala M, Kere J, Petrovic S, Soleimani M. Differential regulation of basolateral Cl[–]/HCO₃[–] exchangers SLC26A7 and AE1 in kidney outer medullary collecting duct. *J Am Soc Nephrol*. 2004; 15:2002–2011. [PubMed: 15284286]
52. Barone S, Amlal H, Xu J, Soleimani M. Deletion of the Cl[–]/HCO₃[–] exchanger pendrin downregulates calcium-transporting proteins in the kidney and causes calcium wasting. *Nephrol Dial Transplant*. 27:1368–1379. [PubMed: 21873623]
53. Barratt LJ, Rector FC Jr, Kokko JP, Seldin DW. Factors governing the transepithelial potential difference across the proximal tubule of the rat kidney. *J Clin Invest*. 1974; 53:454–464. [PubMed: 11344559]
54. Bartolo RC, Harfoot N, Gill M, McLeod BJ, Butt AG. Secretagogues stimulate electrogenic HCO₃[–] secretion in the ileum of the brushtail possum, *Trichosurus vulpecula*: evidence for the role of a Na⁺/HCO₃[–] cotransporter. *J Exp Biol*. 2009; 212:2645–2655. [PubMed: 19648410]
55. Bastani B, Purcell H, Hemken P, Trigg D, Gluck S. Expression and distribution of renal vacuolar proton-translocating adenosine triphosphatase in response to chronic acid and alkali loads in the rat. *The Journal of clinical investigation*. 1991; 88:126–136. [PubMed: 1829094]
56. Bates CM, Merenmies JM, Kelly-Spratt KS, Parada LF. Insulin receptor-related receptor expression in non-A intercalated cells in the kidney. *Kidney international*. 1997; 52:674–681. [PubMed: 9291186]

57. Batlle D, Haque SK. Genetic causes and mechanisms of distal renal tubular acidosis. *Nephrol Dial Transplant*. 27:3691–3704. [PubMed: 23114896]
58. Baum M. Axial heterogeneity of rabbit proximal tubule luminal H⁺ and basolateral HCO₃⁻ transport. *Am J Physiol*. 1989; 256:F335–341. [PubMed: 2537027]
59. Baum M. Effect of luminal chloride on cell pH in rabbit proximal tubule. *Am J Physiol*. 1988; 254:F677–683. [PubMed: 3364577]
60. Baum M. Evidence that parallel Na⁺-H⁺ and Cl⁻-HCO₃⁻(OH⁻) antiporters transport NaCl in the proximal tubule. *Am J Physiol*. 1987; 252:F338–345. [PubMed: 3028174]
61. Baum M. Insulin stimulates volume absorption in the rabbit proximal convoluted tubule. *J Clin Invest*. 1987; 79:1104–1109. [PubMed: 3549779]
62. Baum M, Amemiya M, Dwarakanath V, Alpern RJ, Moe OW. Glucocorticoids regulate NHE-3 transcription in OKP cells. *Am J Physiol*. 1996; 270:F164–169. [PubMed: 8769835]
63. Baum M, Biemesderfer D, Gentry D, Aronson PS. Ontogeny of rabbit renal cortical NHE3 and NHE1: effect of glucocorticoids. *Am J Physiol*. 1995; 268:F815–820. [PubMed: 7771509]
64. Baum M, Hays SR. Phorbol myristate acetate and dioctanoylglycerol inhibit transport in rabbit proximal convoluted tubule. *Am J Physiol*. 1988; 254:F9–14. [PubMed: 2827521]
65. Baum M, Moe OW, Gentry DL, Alpern RJ. Effect of glucocorticoids on renal cortical NHE-3 and NHE-1 mRNA. *Am J Physiol*. 1994; 267:F437–442. [PubMed: 8092257]
66. Baum M, Quigley R. Inhibition of proximal convoluted tubule transport by dopamine. *Kidney Int*. 1998; 54:1593–1600. [PubMed: 9844135]
67. Beach RE, Schwab SJ, Brazy PC, Dennis VW. Norepinephrine increases Na⁺-K⁺-ATPase and solute transport in rabbit proximal tubules. *Am J Physiol*. 1987; 252:F215–220. [PubMed: 3028169]
68. Becker AM, Zhang J, Goyal S, Dwarakanath V, Aronson PS, Moe OW, Baum M. Ontogeny of NHE8 in the rat proximal tubule. *Am J Physiol Renal Physiol*. 2007; 293:F255–261. [PubMed: 17429030]
69. Becker HM, Deitmer JW. Carbonic anhydrase II increases the activity of the human electrogenic Na⁺/HCO₃⁻ cotransporter. *The Journal of biological chemistry*. 2007; 282:13508–13521. [PubMed: 17353189]
70. Bell-Reuss E, Trevino DL, Gottschalk CW. Effect of renal sympathetic nerve stimulation on proximal water and sodium reabsorption. *J Clin Invest*. 1976; 57:1104–1107. [PubMed: 947953]
71. Belleannee C, Da Silva N, Shum WW, Brown D, Breton S. Role of purinergic signaling pathways in V-ATPase recruitment to apical membrane of acidifying epididymal clear cells. *Am J Physiol Cell Physiol*. 298:C817–830. [PubMed: 20071692]
72. Bello-Reuss E, Colindres RE, Pastoriza-Munoz E, Mueller RA, Gottschalk CW. Effects of acute unilateral renal denervation in the rat. *J Clin Invest*. 1975; 56:208–217. [PubMed: 1141432]
73. Bello-Reuss E, Higashi Y, Kaneda Y. Dopamine decreases fluid reabsorption in straight portions of rabbit proximal tubule. *Am J Physiol*. 1982; 242:F634–640. [PubMed: 7091317]
74. Bengel HH, McNamara ER, Schwartz JH, Alexander EA. Acidification adaptation along the inner medullary collecting duct. *Am J Physiol*. 1988; 255:F1155–1159. [PubMed: 3144183]
75. Bengel HH, Schwartz JH, McNamara ER, Alexander EA. Chronic metabolic acidosis augments acidification along the inner medullary collecting duct. *Am J Physiol*. 1986; 250:F690–694. [PubMed: 3963206]
76. Bernstein H, Atherton LJ, Deen WM. Axial heterogeneity and filtered-load dependence of proximal bicarbonate reabsorption. *Biophys J*. 1986; 50:239–252. [PubMed: 3017467]
77. Bezerra CN, Girardi AC, Carraro-Lacroix LR, Reboucas NA. Mechanisms underlying the long-term regulation of NHE3 by parathyroid hormone. *Am J Physiol Renal Physiol*. 2008; 294:F1232–1237. [PubMed: 18322024]
78. Biagi B, Kubota T, Sohtell M, Giebisch G. Intracellular potentials in rabbit proximal tubules perfused in vitro. *Am J Physiol*. 1981; 240:F200–210. [PubMed: 7212067]
79. Biagi BA. Effects of the anion transport inhibitor, SITS, on the proximal straight tubule of the rabbit perfused in vitro. *J Membr Biol*. 1985; 88:25–31. [PubMed: 4093954]

80. Biagi BA, Sohtell M. Electrophysiology of basolateral bicarbonate transport in the rabbit proximal tubule. *Am J Physiol*. 1986; 250:F267–272. [PubMed: 3946604]
81. Bichara M, Mercier O, Borensztein P, Paillard M. Acute metabolic acidosis enhances circulating parathyroid hormone, which contributes to the renal response against acidosis in the rat. *J Clin Invest*. 1990; 86:430–443. [PubMed: 2166755]
82. Bichara M, Mercier O, Houillier P, Paillard M, Leviel F. Effects of antidiuretic hormone on urinary acidification and on tubular handling of bicarbonate in the rat. *The Journal of clinical investigation*. 1987; 80:621–630. [PubMed: 3624481]
83. Bichara M, Mercier O, Paillard M, Leviel F. Effects of parathyroid hormone on urinary acidification. *The American journal of physiology*. 1986; 251:F444–453. [PubMed: 3752255]
84. Bichara M, Paillard M, Corman B, de Rouffignac C, Leviel F. Volume expansion modulates NaHCO_3 and NaCl transport in the proximal tubule and Henle's loop. *Am J Physiol*. 1984; 247:F140–150. [PubMed: 6331198]
85. Biemesderfer D, DeGray B, Aronson PS. Active (9.6 s) and inactive (21 s) oligomers of NHE3 in microdomains of the renal brush border. *J Biol Chem*. 2001; 276:10161–10167. [PubMed: 11120742]
86. Biemesderfer D, Nagy T, DeGray B, Aronson PS. Specific association of megalin and the Na^+/H^+ exchanger isoform NHE3 in the proximal tubule. *J Biol Chem*. 1999; 274:17518–17524. [PubMed: 10364184]
87. Biemesderfer D, Pizzonia J, Abu-Alfa A, Exner M, Reilly R, Igarashi P, Aronson PS. NHE3: a Na^+/H^+ exchanger isoform of renal brush border. *Am J Physiol*. 1993; 265:F736–742. [PubMed: 8238556]
88. Biemesderfer D, Reilly RF, Exner M, Igarashi P, Aronson PS. Immunocytochemical characterization of $\text{Na}^+(\text{+})\text{-H}^+$ exchanger isoform NHE-1 in rabbit kidney. *Am J Physiol*. 1992; 263:F833–840. [PubMed: 1279986]
89. Biemesderfer D, Rutherford PA, Nagy T, Pizzonia JH, Abu-Alfa AK, Aronson PS. Monoclonal antibodies for high-resolution localization of NHE3 in adult and neonatal rat kidney. *The American journal of physiology*. 1997; 273:F289–299. [PubMed: 9277590]
90. Birkenhager R, Otto E, Schurmann MJ, Vollmer M, Ruf EM, Maier-Lutz I, Beekmann F, Fekete A, Omran H, Feldmann D, Milford DV, Jeck N, Konrad M, Landau D, Knoers NV, Antignac C, Sudbrak R, Kispert A, Hildebrandt F. Mutation of BSND causes Bartter syndrome with sensorineural deafness and kidney failure. *Nat Genet*. 2001; 29:310–314. [PubMed: 11687798]
91. Bishop JM, Lee HW, Handlogten ME, Han KH, Verlander JW, Weiner ID. Intercalated cell-specific Rh B glycoprotein deletion diminishes renal ammonia excretion response to hypokalemia. *American journal of physiology Renal physiology*. 2013; 304:F422–431. [PubMed: 23220726]
92. Bishop JM, Verlander JW, Lee HW, Nelson RD, Weiner AJ, Handlogten ME, Weiner ID. Role of the Rhesus glycoprotein, Rh B glycoprotein, in renal ammonia excretion. *Am J Physiol Renal Physiol*. 299:F1065–1077. [PubMed: 20719974]
93. Blanchard A, leviel F, Bichara M, Podevin RA, Paillard M. Interactions of external and internal K^+ with $\text{K}^+(\text{+})\text{-HCO}_3^-$ cotransporter of rat medullary thick ascending limb. *Am J Physiol*. 1996; 271:C218–225. [PubMed: 8760049]
94. Bloch RD, Zikos D, Fisher KA, Schleicher L, Oyama M, Cheng JC, Skopicki HA, Sukowski EJ, Cragoe EJ Jr. Peterson DR. Activation of proximal tubular $\text{Na}^+(\text{+})\text{-H}^+$ exchange by angiotensin II. *Am J Physiol*. 1992; 263:F135–143. [PubMed: 1322054]
95. Blomqvist SR, Vidarsson H, Fitzgerald S, Johansson BR, Ollerstam A, Brown R, Persson AE, Bergstrom GG, Enerback S. Distal renal tubular acidosis in mice that lack the forkhead transcription factor Foxi1. *J Clin Invest*. 2004; 113:1560–1570. [PubMed: 15173882]
96. Bobulescu IA, Di Sole F, Moe OW. Na^+/H^+ exchangers: physiology and link to hypertension and organ ischemia. *Curr Opin Nephrol Hypertens*. 2005; 14:485–494. [PubMed: 16046909]
97. Bobulescu IA, Dwarakanath V, Zou L, Zhang J, Baum M, Moe OW. Glucocorticoids acutely increase cell surface Na^+/H^+ exchanger-3 (NHE3) by activation of NHE3 exocytosis. *Am J Physiol Renal Physiol*. 2005; 289:F685–691. [PubMed: 15942046]
98. Bobulescu IA, Moe OW. Luminal $\text{Na}^+(\text{+})/\text{H}^+(\text{+})$ exchange in the proximal tubule. *Pflugers Arch*. 2009; 458:5–21. [PubMed: 18853182]

99. Bobulescu IA, Moe OW. Na⁺/H⁺ exchangers in renal regulation of acid-base balance. *Semin Nephrol.* 2006; 26:334–344. [PubMed: 17071327]
100. Boettger T, Hubner CA, Maier H, Rust MB, Beck FX, Jentsch TJ. Deafness and renal tubular acidosis in mice lacking the K-Cl co-transporter Kcc4. *Nature.* 2002; 416:874–878. [PubMed: 11976689]
101. Bogusky RT, Dietrich RL. Effect of acute metabolic acidosis on ammonia metabolism in kidney. *Am J Physiol.* 1989; 256:F321–328. [PubMed: 2916664]
102. Bonar P, Schneider HP, Becker HM, Deitmer JW, Casey JR. Three-dimensional model for the human Cl[−]/HCO₃[−] exchanger, AE1, by homology to the E. coli ClC protein. *J Mol Biol.* 425:2591–2608. [PubMed: 23583773]
103. Boron WF, Boulpaep EL. Intracellular pH regulation in the renal proximal tubule of the salamander. Basolateral HCO₃[−] transport. *J Gen Physiol.* 1983; 81:53–94. [PubMed: 6833997]
104. Boron WF, Boulpaep EL. Intracellular pH regulation in the renal proximal tubule of the salamander. Na-H exchange. *J Gen Physiol.* 1983; 81:29–52. [PubMed: 6833996]
105. Bourgeois S, Bounoure L, Christensen EI, Ramakrishnan SK, Houillier P, Devuyst O, Wagner CA. Haploinsufficiency of the ammonia transporter Rhcg predisposes to chronic acidosis: Rhcg is critical for apical and basolateral ammonia transport in the mouse collecting duct. *The Journal of biological chemistry.* 2013; 288:5518–5529. [PubMed: 23281477]
106. Bourgeois S, Masse S, Paillard M, Houillier P. Basolateral membrane Cl(−), Na(+), and K(+)-coupled base transport mechanisms in rat MTALH. *Am J Physiol Renal Physiol.* 2002; 282:F655–668. [PubMed: 11880327]
107. Bourgeois S, Meer LV, Wootla B, Bloch-Faure M, Chambrey R, Shull GE, Gawenis LR, Houillier P. NHE4 is critical for the renal handling of ammonia in rodents. *J Clin Invest.* 120:1895–1904. [PubMed: 20484819]
108. Braam B, Mitchell KD, Fox J, Navar LG. Proximal tubular secretion of angiotensin II in rats. *Am J Physiol.* 1993; 264:F891–898. [PubMed: 8388654]
109. Brenes LG, Brenes JN, Hernandez MM. Familial proximal renal tubular acidosis. A distinct clinical entity. *Am J Med.* 1977; 63:244–252. [PubMed: 888846]
110. Breton S, Alper SL, Gluck SL, Sly WS, Barker JE, Brown D. Depletion of intercalated cells from collecting ducts of carbonic anhydrase II-deficient (CAR2 null) mice. *The American journal of physiology.* 1995; 269:F761–774. [PubMed: 8594870]
111. Breton S, Brown D. Regulation of luminal acidification by the V-ATPase. *Physiology (Bethesda).* 28:318–329. [PubMed: 23997191]
112. Breyer MD, Jacobson HR. Regulation of rabbit medullary collecting duct cell pH by basolateral Na⁺/H⁺ and Cl[−]/base exchange. *J Clin Invest.* 1989; 84:996–1004. [PubMed: 2547843]
113. Breyer MD, Kokko JP, Jacobson HR. Regulation of net bicarbonate transport in rabbit cortical collecting tubule by peritubular pH, carbon dioxide tension, and bicarbonate concentration. *The Journal of clinical investigation.* 1986; 77:1650–1660. [PubMed: 3084564]
114. Brosius FC 3rd, Alper SL, Garcia AM, Lodish HF. The major kidney band 3 gene transcript predicts an amino-terminal truncated band 3 polypeptide. *J Biol Chem.* 1989; 264:7784–7787. [PubMed: 2542243]
115. Brown D, Breton S. H(+)V-ATPase-dependent luminal acidification in the kidney collecting duct and the epididymis/vas deferens: vesicle recycling and transcytotic pathways. *The Journal of experimental biology.* 2000; 203:137–145. [PubMed: 10600682]
116. Brown D, Breton S, Ausiello DA, Marshansky V. Sensing, signaling and sorting events in kidney epithelial cell physiology. *Traffic.* 2009; 10:275–284. [PubMed: 19170982]
117. Brown D, Hirsch S, Gluck S. An H⁺-ATPase in opposite plasma membrane domains in kidney epithelial cell subpopulations. *Nature.* 1988; 331:622–624. [PubMed: 2893294]
118. Brown D, Hirsch S, Gluck S. Localization of a proton-pumping ATPase in rat kidney. *J Clin Invest.* 1988; 82:2114–2126. [PubMed: 2904451]
119. Brown D, Kumpulainen T, Roth J, Orci L. Immunohistochemical localization of carbonic anhydrase in postnatal and adult rat kidney. *The American journal of physiology.* 1983; 245:F110–118. [PubMed: 6408932]

120. Brown D, Paunescu TG, Breton S, Marshansky V. Regulation of the V ATPase in kidney epithelial cells: dual role in acid-base homeostasis and vesicle trafficking. *J Exp Biol.* 2009; 212:1762–1772. [PubMed: 19448085]
121. Brown D, Roth J, Kumpulainen T, Orci L. Ultrastructural immunocytochemical localization of carbonic anhydrase. Presence in intercalated cells of the rat collecting tubule. *Histochemistry.* 1982; 75:209–213. [PubMed: 6813300]
122. Brown D, Zhu XL, Sly WS. Localization of membrane-associated carbonic anhydrase type IV in kidney epithelial cells. *Proc Natl Acad Sci U S A.* 1990; 87:7457–7461. [PubMed: 2120700]
123. Brown JC, Packer RK, Knepper MA. Role of organic anions in renal response to dietary acid and base loads. *Am J Physiol.* 1989; 257:F170–176. [PubMed: 2548397]
124. Bruce LJ, Cope DL, Jones GK, Schofield AE, Burley M, Povey S, Unwin RJ, Wrong O, Tanner MJ. Familial distal renal tubular acidosis is associated with mutations in the red cell anion exchanger (Band 3, AE1) gene. *J Clin Invest.* 1997; 100:1693–1707. [PubMed: 9312167]
125. Bruce LJ, Wrong O, Toye AM, Young MT, Ogle G, Ismail Z, Sinha AK, McMaster P, Hwaihwanje I, Nash GB, Hart S, Lavu E, Palmer R, Othman A, Unwin RJ, Tanner MJ. Band 3 mutations, renal tubular acidosis and South-East Asian ovalocytosis in Malaysia and Papua New Guinea: loss of up to 95% band 3 transport in red cells. *Biochem J.* 2000; 350(Pt 1):41–51. [PubMed: 10926824]
126. Buerkert J, Martin D, Trigg D. Segmental analysis of the renal tubule in buffer production and net acid formation. *The American journal of physiology.* 1983; 244:F442–454. [PubMed: 6837741]
127. Buffin-Meyer B, Younes-Ibrahim M, Barlet-Bas C, Cheval L, Marsy S, Doucet A. K depletion modifies the properties of Sch-28080-sensitive K-ATPase in rat collecting duct. *Am J Physiol.* 1997; 272:F124–131. [PubMed: 9039058]
128. Burckhardt BC, Sato K, Fromter E. Electrophysiological analysis of bicarbonate permeation across the peritubular cell membrane of rat kidney proximal tubule. I. Basic observations. *Pflugers Arch.* 1984; 401:34–42. [PubMed: 6089090]
129. Burg M, Green N. Bicarbonate transport by isolated perfused rabbit proximal convoluted tubules. *Am J Physiol.* 1977; 233:F307–314. [PubMed: 910955]
130. Burg MB, Orloff J. Control of fluid absorption in the renal proximal tubule. *J Clin Invest.* 1968; 47:2016–2024. [PubMed: 4300070]
131. Burnham C, Munzesheimer C, Rabon E, Sachs G. Ion pathways in renal brush border membranes. *Biochim Biophys Acta.* 1982; 685:260–272. [PubMed: 7066312]
132. Callaghan JM, Tan SS, Khan MA, Curran KA, Campbell WG, Smolka AJ, Toh BH, Gleeson PA, Wingo CS, Cain BD, et al. Renal expression of the gene encoding the gastric H(+)-K(+)-ATPase beta-subunit. *Am J Physiol.* 1995; 268:F363–374. [PubMed: 7900835]
133. Cano A, Baum M, Moe OW. Thyroid hormone stimulates the renal Na/H exchanger NHE3 by transcriptional activation. *Am J Physiol.* 1999; 276:C102–108. [PubMed: 9886925]
134. Cano A, Miller RT, Alpern RJ, Preisig PA. Angiotensin II stimulation of Na-H antiporter activity is cAMP independent in OKP cells. *Am J Physiol.* 1994; 266:C1603–1608. [PubMed: 8023891]
135. Cantone A, Wang T, Pica A, Simeoni M, Capasso G. Use of transgenic mice in acid-base balance studies. *J Nephrol.* 2006; 19(Suppl 9):S121–127. [PubMed: 16736435]
136. Capasso G, Jaeger P, Giebisch G, Guckian V, Malnic G. Renal bicarbonate reabsorption in the rat. II. Distal tubule load dependence and effect of hypokalemia. *J Clin Invest.* 1987; 80:409–414. [PubMed: 3112185]
137. Capasso G, Kinne R, Malnic G, Giebisch G. Renal bicarbonate reabsorption in the rat. I. Effects of hypokalemia and carbonic anhydrase. *J Clin Invest.* 1986; 78:1558–1567. [PubMed: 3097074]
138. Capasso G, Unwin R, Ciani F, De Santo NG, De Tommaso G, Russo F, Giebisch G. Bicarbonate transport along the loop of Henle. II. Effects of acid-base, dietary, and neurohumoral determinants. *The Journal of clinical investigation.* 1994; 94:830–838. [PubMed: 8040339]
139. Capasso G, Unwin R, Rizzo M, Pica A, Giebisch G. Bicarbonate transport along the loop of Henle: molecular mechanisms and regulation. *Journal of nephrology.* 2002; 15(Suppl 5):S88–96. [PubMed: 12027225]
140. Carraro-Lacroix LR, Girardi AC, Malnic G. Long-term regulation of vacuolar H(+)-ATPase by angiotensin II in proximal tubule cells. *Pflugers Arch.* 2009; 458:969–979. [PubMed: 19396617]

141. Carraro-Lacroix LR, Lessa LM, Bezerra CN, Pessoa TD, Souza-Menezes J, Morales MM, Girardi AC, Malnic G. Role of CFTR and ClC-5 in modulating vacuolar H⁺-ATPase activity in kidney proximal tubule. *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology*. 2010; 26:563–576.
142. Cassano G, Stieger B, Murer H. Na/H- and Cl/OH-exchange in rat jejunal and rat proximal tubular brush border membrane vesicles. *Studies with acridine orange*. *Pflugers Arch*. 1984; 400:309–317. [PubMed: 6728651]
143. Castelo-Branco RC, Leite-Delova DC, de Mello-Aires M. Dose-dependent effects of angiotensin-(1-7) on the NHE3 exchanger and [Ca(2+)](i) in in vivo proximal tubules. *American journal of physiology Renal physiology*. 2013; 304:F1258–1265. [PubMed: 23515716]
144. Cha B, Kim JH, Hut H, Hogema BM, Nadarja J, Zizak M, Cavet M, Lee-Kwon W, Lohmann SM, Smolenski A, Tse CM, Yun C, de Jonge HR, Donowitz M. cGMP inhibition of Na⁺/H⁺ antiporter 3 (NHE3) requires PDZ domain adapter NHERF2, a broad specificity protein kinase G-anchoring protein. *J Biol Chem*. 2005; 280:16642–16650. [PubMed: 15722341]
145. Cha B, Tse M, Yun C, Kovbasnjuk O, Mohan S, Hubbard A, Arpin M, Donowitz M. The NHE3 juxtamembrane cytoplasmic domain directly binds ezrin: dual role in NHE3 trafficking and mobility in the brush border. *Mol Biol Cell*. 2006; 17:2661–2673. [PubMed: 16540524]
146. Chaillet JR, Lopes AG, Boron WF. Basolateral Na-H exchange in the rabbit cortical collecting tubule. *J Gen Physiol*. 1985; 86:795–812. [PubMed: 3001217]
147. Chalmers, RA.; Lawson, AM. *Organic acids in man*. Methuen; New York: 1982. p. xiip. 523
148. Chalumeau C, du Cheyron D, Defontaine N, Kellermann O, Paillard M, Poggioli J. NHE3 activity and trafficking depend on the state of actin organization in proximal tubule. *Am J Physiol Renal Physiol*. 2001; 280:F283–290. [PubMed: 11208604]
149. Chambrey R, Warnock DG, Podevin RA, Bruneval P, Mandet C, Belair MF, Bariety J, Paillard M. Immunolocalization of the Na⁺/H⁺ exchanger isoform NHE2 in rat kidney. *Am J Physiol*. 1998; 275:F379–386. [PubMed: 9729510]
150. Chan YL. Adrenergic control of bicarbonate absorption in the proximal convoluted tubule of the rat kidney. *Pflugers Arch*. 1980; 388:159–164. [PubMed: 7192853]
151. Chan YL, Biagi B, Giebisch G. Control mechanisms of bicarbonate transport across the rat proximal convoluted tubule. *Am J Physiol*. 1982; 242:F532–543. [PubMed: 6282141]
152. Chan YL, Giebisch G. Relationship between sodium and bicarbonate transport in the rat proximal convoluted tubule. *Am J Physiol*. 1981; 240:F222–230. [PubMed: 7212069]
153. Chan YL, Malnic G, Giebisch G. Passive driving forces of proximal tubular fluid and bicarbonate transport: gradient dependence of H⁺ secretion. *Am J Physiol*. 1983; 245:F622–633. [PubMed: 6638182]
154. Chan YL, Malnic G, Giebisch G. Renal bicarbonate reabsorption in the rat. III. Distal tubule perfusion study of load dependence and bicarbonate permeability. *J Clin Invest*. 1989; 84:931–938. [PubMed: 2760220]
155. Chantrelle B, Cogan MG, Rector FC Jr. Evidence for coupled sodium/hydrogen exchange in the rat superficial proximal convoluted tubule. *Pflugers Arch*. 1982; 395:186–189. [PubMed: 6818517]
156. Chen PY, Illsley NP, Verkman AS. Renal brush-border chloride transport mechanisms characterized using a fluorescent indicator. *Am J Physiol*. 1988; 254:F114–120. [PubMed: 3337241]
157. Chen PY, Verkman AS. Sodium-dependent chloride transport in basolateral membrane vesicles isolated from rabbit proximal tubule. *Biochemistry*. 1988; 27:655–660. [PubMed: 3349054]
158. Chen SH, Bubb MR, Yarmola EG, Zuo J, Jiang J, Lee BS, Lu M, Gluck SL, Hurst IR, Holliday LS. Vacuolar H⁺-ATPase binding to microfilaments: regulation in response to phosphatidylinositol 3-kinase activity and detailed characterization of the actin-binding site in subunit B. *J Biol Chem*. 2004; 279:7988–7998. [PubMed: 14662773]
159. Chen Y, Cann MJ, Litvin TN, Iourgenko V, Sinclair ML, Levin LR, Buck J. Soluble adenylyl cyclase as an evolutionarily conserved bicarbonate sensor. *Science*. 2000; 289:625–628. [PubMed: 10915626]

160. Chernova MN, Jiang L, Friedman DJ, Darman RB, Lohi H, Kere J, Vondorp DH, Alper SL. Functional comparison of mouse slc26a6 anion exchanger with human SLC26A6 polypeptide variants: differences in anion selectivity, regulation, and electrogenicity. *J Biol Chem*. 2005; 280:8564–8580. [PubMed: 15548529]
161. Choi JW, Lee-Kwon W, Jeon ES, Kang YJ, Kawano K, Kim HS, Suh PG, Donowitz M, Kim JH. Lysophosphatidic acid induces exocytic trafficking of Na(+)/H(+) exchanger 3 by E3KARP-dependent activation of phospholipase C. *Biochim Biophys Acta*. 2004; 1683:59–68. [PubMed: 15238220]
162. Choi JY, Shah M, Lee MG, Schultheis PJ, Shull GE, Muallem S, Baum M. Novel amiloride-sensitive sodium-dependent proton secretion in the mouse proximal convoluted tubule. *J Clin Invest*. 2000; 105:1141–1146. [PubMed: 10772659]
163. Chu CY, King JC, Berrini M, Alexander RT, Cordat E. Functional rescue of a kidney anion exchanger 1 trafficking mutant in renal epithelial cells. *PloS one*. 2013; 8:e57062. [PubMed: 23460825]
164. Chu TS, Peng Y, Cano A, Yanagisawa M, Alpern RJ. Endothelin(B) receptor activates NHE-3 by a Ca²⁺-dependent pathway in OKP cells. *J Clin Invest*. 1996; 97:1454–1462. [PubMed: 8617878]
165. Chu TS, Tsuganezawa H, Peng Y, Cano A, Yanagisawa M, Alpern RJ. Role of tyrosine kinase pathways in ETB receptor activation of NHE3. *Am J Physiol*. 1996; 271:C763–771. [PubMed: 8843705]
166. Cipriano DJ, Wang Y, Bond S, Hinton A, Jefferies KC, Qi J, Forgac M. Structure and regulation of the vacuolar ATPases. *Biochim Biophys Acta*. 2008; 1777:599–604. [PubMed: 18423392]
167. Clapp WL, Madsen KM, Verlander JW, Tisher CC. Morphologic heterogeneity along the rat inner medullary collecting duct. *Lab Invest*. 1989; 60:219–230. [PubMed: 2915516]
168. Cogan MG. Chronic hypercapnia stimulates proximal bicarbonate reabsorption in the rat. *J Clin Invest*. 1984; 74:1942–1947. [PubMed: 6439738]
169. Cogan MG. Effects of acute alterations in PCO₂ on proximal HCO₃⁻, Cl⁻, and H₂O reabsorption. *Am J Physiol*. 1984; 246:F21–26. [PubMed: 6421173]
170. Cogan MG, Maddox DA, Lucci MS, Rector FC Jr. Control of proximal bicarbonate reabsorption in normal and acidotic rats. *J Clin Invest*. 1979; 64:1168–1180. [PubMed: 500804]
171. Cogan MG, Maddox DA, Warnock DG, Lin ET, Rector FC Jr. Effect of acetazolamide on bicarbonate reabsorption in the proximal tubule of the rat. *Am J Physiol*. 1979; 237:F447–454. [PubMed: 517658]
172. Cohen RM, Feldman GM, Fernandez PC. The balance of acid, base and charge in health and disease. *Kidney international*. 1997; 52:287–293. [PubMed: 9263983]
173. Cohn DE, Hruska KA, Klahr S, Hammerman MR. Increased Na⁺-H⁺ exchange in brush border vesicles from dogs with renal failure. *Am J Physiol*. 1982; 243:F293–299. [PubMed: 7114259]
174. Cohn DE, Klahr S, Hammerman MR. Metabolic acidosis and parathyroidectomy increase Na⁺-H⁺ exchange in brush border vesicles. *Am J Physiol*. 1983; 245:F217–222. [PubMed: 6309012]
175. Collazo R, Fan L, Hu MC, Zhao H, Wiederkehr MR, Moe OW. Acute regulation of Na⁺/H⁺ exchanger NHE3 by parathyroid hormone via NHE3 phosphorylation and dynamin-dependent endocytosis. *J Biol Chem*. 2000; 275:31601–31608. [PubMed: 10866993]
176. Coppola S, Fromter E. An electrophysiological study of angiotensin II regulation of Na-HCO₃ cotransport and K conductance in renal proximal tubules. I. Effect of picomolar concentrations. *Pflügers Archiv : European journal of physiology*. 1994; 427:143–150. [PubMed: 8058463]
177. Coppola S, Fromter E. An electrophysiological study of angiotensin II regulation of Na-HCO₃ cotransport and K conductance in renal proximal tubules. II. Effect of micromolar concentrations. *Pflügers Archiv : European journal of physiology*. 1994; 427:151–156. [PubMed: 8058464]
178. Cougnon M, Bouyer P, Planelles G, Jaisser F. Does the colonic H,K ATPase also act as an Na,K-ATPase? *Proc Natl Acad Sci U S A*. 1998; 95:6516–6520. [PubMed: 9600998]
179. Crambert G. H,K-ATPase type 2: Relevance for renal physiology and beyond. *Am J Physiol Renal Physiol*.

180. Cruciat CM, Ohkawara B, Acebron SP, Karaulanov E, Reinhard C, Ingelfinger D, Boutros M, Niehrs C. Requirement of prorenin receptor and vacuolar H⁺-ATPase-mediated acidification for Wnt signaling. *Science*. 327:459–463. [PubMed: 20093472]
181. Cunningham R, Biswas R, Steplock D, Shenolikar S, Weinman E. Role of NHERF and scaffolding proteins in proximal tubule transport. *Urol Res*. 38:257–262. [PubMed: 20632170]
182. Cunningham R, Steplock D, Wang F, Huang H, E X, Shenolikar S, Weinman EJ. Defective parathyroid hormone regulation of NHE3 activity and phosphate adaptation in cultured NHERF-1^{-/-} renal proximal tubule cells. *J Biol Chem*. 2004; 279:37815–37821. [PubMed: 15218020]
183. Dass PD, Kurtz I. Renal ammonia and bicarbonate production in chronic renal failure. *Miner Electrolyte Metab*. 1990; 16:308–314. [PubMed: 1980935]
184. Dennis VW. Influence of bicarbonate on parathyroid hormone-induced changes in fluid absorption by the proximal tubule. *Kidney Int*. 1976; 10:373–380. [PubMed: 1003728]
185. Deyev IE, Sohet F, Vassilenko KP, Serova OV, Popova NV, Zozulya SA, Burova EB, Houillier P, Rzhovsky DI, Berchatova AA, Murashev AN, Chugunov AO, Efremov RG, Nikol'sky NN, Bertelli E, Eladari D, Petrenko AG. Insulin receptor-related receptor as an extracellular alkali sensor. *Cell metabolism*. 2011; 13:679–689. [PubMed: 21641549]
186. Dherbecourt O, Cheval L, Bloch-Faure M, Meneton P, Doucet A. Molecular identification of Sch28080-sensitive K-ATPase activities in the mouse kidney. *Pflugers Arch*. 2006; 451:769–775. [PubMed: 16208521]
187. Di Sole F, Babich V, Moe OW. The calcineurin homologous protein-1 increases Na⁽⁺⁾/H⁽⁺⁾ -exchanger 3 trafficking via ezrin phosphorylation. *J Am Soc Nephrol*. 2009; 20:1776–1786. [PubMed: 19556366]
188. Diakov TT, Kane PM. Regulation of vacuolar proton-translocating ATPase activity and assembly by extracellular pH. *The Journal of biological chemistry*. 2010; 285:23771–23778. [PubMed: 20511227]
189. Dixit MP, Xu L, Xu H, Bai L, Collins JF, Ghishan FK. Effect of angiotensin-II on renal Na⁺/H⁺-exchanger-NHE3 and NHE2. *Biochim Biophys Acta*. 2004; 1664:38–44. [PubMed: 15238256]
190. Dolson GM, Hise MK, Weinman EJ. Relationship among parathyroid hormone, cAMP, and calcium on proximal tubule sodium transport. *Am J Physiol*. 1985; 249:F409–416. [PubMed: 2994494]
191. Donowitz M, Ming Tse C, Fuster D. SLC9/NHE gene family, a plasma membrane and organellar family of Na⁽⁺⁾/H⁽⁺⁾ exchangers. *Mol Aspects Med*. 34:236–251. [PubMed: 23506868]
192. Dossena S, Rodighiero S, Vezzoli V, Nofziger C, Salvioni E, Boccazzi M, Grabmayer E, Botta G, Meyer G, Fugazzola L, Beck-Peccoz P, Paulmichl M. Functional characterization of wild-type and mutated pendrin (SLC26A4), the anion transporter involved in Pendred syndrome. *J Mol Endocrinol*. 2009; 43:93–103. [PubMed: 19608655]
193. Doucet A, Marsy S. Characterization of K-ATPase activity in distal nephron: stimulation by potassium depletion. *Am J Physiol*. 1987; 253:F418–423. [PubMed: 2957926]
194. Drumm K, Kress TR, Gassner B, Krug AW, Gekle M. Aldosterone stimulates activity and surface expression of NHE3 in human primary proximal tubule epithelial cells (RPTEC). *Cell Physiol Biochem*. 2006; 17:21–28. [PubMed: 16543718]
195. Du Z, Duan Y, Yan Q, Weinstein AM, Weinbaum S, Wang T. Mechanosensory function of microvilli of the kidney proximal tubule. *Proc Natl Acad Sci U S A*. 2004; 101:13068–13073. [PubMed: 15319475]
196. Du Z, Yan Q, Duan Y, Weinbaum S, Weinstein AM, Wang T. Axial flow modulates proximal tubule NHE3 and H-ATPase activities by changing microvillus bending moments. *Am J Physiol Renal Physiol*. 2006; 290:F289–296. [PubMed: 16144961]
197. DuBose TD Jr. Caflisch CR. Effect of selective aldosterone deficiency on acidification in nephron segments of the rat inner medulla. *J Clin Invest*. 1988; 82:1624–1632. [PubMed: 3183058]
198. DuBose TD Jr. Lucci MS, Hogg RJ, Pucacco LR, Kokko JP, Carter NW. Comparison of acidification parameters in superficial and deep nephrons of the rat. *The American journal of physiology*. 1983; 244:F497–503. [PubMed: 6405628]

199. DuBose TD Jr, Pucacco LR, Carter NW. Determination of disequilibrium pH in the rat kidney in vivo: evidence of hydrogen secretion. *Am J Physiol.* 1981; 240:F138–146. [PubMed: 6781364]
200. Duong Van Huyen JP, Cheval L, Bloch-Faure M, Belair MF, Heudes D, Bruneval P, Doucet A. GDF15 triggers homeostatic proliferation of acid-secreting collecting duct cells. *J Am Soc Nephrol.* 2008; 19:1965–1974. [PubMed: 18650486]
201. Dynia DW, Steinmetz AG, Kocinsky HS. NHE3 function and phosphorylation are regulated by a calyculin A-sensitive phosphatase. *Am J Physiol Renal Physiol.* 298:F745–753. [PubMed: 20015946]
202. Edelman A, Bouthier M, Anagnostopoulos T. Chloride distribution in the proximal convoluted tubule of Necturus kidney. *J Membr Biol.* 1981; 62:7–17. [PubMed: 7277477]
203. Eiam-Ong S, Hilden SA, King AJ, Johns CA, Madias NE. Endothelin-1 stimulates the Na⁺/H⁺ and Na⁺/HCO₃[–] transporters in rabbit renal cortex. *Kidney Int.* 1992; 42:18–24. [PubMed: 1321928]
204. Eladari D, Blanchard A, Levieil F, Paillard M, Stuart-Tilley AK, Alper SL, Podgevin RA. Functional and molecular characterization of luminal and basolateral Cl[–]/HCO₃[–] exchangers of rat thick limbs. *Am J Physiol.* 1998; 275:F334–342. [PubMed: 9729505]
205. Ellis B, Li XC, Miguel-Qin E, Gu V, Zhuo JL. Evidence for a functional intracellular angiotensin system in the proximal tubule of the kidney. *Am J Physiol Regul Integr Comp Physiol.* 302:R494–509. [PubMed: 22170616]
206. Emmons C. Halide transport patterns of apical anion exchange in rabbit cortical collecting duct intercalated cells. *Am J Physiol.* 1996; 271:F799–805. [PubMed: 8898009]
207. Emmons C, Kurtz I. Functional characterization of three intercalated cell subtypes in the rabbit outer cortical collecting duct. *The Journal of clinical investigation.* 1994; 93:417–423. [PubMed: 8282814]
208. Erdorf M, Seifert R. Pharmacological characterization of adenylyl cyclase isoforms in rabbit kidney membranes. *Naunyn Schmiedeberg's Arch Pharmacol.* 383:357–372. [PubMed: 21279330]
209. Estevez R, Boettger T, Stein V, Birkenhager R, Otto E, Hildebrandt F, Jentsch TJ. Barttin is a Cl[–] channel beta-subunit crucial for renal Cl[–] reabsorption and inner ear K⁺ secretion. *Nature.* 2001; 414:558–561. [PubMed: 11734858]
210. Everett LA, Glaser B, Beck JC, Idol JR, Buchs A, Heyman M, Adawi F, Hazani E, Nassir E, Baxevas AD, Sheffield VC, Green ED. Pendred syndrome is caused by mutations in a putative sulphate transporter gene (PDS). *Nat Genet.* 1997; 17:411–422. [PubMed: 9398842]
211. Fahlke C, Fischer M. Physiology and pathophysiology of ClC-K/barttin channels. *Front Physiol.* 1:155. [PubMed: 21423394]
212. Fan L, Wiederkehr MR, Collazo R, Wang H, Crowder LA, Moe OW. Dual mechanisms of regulation of Na/H exchanger NHE-3 by parathyroid hormone in rat kidney. *J Biol Chem.* 1999; 274:11289–11295. [PubMed: 10196218]
213. Farwell WR, Taylor EN. Serum bicarbonate, anion gap and insulin resistance in the National Health and Nutrition Examination Survey. *Diabet Med.* 2008; 25:798–804. [PubMed: 18644066]
214. Fathallah DM, Bejaoui M, Lepaslier D, Chater K, Sly WS, Dellagi K. Carbonic anhydrase II (CA II) deficiency in Maghrebians: evidence for founder effect and genomic recombination at the CA II locus. *Hum Genet.* 1997; 99:634–637. [PubMed: 9150731]
215. Fathallah DM, Bejaoui M, Sly WS, Lakhoua R, Dellagi K. A unique mutation underlying carbonic anhydrase II deficiency syndrome in patients of Arab descent. *Hum Genet.* 1994; 94:581–582. [PubMed: 7959703]
216. Fejes-Toth G, Naray-Fejes-Toth A. Immunohistochemical localization of colonic H-K-ATPase to the apical membrane of connecting tubule cells. *Am J Physiol Renal Physiol.* 2001; 281:F318–325. [PubMed: 11457724]
217. Fejes-Toth G, Rusvai E, Cleaveland ES, Naray-Fejes-Toth A. Regulation of AE2 mRNA expression in the cortical collecting duct by acid/base balance. *Am J Physiol.* 1998; 274:F596–601. [PubMed: 9530277]

218. Fejes-Toth G, Rusvai E, Longo KA, Naray-Fejes-Toth A. Expression of colonic H-K-ATPase mRNA in cortical collecting duct: regulation by acid/base balance. *Am J Physiol.* 1995; 269:F551–557. [PubMed: 7485541]
219. Finberg KE, Wagner CA, Bailey MA, Paunescu TG, Breton S, Brown D, Giebisch G, Geibel JP, Lifton RP. The B1-subunit of the H(+) ATPase is required for maximal urinary acidification. *Proc Natl Acad Sci U S A.* 2005; 102:13616–13621. [PubMed: 16174750]
220. Fiori M, Gras EG, Amorena C. Decreased NHE8 isoform expression and defective acidification in proximal convoluted tubules of senile rats. *Age (Dordr).* 2009; 31:77–84. [PubMed: 19234771]
221. Forgac M. Vacuolar ATPases: rotary proton pumps in physiology and pathophysiology. *Nat Rev Mol Cell Biol.* 2007; 8:917–929. [PubMed: 17912264]
222. Frassetto L, Morris RC Jr, Sebastian A. Potassium bicarbonate reduces urinary nitrogen excretion in postmenopausal women. *J Clin Endocrinol Metab.* 1997; 82:254–259. [PubMed: 8989270]
223. Frische S, Kwon TH, Frokiaer J, Madsen KM, Nielsen S. Regulated expression of pendrin in rat kidney in response to chronic NH₄Cl or NaHCO₃ loading. *Am J Physiol Renal Physiol.* 2003; 284:F584–593. [PubMed: 12556366]
224. Frische S, Zolotarev AS, Kim YH, Praetorius J, Alper S, Nielsen S, Wall SM. AE2 isoforms in rat kidney: immunohistochemical localization and regulation in response to chronic NH₄Cl loading. *Am J Physiol Renal Physiol.* 2004; 286:F1163–1170. [PubMed: 14749257]
225. Fromter E, Gessner K. Active transport potentials, membrane diffusion potentials and streaming potentials across rat kidney proximal tubule. *Pflugers Arch.* 1974; 351:85–98. [PubMed: 4473176]
226. Fromter E, Gessner K. Effect of inhibitors and diuretics on electrical potential differences in rat kidney proximal tubule. *Pflugers Arch.* 1975; 357:209–224. [PubMed: 1238986]
227. Fromter E, Gessner K. Free-flow potential profile along rat kidney proximal tubule. *Pflugers Arch.* 1974; 351:69–83. [PubMed: 4472485]
228. Fuster DG, Bobulescu IA, Zhang J, Wade J, Moe OW. Characterization of the regulation of renal Na⁺/H⁺ exchanger NHE3 by insulin. *Am J Physiol Renal Physiol.* 2007; 292:F577–585. [PubMed: 17018843]
229. Fuster DG, Zhang J, Xie XS, Moe OW. The vacuolar-ATPase B1 subunit in distal tubular acidosis: novel mutations and mechanisms for dysfunction. *Kidney Int.* 2008; 73:1151–1158. [PubMed: 18368028]
230. Gao X, Eladari D, Leviel F, Tew BY, Miro-Julia C, Cheema FH, Miller L, Nelson R, Paunescu TG, McKee M, Brown D, Al-Awqati Q. Deletion of hensin/DMBT1 blocks conversion of beta- to alpha-intercalated cells and induces distal renal tubular acidosis. *Proc Natl Acad Sci U S A.* 107:21872–21877. [PubMed: 21098262]
231. Garcia NH, Garvin JL. Angiotensin 1-7 has a biphasic effect on fluid absorption in the proximal straight tubule. *Journal of the American Society of Nephrology : JASN.* 1994; 5:1133–1138. [PubMed: 7849254]
232. Garcia-Austt J, Good DW, Burg MB, Knepper MA. Deoxycorticosterone-stimulated bicarbonate secretion in rabbit cortical collecting ducts: effects of luminal chloride removal and in vivo acid loading. *Am J Physiol.* 1985; 249:F205–212. [PubMed: 3927746]
233. Garvin JL, Burg MB, Knepper MA. Active NH₄⁺ absorption by the thick ascending limb. *Am J Physiol.* 1988; 255:F57–65. [PubMed: 3394813]
234. Gattineni J, Sas D, Dagan A, Dwarakanath V, Baum M. Effect of thyroid hormone on the postnatal renal expression of NHE8. *Am J Physiol Renal Physiol.* 2008; 294:F198–204. [PubMed: 17977906]
235. Gawenis LR, Bradford EM, Prasad V, Lorenz JN, Simpson JE, Clarke LL, Woo AL, Grisham C, Sanford LP, Doetschman T, Miller ML, Shull GE. Colonic anion secretory defects and metabolic acidosis in mice lacking the NBC1 Na⁺/HCO₃[–] cotransporter. *J Biol Chem.* 2007; 282:9042–9052. [PubMed: 17192275]
236. Geibel J, Giebisch G, Boron WF. Angiotensin II stimulates both Na(+)–H⁺ exchange and Na⁺/HCO₃[–] cotransport in the rabbit proximal tubule. *Proc Natl Acad Sci U S A.* 1990; 87:7917–7920. [PubMed: 2172967]

237. Geibel J, Giebisch G, Boron WF. Basolateral sodium-coupled acid-base transport mechanisms of the rabbit proximal tubule. *Am J Physiol.* 1989; 257:F790–797. [PubMed: 2556038]
238. Geibel J, Giebisch G, Boron WF. Effects of acetate on luminal acidification processes in the S3 segment of the rabbit proximal tubule. *Am J Physiol.* 1989; 257:F586–594. [PubMed: 2801961]
239. Giebisch G, Malnic G, De Mello GB, De Mello Aires M. Kinetics of luminal acidification in cortical tubules of the rat kidney. *J Physiol.* 1977; 267:571–599. [PubMed: 17736]
240. Gifford JD, Sharkins K, Work J, Luke RG, Galla JH. Total CO₂ transport in rat cortical collecting duct in chloride-depletion alkalosis. *Am J Physiol.* 1990; 258:F848–853. [PubMed: 2109936]
241. Gildea JJ, Shah IT, Van Sciver RE, Israel JA, Enzensperger C, McGrath HE, Jose PA, Felder RA. The cooperative roles of the dopamine receptors, DR and DR, on the regulation of renal sodium transport. *Kidney Int.*
242. Girardi AC, Degray BC, Nagy T, Biemesderfer D, Aronson PS. Association of Na(+)-H(+) exchanger isoform NHE3 and dipeptidyl peptidase IV in the renal proximal tubule. *J Biol Chem.* 2001; 276:46671–46677. [PubMed: 11590171]
243. Girardi AC, Fukuda LE, Rossoni LV, Malnic G, Reboucas NA. Dipeptidyl peptidase IV inhibition downregulates Na⁺ - H⁺ exchanger NHE3 in rat renal proximal tubule. *Am J Physiol Renal Physiol.* 2008; 294:F414–422. [PubMed: 18077600]
244. Girardi AC, Knauf F, Demuth HU, Aronson PS. Role of dipeptidyl peptidase IV in regulating activity of Na⁺/H⁺ exchanger isoform NHE3 in proximal tubule cells. *Am J Physiol Cell Physiol.* 2004; 287:C1238–1245. [PubMed: 15213057]
245. Gisler SM, Pribanic S, Bacic D, Forrer P, Gantenbein A, Sabourin LA, Tsuji A, Zhao ZS, Manser E, Biber J, Murer H. PDZK1: I. a major scaffold in brush borders of proximal tubular cells. *Kidney Int.* 2003; 64:1733–1745. [PubMed: 14531806]
246. Goldfarb S. Dietary factors in the pathogenesis and prophylaxis of calcium nephrolithiasis. *Kidney international.* 1988; 34:544–555. [PubMed: 3199675]
247. Gong F, Alzamora R, Smolak C, Li H, Naveed S, Neumann D, Hallows KR, Pastor-Soler NM. Vacuolar H⁺-ATPase apical accumulation in kidney intercalated cells is regulated by PKA and AMP-activated protein kinase. *American journal of physiology Renal physiology.* 2010; 298:F1162–1169. [PubMed: 20147366]
248. Good DW. Active absorption of NH₄⁺ by rat medullary thick ascending limb: inhibition by potassium. *Am J Physiol.* 1988; 255:F78–87. [PubMed: 3134821]
249. Good DW. Adaptation of HCO₃⁻ and NH₄⁺ transport in rat MTAL: effects of chronic metabolic acidosis and Na⁺ intake. *The American journal of physiology.* 1990; 258:F1345–1353. [PubMed: 2337153]
250. Good DW. Effects of osmolality on bicarbonate absorption by medullary thick ascending limb of the rat. *The Journal of clinical investigation.* 1992; 89:184–190. [PubMed: 1729270]
251. Good DW. Hyperosmolality inhibits bicarbonate absorption in rat medullary thick ascending limb via a protein-tyrosine kinase-dependent pathway. *The Journal of biological chemistry.* 1995; 270:9883–9889. [PubMed: 7730371]
252. Good DW. Inhibition of bicarbonate absorption by peptide hormones and cyclic adenosine monophosphate in rat medullary thick ascending limb. *The Journal of clinical investigation.* 1990; 85:1006–1013. [PubMed: 2318960]
253. Good DW. Nerve growth factor regulates HCO₃⁻ absorption in thick ascending limb: modifying effects of vasopressin. *The American journal of physiology.* 1998; 274:C931–939. [PubMed: 9575789]
254. Good DW. PGE₂ reverses AVP inhibition of HCO₃⁻ absorption in rat MTAL by activation of protein kinase C. *The American journal of physiology.* 1996; 270:F978–985. [PubMed: 8764317]
255. Good DW. Sodium-dependent bicarbonate absorption by cortical thick ascending limb of rat kidney. *The American journal of physiology.* 1985; 248:F821–829. [PubMed: 2988349]
256. Good DW, Burg MB. Ammonia production by individual segments of the rat nephron. *J Clin Invest.* 1984; 73:602–610. [PubMed: 6323523]
257. Good DW, Caflisch CR, DuBose TD Jr. Transepithelial ammonia concentration gradients in inner medulla of the rat. *The American journal of physiology.* 1987; 252:F491–500. [PubMed: 3826390]

258. Good DW, Di Mari JF, Watts BA 3rd. Hyposmolality stimulates Na(+)/H(+) exchange and HCO₃(-)- absorption in thick ascending limb via PI 3-kinase. *American journal of physiology Cell physiology*. 2000; 279:C1443–1454. [PubMed: 11029292]
259. Good DW, George T. Regulation of HCO₃⁻ absorption by prostaglandin E₂ and G proteins in rat medullary thick ascending limb. *The American journal of physiology*. 1996; 270:F711–717. [PubMed: 8928831]
260. Good DW, George T, Wang DH. Angiotensin II inhibits HCO₃- absorption via a cytochrome P-450-dependent pathway in MTAL. *The American journal of physiology*. 1999; 276:F726–736. [PubMed: 10330055]
261. Good DW, George T, Watts BA 3rd. Aldosterone inhibits HCO₃ absorption via a nongenomic pathway in medullary thick ascending limb. *American journal of physiology Renal physiology*. 2002; 283:F699–706. [PubMed: 12217861]
262. Good DW, George T, Watts BA 3rd. Lipopolysaccharide directly alters renal tubule transport through distinct TLR4-dependent pathways in basolateral and apical membranes. *Am J Physiol Renal Physiol*. 2009; 297:F866–874. [PubMed: 19625374]
263. Good DW, George T, Watts BA 3rd. Nerve growth factor inhibits Na⁺/H⁺ exchange and formula absorption through parallel phosphatidylinositol 3-kinase-mTOR and ERK pathways in thick ascending limb. *The Journal of biological chemistry*. 2008; 283:26602–26611. [PubMed: 18660503]
264. Good DW, George T, Watts BA 3rd. Nongenomic regulation by aldosterone of the epithelial NHE3 Na(+)/H(+) exchanger. *American journal of physiology Cell physiology*. 2006; 290:C757–763. [PubMed: 16251474]
265. Good DW, George T, Watts BA 3rd. Toll-like receptor 2 is required for LPS-induced Toll-like receptor 4 signaling and inhibition of ion transport in renal thick ascending limb. *J Biol Chem*. 287:20208–20220. [PubMed: 22523073]
266. Good DW, George T, Watts BA 3rd. Toll-like receptor 2 mediates inhibition of HCO₃(-)- absorption by bacterial lipoprotein in medullary thick ascending limb. *Am J Physiol Renal Physiol*. 299:F536–544. [PubMed: 20554644]
267. Good DW, Knepper MA, Burg MB. Ammonia and bicarbonate transport by thick ascending limb of rat kidney. *Am J Physiol*. 1984; 247:F35–44. [PubMed: 6742203]
268. Good DW, Watts BA 3rd, George T, Meyer JW, Shull GE. Transepithelial HCO₃⁻ absorption is defective in renal thick ascending limbs from Na⁺/H⁺ exchanger NHE1 null mutant mice. *American journal of physiology Renal physiology*. 2004; 287:F1244–1249. [PubMed: 15292047]
269. Goyal S, Mentone S, Aronson PS. Immunolocalization of NHE8 in rat kidney. *Am J Physiol Renal Physiol*. 2005; 288:F530–538. [PubMed: 15522984]
270. Goyal S, Vanden Heuvel G, Aronson PS. Renal expression of novel Na⁺/H⁺ exchanger isoform NHE8. *Am J Physiol Renal Physiol*. 2003; 284:F467–473. [PubMed: 12409279]
271. Grassl SM, Aronson PS. Na⁺/HCO₃⁻-co-transport in basolateral membrane vesicles isolated from rabbit renal cortex. *J Biol Chem*. 1986; 261:8778–8783. [PubMed: 3013862]
272. Grassl SM, Holohan PD, Ross CR. HCO₃⁻ transport in basolateral membrane vesicles isolated from rat renal cortex. *J Biol Chem*. 1987; 262:2682–2687. [PubMed: 3029092]
273. Greenlee MM, Lynch IJ, Gumz ML, Cain BD, Wingo CS. Mineralocorticoids stimulate the activity and expression of renal H⁺,K⁺-ATPases. *J Am Soc Nephrol*. 22:49–58. [PubMed: 21164026]
274. Greenlee MM, Lynch IJ, Gumz ML, Cain BD, Wingo CS. The renal H,K-ATPases. *Curr Opin Nephrol Hypertens*. 19:478–482. [PubMed: 20616716]
275. Gross E, Hawkins K, Abuladze N, Pushkin A, Cotton CU, Hopfer U, Kurtz I. The stoichiometry of the electrogenic sodium bicarbonate cotransporter NBC1 is cell-type dependent. *J Physiol*. 2001; 531:597–603. [PubMed: 11251043]
276. Gross E, Hawkins K, Pushkin A, Sassani P, Dukkipati R, Abuladze N, Hopfer U, Kurtz I. Phosphorylation of Ser(982) in the sodium bicarbonate cotransporter kNBC1 shifts the HCO₃(-) : Na(+) stoichiometry from 3 : 1 to 2 : 1 in murine proximal tubule cells. *J Physiol*. 2001; 537:659–665. [PubMed: 11744745]

277. Gross E, Pushkin A, Abuladze N, Fedotoff O, Kurtz I. Regulation of the sodium bicarbonate cotransporter kNBC1 function: role of Asp⁹⁸⁶, Asp⁹⁸⁸ and kNBC1-carbonic anhydrase II binding. *The Journal of physiology*. 2002; 544:679–685. [PubMed: 12411514]
278. Gueutin V, Vallet M, Jayat M, Peti-Peterdi J, Corniere N, Leviel F, Sohet F, Wagner CA, Eladari D, Chambrey R. Renal beta-intercalated cells maintain body fluid and electrolyte balance. *J Clin Invest*. 123:4219–4231. [PubMed: 24051376]
279. Guggino WB, Boulpaep EL, Giebisch G. Electrical properties of chloride transport across the necturus proximal tubule. *J Membr Biol*. 1982; 65:185–196. [PubMed: 7062339]
280. Guggino WB, London R, Boulpaep EL, Giebisch G. Chloride transport across the basolateral cell membrane of the Necturus proximal tubule: dependence on bicarbonate and sodium. *J Membr Biol*. 1983; 71:227–240. [PubMed: 6302263]
281. Gumz ML, Lynch IJ, Greenlee MM, Cain BD, Wingo CS. The renal H⁺-K⁺-ATPases: physiology, regulation, and structure. *Am J Physiol Renal Physiol*. 298:F12–21. [PubMed: 19640897]
282. Guntupalli J, DuBose TD Jr. Effects of endothelin on rat renal proximal tubule Na(+)-Pi cotransport and Na⁺/H⁺ exchange. *Am J Physiol*. 1994; 266:F658–666. [PubMed: 8184900]
283. Guntupalli J, Onuigbo M, Wall S, Alpern RJ, DuBose TD Jr. Adaptation to low-K⁺ media increases H(+)-K(+)-ATPase but not H(+)-ATPase-mediated pHi recovery in OMCD1 cells. *Am J Physiol*. 1997; 273:C558–571. [PubMed: 9277353]
284. Haithcock D, Jiao H, Cui XL, Hopfer U, Douglas JG. Renal proximal tubular AT2 receptor: signaling and transport. *J Am Soc Nephrol*. 1999; 10(Suppl 11):S69–74. [PubMed: 9892143]
285. Hall RA, Premont RT, Chow CW, Blitzer JT, Pitcher JA, Claing A, Stoffel RH, Barak LS, Shenolikar S, Weinman EJ, Grinstein S, Lefkowitz RJ. The beta2-adrenergic receptor interacts with the Na⁺/H⁺-exchanger regulatory factor to control Na⁺/H⁺ exchange. *Nature*. 1998; 392:626–630. [PubMed: 9560162]
286. Hallows KR, Alzamora R, Li H, Gong F, Smolak C, Neumann D, Pastor-Soler NM. AMP-activated protein kinase inhibits alkaline pH- and PKA-induced apical vacuolar H⁺-ATPase accumulation in epididymal clear cells. *Am J Physiol Cell Physiol*. 2009; 296:C672–681. [PubMed: 19211918]
287. Halperin ML, Jungas RL. Metabolic production and renal disposal of hydrogen ions. *Kidney Int*. 1983; 24:709–713. [PubMed: 6674669]
288. Halperin ML, Kamel KS. Ammonium metabolism: emphasis on energy considerations. *Miner Electrolyte Metab*. 1990; 16:277–282. [PubMed: 2283991]
289. Hamburger RJ, Lawson NL, Schwartz JH. Response to parathyroid hormone in defined segments of proximal tubule. *Am J Physiol*. 1976; 230:286–290. [PubMed: 1259005]
290. Hamm LL. Renal handling of citrate. *Kidney international*. 1990; 38:728–735. [PubMed: 2232510]
291. Hamm LL, Pucacco LR, Kokko JP, Jacobson HR. Hydrogen ion permeability of the rabbit proximal convoluted tubule. *Am J Physiol*. 1984; 246:F3–11. [PubMed: 6696075]
292. Hamm LL, Simon EE. Roles and mechanisms of urinary buffer excretion. *The American journal of physiology*. 1987; 253:F595–605. [PubMed: 3310662]
293. Han KH, Croker BP, Clapp WL, Werner D, Sahni M, Kim J, Kim HY, Handlogten ME, Weiner ID. Expression of the ammonia transporter, rh C glycoprotein, in normal and neoplastic human kidney. *J Am Soc Nephrol*. 2006; 17:2670–2679. [PubMed: 16928804]
294. Han KH, Lee HW, Handlogten ME, Whitehill F, Osis G, Croker BP, Clapp WL, Verlander JW, Weiner ID. Expression of the ammonia transporter family member, Rh B Glycoprotein, in the human kidney. *Am J Physiol Renal Physiol*. 304:F972–981. [PubMed: 23324176]
295. Han W, Kim KH, Jo MJ, Lee JH, Yang J, Doctor RB, Moe OW, Lee J, Kim E, Lee MG. Shank2 associates with and regulates Na⁺/H⁺ exchanger 3. *J Biol Chem*. 2006; 281:1461–1469. [PubMed: 16293618]
296. Haque SK, Ariceta G, Batlle D. Proximal renal tubular acidosis: a not so rare disorder of multiple etiologies. *Nephrol Dial Transplant*. 27:4273–4287. [PubMed: 23235953]

297. Harris RC, Seifter JL, Brenner BM. Adaptation of Na⁺-H⁺ exchange in renal microvillus membrane vesicles. Role of dietary protein and uninephrectomy. *J Clin Invest.* 1984; 74:1979–1987. [PubMed: 6511911]
298. Haussinger D, Gerok W, Sies H. The effect of urea synthesis on extracellular pH in isolated perfused rat liver. *Biochem J.* 1986; 236:261–265. [PubMed: 3790075]
299. Hayashi M, Yamaji Y, Iyori M, Kitajima W, Saruta T. Effect of isoproterenol on intracellular pH of the intercalated cells in the rabbit cortical collecting ducts. *J Clin Invest.* 1991; 87:1153–1157. [PubMed: 1849143]
300. Hays S, Kokko JP, Jacobson HR. Hormonal regulation of proton secretion in rabbit medullary collecting duct. *J Clin Invest.* 1986; 78:1279–1286. [PubMed: 3021819]
301. Hays SR, Alpern RJ. Apical and basolateral membrane H⁺ extrusion mechanisms in inner stripe of rabbit outer medullary collecting duct. *Am J Physiol.* 1990; 259:F628–635. [PubMed: 2171359]
302. Hays SR, Alpern RJ. Basolateral membrane Na⁽⁺⁾-independent Cl[–]/HCO₃[–] exchange in the inner stripe of the rabbit outer medullary collecting tubule. *J Gen Physiol.* 1990; 95:347–367. [PubMed: 2307960]
303. Hays SR, Baum M, Kokko JP. Effects of protein kinase C activation on sodium, potassium, chloride, and total CO₂ transport in the rabbit cortical collecting tubule. *The Journal of clinical investigation.* 1987; 80:1561–1570. [PubMed: 3680514]
304. Hennings JC, Picard N, Huebner AK, Stauber T, Maier H, Brown D, Jentsch TJ, Vargas-Poussou R, Eladari D, Hubner CA. A mouse model for distal renal tubular acidosis reveals a previously unrecognized role of the V-ATPase a4 subunit in the proximal tubule. *EMBO Mol Med.* 4:1057–1071. [PubMed: 22933323]
305. Hensley CB, Bradley ME, Mircheff AK. Parathyroid hormone-induced translocation of Na-H antiporters in rat proximal tubules. *Am J Physiol.* 1989; 257:C637–645. [PubMed: 2552817]
306. Hentschke M, Hentschke S, Borgmeyer U, Hubner CA, Kurth I. The murine AE4 promoter predominantly drives type B intercalated cell specific transcription. *Histochem Cell Biol.* 2009; 132:405–412. [PubMed: 19544066]
307. Hering-Smith KS, Cragoe EJ Jr, Weiner D, Hamm LL. Inner medullary collecting duct Na⁽⁺⁾-H⁺ exchanger. *Am J Physiol.* 1991; 260:C1300–1307. [PubMed: 1647666]
308. Holliday LS, Lu M, Lee BS, Nelson RD, Solivan S, Zhang L, Gluck SL. The amino-terminal domain of the B subunit of vacuolar H⁺-ATPase contains a filamentous actin binding site. *J Biol Chem.* 2000; 275:32331–32337. [PubMed: 10915794]
309. Holmberg C, Kokko JP, Jacobson HR. Determination of chloride and bicarbonate permeabilities in proximal convoluted tubules. *Am J Physiol.* 1981; 241:F386–394. [PubMed: 7315963]
310. Holthouser KA, Mandal A, Merchant ML, Schelling JR, Delamere NA, Valdes RR Jr, Tyagi SC, Lederer ED, Khundmiri SJ. Ouabain stimulates Na-K-ATPase through a sodium/hydrogen exchanger-1 (NHE-1)-dependent mechanism in human kidney proximal tubule cells. *Am J Physiol Renal Physiol.* 299:F77–90. [PubMed: 20427472]
311. Honegger KJ, Capuano P, Winter C, Bacic D, Stange G, Wagner CA, Biber J, Murer H, Hernando N. Regulation of sodium-proton exchanger isoform 3 (NHE3) by PKA and exchange protein directly activated by cAMP (EPAC). *Proc Natl Acad Sci U S A.* 2006; 103:803–808. [PubMed: 16407144]
312. Horie S, Moe O, Yamaji Y, Cano A, Miller RT, Alpern RJ. Role of protein kinase C and transcription factor AP-1 in the acid-induced increase in Na/H antiporter activity. *Proc Natl Acad Sci U S A.* 1992; 89:5236–5240. [PubMed: 1319056]
313. Horita S, Zheng Y, Hara C, Yamada H, Kunimi M, Taniguchi S, Uwatoko S, Sugaya T, Goto A, Fujita T, Seki G. Biphasic regulation of Na⁺-HCO₃[–] cotransporter by angiotensin II type 1A receptor. *Hypertension.* 2002; 40:707–712. [PubMed: 12411466]
314. Hosokawa H, Dip PV, Merkulova M, Bakulina A, Zhuang Z, Khatri A, Jian X, Keating SM, Bueler SA, Rubinstein JL, Randazzo PA, Ausiello DA, Gruber G, Marshansky V. The N termini of a-subunit isoforms are involved in signaling between vacuolar H⁺-ATPase (V-ATPase) and cytohesin-2. *J Biol Chem.* 288:5896–5913. [PubMed: 23288846]

315. Houillier P, Bourgeois S. More actors in ammonia absorption by the thick ascending limb. *Am J Physiol Renal Physiol.* 302:F293–297. [PubMed: 22088435]
316. Houillier P, Chambrey R, Achard JM, Froissart M, Poggioli J, Paillard M. Signaling pathways in the biphasic effect of angiotensin II on apical Na/H antiport activity in proximal tubule. *Kidney international.* 1996; 50:1496–1505. [PubMed: 8914015]
317. Howlin KJ, Alpern RJ, Rector FC Jr. Amiloride inhibition of proximal tubular acidification. *Am J Physiol.* 1985; 248:F773–778. [PubMed: 2988347]
318. Hruska KA, Moskowitz D, Esbrit P, Civitelli R, Westbrook S, Huskey M. Stimulation of inositol trisphosphate and diacylglycerol production in renal tubular cells by parathyroid hormone. *J Clin Invest.* 1987; 79:230–239. [PubMed: 3025260]
319. Hu PY, Roth DE, Skaggs LA, Venta PJ, Tashian RE, Guibaud P, Sly WS. A splice junction mutation in intron 2 of the carbonic anhydrase II gene of osteopetrosis patients from Arabic countries. *Hum Mutat.* 1992; 1:288–292. [PubMed: 1301935]
320. Iacovitti M, Nash L, Peterson LN, Rochon J, Levine DZ. Distal tubule bicarbonate accumulation in vivo. Effect of flow and transtubular bicarbonate gradients. *The Journal of clinical investigation.* 1986; 78:1658–1665. [PubMed: 3097075]
321. Igarashi T, Inatomi J, Sekine T, Cha SH, Kanai Y, Kunimi M, Tsukamoto K, Satoh H, Shimadzu M, Tozawa F, Mori T, Shiobara M, Seki G, Endou H. Mutations in SLC4A4 cause permanent isolated proximal renal tubular acidosis with ocular abnormalities. *Nat Genet.* 1999; 23:264–266. [PubMed: 10545938]
322. Iino Y, Burg MB. Effect of parathyroid hormone on bicarbonate absorption by proximal tubules in vitro. *Am J Physiol.* 1979; 236:F387–391. [PubMed: 434213]
323. Imai M, Kokko JP. Mechanism of sodium and chloride transport in the thin ascending limb of Henle. *The Journal of clinical investigation.* 1976; 58:1054–1060. [PubMed: 993330]
324. Imai M, Kokko JP. Sodium chloride, urea, and water transport in the thin ascending limb of Henle. Generation of osmotic gradients by passive diffusion of solutes. *The Journal of clinical investigation.* 1974; 53:393–402. [PubMed: 11344552]
325. International Union of Pure and Applied Chemistry. Commission on Equilibrium Data. Serjeant, EP.; Dempsey, B.; International Union of Pure and Applied Chemistry. Commission on Electrochemical Data. Ionisation constants of organic acids in aqueous solution. Pergamon Press; Oxford ; New York: 1979. p. xip. 989
326. Ishibashi K, Sasaki S, Yoshiyama N. Intracellular chloride activity of rabbit proximal straight tubule perfused in vitro. *Am J Physiol.* 1988; 255:F49–56. [PubMed: 3394812]
327. Ishibashi K, Sasaki S, Yoshiyama N, Shiigai T, Takeuchi J. Generation of pH gradient across the rabbit collecting duct segments perfused in vitro. *Kidney Int.* 1987; 31:930–936. [PubMed: 3586499]
328. Ives HE. Proton/hydroxyl permeability of proximal tubule brush border vesicles. *Am J Physiol.* 1985; 248:F78–86. [PubMed: 2982277]
329. Ives HE, Chen PY, Verkman AS. Mechanism of coupling between Cl⁻ and OH⁻ transport in renal brush-border membranes. *Biochim Biophys Acta.* 1986; 863:91–100. [PubMed: 3778914]
330. Ives HE, Yee VJ, Warnock DG. Asymmetric distribution of the Na⁺/H⁺ antiporter in the renal proximal tubule epithelial cell. *J Biol Chem.* 1983; 258:13513–13516. [PubMed: 6315699]
331. Ives HE, Yee VJ, Warnock DG. Mixed type inhibition of the renal Na⁺/H⁺ antiporter by Li⁺ and amiloride. Evidence for a modifier site. *J Biol Chem.* 1983; 258:9710–9716. [PubMed: 6309782]
332. Izumi Y, Hori K, Nakayama Y, Kimura M, Hasuike Y, Nanami M, Kohda Y, Otaki Y, Kuragano T, Obinata M, Kawahara K, Tanoue A, Tomita K, Nakanishi T, Nonoguchi H. Aldosterone requires vasopressin V1a receptors on intercalated cells to mediate acid-base homeostasis. *J Am Soc Nephrol.* 22:673–680. [PubMed: 21415155]
333. Jacobson HR. Effects of CO₂ and acetazolamide on bicarbonate and fluid transport in rabbit proximal tubules. *Am J Physiol.* 1981; 240:F54–62. [PubMed: 6779638]
334. Jacobson HR. Medullary collecting duct acidification. Effects of potassium, HCO₃ concentration, and pCO₂. *The Journal of clinical investigation.* 1984; 74:2107–2114. [PubMed: 6096401]
335. Jacques T, Picard N, Miller RL, Riemony KA, Houillier P, Sohet F, Ramakrishnan SK, Busst CJ, Jayat M, Corniere N, Hassan H, Aronson PS, Hennings JC, Hubner CA, Nelson RD,

- Chambrey R, Eladari D. Overexpression of pendrin in intercalated cells produces chloride-sensitive hypertension. *Journal of the American Society of Nephrology : JASN*. 2013; 24:1104–1113. [PubMed: 23766534]
336. Jarolim P, Palek J, Amato D, Hassan K, Sapak P, Nurse GT, Rubin HL, Zhai S, Sahr KE, Liu SC. Deletion in erythrocyte band 3 gene in malaria-resistant Southeast Asian ovalocytosis. *Proc Natl Acad Sci U S A*. 1991; 88:11022–11026. [PubMed: 1722314]
337. Jehle S, Zanetti A, Muser J, Hulter HN, Krapf R. Partial neutralization of the acidogenic Western diet with potassium citrate increases bone mass in postmenopausal women with osteopenia. *J Am Soc Nephrol*. 2006; 17:3213–3222. [PubMed: 17035614]
338. Jin XH, Siragy HM, Carey RM. Renal interstitial cGMP mediates natriuresis by direct tubule mechanism. *Hypertension*. 2001; 38:309–316. [PubMed: 11566896]
339. Joshua B, Kaplan DM, Raveh E, Lotan D, Anikster Y. Audiometric and imaging characteristics of distal renal tubular acidosis and deafness. *J Laryngol Otol*. 2008; 122:193–198. [PubMed: 17669226]
340. Josifovska T, Nonoguchi H, Machida K, Tomita K. Mechanisms of down-regulation of the renal parathyroid hormone receptor in rats with chronic renal failure. *Nephron Exp Nephrol*. 2003; 93:e141–149. [PubMed: 12759575]
341. Joyner J, Neves LA, Granger JP, Alexander BT, Merrill DC, Chappell MC, Ferrario CM, Davis WP, Brosnihan KB. Temporal-spatial expression of ANG-(1-7) and angiotensin-converting enzyme 2 in the kidney of normal and hypertensive pregnant rats. *American journal of physiology Regulatory, integrative and comparative physiology*. 2007; 293:R169–177.
342. Kahn AM, Dolson GM, Hise MK, Bennett SC, Weinman EJ. Parathyroid hormone and dibutyryl cAMP inhibit Na⁺/H⁺ exchange in renal brush border vesicles. *Am J Physiol*. 1985; 248:F212–218. [PubMed: 2982285]
343. Kane PM. Disassembly and reassembly of the yeast vacuolar H⁺-ATPase in vivo. *J Biol Chem*. 1995; 270:17025–17032. [PubMed: 7622524]
344. Kane PM. Targeting reversible disassembly as a mechanism of controlling V ATPase activity. *Curr Protein Pept Sci*. 13:117–123. [PubMed: 22044153]
345. Kao L, Sassani P, Azimov R, Pushkin A, Abuladze N, Peti-Peterdi J, Liu W, Newman D, Kurtz I. Oligomeric structure and minimal functional unit of the electrogenic sodium bicarbonate cotransporter NBCe1-A. *J Biol Chem*. 2008; 283:26782–26794. [PubMed: 18658147]
346. Karet FE, Finberg KE, Nayir A, Bakaloglu A, Ozen S, Hulton SA, Sanjad SA, Al-Sabban EA, Medina JF, Lifton RP. Localization of a gene for autosomal recessive distal renal tubular acidosis with normal hearing (rdRTA2) to 7q33-34. *Am J Hum Genet*. 1999; 65:1656–1665. [PubMed: 10577919]
347. Karet FE, Finberg KE, Nelson RD, Nayir A, Mocan H, Sanjad SA, Rodriguez-Soriano J, Santos F, Cremers CW, Di Pietro A, Hoffbrand BI, Winiarski J, Bakaloglu A, Ozen S, Dusunsal R, Goodyer P, Hulton SA, Wu DK, Skvorak AB, Morton CC, Cunningham MJ, Jha V, Lifton RP. Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. *Nat Genet*. 1999; 21:84–90. [PubMed: 9916796]
348. Karet FE, Gainza FJ, Gyory AZ, Unwin RJ, Wrong O, Tanner MJ, Nayir A, Alpay H, Santos F, Hulton SA, Bakaloglu A, Ozen S, Cunningham MJ, di Pietro A, Walker WG, Lifton RP. Mutations in the chloride-bicarbonate exchanger gene AE1 cause autosomal dominant but not autosomal recessive distal renal tubular acidosis. *Proc Natl Acad Sci U S A*. 1998; 95:6337–6342. [PubMed: 9600966]
349. Karim ZG, Chambrey R, Chalumeau C, Defontaine N, Warnock DG, Paillard M, Poggioli J. Regulation by PKC isoforms of Na⁺/H⁺ exchanger in luminal membrane vesicles isolated from cortical tubules. *Am J Physiol*. 1999; 277:F773–778. [PubMed: 10564242]
350. Karniski LP, Aronson PS. Anion exchange pathways for Cl[–] transport in rabbit renal microvillus membranes. *Am J Physiol*. 1987; 253:F513–521. [PubMed: 3631282]
351. Karniski LP, Aronson PS. Chloride/formate exchange with formic acid recycling: a mechanism of active chloride transport across epithelial membranes. *Proc Natl Acad Sci U S A*. 1985; 82:6362–6365. [PubMed: 3862136]

352. Karniski LP, Wang T, Everett LA, Green ED, Giebisch G, Aronson PS. Formate-stimulated NaCl absorption in the proximal tubule is independent of the pendrin protein. *Am J Physiol Renal Physiol.* 2002; 283:F952–956. [PubMed: 12372770]
353. Katzir Z, Dinour D, Reznik-Wolf H, Nissenkorn A, Holtzman E. Familial pure proximal renal tubular acidosis--a clinical and genetic study. *Nephrol Dial Transplant.* 2008; 23:1211–1215. [PubMed: 17881426]
354. Kaufman AM, Kahn T. Complementary role of citrate and bicarbonate excretion in acid-base balance in the rat. *Am J Physiol.* 1988; 255:F182–187. [PubMed: 3394809]
355. Kaunisto K, Parkkila S, Rajaniemi H, Waheed A, Grubb J, Sly WS. Carbonic anhydrase XIV: luminal expression suggests key role in renal acidification. *Kidney Int.* 2002; 61:2111–2118. [PubMed: 12028451]
356. Khanna A, Simoni J, Wesson DE. Endothelin-induced increased aldosterone activity mediates augmented distal nephron acidification as a result of dietary protein. *Journal of the American Society of Nephrology : JASN.* 2005; 16:1929–1935. [PubMed: 15872074]
357. Kikeri D, Azar S, Sun A, Zeidel ML, Hebert SC. Na(+)-H+ antiporter and Na(+)-(HCO3-)n symporter regulate intracellular pH in mouse medullary thick limbs of Henle. *Am J Physiol.* 1990; 258:F445–456. [PubMed: 2156445]
358. Kikeri D, Sun A, Zeidel ML, Hebert SC. Cell membranes impermeable to NH3. *Nature.* 1989; 339:478–480. [PubMed: 2725680]
359. Kikeri D, Sun A, Zeidel ML, Hebert SC. Cellular NH4+/K+ transport pathways in mouse medullary thick limb of Henle. Regulation by intracellular pH. *J Gen Physiol.* 1992; 99:435–461. [PubMed: 1588302]
360. Kim GH, Ecelbarger C, Knepper MA, Packer RK. Regulation of thick ascending limb ion transporter abundance in response to altered acid/base intake. *Journal of the American Society of Nephrology : JASN.* 1999; 10:935–942. [PubMed: 10232678]
361. Kim HY, Verlander JW, Bishop JM, Cain BD, Han KH, Igarashi P, Lee HW, Handlogten ME, Weiner ID. Basolateral expression of the ammonia transporter family member Rh C glycoprotein in the mouse kidney. *Am J Physiol Renal Physiol.* 2009; 296:F543–555. [PubMed: 19129254]
362. Kim J, Kim YH, Cha JH, Tisher CC, Madsen KM. Intercalated cell subtypes in connecting tubule and cortical collecting duct of rat and mouse. *Journal of the American Society of Nephrology : JASN.* 1999; 10:1–12. [PubMed: 9890303]
363. Kim JH, Lee-Kwon W, Park JB, Ryu SH, Yun CH, Donowitz M. Ca(2+)-dependent inhibition of Na+/H+ exchanger 3 (NHE3) requires an NHE3-E3KARP- α -actinin-4 complex for oligomerization and endocytosis. *J Biol Chem.* 2002; 277:23714–23724. [PubMed: 11948184]
364. Kim YH, Kwon TH, Christensen BM, Nielsen J, Wall SM, Madsen KM, Frokiaer J, Nielsen S. Altered expression of renal acid-base transporters in rats with lithium-induced NDI. *Am J Physiol Renal Physiol.* 2003; 285:F1244–1257. [PubMed: 12944321]
365. Kim YH, Kwon TH, Frische S, Kim J, Tisher CC, Madsen KM, Nielsen S. Immunocytochemical localization of pendrin in intercalated cell subtypes in rat and mouse kidney. *American journal of physiology Renal physiology.* 2002; 283:F744–754. [PubMed: 12217866]
366. Kim YH, Verlander JW, Matthews SW, Kurtz I, Shin W, Weiner ID, Everett LA, Green ED, Nielsen S, Wall SM. Intercalated cell H+/OH– transporter expression is reduced in Slc26a4 null mice. *Am J Physiol Renal Physiol.* 2005; 289:F1262–1272. [PubMed: 16144965]
367. Kinne-Saffran E, Beauwens R, Kinne R. An ATP-driven proton pump in brush-border membranes from rat renal cortex. *J Membr Biol.* 1982; 64:67–76. [PubMed: 6120239]
368. Kinne-Saffran E, Kinne R. Further evidence for the existence of an intrinsic bicarbonate-stimulated Mg2+-ATPase in brush border membranes isolated from rat kidney cortex. *J Membr Biol.* 1979; 49:235–251. [PubMed: 226712]
369. Kinne-Saffran E, Kinne R. Presence of bicarbonate stimulated ATPase in the brush border microvillus membranes of the proximal tubule. *Proc Soc Exp Biol Med.* 1974; 146:751–753. [PubMed: 4276357]
370. Kinsella J, Cujdik T, Sacktor B. Na+-H+ exchange activity in renal brush border membrane vesicles in response to metabolic acidosis: The role of glucocorticoids. *Proc Natl Acad Sci U S A.* 1984; 81:630–634. [PubMed: 6320201]

371. Kinsella JL, Aronson PS. Amiloride inhibition of the Na⁺-H⁺ exchanger in renal microvillus membrane vesicles. *Am J Physiol.* 1981; 241:F374–379. [PubMed: 7315961]
372. Kinsella JL, Aronson PS. Interaction of NH₄⁺ and Li⁺ with the renal microvillus membrane Na⁺-H⁺ exchanger. *Am J Physiol.* 1981; 241:C220–226. [PubMed: 6272583]
373. Kinsella JL, Aronson PS. Properties of the Na⁺-H⁺ exchanger in renal microvillus membrane vesicles. *Am J Physiol.* 1980; 238:F461–469. [PubMed: 7386626]
374. Klisic J, Hu MC, Nief V, Reyes L, Fuster D, Moe OW, Ambuhl PM. Insulin activates Na⁽⁺⁾/H⁽⁺⁾ exchanger 3: biphasic response and glucocorticoid dependence. *Am J Physiol Renal Physiol.* 2002; 283:F532–539. [PubMed: 12167605]
375. Knauf F, Yang CL, Thomson RB, Mentone SA, Giebisch G, Aronson PS. Identification of a chloride-formate exchanger expressed on the brush border membrane of renal proximal tubule cells. *Proc Natl Acad Sci U S A.* 2001; 98:9425–9430. [PubMed: 11459928]
376. Ko SB, Luo X, Hager H, Rojek A, Choi JY, Licht C, Suzuki M, Muallem S, Nielsen S, Ishibashi K. AE4 is a DIDS-sensitive Cl⁽⁻⁾/HCO⁽⁻⁾(3) exchanger in the basolateral membrane of the renal CCD and the SMG duct. *Am J Physiol Cell Physiol.* 2002; 283:C1206–1218. [PubMed: 12225984]
377. Kobayashi K, Uchida S, Mizutani S, Sasaki S, Marumo F. Intrarenal and cellular localization of CLC-K2 protein in the mouse kidney. *J Am Soc Nephrol.* 2001; 12:1327–1334. [PubMed: 11423561]
378. Kocinsky HS, Dynia DW, Wang T, Aronson PS. NHE3 phosphorylation at serines 552 and 605 does not directly affect NHE3 activity. *Am J Physiol Renal Physiol.* 2007; 293:F212–218. [PubMed: 17409282]
379. Kocinsky HS, Girardi AC, Biemesderfer D, Nguyen T, Mentone S, Orlowski J, Aronson PS. Use of phospho-specific antibodies to determine the phosphorylation of endogenous Na⁺/H⁺ exchanger NHE3 at PKA consensus sites. *Am J Physiol Renal Physiol.* 2005; 289:F249–258. [PubMed: 15687252]
380. Koeppen BM. Conductive properties of the rabbit outer medullary collecting duct: inner stripe. *Am J Physiol.* 1985; 248:F500–506. [PubMed: 3985156]
381. Koeppen BM. Conductive properties of the rabbit outer medullary collecting duct: outer stripe. *Am J Physiol.* 1986; 250:F70–76. [PubMed: 3942227]
382. Koeppen BM. Electrophysiological identification of principal and intercalated cells in the rabbit outer medullary collecting duct. *Pflugers Arch.* 1987; 409:138–141. [PubMed: 3039450]
383. Koeppen BM, Helman SI. Acidification of luminal fluid by the rabbit cortical collecting tubule perfused in vitro. *Am J Physiol.* 1982; 242:F521–531. [PubMed: 7081439]
384. Kokko JP. Sodium chloride and water transport in the descending limb of Henle. *The Journal of clinical investigation.* 1970; 49:1838–1846. [PubMed: 5460286]
385. Kokko JP. Urea transport in proximal tubule and the descending limb of Henle. *The Journal of clinical investigation.* 1972; 51:1999–2008. [PubMed: 5054459]
386. Kolb RJ, Woost PG, Hopfer U. Membrane trafficking of angiotensin receptor type-1 and mechanochemical signal transduction in proximal tubule cells. *Hypertension.* 2004; 44:352–359. [PubMed: 15262908]
387. Kondo Y, Fromter E. Axial heterogeneity of sodium-bicarbonate cotransport in proximal straight tubule of rabbit kidney. *Pflugers Arch.* 1987; 410:481–486. [PubMed: 2829111]
388. Konrad M, Vollmer M, Lemmink HH, van den Heuvel LP, Jeck N, Vargas-Poussou R, Lakings A, Ruf R, Deschenes G, Antignac C, Guay-Woodford L, Knoers NV, Seyberth HW, Feldmann D, Hildebrandt F. Mutations in the chloride channel gene CLCNKB as a cause of classic Bartter syndrome. *J Am Soc Nephrol.* 2000; 11:1449–1459. [PubMed: 10906158]
389. Krahn TA, Aronson PS, Weinstein AM. Weak acid permeability of a villous membrane: formic acid transport across rat proximal tubule. *Bull Math Biol.* 1994; 56:459–490. [PubMed: 8087078]
390. Krahn TA, Weinstein AM. Acid/base transport in a model of the proximal tubule brush border: impact of carbonic anhydrase. *Am J Physiol.* 1996; 270:F344–355. [PubMed: 8779897]
391. Krapf R. Basolateral membrane H⁺/OH⁻/HCO₃⁻ transport in the rat cortical thick ascending limb. Evidence for an electrogenic Na⁺/HCO₃⁻ cotransporter in parallel with a Na⁺/H⁺ antiporter. *J Clin Invest.* 1988; 82:234–241. [PubMed: 2839547]

392. Krapf R, Alpern RJ, Rector FC Jr, Berry CA. Basolateral membrane Na/base cotransport is dependent on CO₂/HCO₃ in the proximal convoluted tubule. *J Gen Physiol.* 1987; 90:833–853. [PubMed: 2831294]
393. Krug AW, Papavassiliou F, Hopfer U, Ullrich KJ, Gekle M. Aldosterone stimulates surface expression of NHE3 in renal proximal brush borders. *Pflugers Arch.* 2003; 446:492–496. [PubMed: 12684793]
394. Kunau RT Jr, Hart JI, Walker KA. Effect of metabolic acidosis on proximal tubular total CO₂ absorption. *Am J Physiol.* 1985; 249:F62–68. [PubMed: 3925794]
395. Kunau RT Jr, Walker KA. Total CO₂ absorption in the distal tubule of the rat. *Am J Physiol.* 1987; 252:F468–473. [PubMed: 3103468]
396. Kunimi M, Seki G, Hara C, Taniguchi S, Uwatoko S, Goto A, Kimura S, Fujita T. Dopamine inhibits renal Na⁺:HCO₃[−] cotransporter in rabbits and normotensive rats but not in spontaneously hypertensive rats. *Kidney Int.* 2000; 57:534–543. [PubMed: 10652030]
397. Kurashima K, D'Souza S, Szaszi K, Ramjeesingh R, Orlowski J, Grinstein S. The apical Na(+)/H(+) exchanger isoform NHE3 is regulated by the actin cytoskeleton. *J Biol Chem.* 1999; 274:29843–29849. [PubMed: 10514464]
398. Kurashima K, Yu FH, Cabado AG, Szabo EZ, Grinstein S, Orlowski J. Identification of sites required for down-regulation of Na⁺/H⁺ exchanger NHE3 activity by cAMP-dependent protein kinase. phosphorylation-dependent and -independent mechanisms. *J Biol Chem.* 1997; 272:28672–28679. [PubMed: 9353335]
399. Kurtz I. Apical and basolateral Na⁺/H⁺ exchange in the rabbit outer medullary thin descending limb of Henle: role in intracellular pH regulation. *The Journal of membrane biology.* 1988; 106:253–260. [PubMed: 2854165]
400. Kurtz I. Apical Na⁺/H⁺ antiporter and glycolysis-dependent H⁺-ATPase regulate intracellular pH in the rabbit S3 proximal tubule. *J Clin Invest.* 1987; 80:928–935. [PubMed: 2888787]
401. Kurtz I. Basolateral membrane Na⁺/H⁺ antiport, Na⁺/base cotransport, and Na⁺-independent Cl[−]/base exchange in the rabbit S3 proximal tubule. *J Clin Invest.* 1989; 83:616–622. [PubMed: 2536402]
402. Kurtz I. NBCe1 as a model carrier for understanding the structure-function properties of Na-coupled SLC4 transporters in health and disease. *Pflugers Archiv : European journal of physiology.* 2014
403. Kurtz I. NBCe1 as a model carrier for understanding the structure-function properties of Na-coupled SLC4 transporters in health and disease. *Pflugers Archiv: Europ J of Physiol.* 2014 [Epub ahead of print].
404. Kurtz I, Dass PD, Cramer S. The importance of renal ammonia metabolism to whole body acid-base balance: a reanalysis of the pathophysiology of renal tubular acidosis. *Miner Electrolyte Metab.* 1990; 16:331–340. [PubMed: 2283996]
405. Kurtz I, Maher T, Hulter HN, Schambelan M, Sebastian A. Effect of diet on plasma acid-base composition in normal humans. *Kidney Int.* 1983; 24:670–680. [PubMed: 6663989]
406. Kurtz I, Nagami G, Yanagawa N, Li L, Emmons C, Lee I. Mechanism of apical and basolateral Na(+)-independent Cl[−]/base exchange in the rabbit superficial proximal straight tubule. *J Clin Invest.* 1994; 94:173–183. [PubMed: 8040258]
407. Kurtz I, Petrasek D, Tatishchev S. Molecular mechanisms of electrogenic sodium bicarbonate cotransport: structural and equilibrium thermodynamic considerations. *J Membr Biol.* 2004; 197:77–90. [PubMed: 15014910]
408. Kurtz I, Star R, Balaban RS, Garvin JL, Knepper MA. Spontaneous luminal disequilibrium pH in S3 proximal tubules. Role in ammonia and bicarbonate transport. *J Clin Invest.* 1986; 78:989–996. [PubMed: 3760195]
409. Kurtzman NA. Renal tubular acidosis syndromes. *South Med J.* 2000; 93:1042–1052. [PubMed: 11095551]
410. Kwon TH, Fulton C, Wang W, Kurtz I, Frokiaer J, Aalkjaer C, Nielsen S. Chronic metabolic acidosis upregulates rat kidney Na-HCO cotransporters NBCn1 and NBC3 but not NBC1. *Am J Physiol Renal Physiol.* 2002; 282:F341–351. [PubMed: 11788449]

411. Kyllonen MS, Parkkila S, Rajaniemi H, Waheed A, Grubb JH, Shah GN, Sly WS, Kaunisto K. Localization of carbonic anhydrase XII to the basolateral membrane of H⁺-secreting cells of mouse and rat kidney. *J Histochem Cytochem*. 2003; 51:1217–1224. [PubMed: 12923247]
412. L'Hoste S, Diakov A, Andrini O, Genete M, Pinelli L, Grand T, Keck M, Paulais M, Beck L, Korbmacher C, Teulon J, Lourdel S. Characterization of the mouse CIC-K1/Barttin chloride channel. *Biochimica et biophysica acta*. 2013; 1828:2399–2409. [PubMed: 23791703]
413. Lacruz RS, Nanci A, White SN, Wen X, Wang H, Zalzal SF, Luong VQ, Schuetter VL, Conti PS, Kurtz I, Paine ML. The sodium bicarbonate cotransporter (NBCe1) is essential for normal development of mouse dentition. *J Biol Chem*. 285:24432–24438. [PubMed: 20529845]
414. Laghmani K, Borensztein P, Ambuhl P, Froissart M, Bichara M, Moe OW, Alpern RJ, Paillard M. Chronic metabolic acidosis enhances NHE-3 protein abundance and transport activity in the rat thick ascending limb by increasing NHE-3 mRNA. *The Journal of clinical investigation*. 1997; 99:24–30. [PubMed: 9011571]
415. Laghmani K, Preisig PA, Moe OW, Yanagisawa M, Alpern RJ. Endothelin-1/endothelin-B receptor-mediated increases in NHE3 activity in chronic metabolic acidosis. *The Journal of clinical investigation*. 2001; 107:1563–1569. [PubMed: 11413164]
416. Lamprecht G, Weinman EJ, Yun CH. The role of NHERF and E3KARP in the cAMP-mediated inhibition of NHE3. *J Biol Chem*. 1998; 273:29972–29978. [PubMed: 9792717]
417. Laradi A, Sakhrani LM, Massry SG. Effect of dopamine on sodium uptake by renal proximal tubule cells of rabbit. *Miner Electrolyte Metab*. 1986; 12:303–307. [PubMed: 3807828]
418. Laroche-Joubert N, Marsy S, Michelet S, Imbert-Teboul M, Doucet A. Protein kinase A-independent activation of ERK and H,K-ATPase by cAMP in native kidney cells: role of Epac I. *J Biol Chem*. 2002; 277:18598–18604. [PubMed: 11897793]
419. Laski ME, Kurtzman NA. Characterization of acidification in the cortical and medullary collecting tubule of the rabbit. *The Journal of clinical investigation*. 1983; 72:2050–2059. [PubMed: 6417167]
420. Laski ME, Kurtzman NA. Collecting tubule adaptation to respiratory acidosis induced in vivo. *Am J Physiol*. 1990; 258:F15–20. [PubMed: 2105660]
421. Lee HW, Verlander JW, Bishop JM, Igarashi P, Handlogten ME, Weiner ID. Collecting duct-specific Rh C glycoprotein deletion alters basal and acidosis-stimulated renal ammonia excretion. *Am J Physiol Renal Physiol*. 2009; 296:F1364–1375. [PubMed: 19321595]
422. Lee HW, Verlander JW, Bishop JM, Nelson RD, Handlogten ME, Weiner ID. Effect of intercalated cell-specific Rh C glycoprotein deletion on basal and metabolic acidosis-stimulated renal ammonia excretion. *Am J Physiol Renal Physiol*. 299:F369–379. [PubMed: 20462967]
423. Lee HW, Verlander JW, Handlogten ME, Han KH, Weiner ID. Effect of collecting duct-specific deletion of both Rh B Glycoprotein (Rhbg) and Rh C Glycoprotein (Rhcg) on renal response to metabolic acidosis. *American journal of physiology Renal physiology*. 2014; 306:F389–400. [PubMed: 24338819]
424. Lee S, Grichtchenko II, Boron WF. Distinguishing HCO₃⁻ from CO₃²⁻ transport by NBCe1-A. *FASEB J*. 2011; 5:656.9.
425. Lee SK, Boron WF, Parker MD. Substrate specificity of the electrogenic sodium/bicarbonate cotransporter NBCe1-A (SLC4A4, variant A) from humans and rabbits. *Am J Physiol Renal Physiol*. 304:F883–899. [PubMed: 23324180]
426. Lee-Kwon W, Kawano K, Choi JW, Kim JH, Donowitz M. Lysophosphatidic acid stimulates brush border Na⁺/H⁺ exchanger 3 (NHE3) activity by increasing its exocytosis by an NHE3 kinase A regulatory protein-dependent mechanism. *J Biol Chem*. 2003; 278:16494–16501. [PubMed: 12595524]
427. Lee-Kwon W, Kim JH, Choi JW, Kawano K, Cha B, Dartt DA, Zoukhri D, Donowitz M. Ca²⁺-dependent inhibition of NHE3 requires PKC alpha which binds to E3KARP to decrease surface NHE3 containing plasma membrane complexes. *American journal of physiology Cell physiology*. 2003; 285:C1527–1536. [PubMed: 12954600]
428. Leite-Dellova DC, Oliveira-Souza M, Malnic G, Mello-Aires M. Genomic and nongenomic dose-dependent biphasic effect of aldosterone on Na⁺/H⁺ exchanger in proximal S3 segment: role of cytosolic calcium. *Am J Physiol Renal Physiol*. 2008; 295:F1342–1352. [PubMed: 18715944]

429. Lemann J Jr, Adams ND, Wilz DR, Brenes LG. Acid and mineral balances and bone in familial proximal renal tubular acidosis. *Kidney Int.* 2000; 58:1267–1277. [PubMed: 10972690]
430. Lemann J Jr, Lennon EJ. Role of diet, gastrointestinal tract and bone in acid-base homeostasis. *Kidney Int.* 1972; 1:275–279. [PubMed: 4599246]
431. Lemann J Jr, Lennon EJ, Brock J. A potential error in the measurement of urinary titratable acid. *J Lab Clin Med.* 1966; 67:906–913. [PubMed: 5916139]
432. Lemann J Jr, Lennon EJ, Goodman AD, Litzow JR, Relman AS. The Net Balance of Acid in Subjects Given Large Loads of Acid or Alkali. *J Clin Invest.* 1965; 44:507–517. [PubMed: 14278167]
433. Leong PK, Devillez A, Sandberg MB, Yang LE, Yip DK, Klein JB, McDonough AA. Effects of ACE inhibition on proximal tubule sodium transport. *Am J Physiol Renal Physiol.* 2006; 290:F854–863. [PubMed: 16263808]
434. Lessa LM, Amorim JB, Fonteles MC, Malnic G. Effect of renoguanil on hydrogen/bicarbonate ion transport in rat renal tubules. *Regul Pept.* 2009; 157:37–43. [PubMed: 19540271]
435. Lessa LM, Carraro-Lacroix LR, Crajoinas RO, Bezerra CN, Dariolli R, Girardi AC, Fonteles MC, Malnic G. Mechanisms underlying the inhibitory effects of uroguanylin on NHE3 transport activity in renal proximal tubule. *Am J Physiol Renal Physiol.* 303:F1399–1408. [PubMed: 22952280]
436. Leviel F, Eladari D, Blanchard A, Poumarat JS, Paillard M, Pódevin RA. Pathways for HCO₃⁻ exit across the basolateral membrane in rat thick limbs. *Am J Physiol.* 1999; 276:F847–856. [PubMed: 10362773]
437. Leviel F, Hubner CA, Houillier P, Morla L, El Moghrabi S, Brideau G, Hassan H, Parker MD, Kurth I, Kougioumtzes A, Sinning A, Pech V, Riemyndy KA, Miller RL, Hummler E, Shull GE, Aronson PS, Doucet A, Wall SM, Chambrey R, Eladari D. The Na⁺-dependent chloride-bicarbonate exchanger SLC4A8 mediates an electroneutral Na⁺ reabsorption process in the renal cortical collecting ducts of mice. *J Clin Invest.* 120:1627–1635. [PubMed: 20389022]
438. Levine DZ. Effect of acute hypercapnia on proximal tubular water and bicarbonate reabsorption. *Am J Physiol.* 1971; 221:1164–1170. [PubMed: 5111258]
439. Levine DZ. An in vivo microperfusion study of distal tubule bicarbonate reabsorption in normal and ammonium chloride rats. *J Clin Invest.* 1985; 75:588–595. [PubMed: 2982915]
440. Levine DZ. Single-nephron studies: implications for acid-base regulation. *Kidney Int.* 1990; 38:744–761. [PubMed: 2232512]
441. Levine DZ, Iacovitti M, Buckman S, Harrison V. In vivo modulation of rat distal tubule net HCO₃⁻ flux by VIP, isoproterenol, angiotensin II, and ADH. *Am J Physiol.* 1994; 266:F878–883. [PubMed: 8023967]
442. Levine DZ, Iacovitti M, Nash L, Vidorpe D. Secretion of bicarbonate by rat distal tubules in vivo. Modulation by overnight fasting. *The Journal of clinical investigation.* 1988; 81:1873–1878. [PubMed: 3384953]
443. Levine DZ, Vidorpe D, Iacovitti M. Luminal chloride modulates rat distal tubule bidirectional bicarbonate flux in vivo. *J Clin Invest.* 1990; 85:1793–1798. [PubMed: 2347913]
444. Lewis SE, Erickson RP, Barnett LB, Venta PJ, Tashian RE. N-ethyl-N-nitrosourea-induced null mutation at the mouse Car-2 locus: an animal model for human carbonic anhydrase II deficiency syndrome. *Proceedings of the National Academy of Sciences of the United States of America.* 1988; 85:1962–1966. [PubMed: 3126501]
445. Li S, Sato S, Yang X, Preisig PA, Alpern RJ. Pyk2 activation is integral to acid stimulation of sodium/hydrogen exchanger 3. *The Journal of clinical investigation.* 2004; 114:1782–1789. [PubMed: 15599403]
446. Li X, Liu Y, Alvarez BV, Casey JR, Fliegel L. A novel carbonic anhydrase II binding site regulates NHE1 activity. *Biochemistry.* 2006; 45:2414–2424. [PubMed: 16475831]
447. Liang M, Knox FG. Production and functional roles of nitric oxide in the proximal tubule. *Am J Physiol Regul Integr Comp Physiol.* 2000; 278:R1117–1124. [PubMed: 10801277]
448. Licht C, Laghmani K, Yanagisawa M, Preisig PA, Alpern RJ. An autocrine role for endothelin-1 in the regulation of proximal tubule NHE3. *Kidney international.* 2004; 65:1320–1326. [PubMed: 15086471]

449. Liedtke CM, Hopfer U. Anion transport in brush border membranes isolated from rat small intestine. *Biochemical and biophysical research communications*. 1976; 76:579–585. [PubMed: 17408]
450. Lin Z, Jin S, Duan X, Wang T, Martini S, Hulamm P, Cha B, Hubbard A, Donowitz M, Guggino SE. Chloride channel (Clc)-5 is necessary for exocytic trafficking of Na⁺/H⁺ exchanger 3 (NHE3). *J Biol Chem*. 286:22833–22845. [PubMed: 21561868]
451. Liu F, Gesek FA. α (1)-Adrenergic receptors activate NHE1 and NHE3 through distinct signaling pathways in epithelial cells. *Am J Physiol Renal Physiol*. 2001; 280:F415–425. [PubMed: 11181403]
452. Liu FY, Cogan MG. Angiotensin II stimulates early proximal bicarbonate absorption in the rat by decreasing cyclic adenosine monophosphate. *J Clin Invest*. 1989; 84:83–91. [PubMed: 2544631]
453. Liu FY, Cogan MG. Angiotensin II: a potent regulator of acidification in the rat early proximal convoluted tubule. *J Clin Invest*. 1987; 80:272–275. [PubMed: 3597776]
454. Liu FY, Cogan MG. Axial heterogeneity in the rat proximal convoluted tubule. I. Bicarbonate, chloride, and water transport. *Am J Physiol*. 1984; 247:F816–821. [PubMed: 6496747]
455. Liu FY, Cogan MG. Kinetics of bicarbonate transport in the early proximal convoluted tubule. *Am J Physiol*. 1987; 253:F912–916. [PubMed: 3688241]
456. Liu Y, Xu JY, Wang DK, Wang L, Chen LM. Cloning and identification of two novel NBCe1 splice variants from mouse reproductive tract tissues: a comparative study of NCBT genes. *Genomics*. 98:112–119. [PubMed: 21600280]
457. Lombard WE, Kokko JP, Jacobson HR. Bicarbonate transport in cortical and outer medullary collecting tubules. *Am J Physiol*. 1983; 244:F289–296. [PubMed: 6829761]
458. Lonnerholm G, Wistrand PJ. Carbonic anhydrase in the human kidney: a histochemical and immunocytochemical study. *Kidney international*. 1984; 25:886–898. [PubMed: 6433100]
459. Lonnerholm G, Wistrand PJ. Membrane-bound carbonic anhydrase CA IV in the human kidney. *Acta physiologica Scandinavica*. 1991; 141:231–234. [PubMed: 1904675]
460. Lopes AG, Siebens AW, Giebisch G, Boron WF. Electrogenic Na/HCO₃ cotransport across basolateral membrane of isolated perfused Necturus proximal tubule. *Am J Physiol*. 1987; 253:F340–350. [PubMed: 3618795]
461. Lu J, Daly CM, Parker MD, Gill HS, Piermarini PM, Pelletier MF, Boron WF. Effect of human carbonic anhydrase II on the activity of the human electrogenic Na/HCO₃ cotransporter NBCe1-A in *Xenopus* oocytes. *The Journal of biological chemistry*. 2006; 281:19241–19250. [PubMed: 16687407]
462. Lu X, Garrelds IM, Wagner CA, Danser AH, Meima ME. (Pro)renin receptor is required for prorenin-dependent and -independent regulation of vacuolar H⁺-ATPase activity in MDCK.C11 collecting duct cells. *American journal of physiology Renal physiology*. 2013; 305:F417–425. [PubMed: 23698123]
463. Lucci MS, Pucacco LR, Carter NW, DuBose TD Jr. Evaluation of bicarbonate transport in rat distal tubule: effects of acid-base status. *Am J Physiol*. 1982; 243:F335–341. [PubMed: 7124948]
464. Lucci MS, Tinker JP, Weiner IM, DuBose TD Jr. Function of proximal tubule carbonic anhydrase defined by selective inhibition. *Am J Physiol*. 1983; 245:F443–449. [PubMed: 6414310]
465. Lucci MS, Warnock DG. Effects of anion-transport inhibitors on NaCl reabsorption in the rat superficial proximal convoluted tubule. *J Clin Invest*. 1979; 64:570–579. [PubMed: 457869]
466. Ludwig J, Kerscher S, Brandt U, Pfeiffer K, Getlawi F, Apps DK, Schagger H. Identification and characterization of a novel 9.2-kDa membrane sector-associated protein of vacuolar proton-ATPase from chromaffin granules. *The Journal of biological chemistry*. 1998; 273:10939–10947. [PubMed: 9556572]
467. Lynch IJ, Greenlee MM, Gumz ML, Rudin A, Xia SL, Wingo CS. Heterogeneity of H-K-ATPase-mediated acid secretion along the mouse collecting duct. *Am J Physiol Renal Physiol*. 298:F408–415. [PubMed: 19923411]
468. Lynch IJ, Rudin A, Xia SL, Stow LR, Shull GE, Weiner ID, Cain BD, Wingo CS. Impaired acid secretion in cortical collecting duct intercalated cells from H-K ATPase-deficient mice: role of HK α isoforms. *Am J Physiol Renal Physiol*. 2008; 294:F621–627. [PubMed: 18057185]

469. Macdonald HM, New SA, Fraser WD, Campbell MK, Reid DM. Low dietary potassium intakes and high dietary estimates of net endogenous acid production are associated with low bone mineral density in premenopausal women and increased markers of bone resorption in postmenopausal women. *Am J Clin Nutr.* 2005; 81:923–933. [PubMed: 15817873]
470. Madsen KM, Clapp WL, Verlander JW. Structure and function of the inner medullary collecting duct. *Kidney Int.* 1988; 34:441–454. [PubMed: 3059025]
471. Madsen KM, Kim J, Tisher CC. Intracellular band 3 immunostaining in type A intercalated cells of rabbit kidney. *The American journal of physiology.* 1992; 262:F1015–1022. [PubMed: 1377872]
472. Madsen KM, Tisher CC. Response of intercalated cells of rat outer medullary collecting duct to chronic metabolic acidosis. *Lab Invest.* 1984; 51:268–276. [PubMed: 6471808]
473. Madsen KM, Tisher CC. Structural-functional relationships along the distal nephron. *Am J Physiol.* 1986; 250:F1–15.
474. Madsen KM, Verlander JW, Kim J, Tisher CC. Morphological adaptation of the collecting duct to acid-base disturbances. *Kidney international Supplement.* 1991; 33:S57–63. [PubMed: 1890802]
475. Malnic G, De Mello Aires M, Giebisch G. Micropuncture study of renal tubular hydrogen ion transport in the rat. *Am J Physiol.* 1972; 222:147–158. [PubMed: 5060213]
476. Malnic G, de Mello-Aires M. Kinetic study of bicarbonate reabsorption in proximal tubule of the rat. *Am J Physiol.* 1971; 220:1759–1767. [PubMed: 5087824]
477. Malnic G, Geibel JP. Cell pH and H(+) secretion by S3 segment of mammalian kidney: role of H(+)-ATPase and Cl(–). *The Journal of membrane biology.* 2000; 178:115–125. [PubMed: 11083900]
478. Matsumura Y, Uchida S, Kondo Y, Miyazaki H, Ko SB, Hayama A, Morimoto T, Liu W, Arisawa M, Sasaki S, Marumo F. Overt nephrogenic diabetes insipidus in mice lacking the CLC-K1 chloride channel. *Nat Genet.* 1999; 21:95–98. [PubMed: 9916798]
479. Matsuzaki K, Schuster VL, Stokes JB. Reduction in sensitivity to Cl[–] channel blockers by HCO₃[–]–CO₂ in rabbit cortical collecting duct. *Am J Physiol.* 1989; 257:C102–109. [PubMed: 2473650]
480. Maunsbach AB, Vorum H, Kwon TH, Nielsen S, Simonsen B, Choi I, Schmitt BM, Boron WF, Aalkjaer C. Immunoelectron microscopic localization of the electrogenic Na/HCO₃ cotransporter in rat and ambystoma kidney. *J Am Soc Nephrol.* 2000; 11:2179–2189. [PubMed: 11095641]
481. May RC, Kelly RA, Mitch WE. Metabolic acidosis stimulates protein degradation in rat muscle by a glucocorticoid-dependent mechanism. *J Clin Invest.* 1986; 77:614–621. [PubMed: 3511100]
482. McAlear SD, Liu X, Williams JB, McNicholas-Bevensee CM, Bevensee MO. Electrogenic Na/HCO₃ cotransporter (NBCe1) variants expressed in *Xenopus* oocytes: functional comparison and roles of the amino and carboxy termini. *The Journal of general physiology.* 2006; 127:639–658. [PubMed: 16735752]
483. McKinney TD, Burg MB. Bicarbonate absorption by rabbit cortical collecting tubules in vitro. *The American journal of physiology.* 1978; 234:F141–145. [PubMed: 623305]
484. McKinney TD, Burg MB. Bicarbonate secretion by rabbit cortical collecting tubules in vitro. *The Journal of clinical investigation.* 1978; 61:1421–1427. [PubMed: 659604]
485. McKinney TD, Burg MB. Bicarbonate transport by rabbit cortical collecting tubules. Effect of acid and alkali loads in vivo on transport in vitro. *The Journal of clinical investigation.* 1977; 60:766–768. [PubMed: 19497]
486. McKinney TD, Burg MB. Bicarbonate and fluid absorption by renal proximal straight tubules. *Kidney international.* 1977; 12:1–8. [PubMed: 894911]
487. McKinney TD, Davidson KK. Bicarbonate transport in collecting tubules from outer stripe of outer medulla of rabbit kidneys. *Am J Physiol.* 1987; 253:F816–822. [PubMed: 3120600]
488. McKinney TD, Davidson KK. Effect of potassium depletion and protein intake in vivo on renal tubular bicarbonate transport in vitro. *Am J Physiol.* 1987; 252:F509–516. [PubMed: 3103470]
489. McKinney TD, Davidson KK. Effects of respiratory acidosis on HCO₃[–] transport by rabbit collecting tubules. *The American journal of physiology.* 1988; 255:F656–665. [PubMed: 2845808]

490. McKinney TD, Myers P. Bicarbonate transport by proximal tubules: effect of parathyroid hormone and dibutyl cyclic AMP. *Am J Physiol.* 1980; 238:F166–174. [PubMed: 6245584]
491. McKinney TD, Myers P. PTH inhibition of bicarbonate transport by proximal convoluted tubules. *Am J Physiol.* 1980; 239:F127–134. [PubMed: 6250409]
492. Mellas J, Hammerman MR. Phorbol ester-induced alkalinization of canine renal proximal tubular cells. *Am J Physiol.* 1986; 250:F451–459. [PubMed: 3953824]
493. Mello Aires M, Malnic G. Peritubular pH and PCO₂ in renal tubular acidification. *Am J Physiol.* 1975; 228:1766–1774. [PubMed: 239600]
494. Meneton P, Schultheis PJ, Greeb J, Nieman ML, Liu LH, Clarke LL, Duffy JJ, Doetschman T, Lorenz JN, Shull GE. Increased sensitivity to K⁺ deprivation in colonic H,K-ATPase-deficient mice. *J Clin Invest.* 1998; 101:536–542. [PubMed: 9449685]
495. Mercier O, Bichara M, Delahousse M, Prigent A, Leviel F, Paillard M. Effects of glucagon on H(+)–HCO₃[–] transport in Henle's loop, distal tubule, and collecting ducts in the rat. *The American journal of physiology.* 1989; 257:F1003–1014. [PubMed: 2574952]
496. Mercier O, Bichara M, Paillard M, Gardin JP, Leviel F. Parathyroid hormone contributes to volume expansion-induced inhibition of proximal reabsorption. *Am J Physiol.* 1985; 248:F100–103. [PubMed: 2982273]
497. Mercier O, Bichara M, Paillard M, Prigent A. Effects of parathyroid hormone and urinary phosphate on collecting duct hydrogen secretion. *Am J Physiol.* 1986; 251:F802–809. [PubMed: 3096150]
498. Merkulova M, Bakulina A, Thaker YR, Gruber G, Marshansky V. Specific motifs of the V-ATPase α 2-subunit isoform interact with catalytic and regulatory domains of ARNO. *Biochim Biophys Acta.* 1997; 1398–1409. [PubMed: 20153292]
499. Miller RT, Pollock AS. Modification of the internal pH sensitivity of the Na⁺/H⁺ antiporter by parathyroid hormone in a cultured renal cell line. *J Biol Chem.* 1987; 262:9115–9120. [PubMed: 3036848]
500. Milton AE, Weiner ID. Intracellular pH regulation in the rabbit cortical collecting duct A-type intercalated cell. *Am J Physiol.* 1997; 273:F340–347. [PubMed: 9321906]
501. Milton AE, Weiner ID. Regulation of B-type intercalated cell apical anion exchange activity by CO₂/HCO₃. *Am J Physiol.* 1998; 274:F1086–1094. [PubMed: 9841500]
502. Miura K, Sekine T, Takahashi K, Takita J, Harita Y, Ohki K, Park MJ, Hayashi Y, Tajima A, Ishihara M, Hisano M, Murai M, Igarashi T. Mutational analyses of the ATP6V1B1 and ATP6V0A4 genes in patients with primary distal renal tubular acidosis. *Nephrol Dial Transplant.* 28:2123–2130. [PubMed: 23729491]
503. Moe OW, Amemiya M, Yamaji Y. Activation of protein kinase A acutely inhibits and phosphorylates Na/H exchanger NHE-3. *J Clin Invest.* 1995; 96:2187–2194. [PubMed: 7593604]
504. Moe OW, Tejedor A, Levi M, Seldin DW, Preisig PA, Alpern RJ. Dietary NaCl modulates Na(+)–H⁺ antiporter activity in renal cortical apical membrane vesicles. *Am J Physiol.* 1991; 260:F130–137. [PubMed: 1847012]
505. Mohebbi N, Kovacicova J, Nowik M, Wagner CA. Thyroid hormone deficiency alters expression of acid-base transporters in rat kidney. *American journal of physiology Renal physiology.* 2007; 293:F416–427. [PubMed: 17409279]
506. Mohebbi N, Mihailova M, Wagner CA. The calcineurin inhibitor FK506 (tacrolimus) is associated with transient metabolic acidosis and altered expression of renal acid-base transport proteins. *Am J Physiol Renal Physiol.* 2009; 297:F499–509. [PubMed: 19439519]
507. Mohebbi N, Perna A, van der Wijst J, Becker HM, Capasso G, Wagner CA. Regulation of two renal chloride transporters, AE1 and pendrin, by electrolytes and aldosterone. *PloS one.* 2013; 8:e55286. [PubMed: 23383138]
508. Morel F. Sites of hormone action in the mammalian nephron. *Am J Physiol.* 1981; 240:F159–164. [PubMed: 6259951]
509. Mori K, Ogawa Y, Ebihara K, Tamura N, Tashiro K, Kuwahara T, Mukoyama M, Sugawara A, Ozaki S, Tanaka I, Nakao K. Isolation and characterization of CA XIV, a novel membrane-bound carbonic anhydrase from mouse kidney. *J Biol Chem.* 1999; 274:15701–15705. [PubMed: 10336468]

510. Moriyama Y, Nelson N. Cold inactivation of vacuolar proton-ATPases. *The Journal of biological chemistry*. 1989; 264:3577–3582. [PubMed: 2521638]
511. Muller-Berger S, Ducoudret O, Diakov A, Fromter E. The renal Na-HCO₃-cotransporter expressed in *Xenopus laevis* oocytes: change in stoichiometry in response to elevation of cytosolic Ca²⁺ concentration. *Pflügers Archiv : European journal of physiology*. 2001; 442:718–728. [PubMed: 11512028]
512. Muller-Berger S, Nesterov VV, Fromter E. Partial recovery of in vivo function by improved incubation conditions of isolated renal proximal tubule. II. Change of Na-HCO₃ cotransport stoichiometry and of response to acetazolamide. *Pflügers Arch*. 1997; 434:383–391. [PubMed: 9211803]
513. Murer H, Hopfer U, Kinne R. Sodium/proton antiport in brush-border-membrane vesicles isolated from rat small intestine and kidney. *Biochem J*. 1976; 154:597–604. [PubMed: 942389]
514. Murtazina R, Kovbasnjuk O, Zachos NC, Li X, Chen Y, Hubbard A, Hogema BM, Steplock D, Seidler U, Hoque KM, Tse CM, De Jonge HR, Weinman EJ, Donowitz M. Tissue-specific regulation of sodium/proton exchanger isoform 3 activity in Na(+)/H(+) exchanger regulatory factor 1 (NHERF1) null mice. cAMP inhibition is differentially dependent on NHERF1 and exchange protein directly activated by cAMP in ileum versus proximal tubule. *J Biol Chem*. 2007; 282:25141–25151. [PubMed: 17580307]
515. Musa-Aziz R, Chen LM, Pelletier MF, Boron WF. Relative CO₂/NH₃ selectivities of AQP1, AQP4, AQP5, AmtB, and RhAG. *Proc Natl Acad Sci U S A*. 2009; 106:5406–5411. [PubMed: 19273840]
516. Muto S, Yasoshima K, Yoshitomi K, Imai M, Asano Y. Electrophysiological identification of alpha- and beta-intercalated cells and their distribution along the rabbit distal nephron segments. *J Clin Invest*. 1990; 86:1829–1839. [PubMed: 2254448]
517. Muzalef A, Alshehri M, Al-Abidi A, Al-Trabolsi HA. Marble brain disease in two Saudi Arabian siblings. *Ann Trop Paediatr*. 2005; 25:213–218. [PubMed: 16156988]
518. Nagami GT, Chang JA, Plato ME, Santamaria R. Acid loading in vivo and low pH in culture increase angiotensin receptor expression: enhanced ammoniagenic response to angiotensin II. *Am J Physiol Renal Physiol*. 2008; 295:F1864–1870. [PubMed: 18945827]
519. Nakhoul NL, Boron WF. Acetate transport in the S3 segment of the rabbit proximal tubule and its effect on intracellular pH. *J Gen Physiol*. 1988; 92:395–412. [PubMed: 3225555]
520. Nakhoul NL, Chen LK, Boron WF. Intracellular pH regulation in rabbit S3 proximal tubule: basolateral Cl-HCO₃ exchange and Na-HCO₃ cotransport. *Am J Physiol*. 1990; 258:F371–381. [PubMed: 2155541]
521. Nakhoul NL, Lopes AG, Chaillet JR, Boron WF. Intracellular pH regulation in the S3 segment of the rabbit proximal tubule in HCO₃⁻-free solutions. *J Gen Physiol*. 1988; 92:369–393. [PubMed: 3225554]
522. Nguyen G. The (pro)renin receptor in health and disease. *Ann Med*. 42:13–18. [PubMed: 20104654]
523. Nguyen G, Delarue F, Burckle C, Bouzahir L, Giller T, Sraer JD. Pivotal role of the renin/prorenin receptor in angiotensin II production and cellular responses to renin. *J Clin Invest*. 2002; 109:1417–1427. [PubMed: 12045255]
524. Nord EP, Hafezi A, Kaunitz JD, Trizna W, Fine LG. pH gradient-dependent increased Na⁺-H⁺ antiport capacity of the rabbit remnant kidney. *Am J Physiol*. 1985; 249:F90–98. [PubMed: 2409819]
525. Nord EP, Howard MJ, Hafezi A, Moradeshagi P, Vaystub S, Insel PA. Alpha 2 adrenergic agonists stimulate Na⁺-H⁺ antiport activity in the rabbit renal proximal tubule. *J Clin Invest*. 1987; 80:1755–1762. [PubMed: 2890661]
526. Norgett EE, Golder ZJ, Lorente-Canovas B, Ingham N, Steel KP, Karet Frankl FE. Atp6v0a4 knockout mouse is a model of distal renal tubular acidosis with hearing loss, with additional extrarenal phenotype. *Proc Natl Acad Sci U S A*. 109:13775–13780. [PubMed: 22872862]
527. O'Connell DP, Botkin SJ, Ramos SI, Sibley DR, Ariano MA, Felder RA, Carey RM. Localization of dopamine D1A receptor protein in rat kidneys. *Am J Physiol*. 1995; 268:F1185–1197. [PubMed: 7611459]

528. O'Neil RG, Hayhurst RA. Functional differentiation of cell types of cortical collecting duct. *Am J Physiol.* 1985; 248:F449–453. [PubMed: 2579583]
529. O'Neil RG, Sansom SC. Characterization of apical cell membrane Na⁺ and K⁺ conductances of cortical collecting duct using microelectrode techniques. *Am J Physiol.* 1984; 247:F14–24. [PubMed: 6331197]
530. Obermuller N, Gretz N, Kriz W, Reilly RF, Witzgall R. The swelling-activated chloride channel CIC-2, the chloride channel CIC-3, and CIC-5, a chloride channel mutated in kidney stone disease, are expressed in distinct subpopulations of renal epithelial cells. *J Clin Invest.* 1998; 101:635–642. [PubMed: 9449697]
531. Orlowski A, De Giusti VC, Morgan PE, Aiello EA, Alvarez BV. Binding of carbonic anhydrase IX to extracellular loop 4 of the NBCe1 Na⁺/HCO₃⁻ cotransporter enhances NBCe1-mediated HCO₃⁻ influx in the rat heart. *American journal of physiology.* 2012; 303:C69–80. [PubMed: 22538240]
532. Orlowski J. Heterologous expression and functional properties of amiloride high affinity (NHE-1) and low affinity (NHE-3) isoforms of the rat Na/H exchanger. *J Biol Chem.* 1993; 268:16369–16377. [PubMed: 8393860]
533. Orlowski J, Kandasamy RA, Shull GE. Molecular cloning of putative members of the Na/H exchanger gene family. cDNA cloning, deduced amino acid sequence, and mRNA tissue expression of the rat Na/H exchanger NHE-1 and two structurally related proteins. *J Biol Chem.* 1992; 267:9331–9339. [PubMed: 1577762]
534. Owen EE, Johnson JH, Tyor MP. The effect of induced hyperammonemia on renal ammonia metabolism. *J Clin Invest.* 1961; 40:215–221. [PubMed: 13731806]
535. Owen EE, Robinson RR. Amino acid extraction and ammonia metabolism by the human kidney during the prolonged administration of ammonium chloride. *J Clin Invest.* 1963; 42:263–276. [PubMed: 13940831]
536. Ozaki K, Takada N, Tsujimoto K, Tsuji N, Kawamura T, Muso E, Ohta M, Itoh N. Localization of insulin receptor-related receptor in the rat kidney. *Kidney international.* 1997; 52:694–698. [PubMed: 9291189]
537. Packer RK, Curry CA, Brown KM. Urinary organic anion excretion in response to dietary acid and base loading. *Journal of the American Society of Nephrology : JASN.* 1995; 5:1624–1629. [PubMed: 7756596]
538. Padia SH, Howell NL, Siragy HM, Carey RM. Renal angiotensin type 2 receptors mediate natriuresis via angiotensin III in the angiotensin II type 1 receptor-blocked rat. *Hypertension.* 2006; 47:537–544. [PubMed: 16380540]
539. Padia SH, Kemp BA, Howell NL, Fournie-Zaluski MC, Roques BP, Carey RM. Conversion of renal angiotensin II to angiotensin III is critical for AT2 receptor-mediated natriuresis in rats. *Hypertension.* 2008; 51:460–465. [PubMed: 18158338]
540. Paillard M, Bichara M. Peptide hormone effects on urinary acidification and acid-base balance: PTH, ADH, and glucagon. *Am J Physiol.* 1989; 256:F973–985. [PubMed: 2660594]
541. Parkkila S, Parkkila AK, Saarnio J, Kivela J, Karttunen TJ, Kaunisto K, Waheed A, Sly WS, Tureci O, Virtanen I, Rajaniemi H. Expression of the membrane-associated carbonic anhydrase isozyme XII in the human kidney and renal tumors. *J Histochem Cytochem.* 2000; 48:1601–1608. [PubMed: 11101628]
542. Pastor-Soler N, Beaulieu V, Litvin TN, Da Silva N, Chen Y, Brown D, Buck J, Levin LR, Breton S. Bicarbonate-regulated adenylyl cyclase (sAC) is a sensor that regulates pH-dependent V-ATPase recycling. *The Journal of biological chemistry.* 2003; 278:49523–49529. [PubMed: 14512417]
543. Pastor-Soler NM, Hallows KR, Smolak C, Gong F, Brown D, Breton S. Alkaline pH- and cAMP-induced V-ATPase membrane accumulation is mediated by protein kinase A in epididymal clear cells. *American journal of physiology Cell physiology.* 2008; 294:C488–494. [PubMed: 18160485]
544. Patel AB, Frindt G, Palmer LG. Feedback inhibition of ENaC during acute sodium loading in vivo. *Am J Physiol Renal Physiol.* 304:F222–232. [PubMed: 23171553]

545. Paunescu TG, Da Silva N, Russo LM, McKee M, Lu HA, Breton S, Brown D. Association of soluble adenylyl cyclase with the V-ATPase in renal epithelial cells. *American journal of physiology Renal physiology*. 2008; 294:F130–138. [PubMed: 17959750]
546. Paunescu TG, Ljubojevic M, Russo LM, Winter C, McLaughlin MM, Wagner CA, Breton S, Brown D. cAMP stimulates apical V-ATPase accumulation, microvillar elongation, and proton extrusion in kidney collecting duct A-intercalated cells. *American journal of physiology Renal physiology*. 2010; 298:F643–654. [PubMed: 20053793]
547. Paunescu TG, Russo LM, Da Silva N, Kovacicova J, Mohebbi N, Van Hoek AN, McKee M, Wagner CA, Breton S, Brown D. Compensatory membrane expression of the V-ATPase B2 subunit isoform in renal medullary intercalated cells of B1-deficient mice. *Am J Physiol Renal Physiol*. 2007; 293:F1915–1926. [PubMed: 17898041]
548. Pech V, Kim YH, Weinstein AM, Everett LA, Pham TD, Wall SM. Angiotensin II increases chloride absorption in the cortical collecting duct in mice through a pendrin-dependent mechanism. *American journal of physiology Renal physiology*. 2007; 292:F914–920. [PubMed: 17077386]
549. Pech V, Pham TD, Hong S, Weinstein AM, Spencer KB, Duke BJ, Walp E, Kim YH, Sutliff RL, Bao HF, Eaton DC, Wall SM. Pendrin modulates ENaC function by changing luminal HCO₃. *J Am Soc Nephrol*. 21:1928–1941. [PubMed: 20966128]
550. Pech V, Zheng W, Pham TD, Verlander JW, Wall SM. Angiotensin II activates H⁺-ATPase in type A intercalated cells. *Journal of the American Society of Nephrology : JASN*. 2008; 19:84–91. [PubMed: 18178800]
551. Pendergrass KD, Averill DB, Ferrario CM, Diz DI, Chappell MC. Differential expression of nuclear AT1 receptors and angiotensin II within the kidney of the male congenic mRen2. Lewis rat. *American journal of physiology Renal physiology*. 2006; 290:F1497–1506. [PubMed: 16403834]
552. Pendergrass KD, Pirro NT, Westwood BM, Ferrario CM, Brosnihan KB, Chappell MC. Sex differences in circulating and renal angiotensins of hypertensive mRen(2). Lewis but not normotensive Lewis rats. *American journal of physiology Heart and circulatory physiology*. 2008; 295:H10–20. [PubMed: 18456730]
553. Peng H, Vijayakumar S, Schiene-Fischer C, Li H, Purkerson JM, Malesevic M, Liebscher J, Al-Awqati Q, Schwartz GJ. Secreted cyclophilin A, a peptidylprolyl cis-trans isomerase, mediates matrix assembly of hensin, a protein implicated in epithelial differentiation. *J Biol Chem*. 2009; 284:6465–6475. [PubMed: 19112104]
554. Peng Y, Amemiya M, Yang X, Fan L, Moe OW, Yin H, Preisig PA, Yanagisawa M, Alpern RJ. ET(B) receptor activation causes exocytic insertion of NHE3 in OKP cells. *Am J Physiol Renal Physiol*. 2001; 280:F34–42. [PubMed: 11133512]
555. Peng Y, Moe OW, Chu T, Preisig PA, Yanagisawa M, Alpern RJ. ETB receptor activation leads to activation and phosphorylation of NHE3. *Am J Physiol*. 1999; 276:C938–945. [PubMed: 10199826]
556. Pergher PS, Leite-Dellova D, de Mello-Aires M. Direct action of aldosterone on bicarbonate reabsorption in in vivo cortical proximal tubule. *Am J Physiol Renal Physiol*. 2009; 296:F1185–1193. [PubMed: 19225053]
557. Perry C, Baker OJ, Reyland ME, Grichtchenko II. PKC $\alpha\beta\gamma$ - and PKC δ -dependent endocytosis of NBCe1-A and NBCe1-B in salivary parotid acinar cells. *American journal of physiology*. 2009; 297:C1409–1423. [PubMed: 19783762]
558. Perry C, Blaine J, Le H, Grichtchenko II. PMA- and ANG II-induced PKC regulation of the renal Na⁺-HCO₃⁻ cotransporter (hkNBCe1). *American journal of physiology Renal physiology*. 2006; 290:F417–427. [PubMed: 16159892]
559. Perry C, Le H, Grichtchenko II. ANG II and calmodulin/CaMKII regulate surface expression and functional activity of NBCe1 via separate means. *American journal of physiology Renal physiology*. 2007; 293:F68–77. [PubMed: 17376763]
560. Petrovic S, Barone S, Wang Z, McDonough AA, Amlal H, Soleimani M. Slc26a6 (PAT1) deletion downregulates the apical Na⁺/H⁺ exchanger in the straight segment of the proximal tubule. *Am J Nephrol*. 2008; 28:330–338. [PubMed: 18046080]

561. Petrovic S, Barone S, Xu J, Conforti L, Ma L, Kujala M, Kere J, Soleimani M. SLC26A7: a basolateral Cl⁻/HCO₃⁻ exchanger specific to intercalated cells of the outer medullary collecting duct. *Am J Physiol Renal Physiol*. 2004; 286:F161–169. [PubMed: 12965893]
562. Petrovic S, Wang Z, Ma L, Soleimani M. Regulation of the apical Cl⁻/HCO₃⁻ 3 exchanger pendrin in rat cortical collecting duct in metabolic acidosis. *Am J Physiol Renal Physiol*. 2003; 284:F103–112. [PubMed: 12388388]
563. Piermarini PM, Kim EY, Boron WF. Evidence against a direct interaction between intracellular carbonic anhydrase II and pure C-terminal domains of SLC4 bicarbonate transporters. *The Journal of biological chemistry*. 2007; 282:1409–1421. [PubMed: 17090540]
564. Planelles G, Thomas SR, Anagnostopoulos T. Change of apparent stoichiometry of proximal-tubule Na(+)-HCO₃⁻ cotransport upon experimental reversal of its orientation. *Proc Natl Acad Sci U S A*. 1993; 90:7406–7410. [PubMed: 8346263]
565. Pohl M, Kaminski H, Castrop H, Bader M, Himmerkus N, Bleich M, Bachmann S, Theilig F. Intrarenal renin angiotensin system revisited: role of megalin-dependent endocytosis along the proximal nephron. *J Biol Chem*. 285:41935–41946. [PubMed: 20966072]
566. Pollock AS, Warnock DG, Strewler GJ. Parathyroid hormone inhibition of Na⁺-H⁺ antiporter activity in a cultured renal cell line. *Am J Physiol*. 1986; 250:F217–225. [PubMed: 2418691]
567. Pradervand S, Vandewalle A, Bens M, Gautschi I, Loffing J, Hummler E, Schild L, Rossier BC. Dysfunction of the epithelial sodium channel expressed in the kidney of a mouse model for Liddle syndrome. *J Am Soc Nephrol*. 2003; 14:2219–2228. [PubMed: 12937297]
568. Preisig PA. The acid-activated signaling pathway: starting with Pyk2 and ending with increased NHE3 activity. *Kidney international*. 2007; 72:1324–1329. [PubMed: 17882150]
569. Preisig PA. Luminal flow rate regulates proximal tubule H-HCO₃ transporters. *Am J Physiol*. 1992; 262:F47–54. [PubMed: 1310229]
570. Preisig PA, Alpern RJ. Chronic metabolic acidosis causes an adaptation in the apical membrane Na/H antiporter and basolateral membrane Na(HCO₃)₃ symporter in the rat proximal convoluted tubule. *J Clin Invest*. 1988; 82:1445–1453. [PubMed: 2844858]
571. Preisig PA, Alpern RJ. Contributions of cellular leak pathways to net NaHCO₃ and NaCl absorption. *J Clin Invest*. 1989; 83:1859–1867. [PubMed: 2542374]
572. Preisig PA, Alpern RJ. Increased Na/H antiporter and Na/3HCO₃ symporter activities in chronic hyperfiltration. A model of cell hypertrophy. *J Gen Physiol*. 1991; 97:195–217. [PubMed: 1849958]
573. Preisig PA, Ives HE, Cragoe EJ Jr. Alpern RJ, Rector FC Jr. Role of the Na⁺/H⁺ antiporter in rat proximal tubule bicarbonate absorption. *J Clin Invest*. 1987; 80:970–978. [PubMed: 2888788]
574. Preisig PA, Rector FC Jr. Role of Na⁺-H⁺ antiport in rat proximal tubule NaCl absorption. *Am J Physiol*. 1988; 255:F461–465. [PubMed: 2843052]
575. Purkerson JM, Schwartz GJ. The role of carbonic anhydrases in renal physiology. *Kidney Int*. 2007; 71:103–115. [PubMed: 17164835]
576. Purkerson JM, Tsuruoka S, Suter DZ, Nakamori A, Schwartz GJ. Adaptation to metabolic acidosis and its recovery are associated with changes in anion exchanger distribution and expression in the cortical collecting duct. *Kidney Int*. 78:993–1005. [PubMed: 20592712]
577. Puschett JB, Zurbach P, Sylk D. Acute effects of parathyroid hormone on proximal bicarbonate transport in the dog. *Kidney Int*. 1976; 9:501–510. [PubMed: 940283]
578. Pushkin A, Abuladze N, Gross E, Newman D, Tatishchev S, Lee I, Fedotoff O, Bondar G, Azimov R, Ngyuen M, Kurtz I. Molecular mechanism of kNBC1-carbonic anhydrase II interaction in proximal tubule cells. *The Journal of physiology*. 2004; 559:55–65. [PubMed: 15218065]
579. Pushkin A, Abuladze N, Newman D, Muronets V, Sassani P, Tatishchev S, Kurtz I. The COOH termini of NBC3 and the 56-kDa H⁺-ATPase subunit are PDZ motifs involved in their interaction. *American journal of physiology Cell physiology*. 2003; 284:C667–673. [PubMed: 12444018]
580. Quan A, Baum M. The renal nerve is required for regulation of proximal tubule transport by intraluminally produced ANG II. *Am J Physiol Renal Physiol*. 2001; 280:F524–529. [PubMed: 11181415]

581. Quan A, Baum M. Renal nerve stimulation augments effect of intraluminal angiotensin II on proximal tubule transport. *Am J Physiol Renal Physiol.* 2002; 282:F1043–1048. [PubMed: 11997320]
582. Quentin F, Chambrey R, Trinh-Trang-Tan MM, Fysekidis M, Cambillau M, Paillard M, Aronson PS, Eladari D. The Cl⁻/HCO₃⁻ exchanger pendrin in the rat kidney is regulated in response to chronic alterations in chloride balance. *Am J Physiol Renal Physiol.* 2004; 287:F1179–1188. [PubMed: 15292050]
583. Quentin F, Eladari D, Frische S, Cambillau M, Nielsen S, Alper SL, Paillard M, Chambrey R. Regulation of the Cl⁻/HCO₃⁻ exchanger AE2 in rat thick ascending limb of Henle's loop in response to changes in acid-base and sodium balance. *J Am Soc Nephrol.* 2004; 15:2988–2997. [PubMed: 15579501]
584. Rebello G, Ramesar R, Vorster A, Roberts L, Ehrenreich L, Oppon E, Gama D, Bardien S, Greenberg J, Bonapace G, Waheed A, Shah GN, Sly WS. Apoptosis-inducing signal sequence mutation in carbonic anhydrase IV identified in patients with the RP17 form of retinitis pigmentosa. *Proceedings of the National Academy of Sciences of the United States of America.* 2004; 101:6617–6622. [PubMed: 15090652]
585. Rector FC Jr, Carter NW, Seldin DW. The Mechanism of Bicarbonate Reabsorption in the Proximal and Distal Tubules of the Kidney. *J Clin Invest.* 1965; 44:278–290. [PubMed: 14260168]
586. Reenstra WW, Warnock DG, Yee VJ, Forte JG. Proton gradients in renal cortex brush-border membrane vesicles. Demonstration of a rheogenic proton flux with acridine orange. *J Biol Chem.* 1981; 256:11663–11666. [PubMed: 7298622]
587. Reilly AM, Harris PJ, Williams DA. Biphasic effect of angiotensin II on intracellular sodium concentration in rat proximal tubules. *Am J Physiol.* 1995; 269:F374–380. [PubMed: 7573486]
588. Relman AS. Endogenous production of fixed acid and the measurement of the net balance of acid in normal subjects. *Journal of the American Society of Nephrology : JASN.* 2000; 11:2155–2164. [PubMed: 11053495]
589. Relman AS, Lennon EJ, Lemann J Jr. Endogenous production of fixed acid and the measurement of the net balance of acid in normal subjects. *J Clin Invest.* 1961; 40:1621–1630. [PubMed: 13740723]
590. Renkema KY, Velic A, Dijkman HB, Verkaart S, van der Kemp AW, Nowik M, Timmermans K, Doucet A, Wagner CA, Bindels RJ, Hoenderop JG. The calcium-sensing receptor promotes urinary acidification to prevent nephrolithiasis. *J Am Soc Nephrol.* 2009; 20:1705–1713. [PubMed: 19470676]
591. Richardson RM, Kunau RT Jr. Bicarbonate reabsorption in the papillary collecting duct: effect of acetazolamide. *The American journal of physiology.* 1982; 243:F74–80. [PubMed: 6807106]
592. Ridderstrale Y, Kashgarian M, Koeppen B, Giebisch G, Stetson D, Ardito T, Stanton B. Morphological heterogeneity of the rabbit collecting duct. *Kidney Int.* 1988; 34:655–670. [PubMed: 2462075]
593. Riquier-Brison AD, Leong PK, Pihakaski-Maunsbach K, McDonough AA. Angiotensin II stimulates trafficking of NHE3, NaPi2, and associated proteins into the proximal tubule microvilli. *Am J Physiol Renal Physiol.* 298:F177–186. [PubMed: 19864301]
594. Robey RB, Ruiz OS, Espiritu DJ, Ibanez VC, Kear FT, Noboa OA, Bernardo AA, Arruda JA. Angiotensin II stimulation of renal epithelial cell Na/HCO₃ cotransport activity: a central role for Src family kinase/classic MAPK pathway coupling. *J Membr Biol.* 2002; 187:135–145. [PubMed: 12029370]
595. Romero MF, Hediger MA, Boulpaep EL, Boron WF. Expression cloning and characterization of a renal electrogenic Na⁺/HCO₃⁻ cotransporter. *Nature.* 1997; 387:409–413. [PubMed: 9163427]
596. Rothenberger F, Velic A, Stehberger PA, Kovacikova J, Wagner CA. Angiotensin II stimulates vacuolar H⁺-ATPase activity in renal acid-secreting intercalated cells from the outer medullary collecting duct. *J Am Soc Nephrol.* 2007; 18:2085–2093. [PubMed: 17561490]
597. Royaux IE, Wall SM, Karniski LP, Everett LA, Suzuki K, Knepper MA, Green ED. Pendrin, encoded by the Pendred syndrome gene, resides in the apical region of renal intercalated cells

- and mediates bicarbonate secretion. *Proc Natl Acad Sci U S A*. 2001; 98:4221–4226. [PubMed: 11274445]
598. Ruiz OS, Arruda JA. Regulation of the renal Na-HCO₃ cotransporter by cAMP and Ca-dependent protein kinases. *Am J Physiol*. 1992; 262:F560–565. [PubMed: 1314505]
 599. Ruiz OS, Qiu YY, Wang LJ, Arruda JA. Regulation of the renal Na-HCO₃ cotransporter: IV. Mechanisms of the stimulatory effect of angiotensin II. *J Am Soc Nephrol*. 1995; 6:1202–1208. [PubMed: 8589287]
 600. Saari S, Hilvo M, Pan P, Gros G, Hanke N, Waheed A, Sly WS, Parkkila S. The most recently discovered carbonic anhydrase, CA XV, is expressed in the thick ascending limb of Henle and in the collecting ducts of mouse kidney. *PLoS One*. 5:e9624. [PubMed: 20224780]
 601. Sabatini S, Laski ME, Kurtzman NA. NEM-sensitive ATPase activity in rat nephron: effect of metabolic acidosis and alkalosis. *The American journal of physiology*. 1990; 258:F297–304. [PubMed: 2137983]
 602. Sabolic I, Burckhardt G. Proton pathways in rat renal brush-border and basolateral membranes. *Biochim Biophys Acta*. 1983; 734:210–220. [PubMed: 6311264]
 603. Saccomani G, Mitchell KD, Navar LG. Angiotensin II stimulation of Na(+)-H+ exchange in proximal tubule cells. *Am J Physiol*. 1990; 258:F1188–1195. [PubMed: 2159715]
 604. Sakamoto H, Sado Y, Naito I, Kwon TH, Inoue S, Endo K, Kawasaki M, Uchida S, Nielsen S, Sasaki S, Marumo F. Cellular and subcellular immunolocalization of ClC-5 channel in mouse kidney: colocalization with H+-ATPase. *The American journal of physiology*. 1999; 277:F957–965. [PubMed: 10600943]
 605. Saleh AM, Rudnick H, Aronson PS. Mechanism of H(+)-coupled formate transport in rabbit renal microvillus membranes. *Am J Physiol*. 1996; 271:F401–407. [PubMed: 8770172]
 606. Sarker R, Gronborg M, Cha B, Mohan S, Chen Y, Pandey A, Litchfield D, Donowitz M, Li X. Casein kinase 2 binds to the C terminus of Na+/H+ exchanger 3 (NHE3) and stimulates NHE3 basal activity by phosphorylating a separate site in NHE3. *Mol Biol Cell*. 2008; 19:3859–3870. [PubMed: 18614797]
 607. Sasaki S, Berry CA. Mechanism of bicarbonate exit across basolateral membrane of the rabbit proximal convoluted tubule. *Am J Physiol*. 1984; 246:F889–896. [PubMed: 6742135]
 608. Sasaki S, Berry CA, Rector FC Jr. Effect of luminal and peritubular HCO₃(-) concentrations and PCO₂ on HCO₃(-) reabsorption in rabbit proximal convoluted tubules perfused in vitro. *J Clin Invest*. 1982; 70:639–649. [PubMed: 6809795]
 609. Sasaki S, Marumo F. Mechanisms of inhibition of proximal acidification by PTH. *The American journal of physiology*. 1991; 260:F833–838. [PubMed: 1647688]
 610. Sasaki S, Shigai T, Takeuchi J. Intracellular pH in the isolated perfused rabbit proximal straight tubule. *Am J Physiol*. 1985; 249:F417–423. [PubMed: 4037093]
 611. Sasaki S, Shiigai T, Yoshiyama N, Takeuchi J. Mechanism of bicarbonate exit across basolateral membrane of rabbit proximal straight tubule. *Am J Physiol*. 1987; 252:F11–18. [PubMed: 3812695]
 612. Sasaki S, Siragy HM, Gildea JJ, Felder RA, Carey RM. Production and role of extracellular guanosine cyclic 3', 5' monophosphate in sodium uptake in human proximal tubule cells. *Hypertension*. 2004; 43:286–291. [PubMed: 14718358]
 613. Sasaki S, Yoshiyama N. Interaction of chloride and bicarbonate transport across the basolateral membrane of rabbit proximal straight tubule. Evidence for sodium coupled chloride/bicarbonate exchange. *J Clin Invest*. 1988; 81:1004–1011. [PubMed: 2450891]
 614. Satlin LM, Schwartz GJ. Cellular remodeling of HCO₃(-)-secreting cells in rabbit renal collecting duct in response to an acidic environment. *J Cell Biol*. 1989; 109:1279–1288. [PubMed: 2549077]
 615. Schiess WA, Ayer JL, Lotspeich WD, Pitts RF, Miner P. The Renal Regulation of Acid-Base Balance in Man. II. Factors Affecting the Excretion of Titratable Acid by the Normal Human Subject. *J Clin Invest*. 1948; 27:57–64.
 616. Schild L, Aronson PS, Giebisch G. Effects of apical membrane Cl(-)-formate exchange on cell volume in rabbit proximal tubule. *Am J Physiol*. 1990; 258:F530–536. [PubMed: 2316663]

617. Schild L, Giebisch G, Karniski L, Aronson PS. Chloride transport in the mammalian proximal tubule. *Pflugers Arch.* 1986; 407(Suppl 2):S156–159. [PubMed: 3822761]
618. Schild L, Giebisch G, Karniski LP, Aronson PS. Effect of formate on volume reabsorption in the rabbit proximal tubule. *J Clin Invest.* 1987; 79:32–38. [PubMed: 3793930]
619. Schlingmann KP, Konrad M, Jeck N, Waldegger P, Reinalter SC, Holder M, Seyberth HW, Waldegger S. Salt wasting and deafness resulting from mutations in two chloride channels. *N Engl J Med.* 2004; 350:1314–1319. [PubMed: 15044642]
620. Schmitt BM, Biemesderfer D, Romero MF, Boulpaep EL, Boron WF. Immunolocalization of the electrogenic Na⁺-HCO₃⁻ cotransporter in mammalian and amphibian kidney. *Am J Physiol.* 1999; 276:F27–38. [PubMed: 9887077]
621. Schofield AE, Martin PG, Spillett D, Tanner MJ. The structure of the human red blood cell anion exchanger (EPB3, AE1, band 3) gene. *Blood.* 1994; 84:2000–2012. [PubMed: 8081002]
622. Schoolwerth AC, Gesek FA. Intramitochondrial pH and ammonium production in rat and dog kidney cortex. *Miner Electrolyte Metab.* 1990; 16:264–269. [PubMed: 2283989]
623. Schueler C, Becker HM, McKenna R, Deitmer JW. Transport activity of the sodium bicarbonate cotransporter NBCe1 is enhanced by different isoforms of carbonic anhydrase. *PLoS One.* 2011; 6:e27167. [PubMed: 22076132]
624. Schultheis PJ, Clarke LL, Meneton P, Miller ML, Soleimani M, Gawenis LR, Riddle TM, Duffy JJ, Doetschman T, Wang T, Giebisch G, Aronson PS, Lorenz JN, Shull GE. Renal and intestinal absorptive defects in mice lacking the NHE3 Na⁺/H⁺ exchanger. *Nat Genet.* 1998; 19:282–285. [PubMed: 9662405]
625. Schuster VL. Cyclic adenosine monophosphate-stimulated anion transport in rabbit cortical collecting duct. Kinetics, stoichiometry, and conductive pathways. *J Clin Invest.* 1986; 78:1621–1630. [PubMed: 3023449]
626. Schuster VL. Cyclic adenosine monophosphate-stimulated bicarbonate secretion in rabbit cortical collecting tubules. *J Clin Invest.* 1985; 75:2056–2064. [PubMed: 2989340]
627. Schuster VL, Bonsib SM, Jennings ML. Two types of collecting duct mitochondria-rich (intercalated) cells: lectin and band 3 cytochemistry. *Am J Physiol.* 1986; 251:C347–355. [PubMed: 3529979]
628. Schuster VL, Fejes-Toth G, Naray-Fejes-Toth A, Gluck S. Colocalization of H⁺-ATPase and band 3 anion exchanger in rabbit collecting duct intercalated cells. *Am J Physiol.* 1991; 260:F506–517. [PubMed: 1849362]
629. Schuster VL, Kokko JP, Jacobson HR. Angiotensin II directly stimulates sodium transport in rabbit proximal convoluted tubules. *The Journal of clinical investigation.* 1984; 73:507–515. [PubMed: 6699174]
630. Schwaderer AL, Vijayakumar S, Al-Awqati Q, Schwartz GJ. Galectin-3 expression is induced in renal beta-intercalated cells during metabolic acidosis. *Am J Physiol Renal Physiol.* 2006; 290:F148–158. [PubMed: 16131647]
631. Schwartz GJ. Absence of Cl⁻ –OH⁻ or Cl⁻ –HCO₃⁻ exchange in the rabbit renal proximal tubule. *Am J Physiol.* 1983; 245:F462–469. [PubMed: 6624909]
632. Schwartz GJ. Na⁺-dependent H⁺ efflux from proximal tubule: evidence for reversible Na⁺-H⁺ exchange. *Am J Physiol.* 1981; 241:F380–385. [PubMed: 7315962]
633. Schwartz GJ. Physiology and molecular biology of renal carbonic anhydrase. *J Nephrol.* 2002; 15(Suppl 5):S61–74. [PubMed: 12027223]
634. Schwartz GJ, Al-Awqati Q. Carbon dioxide causes exocytosis of vesicles containing H⁺ pumps in isolated perfused proximal and collecting tubules. *J Clin Invest.* 1985; 75:1638–1644. [PubMed: 2582000]
635. Schwartz GJ, Al-Awqati Q. Role of hensen in mediating the adaptation of the cortical collecting duct to metabolic acidosis. *Curr Opin Nephrol Hypertens.* 2005; 14:383–388. [PubMed: 15931009]
636. Schwartz GJ, Barasch J, Al-Awqati Q. Plasticity of functional epithelial polarity. *Nature.* 1985; 318:368–371. [PubMed: 2415824]

637. Schwartz GJ, Kittelberger AM, Barnhart DA, Vijayakumar S. Carbonic anhydrase IV is expressed in H(+)-secreting cells of rabbit kidney. *Am J Physiol Renal Physiol.* 2000; 278:F894–904. [PubMed: 10836977]
638. Schwartz GJ, Kittelberger AM, Watkins RH, O'Reilly MA. Carbonic anhydrase XII mRNA encodes a hydratase that is differentially expressed along the rabbit nephron. *American journal of physiology Renal physiology.* 2003; 284:F399–410. [PubMed: 12388401]
639. Schwartz GJ, Tsuruoka S, Vijayakumar S, Petrovic S, Mian A, Al-Awqati Q. Acid incubation reverses the polarity of intercalated cell transporters, an effect mediated by hensin. *The Journal of clinical investigation.* 2002; 109:89–99. [PubMed: 11781354]
640. Sebastian A, Harris ST, Ottaway JH, Todd KM, Morris RC Jr. Improved mineral balance and skeletal metabolism in postmenopausal women treated with potassium bicarbonate. *N Engl J Med.* 1994; 330:1776–1781. [PubMed: 8190153]
641. Seifter JL, Knickelbein R, Aronson PS. Absence of Cl-OH exchange and NaCl cotransport in rabbit renal microvillus membrane vesicles. *Am J Physiol.* 1984; 247:F753–759. [PubMed: 6093589]
642. Seikaly MG, Arant BS Jr, Seney FD Jr. Endogenous angiotensin concentrations in specific intrarenal fluid compartments of the rat. *J Clin Invest.* 1990; 86:1352–1357. [PubMed: 2212017]
643. Seshadri RM, Klein JD, Kozlowski S, Sands JM, Kim YH, Han KH, Handlogten ME, Verlander JW, Weiner ID. Renal expression of the ammonia transporters, Rhbg and Rhcg, in response to chronic metabolic acidosis. *Am J Physiol Renal Physiol.* 2006; 290:F397–408. [PubMed: 16144966]
644. Seshadri RM, Klein JD, Smith T, Sands JM, Handlogten ME, Verlander JW, Weiner ID. Changes in subcellular distribution of the ammonia transporter, Rhcg, in response to chronic metabolic acidosis. *Am J Physiol Renal Physiol.* 2006; 290:F1443–1452. [PubMed: 16434569]
645. Shah GN, Bonapace G, Hu PY, Strisciuglio P, Sly WS. Carbonic anhydrase II deficiency syndrome (osteopetrosis with renal tubular acidosis and brain calcification): novel mutations in CA2 identified by direct sequencing expand the opportunity for genotype-phenotype correlation. *Hum Mutat.* 2004; 24:272. [PubMed: 15300855]
646. Shao J, Gumz ML, Cain BD, Xia SL, Shull GE, van Driel IR, Wingo CS. Pharmacological profiles of the murine gastric and colonic H,K-ATPases. *Biochim Biophys Acta.* 1800:906–911. [PubMed: 20594946]
647. Shenolikar S, Voltz JW, Minkoff CM, Wade JB, Weinman EJ. Targeted disruption of the mouse NHERF-1 gene promotes internalization of proximal tubule sodium-phosphate cotransporter type IIa and renal phosphate wasting. *Proc Natl Acad Sci U S A.* 2002; 99:11470–11475. [PubMed: 12169661]
648. Sheu JN, Quigley R, Baum M. Heterogeneity of chloride/base exchange in rabbit superficial and juxtamedullary proximal convoluted tubules. *Am J Physiol.* 1995; 268:F847–853. [PubMed: 7771513]
649. Shindo T, Spring KR. Chloride movement across the basolateral membrane of proximal tubule cells. *J Membr Biol.* 1981; 58:35–42. [PubMed: 7218331]
650. Shiuan D, Weinstein SW. Evidence for electroneutral chloride transport in rabbit renal cortical brush border membrane vesicles. *The American journal of physiology.* 1984; 247:F837–847. [PubMed: 6093593]
651. Shum WW, Da Silva N, Belleannee C, McKee M, Brown D, Breton S. Regulation of V-ATPase recycling via a RhoA- and ROCKII-dependent pathway in epididymal clear cells. *Am J Physiol Cell Physiol.* 301:C31–43. [PubMed: 21411727]
652. Siebens AW, Boron WF. Effect of electroneutral luminal and basolateral lactate transport on intracellular pH in salamander proximal tubules. *J Gen Physiol.* 1987; 90:799–831. [PubMed: 3440860]
653. Silver RB, Choe H, Frindt G. Low-NaCl diet increases H-K-ATPase in intercalated cells from rat cortical collecting duct. *Am J Physiol.* 1998; 275:F94–102. [PubMed: 9689010]
654. Silver RB, Frindt G, Mennitt P, Satlin LM. Characterization and regulation of H-K-ATPase in intercalated cells of rabbit cortical collecting duct. *J Exp Zool.* 1997; 279:443–455. [PubMed: 9392865]

655. Simon DB, Bindra RS, Mansfield TA, Nelson-Williams C, Mendonca E, Stone R, Schurman S, Nayir A, Alpay H, Bakkaloglu A, Rodriguez-Soriano J, Morales JM, Sanjad SA, Taylor CM, Pilz D, Brem A, Trachtman H, Griswold W, Richard GA, John E, Lifton RP. Mutations in the chloride channel gene, CLCNKB, cause Bartter's syndrome type III. *Nat Genet.* 1997; 17:171–178. [PubMed: 9326936]
656. Simpson DP. Citrate excretion: a window on renal metabolism. *The American journal of physiology.* 1983; 244:F223–234. [PubMed: 6338740]
657. Simpson JE, Schweinfest CW, Shull GE, Gawenis LR, Walker NM, Boyle KT, Soleimani M, Clarke LL. PAT-1 (Slc26a6) is the predominant apical membrane Cl[−]/HCO₃[−] exchanger in the upper villous epithelium of the murine duodenum. *Am J Physiol Gastrointest Liver Physiol.* 2007; 292:G1079–1088. [PubMed: 17170027]
658. Singh V, Lin R, Yang J, Cha B, Sarker R, Tse CM, Donowitz M. AKT and GSK-3 are Necessary for Direct Ezrin Binding to NHE3 as Part of a C-Terminal Stimulatory Complex: Role of a Novel Ser-Rich NHE3 C-terminal Motif in NHE3 Activity and Trafficking. *J Biol Chem.*
659. Sly WS, Whyte MP, Sundaram V, Tashian RE, Hewett-Emmett D, Guibaud P, Vainsel M, Baluarte HJ, Gruskin A, Al-Mosawi M, et al. Carbonic anhydrase II deficiency in 12 families with the autosomal recessive syndrome of osteopetrosis with renal tubular acidosis and cerebral calcification. *N Engl J Med.* 1985; 313:139–145. [PubMed: 3925334]
660. Smith AN, Skaug J, Choate KA, Nayir A, Bakkaloglu A, Ozen S, Hulton SA, Sanjad SA, Al-Sabban EA, Lifton RP, Scherer SW, Karet FE. Mutations in ATP6N1B, encoding a new kidney vacuolar proton pump 116-kD subunit, cause recessive distal renal tubular acidosis with preserved hearing. *Nat Genet.* 2000; 26:71–75. [PubMed: 10973252]
661. Soleimani M, Aronson PS. Effects of acetazolamide on Na⁺-HCO₃[−] cotransport in basolateral membrane vesicles isolated from rabbit renal cortex. *J Clin Invest.* 1989; 83:945–951. [PubMed: 2921327]
662. Soleimani M, Aronson PS. Ionic mechanism of Na⁺-HCO₃[−] cotransport in rabbit renal basolateral membrane vesicles. *J Biol Chem.* 1989; 264:18302–18308. [PubMed: 2509453]
663. Sonalker PA, Tofovic SP, Bastacky SI, Jackson EK. Chronic noradrenaline increases renal expression of NHE-3, NBC-1, BSC-1 and aquaporin-2. *Clin Exp Pharmacol Physiol.* 2008; 35:594–600. [PubMed: 18177483]
664. Sonalker PA, Tofovic SP, Jackson EK. Increased expression of the sodium transporter BSC-1 in spontaneously hypertensive rats. *J Pharmacol Exp Ther.* 2004; 311:1052–1061. [PubMed: 15340004]
665. Spicer SS, Sens MA, Tashian RE. Immunocytochemical demonstration of carbonic anhydrase in human epithelial cells. *The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society.* 1982; 30:864–873. [PubMed: 6813372]
666. Spicer Z, Miller ML, Andringa A, Riddle TM, Duffy JJ, Doetschman T, Shull GE. Stomachs of mice lacking the gastric H,K-ATPase alpha-subunit have achlorhydria, abnormal parietal cells, and ciliated metaplasia. *J Biol Chem.* 2000; 275:21555–21565. [PubMed: 10764766]
667. Stanton BA. Characterization of apical and basolateral membrane conductances of rat inner medullary collecting duct. *Am J Physiol.* 1989; 256:F862–868. [PubMed: 2719119]
668. Star RA. Basolateral membrane sodium-independent Cl[−]/HCO₃[−] exchanger in rat inner medullary collecting duct cell. *J Clin Invest.* 1990; 85:1959–1966. [PubMed: 2347919]
669. Star RA, Burg MB, Knepper MA. Bicarbonate secretion and chloride absorption by rabbit cortical collecting ducts. Role of chloride/bicarbonate exchange. *J Clin Invest.* 1985; 76:1123–1130. [PubMed: 3930570]
670. Star RA, Burg MB, Knepper MA. Luminal disequilibrium pH and ammonia transport in outer medullary collecting duct [corrected and issued with original paging in *Am J Physiol* 1987 Aug; 253(2 Pt 2)]. *Am J Physiol.* 1987; 252:F1148–1157. [PubMed: 3109255]
671. Star RA, Kurtz I, Mejia R, Burg MB, Knepper MA. Disequilibrium pH and ammonia transport in isolated perfused cortical collecting ducts. *The American journal of physiology.* 1987; 253:F1232–1242. [PubMed: 3122587]
672. Stetson DL, Wade JB, Giebisch G. Morphologic alterations in the rat medullary collecting duct following potassium depletion. *Kidney Int.* 1980; 17:45–56. [PubMed: 7374020]

673. Stettner P, Bourgeois S, Marsching C, Traykova-Brauch M, Porubsky S, Nordstrom V, Hopf C, Koesters R, Sandhoff R, Wiegandt H, Wagner CA, Grone HJ, Jennemann R. Sulfatides are required for renal adaptation to chronic metabolic acidosis. *Proceedings of the National Academy of Sciences of the United States of America*. 2013; 110:9998–10003. [PubMed: 23716689]
674. Stone DK, Seldin DW, Kokko JP, Jacobson HR. Mineralocorticoid modulation of rabbit medullary collecting duct acidification. A sodium-independent effect. *J Clin Invest*. 1983; 72:77–83. [PubMed: 6874954]
675. Stover EH, Borthwick KJ, Bavalia C, Eady N, Fritz DM, Rungroj N, Giersch AB, Morton CC, Axon PR, Akil I, Al-Sabban EA, Baguley DM, Bianca S, Bakkaloglu A, Bircan Z, Chauveau D, Clermont MJ, Guala A, Hulton SA, Kroes H, Li Volti G, Mir S, Mocan H, Nayir A, Ozen S, Rodriguez Soriano J, Sanjad SA, Tasic V, Taylor CM, Topaloglu R, Smith AN, Karet FE. Novel ATP6V1B1 and ATP6V0A4 mutations in autosomal recessive distal renal tubular acidosis with new evidence for hearing loss. *J Med Genet*. 2002; 39:796–803. [PubMed: 12414817]
676. Stuart-Tilley AK, Shmukler BE, Brown D, Alper SL. Immunolocalization and tissue-specific splicing of AE2 anion exchanger in mouse kidney. *J Am Soc Nephrol*. 1998; 9:946–959. [PubMed: 9621277]
677. Su Y, Blake-Palmer KG, Sorrell S, Javid B, Bowers K, Zhou A, Chang SH, Qamar S, Karet FE. Human H⁺ATPase α 4 subunit mutations causing renal tubular acidosis reveal a role for interaction with phosphofructokinase-1. *Am J Physiol Renal Physiol*. 2008; 295:F950–958. [PubMed: 18632794]
678. Sun AM, Liu Y, Dworkin LD, Tse CM, Donowitz M, Yip KP. Na⁺/H⁺ exchanger isoform 2 (NHE2) is expressed in the apical membrane of the medullary thick ascending limb. *The Journal of membrane biology*. 1997; 160:85–90. [PubMed: 9351894]
679. Sun X, Soleimani M, Petrovic S. Decreased expression of Slc26a4 (Pendrin) and Slc26a7 in the kidneys of carbonic anhydrase II-deficient mice. *Cell Physiol Biochem*. 2008; 21:95–108. [PubMed: 18209476]
680. Suzuki M, Van Paesschen W, Stalmans I, Horita S, Yamada H, Bergmans BA, Legius E, Riant F, De Jonghe P, Li Y, Sekine T, Igarashi T, Fujimoto I, Mikoshiba K, Shimadzu M, Shiohara M, Braverman N, Al-Gazali L, Fujita T, Seki G. Defective membrane expression of the Na⁺-HCO₃⁻ cotransporter NBCe1 is associated with familial migraine. *Proc Natl Acad Sci U S A*. 2007; 104:15963–15968. [PubMed: 17800355]
681. Szaszi K, Kurashima K, Kaibuchi K, Grinstein S, Orlowski J. Role of the cytoskeleton in mediating cAMP-dependent protein kinase inhibition of the epithelial Na⁺/H⁺ exchanger NHE3. *J Biol Chem*. 2001; 276:40761–40768. [PubMed: 11522794]
682. Tago K, Schuster VL, Stokes JB. Regulation of chloride self exchange by cAMP in cortical collecting tubule. *Am J Physiol*. 1986; 251:F40–48. [PubMed: 3014899]
683. Tago K, Schuster VL, Stokes JB. Stimulation of chloride transport by HCO₃-CO₂ in rabbit cortical collecting tubule. *Am J Physiol*. 1986; 251:F49–56. [PubMed: 3089028]
684. Takito J, Hikita C, Al-Awqati Q. Hensin, a new collecting duct protein involved in the in vitro plasticity of intercalated cell polarity. *J Clin Invest*. 1996; 98:2324–2331. [PubMed: 8941650]
685. Teng-umnuay P, Verlander JW, Yuan W, Tisher CC, Madsen KM. Identification of distinct subpopulations of intercalated cells in the mouse collecting duct. *Journal of the American Society of Nephrology : JASN*. 1996; 7:260–274. [PubMed: 8785396]
686. Tizianello A, Deferrari G, Garibotto G, Robaudo C, Acquarone N, Ghiggeri GM. Renal ammoniogenesis in an early stage of metabolic acidosis in man. *J Clin Invest*. 1982; 69:240–250. [PubMed: 7054241]
687. Tizianello A, Deferrari G, Garibotto G, Robaudo C, Bruzzzone M, Passerone GC. Renal ammoniogenesis during the adaptation to metabolic acidosis in man. *Contrib Nephrol*. 1982; 31:40–46. [PubMed: 7105750]
688. Tokonami N, Morla L, Centeno G, Mordasini D, Ramakrishnan SK, Nikolaeva S, Wagner CA, Bonny O, Houillier P, Doucet A, Firsov D. α -Ketoglutarate regulates acid-base balance through an intrarenal paracrine mechanism. *J Clin Invest*. 2005; 115:3166–3171. [PubMed: 16134124]

689. Tomita K, Pisano JJ, Burg MB, Knepper MA. Effects of vasopressin and bradykinin on anion transport by the rat cortical collecting duct. Evidence for an electroneutral sodium chloride transport pathway. *J Clin Invest.* 1986; 77:136–141. [PubMed: 3080471]
690. Tsai CJ, Ives HE, Alpern RJ, Yee VJ, Warnock DG, Rector FC Jr. Increased V_{max} for Na^+/H^+ -antiporter activity in proximal tubule brush border vesicles from rabbits with metabolic acidosis. *Am J Physiol.* 1984; 247:F339–343. [PubMed: 6087681]
691. Tse CM, Brant SR, Walker MS, Pouyssegur J, Donowitz M. Cloning and sequencing of a rabbit cDNA encoding an intestinal and kidney-specific Na^+/H^+ exchanger isoform (NHE-3). *J Biol Chem.* 1992; 267:9340–9346. [PubMed: 1374392]
692. Tse CM, Levine SA, Yun CH, Brant SR, Pouyssegur J, Montrose MH, Donowitz M. Functional characteristics of a cloned epithelial Na^+/H^+ exchanger (NHE3): resistance to amiloride and inhibition by protein kinase C. *Proc Natl Acad Sci U S A.* 1993; 90:9110–9114. [PubMed: 8415663]
693. Tsuganezawa H, Kobayashi K, Iyori M, Araki T, Koizumi A, Watanabe S, Kaneko A, Fukao T, Monkawa T, Yoshida T, Kim DK, Kanai Y, Endou H, Hayashi M, Saruta T. A new member of the HCO_3^- transporter superfamily is an apical anion exchanger of beta-intercalated cells in the kidney. *J Biol Chem.* 2001; 276:8180–8189. [PubMed: 11102437]
694. Tsuruoka S, Kittelberger AM, Schwartz GJ. Carbonic anhydrase II and IV mRNA in rabbit nephron segments: stimulation during metabolic acidosis. *The American journal of physiology.* 1998; 274:F259–267. [PubMed: 9486220]
695. Tsuruoka S, Schwartz GJ. Adaptation of rabbit cortical collecting duct HCO_3^- transport to metabolic acidosis in vitro. *J Clin Invest.* 1996; 97:1076–1084. [PubMed: 8613531]
696. Tsuruoka S, Schwartz GJ. Mechanisms of HCO_3^- secretion in the rabbit connecting segment. *Am J Physiol.* 1999; 277:F567–574. [PubMed: 10516281]
697. Tsuruoka S, Schwartz GJ. Metabolic acidosis stimulates H^+ secretion in the rabbit outer medullary collecting duct (inner stripe) of the kidney. *J Clin Invest.* 1997; 99:1420–1431. [PubMed: 9077552]
698. Tsuruoka S, Swenson ER, Petrovic S, Fujimura A, Schwartz GJ. Role of basolateral carbonic anhydrase in proximal tubular fluid and bicarbonate absorption. *Am J Physiol Renal Physiol.* 2001; 280:F146–154. [PubMed: 11133524]
699. Tsuruoka S, Watanabe S, Purkerson JM, Fujimura A, Schwartz GJ. Endothelin and nitric oxide mediate adaptation of the cortical collecting duct to metabolic acidosis. *Am J Physiol Renal Physiol.* 2006; 291:F866–873. [PubMed: 16705153]
700. Turban S, Beutler KT, Morris RG, Masilamani S, Fenton RA, Knepper MA, Packer RK. Long-term regulation of proximal tubule acid-base transporter abundance by angiotensin II. *Kidney Int.* 2006; 70:660–668. [PubMed: 16807546]
701. Twombly K, Gattineni J, Bobulescu IA, Dwarakanath V, Baum M. Effect of metabolic acidosis on neonatal proximal tubule acidification. *Am J Physiol Regul Integr Comp Physiol.* 299:R1360–1368. [PubMed: 20720175]
702. Ullrich KJ, Papavassiliou F. Bicarbonate reabsorption in the papillary collecting duct of rats. *Pflugers Archiv : European journal of physiology.* 1981; 389:271–275. [PubMed: 6453328]
703. Ullrich KJ, Rumrich G, Baumann K. Renal proximal tubular buffer-(glycodiazine) transport. Inhomogeneity of local transport rate, dependence on sodium, effect of inhibitors and chronic adaptation. *Pflugers Arch.* 1975; 357:149–163. [PubMed: 127986]
704. Unwin R, Capasso G, Giebisch G. Bicarbonate transport along the loop of Henle effects of adrenal steroids. *The American journal of physiology.* 1995; 268:F234–239. [PubMed: 7864161]
705. van Adelsberg J, Edwards JC, Takito J, Kiss B, al-Awqati Q. An induced extracellular matrix protein reverses the polarity of band 3 in intercalated epithelial cells. *Cell.* 1994; 76:1053–1061. [PubMed: 8137422]
706. Vandewalle A, Cluzeaud F, Bens M, Kieferle S, Steinmeyer K, Jentsch TJ. Localization and induction by dehydration of $ClC-K$ chloride channels in the rat kidney. *Am J Physiol.* 1997; 272:F678–688. [PubMed: 9176380]
707. Vargas-Poussou R, Houillier P, Le Pottier N, Strompf L, Loirat C, Baudouin V, Macher MA, Dechaux M, Ulinski T, Nobili F, Eckart P, Novo R, Cailliez M, Salomon R, Nivet H, Cochat P,

- Tack I, Fargeot A, Bouissou F, Kesler GR, Lorotte S, Godefroid N, Layet V, Morin G, Jeunemaitre X, Blanchard A. Genetic investigation of autosomal recessive distal renal tubular acidosis: evidence for early sensorineural hearing loss associated with mutations in the ATP6V0A4 gene. *J Am Soc Nephrol*. 2006; 17:1437–1443. [PubMed: 16611712]
708. Velazquez H, Silva T. Cloning and localization of KCC4 in rabbit kidney: expression in distal convoluted tubule. *Am J Physiol Renal Physiol*. 2003; 285:F49–58. [PubMed: 12709395]
709. Verlander JW, Hassell KA, Royaux IE, Glapion DM, Wang ME, Everett LA, Green ED, Wall SM. Deoxycorticosterone upregulates PDS (Slc26a4) in mouse kidney: role of pendrin in mineralocorticoid-induced hypertension. *Hypertension*. 2003; 42:356–362. [PubMed: 12925556]
710. Verlander JW, Kim YH, Shin W, Pham TD, Hassell KA, Beierwaltes WH, Green ED, Everett L, Matthews SW, Wall SM. Dietary Cl(–) restriction upregulates pendrin expression within the apical plasma membrane of type B intercalated cells. *Am J Physiol Renal Physiol*. 2006; 291:F833–839. [PubMed: 16670435]
711. Verlander JW, Madsen KM, Galla JH, Luke RG, Tisher CC. Response of intercalated cells to chloride depletion metabolic alkalosis. *Am J Physiol*. 1992; 262:F309–319. [PubMed: 1531735]
712. Verlander JW, Madsen KM, Low PS, Allen DP, Tisher CC. Immunocytochemical localization of band 3 protein in the rat collecting duct. *Am J Physiol*. 1988; 255:F115–125. [PubMed: 3394806]
713. Verlander JW, Madsen KM, Stone DK, Tisher CC. Ultrastructural localization of H⁺ATPase in rabbit cortical collecting duct. *Journal of the American Society of Nephrology : JASN*. 1994; 4:1546–1557. [PubMed: 8025228]
714. Verlander JW, Madsen KM, Tisher CC. Effect of acute respiratory acidosis on two populations of intercalated cells in rat cortical collecting duct. *The American journal of physiology*. 1987; 253:F1142–1156. [PubMed: 3425724]
715. Verlander JW, Madsen KM, Tisher CC. Structural and functional features of proton and bicarbonate transport in the rat collecting duct. *Semin Nephrol*. 1991; 11:465–477. [PubMed: 1658904]
716. Verlander JW, Miller RT, Frank AE, Royaux IE, Kim YH, Weiner ID. Localization of the ammonium transporter proteins RhBG and RhCG in mouse kidney. *Am J Physiol Renal Physiol*. 2003; 284:F323–337. [PubMed: 12388412]
717. Verlander JW, Moudy RM, Campbell WG, Cain BD, Wingo CS. Immunohistochemical localization of H-K-ATPase alpha(2c)-subunit in rabbit kidney. *Am J Physiol Renal Physiol*. 2001; 281:F357–365. [PubMed: 11457728]
718. Vidarsson H, Westergren R, Heglund M, Blomqvist SR, Breton S, Enerback S. The forkhead transcription factor Foxi1 is a master regulator of vacuolar H-ATPase proton pump subunits in the inner ear, kidney and epididymis. *PLoS One*. 2009; 4:e4471. [PubMed: 19214237]
719. Vieira FL, Malnic G. Hydrogen ion secretion by rat renal cortical tubules as studied by an antimony microelectrode. *Am J Physiol*. 1968; 214:710–718. [PubMed: 4966812]
720. Vitavska O, Wieczorek H, Merzendorfer H. A novel role for subunit C in mediating binding of the H⁺-V-ATPase to the actin cytoskeleton. *J Biol Chem*. 2003; 278:18499–18505. [PubMed: 12606563]
721. Vorum H, Kwon TH, Fulton C, Simonsen B, Choi I, Boron W, Maunsbach AB, Nielsen S, Aalkjaer C. Immunolocalization of electroneutral Na-HCO₃(–) cotransporter in rat kidney. *Am J Physiol Renal Physiol*. 2000; 279:F901–909. [PubMed: 11053051]
722. Voss M, Schmidt R, Walz B, Baumann O. Stimulus-induced translocation of the protein kinase A catalytic subunit to the apical membrane in blowfly salivary glands. *Cell and tissue research*. 2009; 335:657–662. [PubMed: 18766382]
723. Wade JB, Liu J, Coleman RA, Cunningham R, Steplock DA, Lee-Kwon W, Pallone TL, Shenolikar S, Weinman EJ. Localization and interaction of NHERF isoforms in the renal proximal tubule of the mouse. *Am J Physiol Cell Physiol*. 2003; 285:C1494–1503. [PubMed: 12917102]
724. Wade JB, Welling PA, Donowitz M, Shenolikar S, Weinman EJ. Differential renal distribution of NHERF isoforms and their colocalization with NHE3, ezrin, and ROMK. *Am J Physiol Cell Physiol*. 2001; 280:C192–198. [PubMed: 11121391]

725. Wagner CA, Finberg KE, Breton S, Marshansky V, Brown D, Geibel JP. Renal vacuolar H⁺-ATPase. *Physiol Rev.* 2004; 84:1263–1314. [PubMed: 15383652]
726. Wagner CA, Finberg KE, Stehberger PA, Lifton RP, Giebisch GH, Aronson PS, Geibel JP. Regulation of the expression of the Cl[−]/anion exchanger pendrin in mouse kidney by acid-base status. *Kidney international.* 2002; 62:2109–2117. [PubMed: 12427135]
727. Wagner CA, Giebisch G, Lang F, Geibel JP. Angiotensin II stimulates vesicular H⁺-ATPase in rat proximal tubular cells. *Proc Natl Acad Sci U S A.* 1998; 95:9665–9668. [PubMed: 9689138]
728. Wagner CA, Mohebbi N, Uhlig U, Giebisch GH, Breton S, Brown D, Geibel JP. Angiotensin II stimulates H⁽⁺⁾-ATPase activity in intercalated cells from isolated mouse connecting tubules and cortical collecting ducts. *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology.* 2011; 28:513–520.
729. Wakabayashi S, Bertrand B, Ikeda T, Pouyssegur J, Shigekawa M. Mutation of calmodulin-binding site renders the Na⁺/H⁺ exchanger (NHE1) highly H⁽⁺⁾-sensitive and Ca²⁺ regulation-defective. *J Biol Chem.* 1994; 269:13710–13715. [PubMed: 8175807]
730. Wakabayashi S, Bertrand B, Shigekawa M, Fafournoux P, Pouyssegur J. Growth factor activation and “H⁽⁺⁾-sensing” of the Na⁺/H⁺ exchanger isoform 1 (NHE1). Evidence for an additional mechanism not requiring direct phosphorylation. *J Biol Chem.* 1994; 269:5583–5588. [PubMed: 8119893]
731. Wakabayashi S, Ikeda T, Noel J, Schmitt B, Orlowski J, Pouyssegur J, Shigekawa M. Cytoplasmic domain of the ubiquitous Na⁺/H⁺ exchanger NHE1 can confer Ca²⁺ responsiveness to the apical isoform NHE3. *J Biol Chem.* 1995; 270:26460–26465. [PubMed: 7592862]
732. Waldegger S, Jeck N, Barth P, Peters M, Vitzthum H, Wolf K, Kurtz A, Konrad M, Seyberth HW. Barttin increases surface expression and changes current properties of ClC-K channels. *Pflugers Arch.* 2002; 444:411–418. [PubMed: 12111250]
733. Wall SM, Fischer MP, Glapion DM, De La Calzada M. ANG II reduces net acid secretion in rat outer medullary collecting duct. *Am J Physiol Renal Physiol.* 2003; 285:F930–937. [PubMed: 12851254]
734. Wall SM, Flessner MF, Knepper MA. Distribution of luminal carbonic anhydrase activity along rat inner medullary collecting duct. *Am J Physiol.* 1991; 260:F738–748. [PubMed: 1903600]
735. Wall SM, Kim YH, Stanley L, Glapion DM, Everett LA, Green ED, Verlander JW. NaCl restriction upregulates renal Slc26a4 through subcellular redistribution: role in Cl[−] conservation. *Hypertension.* 2004; 44:982–987. [PubMed: 15477386]
736. Wall SM, Sands JM, Flessner MF, Nonoguchi H, Spring KR, Knepper MA. Net acid transport by isolated perfused inner medullary collecting ducts. *Am J Physiol.* 1990; 258:F75–84. [PubMed: 2301597]
737. Wall SM, Truong AV, DuBose TD Jr. H⁽⁺⁾-K⁽⁺⁾-ATPase mediates net acid secretion in rat terminal inner medullary collecting duct. *Am J Physiol.* 1996; 271:F1037–1044. [PubMed: 8945998]
738. Wall SM, Weinstein AM. Cortical distal nephron Cl^(−) transport in volume homeostasis and blood pressure regulation. *Am J Physiol Renal Physiol.* 305:F427–438. [PubMed: 23637202]
739. Wang D, Sun H, Lang F, Yun CC. Activation of NHE3 by dexamethasone requires phosphorylation of NHE3 at Ser663 by SGK1. *Am J Physiol Cell Physiol.* 2005; 289:C802–810. [PubMed: 15888551]
740. Wang D, Zhang H, Lang F, Yun CC. Acute activation of NHE3 by dexamethasone correlates with activation of SGK1 and requires a functional glucocorticoid receptor. *Am J Physiol Cell Physiol.* 2007; 292:C396–404. [PubMed: 16971495]
741. Wang G, Li C, Kim SW, Ring T, Wen J, Djurhuus JC, Wang W, Nielsen S, Frokiaer J. Ureter obstruction alters expression of renal acid-base transport proteins in rat kidney. *Am J Physiol Renal Physiol.* 2008; 295:F497–506. [PubMed: 18508879]
742. Wang T. Flow-activated transport events along the nephron. *Curr Opin Nephrol Hypertens.* 2006; 15:530–536. [PubMed: 16914967]
743. Wang T. Nitric oxide regulates HCO₃[−] and Na⁺ transport by a cGMP-mediated mechanism in the kidney proximal tubule. *Am J Physiol.* 1997; 272:F242–248. [PubMed: 9124402]

744. Wang T, Chan YL. Time- and dose-dependent effects of protein kinase C on proximal bicarbonate transport. *J Membr Biol.* 1990; 117:131–139. [PubMed: 2120446]
745. Wang T, Egbert AL Jr, Abbiati T, Aronson PS, Giebisch G. Mechanisms of stimulation of proximal tubule chloride transport by formate and oxalate. *Am J Physiol.* 1996; 271:F446–450. [PubMed: 8770178]
746. Wang T, Giebisch G. Effects of angiotensin II on electrolyte transport in the early and late distal tubule in rat kidney. *Am J Physiol.* 1996; 271:F143–149. [PubMed: 8760255]
747. Wang T, Giebisch G, Aronson PS. Effects of formate and oxalate on volume absorption in rat proximal tubule. *Am J Physiol.* 1992; 263:F37–42. [PubMed: 1636743]
748. Wang T, Hropot M, Aronson PS, Giebisch G. Role of NHE isoforms in mediating bicarbonate reabsorption along the nephron. *American journal of physiology Renal physiology.* 2001; 281:F1117–1122. [PubMed: 11704563]
749. Wang T, Inglis FM, Kalb RG. Defective fluid and HCO_3^- absorption in proximal tubule of neuronal nitric oxide synthase-knockout mice. *Am J Physiol Renal Physiol.* 2000; 279:F518–524. [PubMed: 10966931]
750. Wang T, Malnic G, Giebisch G, Chan YL. Renal bicarbonate reabsorption in the rat. IV. Bicarbonate transport mechanisms in the early and late distal tubule. *J Clin Invest.* 1993; 91:2776–2784. [PubMed: 8390489]
751. Wang T, Yang CL, Abbiati T, Schultheis PJ, Shull GE, Giebisch G, Aronson PS. Mechanism of proximal tubule bicarbonate absorption in NHE3 null mice. *Am J Physiol.* 1999; 277:F298–302. [PubMed: 10444585]
752. Wang T, Yang CL, Abbiati T, Shull GE, Giebisch G, Aronson PS. Essential role of NHE3 in facilitating formate-dependent NaCl absorption in the proximal tubule. *Am J Physiol Renal Physiol.* 2001; 281:F288–292. [PubMed: 11457720]
753. Wang X, Kurtz I. H⁺/base transport in principal cells characterized by confocal fluorescence imaging. *Am J Physiol.* 1990; 259:C365–373. [PubMed: 2382707]
754. Wang Z, Wang T, Petrovic S, Tuo B, Riederer B, Barone S, Lorenz JN, Seidler U, Aronson PS, Soleimani M. Renal and intestinal transport defects in Slc26a6-null mice. *American journal of physiology Cell physiology.* 2005; 288:C957–965. [PubMed: 15574486]
755. Warnock DG, Reenstra WW, Yee VJ. Na⁺/H⁺ antiporter of brush border vesicles: studies with acridine orange uptake. *Am J Physiol.* 1982; 242:F733–739. [PubMed: 6283903]
756. Warnock DG, Yee VJ. Chloride uptake by brush border membrane vesicles isolated from rabbit renal cortex. Coupling to proton gradients and K⁺ diffusion potentials. *The Journal of clinical investigation.* 1981; 67:103–115. [PubMed: 7451645]
757. Warth R, Barriere H, Meneton P, Bloch M, Thomas J, Tauc M, Heitzmann D, Romeo E, Verrey F, Mengual R, Guy N, Bendahhou S, Lesage F, Poujeol P, Barhanin J. Proximal renal tubular acidosis in TASK2 K⁺ channel-deficient mice reveals a mechanism for stabilizing bicarbonate transport. *Proc Natl Acad Sci U S A.* 2004; 101:8215–8220. [PubMed: 15141089]
758. Watts BA 3rd, George T, Good DW. Aldosterone inhibits apical NHE3 and HCO_3^- absorption via a nongenomic ERK-dependent pathway in medullary thick ascending limb. *American journal of physiology Renal physiology.* 2006; 291:F1005–1013. [PubMed: 16757729]
759. Watts BA 3rd, George T, Good DW. The basolateral NHE1 Na⁺/H⁺ exchanger regulates transepithelial HCO_3^- absorption through actin cytoskeleton remodeling in renal thick ascending limb. *The Journal of biological chemistry.* 2005; 280:11439–11447. [PubMed: 15644322]
760. Watts BA 3rd, George T, Good DW. Lumen LPS inhibits HCO_3^- absorption in the medullary thick ascending limb through TLR4-PI3K-Akt-mTOR-dependent inhibition of basolateral Na⁺/H⁺ exchange. *Am J Physiol Renal Physiol.* 305:F451–462. [PubMed: 23698118]
761. Watts BA 3rd, George T, Sherwood ER, Good DW. Basolateral LPS inhibits NHE3 and HCO_3^- absorption through TLR4/MyD88-dependent ERK activation in medullary thick ascending limb. *Am J Physiol Cell Physiol.* 301:C1296–1306. [PubMed: 21881005]
762. Watts BA 3rd, George T, Sherwood ER, Good DW. A two-hit mechanism for sepsis-induced impairment of renal tubule function. *Am J Physiol Renal Physiol.* 304:F863–874. [PubMed: 23324175]

763. Watts BA 3rd, Good DW. An apical K(+)-dependent HCO₃⁻ transport pathway opposes transepithelial HCO₃⁻ absorption in rat medullary thick ascending limb. *Am J Physiol Renal Physiol.* 2004; 287:F57–63. [PubMed: 15026301]
764. Watts BA 3rd, Good DW. Effects of ammonium on intracellular pH in rat medullary thick ascending limb: mechanisms of apical membrane NH₄⁺ transport. *J Gen Physiol.* 1994; 103:917–936. [PubMed: 8035168]
765. Watts BA 3rd, Good DW. ERK mediates inhibition of Na(+)/H(+) exchange and HCO₃(-)-absorption by nerve growth factor in MTAL. *American journal of physiology Renal physiology.* 2002; 282:F1056–1063. [PubMed: 11997322]
766. Watts BA 3rd, Good DW. Hyposmolality stimulates apical membrane Na(+)/H(+) exchange and HCO₃(-)-absorption in renal thick ascending limb. *The Journal of clinical investigation.* 1999; 104:1593–1602. [PubMed: 10587523]
767. Weiner ID, Hamm LL. Regulation of Cl⁻/HCO₃⁻ exchange in the rabbit cortical collecting tubule. *J Clin Invest.* 1991; 87:1553–1558. [PubMed: 2022727]
768. Weiner ID, Hamm LL. Regulation of intracellular pH in the rabbit cortical collecting tubule. *J Clin Invest.* 1990; 85:274–281. [PubMed: 2153152]
769. Weiner ID, New AR, Milton AE, Tisher CC. Regulation of luminal alkalization and acidification in the cortical collecting duct by angiotensin II. *The American journal of physiology.* 1995; 269:F730–738. [PubMed: 7503240]
770. Weiner ID, Verlander JW. Ammonia Transport in the Kidney by Rhesus Glycoproteins. *American journal of physiology Renal physiology.* 2014
771. Weiner ID, Verlander JW. Renal ammonia metabolism and transport. *Compr Physiol.* 3:201–220. [PubMed: 23720285]
772. Weiner ID, Weill AE, New AR. Distribution of Cl⁻/HCO₃⁻ exchange and intercalated cells in rabbit cortical collecting duct. *The American journal of physiology.* 1994; 267:F952–964. [PubMed: 7810703]
773. Weinman EJ, Minkoff C, Shenolikar S. Signal complex regulation of renal transport proteins: NHERF and regulation of NHE3 by PKA. *Am J Physiol Renal Physiol.* 2000; 279:F393–399. [PubMed: 10966919]
774. Weinman EJ, Shenolikar S. Protein kinase C activates the renal apical membrane Na⁺/H⁺ exchanger. *J Membr Biol.* 1986; 93:133–139. [PubMed: 3027347]
775. Weinman EJ, Shenolikar S, Kahn AM. cAMP-associated inhibition of Na⁺-H⁺ exchanger in rabbit kidney brush-border membranes. *Am J Physiol.* 1987; 252:F19–25. [PubMed: 3028154]
776. Weinman EJ, Steplock D, Donowitz M, Shenolikar S. NHERF associations with sodium-hydrogen exchanger isoform 3 (NHE3) and ezrin are essential for cAMP-mediated phosphorylation and inhibition of NHE3. *Biochemistry.* 2000; 39:6123–6129. [PubMed: 10821685]
777. Weinman EJ, Steplock D, Shenolikar S. Acute regulation of NHE3 by protein kinase A requires a multiprotein signal complex. *Kidney Int.* 2001; 60:450–454. [PubMed: 11473625]
778. Weinman EJ, Steplock D, Shenolikar S. CAMP-mediated inhibition of the renal brush border membrane Na⁺-H⁺ exchanger requires a dissociable phosphoprotein cofactor. *J Clin Invest.* 1993; 92:1781–1786. [PubMed: 8408631]
779. Weinman EJ, Steplock D, Shenolikar S. NHERF-1 uniquely transduces the cAMP signals that inhibit sodium-hydrogen exchange in mouse renal apical membranes. *FEBS Lett.* 2003; 536:141–144. [PubMed: 12586353]
780. Weinman EJ, Steplock D, Wade JB, Shenolikar S. Ezrin binding domain-deficient NHERF attenuates cAMP-mediated inhibition of Na(+)/H(+) exchange in OK cells. *Am J Physiol Renal Physiol.* 2001; 281:F374–380. [PubMed: 11457730]
781. Weinman EJ, Steplock D, Wang Y, Shenolikar S. Characterization of a protein cofactor that mediates protein kinase A regulation of the renal brush border membrane Na(+)-H⁺ exchanger. *J Clin Invest.* 1995; 95:2143–2149. [PubMed: 7738182]
782. Weinman EJ, Wang Y, Wang F, Greer C, Steplock D, Shenolikar S. A C-terminal PDZ motif in NHE3 binds NHERF-1 and enhances cAMP inhibition of sodium-hydrogen exchange. *Biochemistry.* 2003; 42:12662–12668. [PubMed: 14580213]

783. Weinstein AM. Chloride transport in a mathematical model of the rat proximal tubule. *Am J Physiol.* 1992; 263:F784–798. [PubMed: 1443169]
784. Welsh-Bacic D, Nowik M, Kaissling B, Wagner CA. Proliferation of acid-secretory cells in the kidney during adaptive remodelling of the collecting duct. *PLoS One.* 6:e25240. [PubMed: 22039408]
785. Wesson DE. Na/H exchange and H-K ATPase increase distal tubule acidification in chronic alkalosis. *Kidney international.* 1998; 53:945–951. [PubMed: 9551402]
786. Wesson DE. Prostacyclin increases distal tubule HCO₃ secretion in the rat. *Am J Physiol.* 1996; 271:F1183–1192. [PubMed: 8997392]
787. Wesson DE, Dolson GM. Endothelin-1 increases rat distal tubule acidification in vivo. *Am J Physiol.* 1997; 273:F586–594. [PubMed: 9362336]
788. Wesson DE, Simoni J, Green DF. Reduced extracellular pH increases endothelin-1 secretion by human renal microvascular endothelial cells. *J Clin Invest.* 1998; 101:578–583. [PubMed: 9449690]
789. Wiederkehr MR, Zhao H, Moe OW. Acute regulation of Na/H exchanger NHE3 activity by protein kinase C: role of NHE3 phosphorylation. *Am J Physiol.* 1999; 276:C1205–1217. [PubMed: 10329970]
790. Wingo CS, Madsen KM, Smolka A, Tisher CC. H-K-ATPase immunoreactivity in cortical and outer medullary collecting duct. *Kidney Int.* 1990; 38:985–990. [PubMed: 2176257]
791. Winkler CA, Kittelberger AM, Schwartz GJ. Expression of carbonic anhydrase IV mRNA in rabbit kidney: stimulation by metabolic acidosis. *The American journal of physiology.* 1997; 272:F551–560. [PubMed: 9140058]
792. Winter C, Kampik NB, Vedovelli L, Rothenberger F, Paunescu TG, Stehberger PA, Brown D, John H, Wagner CA. Aldosterone stimulates vacuolar H(+)-ATPase activity in renal acid-secretory intercalated cells mainly via a protein kinase C-dependent pathway. *Am J Physiol Cell Physiol.* 301:C1251–1261. [PubMed: 21832245]
793. Winter C, Schulz N, Giebisch G, Geibel JP, Wagner CA. Nongenomic stimulation of vacuolar H⁺-ATPases in intercalated renal tubule cells by aldosterone. *Proc Natl Acad Sci U S A.* 2004; 101:2636–2641. [PubMed: 14983061]
794. Woodcock EA, Johnston CI. Inhibition of adenylate cyclase by angiotensin II in rat renal cortex. *Endocrinology.* 1982; 111:1687–1691. [PubMed: 7128530]
795. Wright PA, Knepper MA. Glutamate dehydrogenase activities in microdissected rat nephron segments: effects of acid-base loading. *Am J Physiol.* 1990; 259:F53–59. [PubMed: 2375392]
796. Wright PA, Knepper MA. Phosphate-dependent glutaminase activity in rat renal cortical and medullary tubule segments. *Am J Physiol.* 1990; 259:F961–970. [PubMed: 2175563]
797. Wu J, McNicholas CM, Bevensee MO. Phosphatidylinositol 4,5-bisphosphate (PIP₂) stimulates the electrogenic Na/HCO₃ cotransporter NBCe1-A expressed in *Xenopus* oocytes. *Proceedings of the National Academy of Sciences of the United States of America.* 2009; 106:14150–14155. [PubMed: 19667194]
798. Wu MS, Biemesderfer D, Giebisch G, Aronson PS. Role of NHE3 in mediating renal brush border Na⁺-H⁺ exchange. Adaptation to metabolic acidosis. *J Biol Chem.* 1996; 271:32749–32752. [PubMed: 8955109]
799. Xie Q, Welch R, Mercado A, Romero MF, Mount DB. Molecular characterization of the murine Slc26a6 anion exchanger: functional comparison with Slc26a1. *Am J Physiol Renal Physiol.* 2002; 283:F826–838. [PubMed: 12217875]
800. Xu J, Barone S, Petrovic S, Wang Z, Seidler U, Riederer B, Ramaswamy K, Dudeja PK, Shull GE, Soleimani M. Identification of an apical Cl[−]/HCO₃[−] exchanger in gastric surface mucous and duodenal villus cells. *Am J Physiol Gastrointest Liver Physiol.* 2003; 285:G1225–1234. [PubMed: 12893630]
801. Xu J, Worrell RT, Li HC, Barone SL, Petrovic S, Amlal H, Soleimani M. Chloride/bicarbonate exchanger SLC26A7 is localized in endosomes in medullary collecting duct cells and is targeted to the basolateral membrane in hypertonicity and potassium depletion. *J Am Soc Nephrol.* 2006; 17:956–967. [PubMed: 16524946]

802. Xu L, Dixit MP, Nullmeyer KD, Xu H, Kiela PR, Lynch RM, Ghishan FK. Regulation of Na⁺/H⁺ exchanger-NHE3 by angiotensin-II in OKP cells. *Biochim Biophys Acta*. 2006; 1758:519–526. [PubMed: 16603121]
803. Yamada H, Horita S, Suzuki M, Fujita T, Seki G. Functional role of a putative carbonic anhydrase II-binding domain in the electrogenic Na⁺-HCO₃⁻ cotransporter NBCe1 expressed in *Xenopus* oocytes. *Channels (Austin)*. 2011; 5:106–109. [PubMed: 21224720]
804. Yamazaki O, Yamada H, Suzuki M, Horita S, Shirai A, Nakamura M, Satoh N, Fujita T, Seki G. Identification of dominant negative effect of L522P mutation in the electrogenic Na⁺(+)-HCO₃(3) (-) cotransporter NBCe1. *Pflugers Arch*. 465:1281–1291. [PubMed: 23559100]
805. Yan Y, Deneff N, Schupbach T. The vacuolar proton pump, V-ATPase, is required for notch signaling and endosomal trafficking in *Drosophila*. *Dev Cell*. 2009; 17:387–402. [PubMed: 19758563]
806. Yang LE, Maunsbach AB, Leong PK, McDonough AA. Differential traffic of proximal tubule Na⁺ + transporters during hypertension or PTH: NHE3 to base of microvilli vs. NaPi2 to endosomes. *Am J Physiol Renal Physiol*. 2004; 287:F896–906. [PubMed: 15265767]
807. Yang LE, Sandberg MB, Can AD, Pihakaski-Maunsbach K, McDonough AA. Effects of dietary salt on renal Na⁺ transporter subcellular distribution, abundance, and phosphorylation status. *Am J Physiol Renal Physiol*. 2008; 295:F1003–1016. [PubMed: 18653479]
808. Yang Q, Li G, Singh SK, Alexander EA, Schwartz JH. Vacuolar H⁺ - ATPase B1 subunit mutations that cause inherited distal renal tubular acidosis affect proton pump assembly and trafficking in inner medullary collecting duct cells. *J Am Soc Nephrol*. 2006; 17:1858–1866. [PubMed: 16769747]
809. Yenchitsomanus PT, Kittanakom S, Rungroj N, Cordat E, Reithmeier RA. Molecular mechanisms of autosomal dominant and recessive distal renal tubular acidosis caused by SLC4A1 (AE1) mutations. *J Mol Genet Med*. 2005; 1:49–62. [PubMed: 19565014]
810. Yingst DR, Massey KJ, Rossi NF, Mohanty MJ, Mattingly RR. Angiotensin II directly stimulates activity and alters the phosphorylation of Na-K-ATPase in rat proximal tubule with a rapid time course. *Am J Physiol Renal Physiol*. 2004; 287:F713–721. [PubMed: 15161602]
811. Yip KP, Tsuruoka S, Schwartz GJ, Kurtz I. Apical H(+)/base transporters mediating bicarbonate absorption and pH(i) regulation in the OMCD. *Am J Physiol Renal Physiol*. 2002; 283:F1098–1104. [PubMed: 12372786]
812. Yoshitomi K, Burckhardt BC, Fromter E. Rheogenic sodium-bicarbonate cotransport in the peritubular cell membrane of rat renal proximal tubule. *Pflugers Arch*. 1985; 405:360–366. [PubMed: 3936018]
813. Yoshitomi K, Fromter E. Cell pH of rat renal proximal tubule in vivo and the conductive nature of peritubular HCO₃⁻ (OH⁻) exit. *Pflugers Arch*. 1984; 402:300–305. [PubMed: 6441147]
814. Younes-Ibrahim M, Barlet-Bas C, Buffin-Meyer B, Cheval L, Rajerison R, Doucet A. Ouabain-sensitive and -insensitive K-ATPases in rat nephron: effect of K depletion. *Am J Physiol*. 1995; 268:F1141–1147. [PubMed: 7611455]
815. Yu H, Riederer B, Stieger N, Boron WF, Shull GE, Manns MP, Seidler UE, Bachmann O. Secretagogue stimulation enhances NBCe1 (electrogenic Na⁺/HCO₃⁻ cotransporter) surface expression in murine colonic crypts. *Am J Physiol Gastrointest Liver Physiol*. 2009; 297:G1223–1231. [PubMed: 19779011]
816. Yuge S, Inoue K, Hyodo S, Takei Y. A novel guanylin family (guanylin, uroguanylin, and renoguanylin) in eels: possible osmoregulatory hormones in intestine and kidney. *J Biol Chem*. 2003; 278:22726–22733. [PubMed: 12684514]
817. Yun CH, Gurubhagavatula S, Levine SA, Montgomery JL, Brant SR, Cohen ME, Cragoe EJ Jr. Pouyssegur J, Tse CM, Donowitz M. Glucocorticoid stimulation of ileal Na⁺ absorptive cell brush border Na⁺/H⁺ exchange and association with an increase in message for NHE-3, an epithelial Na⁺/H⁺ exchanger isoform. *J Biol Chem*. 1993; 268:206–211. [PubMed: 8380155]
818. Yun CH, Lamprecht G, Forster DV, Sidor A. NHE3 kinase A regulatory protein E3KARP binds the epithelial brush border Na⁺/H⁺ exchanger NHE3 and the cytoskeletal protein ezrin. *J Biol Chem*. 1998; 273:25856–25863. [PubMed: 9748260]

819. Yun CH, Oh S, Zizak M, Steplock D, Tsao S, Tse CM, Weinman EJ, Donowitz M. cAMP-mediated inhibition of the epithelial brush border Na⁺/H⁺ exchanger, NHE3, requires an associated regulatory protein. *Proc Natl Acad Sci U S A*. 1997; 94:3010–3015. [PubMed: 9096337]
820. Zhang D, Kiyatkin A, Bolin JT, Low PS. Crystallographic structure and functional interpretation of the cytoplasmic domain of erythrocyte membrane band 3. *Blood*. 2000; 96:2925–2933. [PubMed: 11049968]
821. Zhang Y, Norian JM, Magyar CE, Holstein-Rathlou NH, Mircheff AK, McDonough AA. In vivo PTH provokes apical NHE3 and NaPi2 redistribution and Na-K ATPase inhibition. *Am J Physiol*. 1999; 276:F711–719. [PubMed: 10330053]
822. Zhao H, Wiederkehr MR, Fan L, Collazo RL, Crowder LA, Moe OW. Acute inhibition of Na/H exchanger NHE-3 by cAMP. Role of protein kinase a and NHE-3 phosphoserines 552 and 605. *J Biol Chem*. 1999; 274:3978–3987. [PubMed: 9933588]
823. Zheng Y, Horita S, Hara C, Kunimi M, Yamada H, Sugaya T, Goto A, Fujita T, Seki G. Biphasic regulation of renal proximal bicarbonate absorption by luminal AT(1A) receptor. *Journal of the American Society of Nephrology : JASN*. 2003; 14:1116–1122. [PubMed: 12707382]
824. Zhou Y, Boron WF. Role of endogenously secreted angiotensin II in the CO₂-induced stimulation of HCO₃⁻ reabsorption by renal proximal tubules. *American journal of physiology Renal physiology*. 2008; 294:F245–252. [PubMed: 17913836]
825. Zhou Y, Bouyer P, Boron WF. Role of a tyrosine kinase in the CO₂-induced stimulation of HCO₃⁻ reabsorption by rabbit S2 proximal tubules. *Am J Physiol Renal Physiol*. 2006; 291:F358–367. [PubMed: 16705143]
826. Zhou Y, Zhao J, Bouyer P, Boron WF. Evidence from renal proximal tubules that HCO₃⁻ and solute reabsorption are acutely regulated not by pH but by basolateral HCO₃⁻ and CO₂. *Proc Natl Acad Sci U S A*. 2005; 102:3875–3880. [PubMed: 15728388]
827. Zhu Q, Kao L, Azimov R, Abuladze N, Newman D, Pushkin A, Liu W, Chang C, Kurtz I. Structural and functional characterization of the C-terminal transmembrane region of NBCe1-A. *J Biol Chem*. 285:37178–37187. [PubMed: 20837482]
828. Zhu Q, Kao L, Azimov R, Newman D, Liu W, Pushkin A, Abuladze N, Kurtz I. Topological location and structural importance of the NBCe1-A residues mutated in proximal renal tubular acidosis. *J Biol Chem*. 285:13416–13426. [PubMed: 20197274]
829. Zhu Q, Shao XM, Kao L, Azimov R, Weinstein AM, Newman D, Liu W, Kurtz I. Missense mutation T485S alters NBCe1-A electrogenicity causing proximal renal tubular acidosis. *Am J Physiol Cell Physiol*. 305:C392–405. [PubMed: 23636456]
830. Zimolo Z, Montrose MH, Murer H. H⁺ extrusion by an apical vacuolar-type H⁽⁺⁾-ATPase in rat renal proximal tubules. *The Journal of membrane biology*. 1992; 126:19–26. [PubMed: 1317456]
831. Zizak M, Lamprecht G, Steplock D, Tariq N, Shenolikar S, Donowitz M, Yun CH, Weinman EJ. cAMP-induced phosphorylation and inhibition of Na⁽⁺⁾/H⁽⁺⁾ exchanger 3 (NHE3) are dependent on the presence but not the phosphorylation of NHE regulatory factor. *J Biol Chem*. 1999; 274:24753–24758. [PubMed: 10455146]