Published in final edited form as:

Curr Opin Psychol. 2015 August 1; 4: 93–97. doi:10.1016/j.copsyc.2014.12.017.

Frontal EEG Asymmetry as a Promising Marker of Depression Vulnerability: Summary and Methodological Considerations

John J.B. Allen and Samantha J. Reznik

Department of Psychology, University of Arizona

Abstract

Frontal EEG asymmetry is a promising neurophysiological marker of depression risk. It predicts emotional response and negative affect hours to years later. Yet, inconsistencies in the literature may be due to differing methodological approaches between research groups. Within the past two years, a number of studies have shown this line of research may be strengthened by augmenting resting assessments with emotionally evocative tasks, utilizing optimal recording montages, and taking an integrative neuroscience approach that links frontal asymmetry to other indices of neural function. This review will focus on recent work in frontal asymmetry and depression with a particular focus on promising future directions and methodological considerations that may increase consistency between research groups.

Introduction

Investigating neurophysiological indices of depression may facilitate the identification of those at risk for the disorder, provide a metric to assess preventative measures, and lead to the development of biologically driven diagnosis and treatment. In particular, a large body of research suggests that frontal asymmetry may be a neurophysiological biomarker of depression risk.

Frontal asymmetry is a relative measure of the difference in electroencephalogram (EEG) alpha power between the right and left frontal regions and has high internal consistency reliability and good test-retest reliability over intervals from weeks to months [1,2]. As alpha activity has an inhibitory influence on cortical network activity, lower frontal asymmetry scores (right minus left alpha) reflect relatively less left than right cortical activity [1]. Less relative left frontal activity during the resting state has been associated with depressive symptoms [3] and current and past depression status [4–7]. This pattern of frontal asymmetry has also been found in infants of depressed mothers [8–10] and is related to mother's depression score [10]. Less relative left frontal activity prospectively predicts first

Corresponding author: John J.B. Allen, University of Arizona, P.O. Box 210068, Tucson, AZ, USA, 85721, John.JB.Allen@arizona.edu, **Phone: 520-621-4992, FAX: 520-844-8270.**

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

depressive episode in college students [11]. This line of research suggests frontal asymmetry may indicate vulnerability to depression [12].

Although frontal asymmetry has emerged as a promising biomarker for depression risk, inconsistencies in this literature stem from differences in methodology between investigators. The present article will review recent work in frontal asymmetry and depression, focusing on promising future directions and methodological considerations that may increase consistency between research groups. In particular, frontal asymmetry has utility as a predictor of depression onset, but this line of research may be strengthened by: augmenting resting assessments with emotionally evocative tasks (i.e. replacing a trait model with a capability model); utilizing optimal recording montages that isolate local activity; and taking an integrative neuroscience approach that links frontal asymmetry to other indices of neural function.

The predictive power of frontal asymmetry

Recent findings indicate frontal asymmetry is a predictor of depressive symptoms and treatment response. Frontal asymmetry predicts emotional response and negative affect on a short-term scale. Less relative left frontal activity during rest predicted decreased neurophysiological change on subsequent emotional tasks [13], and less relative left frontal activity during negative films predicted mood deterioration in the next week [14]. Additionally, frontal asymmetry has predictive power over longer time frames: less relative left frontal activity predicted depressive symptoms in adolescent boys a year later [15] and also predicted self-reported homesickness among college freshman [16].

A recent review focused on a number of EEG indices as biomarkers of depression treatment response. Although alpha asymmetry in regions other than the frontal sites predicted treatment response for participants taking tricyclic antidepressants and SSRIs, frontal asymmetry did not differentiate responders from non-responders [17–19]. A recent literature review of music therapy cited increased relative left frontal activity after music exposure as support for the effectiveness of music therapy [20], although the longer-term impact of music therapy must be examined. In a behavioral activation treatment study, absolute magnitude of frontal asymmetry (in either direction) predicted negative affect after treatment but not treatment response [21]. The extent to which frontal asymmetry may mark treatment response more broadly – for psychotherapy, pharmacological, and deep-brain stimulation treatments – remains largely unexplored (as research has focused on alternate alpha measures) and is an important area for future research.

Research also suggests the possibility of treating depression by changing frontal asymmetry. Inpatients with treatment-resistant depression had reduced depressive symptoms after repetitive transcranial magnetic stimulation (rTMS) targeted to the left prefrontal cortex [22,23]. A less invasive neurofeedback procedure provides auditory feedback based on frontal asymmetry scores and can alter frontal asymmetry. Five-weeks of neurofeedback training increased relative left frontal asymmetry and decreased depressive symptoms in 24 participants with current depression [24]. A one-session manipulation of frontal asymmetry in individuals without history of psychopathology, however, created mood changes [25].

Thus repeated neurofeedback may be a tenable intervention for depression, directly targeting the neural systems indexed by frontal asymmetry. Further research should examine the duration and course of neurofeedback necessary for efficacy on various populations. Additionally, tonic stimulation with weak direct currents (tDCS) changes cortical excitability and has been investigated as a potential non-invasive measure for depression [26]. Transcranial ultrasound (TUS) over lateral prefrontal cortex improves mood acutely [27], although its longer-term impact on mood remains to be assessed. The impact of these new stimulation approaches, on frontal EEG asymmetry, however, has not been investigated and is an important direction for future research. Finally, transcranial alternating current stimulation (tACS) directly interferes with cortical rhythms but its potential to alter prefrontal asymmetry has never been investigated [28].

Resting assessments versus emotionally evocative tasks: Support for the capability model

Given inconsistencies in the literature and the potential of frontal asymmetry to help identify those at risk for depression, finding the conditions under which frontal asymmetry can best serve as a marker of depression risk remains of great importance. The capability model proposes that frontal asymmetry may be best understood as the interaction between individual differences in emotion regulation and the emotional salience of a particular situation [29]. The major hypothesis of the capability model is that individual differences in frontal asymmetry will be more pronounced during emotionally evocative tasks than during rest. A growing body of research supports this hypothesis. A study of 306 currently, previously, and never depressed participants found that depression status was more strongly predicted by frontal asymmetry during a directed facial action task than frontal asymmetry during rest for Cz, linked mastoids, and average references [30].

Several recent studies found less relative left frontal activity associated with depressive affect during emotionally evocative tasks but not during rest. For example, frontal asymmetry did not differ between individuals with major depressive disorder and those with borderline personality disorder during rest, yet less relative left frontal activity was found in depressed but not borderline participants after a rejection task [31]. Relatedly, women with premenstrual dysphoric disorder had less relative left frontal activity during both depression induction and relaxation tasks but not during rest [32]. Furthermore, 27 mothers of infants 5–7 months old had EEG recorded during rest and during videos of their infants expressing emotions. Self-reported experiential sadness of the mothers related to less relative left frontal activity during videos of their infants expressing emotion but not rest [33].

Similar findings derive from recent studies of children and older adults. Children ages 6–13 whose mothers have a history of depression had less relative left frontal activity in comparison to their peers during emotionally evocative films but not during rest [34]. In a small sample of adults over 65, no relationship was found between frontal asymmetry during rest and depressive symptoms or depression status. The authors suggested that frontal asymmetry may not have the same relationship in older adults [35]. Although such an interpretation is possible, these results are consistent with the hypothesis that individual differences in frontal asymmetry are more pronounced during emotionally evocative tasks

than at rest, thus supporting the capability model. Moreover, a linked mastoids reference montage may limit the ability to detect meaningful relationships between prefrontal asymmetry and depression (see below). A comprehensive assessment of frontal asymmetry throughout the lifespan should include emotionally evocative tasks as well as resting state data.

Additionally, consideration must be given to the type of emotionally evocative tasks used. It might be reasonable to expect that tasks that call for emotion regulation that is partially dependent on lateral prefrontal activity would likely use systems that may be altered in those at risk for depression. For example, Meyer et. al (2014) found that frontal asymmetry was moderated by the extent to which their emotionally evocative videos allowed cognitive emotion regulation [36]. Significant interactions between frontal asymmetry, electrodermal reaction, and internalizing and externalizing symptoms were found during sad and happy videos but not fearful or angry videos [37].

The importance of recording montage

Further consideration should be given to recording montage in frontal asymmetry research. Electrical signals recorded at any given electrode are highly dependent on the reference site used. For over three decades, researchers have examined effects of several recording montages, including: average (the average of all sites); CZ (vertex); and, averaged (“linked”) mastoids reference [38,39]. These traditional references reflect not only frontal alpha but also substantial alpha power originating in other regions (e.g., the Occipital cortex). The reference-free current-source density (CSD) transformation, however, uses an estimate of current sources and sinks on the scalp and attenuates the contribution of distal sources, thus highlighting the contribution of sources nearer the recording electrode. Some recent work suggests that CSD transformation may provide a more sensitive index of individual variations in frontal asymmetry and reduce non-frontal contributions to frontal asymmetry compared to these typical recording montages [40]. Stewart, Coan, Towers, & Allen (2014) found that only CSD transformed frontal asymmetry – but not average, nor linked mastoids, nor CZ referenced asymmetry – differed as a function of lifetime history of depression during the resting state for 306 participants [30,41]. Another study found that the amount of time since a participant had woken up at EEG recording differed more for average and linked mastoids references than for CSD [40]. Thus on both empirical and rational grounds, the CSD transformation may be the preferred approach for assessing the relationship of frontal asymmetry to depression, and for analyzing the asymmetrical activity of specifically frontal systems.

Using integrative neuroscience approaches to link frontal asymmetry to underlying neural systems

Despite considerable research indicating frontal asymmetry may be a marker of depression risk, few studies have linked frontal asymmetry to other neural systems. Functional magnetic resonance (fMRI) studies suggest the importance of lateral frontal systems in depression and depression risk [42,43]. Abnormalities in the cognitive control network (CCN) characterize depression [44]; the CCN serves basic cognitive control mechanisms

(e.g. attentional control) and is located primarily in frontal and parietal regions. Individuals with depression have increased connectivity between the CCN and the dorsal nexus, an area in the bilateral dorsal medial prefrontal cortex region [45] and higher dorsal to rostral anterior cingulate cortex connectivity [46]. Although frontal regions are often implicated in fMRI activation and connectivity studies, frontal asymmetry has seldom been tested explicitly in this work.

One recent study used simultaneous CSD-transformed EEG and fMRI during rest to examine correspondence between frontal asymmetry and connectivity assessed with fMRI. Less relative left frontal activity was associated with increased connectivity between the left inferior frontal gyrus and two anterior cingulate cortex resting networks [47]. This work suggests frontal asymmetry may relate to a hyper connected emotional verbalization network that may reflect ineffective attempts at emotion regulation such as rumination [48]. These results provide one of the few links between resting frontal EEG asymmetry and brain connectivity. These findings, however, reflect between-subjects correlations, and future work should examine whether fMRI connectivity is moderated within each participant by changes in asymmetry throughout the recording session. This research can provide clues about how frontal EEG asymmetry may influence network organization in ways that promote depression risk.

Additionally, linking frontal asymmetry to underlying neural network organization holds the promise for the development of neurally-informed treatments by suggesting specific regional targets for intervention, for example using brain stimulation approaches such as rTMS, tDCS, TUS, and tACS.

Conclusion

Frontal asymmetry may serve as a prospective predictor of depression onset and, as such, has wide-ranging theoretical and translational applications. Frontal asymmetry can potentially identify those at risk for depression, although a large-scale prospective study is needed. Additionally, because frontal asymmetry may index the activity of neural systems that promote depression risk, research in this area can inform the development of neurally-informed, individualized treatments, and allow for an assessment of how well treatments (including psychotherapy) alter activity in within these neural systems. This important line of research may benefit from utilizing EEG not only during rest but also during emotionally evocative tasks (capability model), using the CSD-transformation to highlight specifically frontal activity, and employing multi-modal imaging to link an easily-obtained and widely-studied measure frontal asymmetry to other measures of brain activity known to be involved in depression.

References

1. Towers DN, Allen JJB. A better estimate of the internal consistency reliability of frontal EEG asymmetry scores. *Psychophysiology*. 2009; 46:132–142. [PubMed: 19055503]
2. Sutton SK, Davidson RJ, Science SP, May N. Prefrontal brain asymmetry: A biological substrate of the behavioral approach and inhibition systems. *Psychol Sci*. 2014; 8:204–210.

3. Schaffer C, Davidson R, Saron C. Frontal and parietal electroencephalogram asymmetry in depressed and nondepressed subject. *Biol Psychiatry*. 1983; 18:753–762. [PubMed: 6615936]
4. Gotlib IH, Ranganath C, Rosenfeld P. Frontal EEG alpha asymmetry, depression, and cognitive functioning. *Cogn Emot*. 1998; 12:449–478.
5. Henriques JB, Davidson RJ. Left frontal hypoactivation in depression. *J Abnorm Psychol*. 1991; 100:535–545. [PubMed: 1757667]
6. Thibodeau R, Jorgensen RS, Kim S. Depression, anxiety, and resting frontal EEG asymmetry: a meta-analytic review. *J Abnorm Psychol*. 2006; 115:715–729. [PubMed: 17100529] ** The first meta-analysis of frontal EEG asymmetry and its relationship to depression and anxiety supports the conclusion that this is a moderate effect size.
7. Davidson RJ. Cerebral asymmetry and emotion: Conceptual and methodological conundrums. *Cogn Emot*. 1993; 7:115–138.
8. Dawson G, Frey K, Panagiotides H, Yamada E, Hessel D, Osterling J. Infants of depressed mothers exhibit atypical frontal electrical brain activity during interactions with mother and with a familiar, nondepressed adult. *Child Dev*. 1999; 70:1058–1066. [PubMed: 10546334]
9. Field T, Fox NA, Pickens J, Nawrocki T. Relative right frontal EEG activation in 3- to 6-month-old infants of “depressed” mothers. *Dev Psychol*. 1995; 31:358–363.
10. Lusby CM, Goodman SH, Bell MA, Newport DJ. Electroencephalogram patterns in infants of depressed mothers. *Dev Psychobiol*. 2013; 56:459–473. [PubMed: 23852456] * Infant EEG (3 and 6 months) was predicted by prenatal maternal depressive severity in a sample of 83 mother-infant dyads, but only among those women who also had high post-partum depressive symptoms. Data were averaged referenced and may benefit from further consideration of recording montage.
11. Nusslock R, Shackman AJ, Harmon-Jones E, Alloy LB, Coan JA, Abramson LY. Cognitive vulnerability and frontal brain asymmetry: common predictors of first prospective depressive episode. *J Abnorm Psychol*. 2011; 120:497–503. [PubMed: 21381804] ** Provided the first demonstration of the predictive value of frontal asymmetry for future onset of a depressive episode.
12. Coan JA, Allen JJB. Frontal EEG asymmetry as a moderator and mediator of emotion. *Biol Psychol*. 2004; 67:7–49. [PubMed: 15130524]
13. Papousek I, Reiser EM, Weber B, Freudenthaler HH, Schuster G. Frontal brain asymmetry and affective flexibility in an emotional contagion paradigm. *Psychophysiology*. 2012; 49:489–498. [PubMed: 22176666] * Evaluated emotional responsivity to sounds in 80 participants. Relative less left frontal activity during rest with averaged ears reference predicted increased change in frontal asymmetry during sounds. This study supports predictive power of frontal asymmetry and capability model.
14. Papousek I, Weiss EM, Schuster G, Fink A, Reiser EM, Lackner HK. Prefrontal EEG alpha asymmetry changes while observing disaster happening to other people: Cardiac correlates and prediction of emotional impact. *Biol Psychol*. 2014; 103:184–194. [PubMed: 25224180] * In 148 undergraduates, relative less left frontal activity while viewing emotional videos predicted sadness and hopelessness in the following week. Data were averaged ears referenced and may benefit from further consideration of recording montage. This study supports short-term predictive power of frontal asymmetry during emotionally evocative tasks but not during rest.
15. Mitchell AM, Pössel P. Frontal brain activity pattern predicts depression in adolescent boys. *Biol Psychol*. 2012; 89:525–527. [PubMed: 22200655] * Relative less left frontal activity during rest in 50 adolescent boys (mean age =13.93) with previous or current diagnosis of depression predicted depressive symptoms one year later. A nose reference was used, so further consideration of recording montage may be useful. This study supports the power of prefrontal asymmetry to predict depression severity.
16. Steiner ARW, Coan JA. Prefrontal asymmetry predicts affect, but not beliefs about affect. *Biol Psychol*. 2011; 88:65–71. [PubMed: 21741433]
17. Bruder GE, Sedoruk JP, Stewart JW, McGrath PJ, Quitkin FM, Tenke CE. Electroencephalographic alpha measures predict therapeutic response to a selective serotonin reuptake inhibitor antidepressant: pre- and post-treatment findings. *Biol Psychol*. 2008; 63:1171–1177.

18. Bruder GE, Stewart JW, Tenke CE, McGrath PJ, Leite P, Bhattacharya N, Quitkin FM. Electroencephalographic and perceptual asymmetry differences between responders and nonresponders to an SSRI antidepressant. *Biol Psychol.* 2001; 49:416–425.
19. Baskaran A, Milev R, McIntyre RS. The neurobiology of the EEG biomarker as a predictor of treatment response in depression. *Neuropharmacology.* 2012; 63:507–513. [PubMed: 22569197]
20. Fachner J, Gold C, Erkkilä J. Music therapy modulates fronto-temporal activity in rest-EEG in depressed clients. *Brain Topogr.* 2013; 26:338–354. [PubMed: 22983820] * This review focuses on the effect of music and music therapy on various EEG indices. Frontal asymmetry as a potential measure of treatment response to music therapy is discussed.
21. Gollan JK, Hoxha D, Chihade D, Pflieger ME, Rosebrock L, Cacioppo J. Frontal alpha EEG asymmetry before and after behavioral activation treatment for depression. *Biol Psychol.* 2014; 99:198–208. [PubMed: 24674708] * Thirty seven currently depressed participants completed 16 sessions of behavioral activation therapy over the course of 16 weeks, and 35 healthy control participants were followed prospectively for 16 weeks. The authors examined absolute magnitude of frontal asymmetry (in either direction, by simply taking the absolute value). This novel metric acquired before treatment did not predict treatment response as expected but was related to negative affect post-treatment. Given methodological differences in measuring prefrontal asymmetry in this study (absolute difference rather than directional difference between $\ln[\text{right}]$ minus $\ln[\text{left}]$), further consideration of traditional measures of frontal asymmetry as a predictor of treatment response may be beneficial.
22. George MS, Wassermann EM, Williams WA, Callahan A, Ketter TA, Basser P, Hallett M, Post RM. Daily repetitive transcranial magnetic stimulation (rTMS) improves mood in depression. *Neuroreport.* 1995; 6:1853–1856. [PubMed: 8547583]
23. Pascual-leone A, Rubio B, Pallardó F, Catalá MD. Rapid-rate transcranial magnetic stimulation of left dorsolateral prefrontal cortex in drug-resistant depression. *Lancet.* 1996; 348:233–237. [PubMed: 8684201]
24. Choi SW, Chi SE, Chung SY, Kim JW, Ahn CY, Kim HT. Is alpha wave neurofeedback effective with randomized clinical trials in depression? A pilot study. *Neuropsychobiology.* 2011; 63:43–51. [PubMed: 21063132]
25. Peeters F, Ronner J, Bodar L, van Os J, Lousberg R. Validation of a neurofeedback paradigm: manipulating frontal EEG alpha-activity and its impact on mood. *Int J Psychophysiol.* 2014; 93:116–120. [PubMed: 23773999] ** 40 women with no history of psychopathology were able to use neurofeedback to upregulate either relative right or left activity, but no changes in mood were observed. Data were averaged ears referenced, and may benefit from consideration of recording montage and increased sample size.
26. Nitsche MA, Boggio PS, Fregni F, Pascual-Leone A. Treatment of depression with transcranial direct current stimulation (tDCS): a review. *Exp Neurol.* 2009; 219:14–19. [PubMed: 19348793]
27. Sanguinetti JL, Smith EE, Dieckman L, Vanuk J, Hameroff S, Allen JJB. Noninvasive transcranial ultrasound (TUS) for brain stimulation: Effects on mood in a pilot study. *Psychophysiology.* 2012; 50:S36.
28. Antal A, Paulus W. Transcranial alternating current stimulation (tACS). *Front Hum Neurosci.* 2013; 7:317. [PubMed: 23825454]
29. Coan JA, Allen JJB, McKnight PE. A capability model of individual differences in frontal EEG asymmetry. *Biol Psychol.* 2006; 72:198–207. [PubMed: 16316717]
30. Stewart JL, Coan JA, Towers DN, Allen JJB. Frontal EEG asymmetry during emotional challenge differentiates individuals with and without lifetime major depressive disorder. *J Affect Disord.* 2011; 129:167–174. [PubMed: 20870293]
31. Beeney JE, Levy KN, Gatzke-Kopp LM, Hallquist MN. EEG asymmetry in borderline personality disorder and depression following rejection. *Personal Disord.* 2014; 5:178–185. [PubMed: 24364503] ** Frontal asymmetry with average reference was measured during resting state and after rejection task for healthy controls (n=21), individuals with major depressive disorder (MDD, n=13) and borderline personality disorder (BPD, n=23). MDD and BPD individuals showed opposite patterns of frontal asymmetry after rejection task but not during rest, supporting the capability model. A limitation of this study was the small number of individuals in MDD group, and further work may benefit from expanded sample size and additional recording montages.

32. Lin IM, Tsai YC, Peper E, Yen CF. Depressive mood and frontal alpha asymmetry during the luteal phase in premenstrual dysphoric disorder. *J Obstet Gynaecol Res.* 2013; 39:998–1006. [PubMed: 23551544] *12 women with PMDD were found to have relative less left frontal activity during the luteal phase of the menstrual cycle during a depression induction and relaxation task but not during rest. Data were CZ-referenced. Although this study supports the capability model, larger sample size may strengthen the conclusions.
33. Killeen LA, Teti DM. Mothers' frontal EEG asymmetry in response to infant emotion states and mother-infant emotional availability, emotional experience, and internalizing symptoms. *Dev Psychopathol.* 2012; 24:9–21. [PubMed: 22292990] * Frontal asymmetry (re-referenced offline to a variant of the average reference) predicted self-report emotional experience for 27 mothers while watching emotional videos of their infants but not during rest. This work supports the capability model and has important implications for identifying parenting risk.
34. Lopez-Duran NL, Nusslock R, George C, Kovacs M. Frontal EEG asymmetry moderates the effects of stressful life events on internalizing symptoms in children at familial risk for depression. *Psychophysiology.* 2012; 49:510–521. [PubMed: 22220930] * Average referenced frontal asymmetry during emotionally evocative films but not during rest differentiated between 135 children at high and low-risk for depression. This study supports the capability model and frontal asymmetry as biomarker of depression risk.
35. Carvalho A, Moraes H, Silveira H, Ribeiro P, Piedade RAM, Deslandes AC, Laks J, Versiani M. EEG frontal asymmetry in the depressed and remitted elderly: is it related to the trait or to the state of depression? *J Affect Disord.* 2011; 129:143–148. [PubMed: 20870292]
36. Meyer T, Quaedflieg CWEM, Giesbrecht T, Meijer EH, Abiad S, Smeets T. Frontal EEG asymmetry as predictor of physiological responses to aversive memories. *Psychophysiology.* 2014; 51:853–865. [PubMed: 24779522] * Less relative right-sided activity during road accident video predicted dampened startle response to image reminders of the video. Less relative right-sided activity during genocide video predicted increased startle response to image reminders. This study referenced data to A1 and A2 and used 64 participants. It suggests type of media may influence frontal asymmetry.
37. Gatzke-Kopp LM, Jetha MK, Segalowitz SJ. The role of resting frontal EEG asymmetry in psychopathology: afferent or efferent filter? *Dev Psychobiol.* 2014; 56:73–85. [PubMed: 23168718] * Correlation between less relative left frontal activity and internalizing symptoms was found during rest for 209 kindergarten children using linked mastoids reference. This study has important implications for predicting internalizing and externalizing symptoms in children.
38. Reid SA, Duke LM, Allen JJ. Resting frontal electroencephalographic asymmetry in depression: inconsistencies suggest the need to identify mediating factors. *Psychophysiology.* 1998; 35:389–404. [PubMed: 9643053]
39. Tomarken AJ, Dichter GS, Garber J, Simien C. Resting frontal brain activity: linkages to maternal depression and socio-economic status among adolescents. 2004; 67:77–102.
40. Velo JR, Stewart JL, Hasler BP, Towers DN, Allen JJB. Should it matter when we record? Time of year and time of day as factors influencing frontal EEG asymmetry. *Biol Psychol.* 2012; 91:283–291. [PubMed: 22867950]
41. Stewart JL, Bismark AW, Towers DN, Coan JA, Allen JJB. Resting frontal EEG asymmetry as an endophenotype for depression risk: sex-specific patterns of frontal brain asymmetry. *J Abnorm Psychol.* 2010; 119:502–512. [PubMed: 20677839] ** The largest study of whether frontal EEG asymmetry may index risk for depression, by examining 306 participants: currently depressed, previously depressed but euthymic, and never depressed; Using CSD-transformed frontal asymmetry, less relative left frontal activity characterized those with any history of depression (current or past) but not never-depressed controls. This pattern is consistent with the idea that EEG asymmetry is not merely a correlate of the current depressive state, but may serve as a trait marker of risk for depression.
42. Pizzagalli DA. Frontocingulate dysfunction in depression: toward biomarkers of treatment response. *Neuropsychopharmacology.* 2011; 36:183–206. [PubMed: 20861828]
43. Gotlib IH, Hamilton JP. Neuroimaging and depression: Current status and unresolved issues. *Curr Dir Psychol Sci.* 2014; 17:159–163.

44. Beck AT, Steer RA, Carbin MG. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clin Psychol Rev.* 1988; 8:77–100.
45. Sheline YI, Price JL, Yan Z, Mintun MA. Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. *Proc Natl Acad Sci USA.* 2010; 107:11020–11025. [PubMed: 20534464]
46. Schlösser RGM, Wagner G, Koch K, Dahnke R, Reichenbach JR, Sauer H. Fronto-cingulate effective connectivity in major depression: a study with fMRI and dynamic causal modeling. *Neuroimage.* 2008; 43:645–655. [PubMed: 18761094]
47. Allen JJB, Hewig JS, Hect H, Miltner WHR, Schneyer DM. Linking resting state EEG asymmetry to resting state fMRI with simultaneous recordings. *Psychophysiology.* 2013; 50:S35.
48. Cooney RE, Joormann J, Eugène F, Dennis EL, Gotlib IH. Neural correlates of rumination in depression. *Cogn Affect Behav Neurosci.* 2010; 10:470–478. [PubMed: 21098808]