Published in final edited form as:

J Am Chem Soc. 2015 September 9; 137(35): 11491–11497. doi:10.1021/jacs.5b07201.

Catalytic Kinetic Resolution of Disubstituted Piperidines by Enantioselective Acylation: Synthetic Utility and Mechanistic Insights

Benedikt Wanner[†], Imants Kreituss[†], Osvaldo Gutierrez[‡], Marisa C. Kozlowski^{*,‡}, and Jeffrey W. Bode^{*,†}

[†]Laboratorium für Organische Chemie, Department of Chemistry and Applied Biosciences, ETH-Zürich, 8093 Zürich, Switzerland [‡]Department of Chemistry, Roy and Diana Vagelos Laboratories, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

Abstract

The catalytic kinetic resolution of cyclic amines with achiral N-heterocyclic carbenes (NHC) and chiral hydroxamic acids has emerged as a promising method to obtain enantioenriched amines with high selectivity factors. In this report, we describe the catalytic kinetic resolution of disubstituted piperidines with practical selectivity factors (*s* up to 52) in which we uncovered an unexpected and pronounced conformational effect resulting in disparate reactivity and selectivity between the *cis* and *trans*-substituted piperidine isomers. Detailed experimental and computational (DFT) studies of the kinetic resolution of various disubstituted piperidines revealed a strong preference for the acylation of conformers in which the α -substituent occupies the axial position. This work provides further experimental and computational support for the concerted 7-member transition state model for acyl transfer reagents and expands the scope and functional group tolerance of the secondary amine kinetic resolution.

INTRODUCTION

Disubstituted piperidines are increasingly attractive platforms for drug development, as the precise placement of substituents about the basic, low-molecular weight, three-dimensional scaffold is ideally suited for structure-activity relationship studies.¹ Although the synthesis of such saturated N-heterocycles remains challenging,² recent advances including diastereoselective metalation/cross-coupling,³ conjugate additions of dihydropyridinones,⁴ hydrogenation of disubstituted pyridines⁵ and cyclization methods⁶ are bringing these once exotic building blocks into the mainstream. Unfortunately, most of the available methods for

*Corresponding Author: bode@org.chem.ethz.ch, marisa@sas.upenn.edu.

Supporting Information

Experimental procedures and spectroscopic data for all new compounds. Transition state structures of all minimized TSs. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

The authors declare no competing financial interest.

the preparation of multisubstituted N-heterocycles deliver racemic products and prospects for enantioselective approaches to many substitution patterns are limited.⁷

For such challenging cases, kinetic resolution can be an efficient and powerful alternative to enantioselective synthesis of chiral N-heterocycles.⁸ Although kinetic resolution has the disadvantage of a maximum yield of 50% of a desired stereoisomer, for early stage applications it can be a reliable, effective, and easily implemented tool for the production of enantiopure material. Provided that selectivity factors (*s*) greater than 20 can be obtained, kinetic resolution can be used to separate enantiomers with acceptable yield and outstanding enantiopurity.⁹

In 2011, our group reported a new approach to the catalytic, kinetic resolution of saturated N-heterocycles consisting of two catalysts working synergistically.¹⁰ Our key advance was the use of a chiral hydroxamic acid, employed in either catalytic or stoichiometric amounts, as highly enantio- and chemoselective acylating agent, which operates by a unique mechanism involving concerted acyl and proton transfer via a 7-member transition state.^{11,12} To date, our studies on the kinetic resolution of chiral N-heterocycles using chiral hydroxamic acids have been limited to mono-, α -substituted N-heterocyclic substrates. In contrast to the resolution of α -monosubstituted N-heterocycles, for which only a single enantiomeric pair exists, the resolution of disubstituted piperidines requires an enantioselective catalyst to discriminate among several different ring conformations (Scheme 1).

In this report, we examine the reactivity and selectivity of disubstituted piperidines in the kinetic resolution by means of enantioselective acylation with a chiral hydroxamic acid. Kinetic resolution studies and density functional theory (DFT) calculations exposed remarkable conformational preferences in both reactivity and selectivity, revealing a requirement for axial- α substitution on the reactive form of the N-heterocycle for effective resolution. This preference extends even to substitution patterns where the substituents must occupy bis-axial positions. This work provides, 1) access to valuable enantioenriched disubstituted piperidines, 2) general guidelines for selecting suitable N-heterocycles as substrates for the kinetic resolution via chiral hydroxamic acids, and 3) affirmation of the transition state model involving a high energy conformation of the substrate that we had previously developed on the basis of DFT calculations.

RESULTS AND DISCUSSION

As depicted in Scheme 1, there are four types of disubstituted piperidines that could serve as target substrates for our study. Piperidines without α -substituents, such as 3,5-disubstituted piperidines, undergo acylation in an unselective manner. Further, 2,6-disubstituted substrates were found to be unreactive and were excluded from further studies.

Our work therefore targeted the six isomers arising from 2,3-, 2,4-, and 2,5-disubstitution of piperidine, with *cis* and *trans* stereoisomers in each case. Further disubstituted compounds with an exocyclic 4-methylene group were also prepared and evaluated. The results of the catalytic kinetic resolution are summarized in Table 1. Surprisingly, we observed substantial differences in the reactivity and selectivity between the *cis*- and *trans*-isomers of the

otherwise identical positional isomers. For the 2,3- (Table 1, entries 1 and 2, entries 7 and 8) and 2,5- (Table 1, entries 5 and 6) disubstituted piperidines the *cis*-isomer underwent faster reactions with higher selectivity, while the *trans*-isomer gave rise to superior outcomes for the 2,4- (Table 1, entries 3 and 4) disubstituted compounds. The decahydroquinoline isomers were particularly interesting, as the *trans* isomer is conformationally locked. These results already hinted at strict stereochemical requirements for effective enantioselective acylation.

In light of these initially confusing results, we undertook an extensive study to assess the role of the positional isomers and their stereoisomers. Our goal was to reliably predict whether a given substrate would undergo a selective kinetic resolution. We elected to evaluate at least three substrate pairs (*cis/trans*) for each of the three positional isomers, giving 20 substrates in total. Considerations for compound selection included ease of synthesis, the presence of groups for further functionalization of the products, and ready access to diastereomerically pure forms of the targets (See Supporting Information for experimental procedures and characterization data).

To rule out selectivity arising from the NHC-catalyst in the kinetic resolution, we used our previously reported stoichiometric variant in the resolution of these N-heterocycles.¹⁰ Slightly higher reaction rates and selectivities are observed with this reagent and a recyclable, resin supported variant makes it an attractive option for practical amine resolution.¹³ We first examined the kinetic resolution of the compounds in the 2,3-disubstitution series (Table 2). The data presented in Table 1 implied that substrates with a *cis* configuration undergo faster reaction with much higher selectivities than those observed for the *trans* isomers. With *cis*-2-phenylpiperidin-3-ol (Table 2, entry 1) an *s*-factor of 24 and a conversion of 33% could be obtained after 48 hours, whereas the *trans* diastereomer (Table 2, entry 2) reached only 14% conversion after 72 hours with essentially no selectivity. This trend could also be observed for the other two sets of substrates (Table 2, entries 3 and 4, entries 5 and 6). Decahydroquinolines showed a similar behavior in terms of reaction rate, although acceptable selectivity was obtained in both cases. In the resolution of *cis*-decahydroquinoline (Table 2, entry 7), 65% conversion was obtained after 15 hours whereas the conformationally locked *trans*-decahydroquinoline (Table 2, entry 8) reacted much more slowly. Notably, the opposite sense of induction was observed with this substrate, which contradicted a simple stereochemical model based only on the stereochemistry of the acyl transfer reagent (*vide infra*). The kinetic resolution reactions were stopped once the stoichiometric reagent was fully consumed.¹⁴

For the 2,4-disubstituted piperidines (Table 3), the trend was reversed in comparison to that observed for the 2,3-disubstituted examples in Table 2. For this series, the *cis*-diastereomers exhibit poor selectivities (*s* = 3–7) and low reactivity with reaction times between 72 and 120 hours (Table 3, entries 1, 3, and 5). In contrast, the *trans*-diastereomers reached full conversion in 20 hours or less with high selectivities (*s* = 10–29) (Table 3, entries 2, 4, and 6).

In the case of the 2,5-disubstituted piperidines (Table 4) we observed a similar trend as with the 2,3-disubstituted substrates. Namely, higher *s*-factors were observed for the *cis*-diastereomers as well as higher reaction rates compared to the corresponding *trans*-

diastereomers. The biggest contrast with regards to *s*-factors can be noticed in entry 5; for *cis*-6-propylpiperidin-3-ol an *s*-factor of 52 was obtained while the *trans*-isomer (Table 4, entry 6) afforded an *s*-factor of only 4. Such large differences in selectivity were not evident in entries 1–4; however, the data supports the conjecture that *cis* isomers in this series are superior to the *trans* isomers. These substrates also demonstrate the chemoselectivity of the kinetic resolution for N-heterocycles bearing a variety of functional groups suitable for further functionalization.

The divergent reactivity of the diastereomeric pairs in the different series can be rationalized on the basis of our recent DFT calculations of the transition states of α -methyl piperidine reacting with the chiral acylating agent.¹² These studies concluded that acylation occurred via a concerted 7-membered transition state with concomitant proton transfer guided by the hydroxamic acid and with the α -substituent placed in the axial position. In contrast, other work from our group on the resolution of morpholines with stoichiometric chiral acyl donors and achiral hydroxamic acids was most consistent with a stereochemical model featuring an equatorial substituent of a morpholine.¹⁶ However, no computational studies were performed on this system and significant conformational differences between morpholine and piperidines can be expected. Both models suggest that the stereoselective step involves a seven-membered transition state with the lone pair of the nitrogen in the equatorial position. This mechanism of acyl transfer was found to be general for a variety of widely used acyl transfer reagents including *N*-hydroxysuccinimide, and HOAt.

In order to reconcile these two models with the results obtained for the resolution of disubstituted piperidines, we undertook detailed DFT studies on the relevant transition states using the *cis* and *trans* geometric isomers of 2,3-dimethylpiperidine as a model system and the chiral hydroxamic acid.¹⁷ All calculations were carried out using the same computational methods previously employed in related systems using Gaussian 09.¹⁸ Structures were optimized with B3LYP/6-31G(d).^{19,20} Single-point energy calculations in the condensed phase (CH₂Cl₂; ϵ = 8.93) were undertaken using the IEPCM solvation model with M06-2X/6-311+G(d,p).²¹ Scheme 2 shows the lowest energy transition states for acyl transfer to each enantiomer of *cis* and *trans* 2,3-dimethyl piperidine. As anticipated from our previous work, the concerted 7-membered transition states feature an intramolecular proton shuttle between the secondary amine and the hydroxamate carbonyl. The calculations are in agreement with experimental rates and selectivity trends listed in Table 2 (entries 1–6) in which the *cis* 2,3-disubstituted piperidines showed significantly faster rates and superior *s*-factors (*s* = up to 24) in comparison to the *trans* 2,3-disubstituted piperidines. For example, the lowest transition state free energy for the **II-cis** (25.8 kcal/mol gas phase) is approximately 3 kcal/mol lower in energy than the lowest for **II-trans** (29.1 kcal/mol gas phase), which is consistent with the relative rates of these two amine diastereomers.

This finding is in line with our postulate that the lower energy transition states for acyl transfer feature an axial α -substituent, which requires the *trans*-2,3-dimethyl piperidine to adopt an unfavorable di-axial conformation. In contrast, the chair conformation of the *cis*-2,3-dimethylpiperidine with an α -axial substituent is the preferred ground state conformation. Further, our calculations correctly predict the fastest reacting enantiomer via **II-trans-TS** and **II-cis-TS**, respectively. In agreement with experiment, the quantum

mechanical calculations indicate a larger gap between the *cis*-1,2-disubstituted systems (here modeled as 2,3-dimethyl piperidine **II-cis** and *ent*-**II-cis**) than the *trans*-1,2-disubstituted systems, which reflects the significant difference in selectivity between *cis* and *trans* diastereomers.²² Notably, the lowest energy diastereomeric transition states place the α -substituent away from both the aromatic ring of the acyl transfer agent and the acyl chain. Exhaustive conformational analysis, including other chair conformations, revealed that all other transition states are much higher in energy (see Figure 1 and Supporting Information). For kinetic resolution, the energy difference between the diastereomeric acylation transition states determines the selectivity. As shown in Figure 1, the significant energy difference (3.7 kcal/mol, $s = 19$ –24) between **II-cis-TS** and *ent*-**II-cis-TS** arises from the additional unfavorable *gauche* interactions between α -substituent in the equatorial position and the carbonyl group. In addition, *ent*-**II-cis-TS** has an further interaction between the axial β -methyl group and the hydroxamate carbonyl.

In contrast, the energy difference between **II-trans-TS** and *ent*-**II-trans-TS** is small at 0.5 kcal/mol, which is consistent with the poor s -factors ($s = 1$ –4) observed for the *trans* substituted systems. It is remarkable, however, that the preferred conformation for the lowest energy transition state is the *trans*-2,3-bis-axial, which demonstrates again the strong preference for α -axial substituents in the acylation step and offers an excellent example of the Curtin-Hammett principle (Figure 1, bottom). The dimethyl groups are in the axial position in **II-trans-TS** and in the equatorial position in *ent*-**II-trans-TS**; in this case, the additional 1,3-diaxial interactions in **II-trans-TS** raise its energy leading to a smaller energy gap between the two enantiomers and smaller selectivity values.

The disparate results seen for the decahydroquinolines serve to further confirm this model. The *trans*-decahydroquinoline leads to the opposite sense of induction relative to the *cis*-decahydroquinoline even though the same configuration of the chiral acylating reagent is employed. Since the *trans*-decahydroquinoline cannot react in the conformation where the α -substituent is axial, the only transition states that are accessible have the α -substituent in equatorial position (Figure 1, lower right). When only these two transition states are considered, selectivity is expected for the *opposite enantiomer* due to the large calculated energy gap. Further, the reaction rate of *trans*-decahydroquinoline is expected to be slower, as observed experimentally, since the overall, energy of this pathway is higher.

The same results can be applied to 2,5-disubstituted piperidines. Just as with the 2,3-disubstituted compounds, the *cis*-diastereomer consists of two conformers that differ only slightly in energy whereas the *trans*-diastereomer exists primarily in the diequatorial conformer rather than the higher energy diaxial conformation. An analogous situation occurs for the 2,4-disubstituted compounds; however, it is the *trans* isomer that always has one axial substituent in the lowest energy conformations, and is more selective. The *cis*-2,4-disubstituted isomers, in contrast, exist almost exclusively in the more favorable diequatorial conformation and gives rise to lower selectivity.

This analysis suggests that disubstituted piperidines with a single chiral center and no large conformational preferences between the axial and equatorial α -substituents should be good substrates for the resolution. This hypothesis is confirmed by the excellent results obtained

with substrates bearing an exocyclic 4-methylene and a single α -substituent (Table 5, entries 1–3). In addition 2,4,4-trisubstituted piperidines containing a cyclic ketal were also resolved with modest selectivity (Table 5, entries 4 and 5). All five substrates evaluated show fast reaction rates and useful selectivity, indicating no adverse affect on the resolution of substrates containing additional achiral substitution. The alkene or protected carbonyl in the products provide a means for post-resolution derivatization.

In summary we have examined the kinetic resolution of disubstituted piperidines using a chiral hydroxamic acid catalyst and its corresponding stoichiometric reagent. For most cases, synthetically useful relative rates and outstanding functional group tolerance were observed. The exceptions – stereoisomeric substrates that gave poor reactivity and selectivity – revealed a pronounced conformational preference in which the lowest energy transition states require that the α -substituent populates the axial position. This conjecture was fully supported by DFT calculations on the transition states for all of the relevant stereoisomers and ring conformations. These finding both rationalized the initially unexpected differences in reactivity and selectivity between *cis* and *trans* stereoisomers and strengthen our stereochemical model.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the European Research Council (ERC Starting Grant No. 306793 – CASAA) and National Institutes of Health (GM-112684 to M.C.K.) and the National Science Foundation (CHE1464778 to M.C.K.). Computational support was provided by XSEDE on SDSC Gordon (TG-CHE120052). Computational support was provided by the XSEDE on SDSC Gordon (TG-CHE120052). We thank LOC MS Service (ETH Zürich) for analyses and Dr. Nils Trapp and Dr. Bernd Schweizer for the acquisition of X-ray structures. We are also grateful to Dr. Kimberly Geoghegan, Sheng-Ying Hsieh and Dr. Hidetoshi Noda for helpful discussions.

References

1. Lovering F, Bikker J, Humblet C. J Med Chem. 2009; 52:6752–6756. [PubMed: 19827778]
2. For reviews on the a functionalisation of unsubstituted heterocycles see:(a) Mitchell EA, Peschiulli A, Lefevre N, Meerpoal L, Maes BUW. Chem Eur J. 2012; 18:10092–10142. [PubMed: 22829434] (b) Campos KR. Chem Soc Rev. 2007; 36:1069–1084. [PubMed: 17576475] (c) Vo CT, Bode JW. J Org Chem. 2014; 79:2809–2815. [PubMed: 24617516]
3. (a) Millet A, Larini P, Clot E, Baudoin O. Chem Sci. 2013; 4:2241–2247.(b) Cordier CJ, Lundgren RJ, Fu GC. J Am Chem Soc. 2013; 135:10946–10949. [PubMed: 23869442] (c) McNally A, Prier CK, MacMillan DWC. Science. 2011; 334:1114–1117. [PubMed: 22116882]
4. (a) Comins DL, Abdullah AH. J Org Chem. 1982; 47:4315–4319.(b) Chau ST, Lutz JP, Wu K, Doyle AG. Angew Chem Int Ed. 2013; 52:9153–9156.
5. (a) Cai W, Colony JL, Frost H, Hudspeth JP, Kendall PM, Krishnan AM, Maskowski T, Mazur DJ, Phillips J, Brown Ripin DH, Gut Ruggeri S, Stearns JF, White TD. Org Process Res Dev. 2005; 9:51–56.(b) Ye Z-S, Chen M-W, Chen Q-A, Shi L, Duan Y, Zhou Y-G. Angew Chem Int Ed. 2012; 51:10181–10184.(c) Wang D-S, Chen Q-A, Lu S-M, Zhou Y-G. Chem Rev. 2012; 112:2557–2590. [PubMed: 22098109] (d) Imuro A, Yamaji K, Kandula S, Nagano T, Kita Y, Mashima K. Angew Chem Int Ed. 2013; 52:2046–2050.(e) Romanov-Michailidis F, Sedillo KF, Neely JM, Rovis T. J Am Chem Soc. 2015; 137:8892–8895. [PubMed: 26154248]
6. (a) Pablo O, Guijarro D, Yus M. J Org Chem. 2013; 78:9181–9189. [PubMed: 23952598] (b) Launay GG, Slawin AMZ, O'Hagan D. Beilstein J Org Chem. 2010; 6:41. [PubMed: 20502655]

7. (a) Chen S, Mercado BQ, Bergman RG, Ellman JA. *J Org Chem*. 2015; 80:6660–6668. [PubMed: 26098485] (b) Duttwyler S, Chen S, Lu C, Mercado BQ, Bergman RG, Ellman JA. *Angew Chem Int Ed*. 2014; 53:3877–3880. (c) Ischay M, Takase MK, Bergman RG, Ellman JA. *J Am Chem Soc*. 2013; 135:2478–2481. [PubMed: 23398467] (d) Duttwyler S, Chen S, Takase MK, Wiberg KB, Bergman RG, Ellman JA. *Science*. 2013; 339:678–682. [PubMed: 23393259]
8. (a) Mittal N, Lippert KM, Kanta De C, Klauber EG, Emge TJ, Schreiner PR, Seidel D. *J Am Chem Soc*. 2015; 137:5748–5758. [PubMed: 25871925] (b) Kanta De C, Klauber EG, Seidel D. *J Am Chem Soc*. 2009; 131:17060–17061. [PubMed: 19929016] (c) Klauber EG, Kanta De C, Shah TK, Seidel D. *J Am Chem Soc*. 2010; 132:13624–13626. [PubMed: 20843041] (d) Yang X, Bumbu VD, Liu P, Li X, Jiang H, Uffman EW, Guo L, Zhang W, Jiang X, Houk KN, Birman VB. *J Am Chem Soc*. 2012; 134:17605–17612. [PubMed: 23030692] (e) Fu GC. *Acc Chem Res*. 2000; 33:412–420. [PubMed: 10891059] (f) Fu GC. *Acc Chem Res*. 2004; 37:542–547. [PubMed: 15311953] (g) Fowler BS, Mikochick PJ, Miller SJ. *J Am Chem Soc*. 2010; 132:2870–2871. [PubMed: 20158213] (h) Birman VB, Uffman EW, Jiang H, Li X, Kilbane CJ. *J Am Chem Soc*. 2004; 126:12226–12227. [PubMed: 15453730] (i) Li X, Liu P, Houk KN, Birman VB. *J Am Chem Soc*. 2008; 130:13836–13837. [PubMed: 18817392] (j) Brown MK, Blewett MM, Colombe JR, Corey EJ. *J Am Chem Soc*. 2010; 132:11165–11170. [PubMed: 20666410] (k) Larionov E, Mahesh M, Spivey AC, Wei Y, Zipse H. *J Am Chem Soc*. 2012; 134:9390–9399. [PubMed: 22568686]
9. (a) Keith JM, Larrow JF, Jacobsen EN. *Adv Synth Catal*. 2001; 343:5–26. (b) Breuer M, Dittrich K, Habicher T, Hauer B, Keßeler M, Stürmer R, Zelinski T. *Angew Chem*. 2004; 116:806–843. *Angew. Chem. Int. Ed* **2004**, 43, 788–824.
10. Binanzer M, Hsieh SY, Bode JW. *J Am Chem Soc*. 2011; 133:19698–19701. [PubMed: 22082205]
11. Hsieh S-Y, Binanzer M, Kreituss I, Bode JW. *Chem Commun*. 2012; 48:8892–8894.
12. Allen SE, Hsieh S-Y, Gutierrez O, Bode JW, Kozlowski MC. *J Am Chem Soc*. 2014; 136:11783–11791. [PubMed: 25050843]
13. Kreituss I, Murakami Y, Binanzer M, Bode JW. *Angew Chem Int Ed*. 2012; 51:10660–10663.
14. For the purposes of method development and study we performed the kinetic resolutions under conditions that allow for accurate calculations of the relative rates. At longer reaction times, the hydroxyenone and stoichiometric reagent can undergo side reactions. This protocol explains the low conversion even with full consumption of the reagent with slow-reacting substrates. For preparative work, the resolutions are allowed to proceed to higher conversion, which affords virtually enantiopure recovered amines. As a guideline, $s = 20$ can deliver >99% ee at 62% conversion. See Vedejs E, Jure M. *Angew Chem*. 2005; 117:4040–4069. *Angew. Chem. Int. Ed*. **2005** **44**, 3974–4001 for a discussion.
15. Calculated according to: Kagan HB, Fiaud JC. *Topic in Stereochemistry*. 1988; 18:249–330.
16. Hsieh S-Y, Wanner B, Wheeler P, Beauchemin AM, Rovis T, Bode JW. *Chem Eur J*. 2014; 20:7228–7231. [PubMed: 24839065]
17. All calculations were carried out using Gaussian 09, Revision D.01
18. Frisch, MJ. Gaussian 09, Revision D.01. Gaussian, Inc; Wallingford CT: 2013. Gaussian, Inc.; Wallingford, CT, 2013
19. (a) Becke AD. *J Chem Phys*. 1993; 98:5648–5652. (b) Lee C, Yang W, Parr RG. *Phys Rev B*. 1988; 37:785–789. (c) Vosko SH, Wilk L, Nusair M. *Can J Phys*. 1980; 58:1200.
20. (a) Ditchfield R, Hehre WJ, Pople JA. *J Chem Phys*. 1971; 54:724–728. (b) Frisch MJ, Pople JA, Binkley JS. *J Chem Phys*. 1984; 80:3265–3269.
21. (a) Cancès E, Mennucci B, Tomasi J. *J Chem Phys*. 1997; 107:3032–3041. (b) Zhao Y, Truhlar DG. *Theor Chem Acc*. 2008; 120:215–241.
22. Additional calculations using *cis* and *trans*-2-phenyl-3-hydroxypiperidine showed a similar trend as the model, *cis* and *trans* 2,3-dimethylpiperidine. For instance the enthalpic energy difference (in solvent) between *cis*-2-phenylpiperidin-3-ol was computed to be 1.7 kcal/mol vs only 0.4 kcal/mol for the corresponding *trans* diastereomer. This calculation is in excellent agreement with experiment for kinetic resolution of the *cis* and *trans*-2-phenylpiperidin-3-ol systems (Table 2, entries 1 and 2) which exhibited a s -values of 24 and 1, respectively.

Figure 1.

Proposed reaction pathways for possible conformers/enantiomers in the kinetic resolution. Relative acyl transfer reaction barriers (free energies are in kcal/mol; gas phase energetics calculated using B3LYP/6-31G(d,p)/(in parenthesis IEPCM-CH₂Cl₂-M06-2X/6-311+G(d,p)) for the competing diastereomeric transition states between *cis* and *trans* 2,3-dimethyl substituted piperidines and the acyl transfer agent.

Scheme 1.

Four possible positional isomers of disubstituted piperidines that serve as targets for our study.

Scheme 2. Relative acyl transfer reaction barriers

Relative acyl transfer reaction barriers (free energies are in kcal/mol; gas phase energetics calculated using B3LYP/6-31G(d,p)//(in parenthesis IEPCM-CH₂Cl₂-M06-2X/6-311+G(d,p)) for the competing diastereomeric transition states between *cis* and *trans* 2,3-dimethyl substituted piperidines and the acyl transfer agent.

Table 1

Catalytic Kinetic resolution of disubstituted piperidines.

Entry	Substrate	<i>s</i> ^a	Conv. ^b (%)	reaction time ^c (h)	er amine ^d yield ^e (%)	er amide ^d yield ^e (%)
1		11	28	72	64:36 (43)	89:11 (25)
2		2	3	72	51:49 (58)	67:33 (5)
3		3	4	48	51:49 (43)	78:22 (4)
4		21	59	20	99:1 (28)	83:17 (26)
5		19	17	20	59:41 (35)	94:6 (20)
6		8	5	48	52:48 (54)	89:11 (9)
7		18	63	48	99:1 (20)	79:21 (40)

Entry	Substrate	<i>s</i> ^a	Conv. ^b (%)	reaction time ^c (h)	er amine ^d yield ^e (%)	er amide ^d yield ^e (%)
8 ^f		9	11	65	45:55 (61)	10:90 (9)
9		19	43	20	81:19 (40)	91:9 (42)
10		14	41	20	77:23 (43)	89:11 (40)
11		21	55	20	96:4 (42)	87:13 (43)

^a Calculated selectivity.¹⁵^b Calculated conversion.¹⁵^c Reaction time until the α-hydroxyketone is fully consumed.^d Determined by SFC or HPLC on a chiral support.^e Isolated as the Cbz-derivative. Yield after column chromatography.^f Opposite sense of induction observed. RMesClO₄ = (2,4,6-trimethylphenyl)-2,5,6,7-tetrahydro-pyrrolo[2,1-c][1,2,4]triazol-4-ylm perchlorate.

Table 2

Kinetic resolution of 2,3-disubstituted piperidines with stoichiometric reagent 1.

Entry	Substrate	<i>s</i> ^a	Conv. <i>b</i> (%)	reaction time ^c (h)	er amine ^d yield ^e (%)	er amide ^d yield ^e (%)
1		24	33	48	72:28 (31)	94:6 (30)
2		1	14	72	51:49 (32)	56:44 (5)
3		19	31	72	69:31 (34)	93:7 (28)
4		2	46	96	56:44 (39)	57:43 (7)
5		23	50	72	90:10 (39)	91:9 (50)
6		4	26	96	59:41 (40)	75:25 (21)
7 ^f		20	65	15	99:1 (17)	73:27 (50)
8 ^g		20	36	48	24:76 (25)	7:93 (30)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^a Calculated selectivity.^b Calculated conversion.^c Reaction time until stoichiometric reagent is fully consumed.^d Determined by SFC or HPLC on a chiral support.^e Isolated as the Cbz-derivative. Yield after column chromatography.^f 0.6 equiv of stoichiometric reagent **1** were used.^g Opposite sense of induction observed.

Table 3

Kinetic resolution of 2,4-disubstituted piperidines with stoichiometric reagent 1.

Entry	Substrate	<i>s</i> ^a	Conv. ^b (%)	reaction time ^c (h)	er amine ^d yield ^e (%)	er amide ^d yield ^e (%)
1		3	22	120	57:43 (39)	78:22 (4)
2f		10	65	20	97:3 (15)	75:25 (43)
3		7	32	72	66:34 (56)	84:16 (29)
4		29	52	20	94:6 (46)	91:9 (45)
5		6	38	96	68:32 (46)	80:20 (34)
6f		15	62	20	98:2 (19)	80:20 (40)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^a Calculated selectivity.^b Calculated conversion.^c Reaction time until stoichiometric reagent is fully consumed.^d Determined by SFC or HPLC on a chiral support.^e Isolated as the Cbz-derivative. Yield after column chromatography.^f 0.6 equiv of stoichiometric reagent **1** were used.

Table 4

Kinetic resolution of 2,5-disubstituted piperidines with stoichiometric reagent 1.

Entry	Substrate	<i>s</i> ^a	Conv. ^b (%)	reaction time ^c (h)	er amine ^d yield ^e (%)	er amide ^d yield ^e (%)
1		22	42	24	81:19 (44)	92:8 (33)
2		20	40	96	78:22 (47)	92:8 (33)
3		13	54	72	90:10 (22)	84:16 (28)
4		9	29	72	65:35 (45)	87:13 (19)
5		52	54	20	99:1 (25)	92:2 (38)
6		4	53	48	74:26 (29)	71:29(51)

^a Calculated selectivity.^b Calculated conversion.^c Reaction time until stoichiometric reagent is fully consumed.^d Determined by SFC or HPLC on a chiral support.

Isolated as the Cbz-derivative. Yield after column chromatography.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 5

Kinetic resolution of 2,4,4-trisubstituted piperidines with stoichiometric reagent 1.

Entry	Substrate	<i>s^a</i>	Conv. <i>b</i> (%)	reaction time ^c (h)	er amine ^d yield ^e (%)	er amide ^d yield ^e (%)
1		27	51	20	92:8 (45)	91:9 (43)
2		18	49	20	88:12 (46)	89:11 (43)
3 ^f		18	56	20	95:5 (39)	86:14 (54)
4		8	48	20	79:21 (25)	81:19 (44)
5 ^f		9	64	20	94:6 (23)	75:25 (21)

^a Calculated selectivity.^b Calculated conversion.^c Reaction time until stoichiometric reagent is fully consumed.^d Determined by SFC or HPLC on a chiral support.

Isolated as the Cbz-derivative. Yield after column chromatography,

0.6 equiv of stoichiometric reagent **1** were used.