Published in final edited form as:

Org Lett. 2007 April 26; 9(9): 1737–1740. doi:10.1021/ol070435s.

Fischer carbene catalysis of alkynol cycloisomerization: Application to the synthesis of the altromycin B disaccharide

BonSuk Koo and Frank E. McDonald

Department of Chemistry, Emory University, Atlanta, GA 30322

Frank E. McDonald: fmcdona@emory.edu

Abstract

The tungsten-catalyzed cycloisomerization of alkynyl alcohols can be conducted without using photochemistry, using a stable tungsten Fischer carbene as the precatalyst for this transformation. A variety of alkynyl alcohols undergo cycloisomerization under these conditions to provide endocyclic enol ethers of 5, 6, and 7-membered ring sizes. The utility of this method is further demonstrated in the stereoselective synthesis of the disaccharide substructure of altromycin B.

The importance of cyclic enol ethers in synthetic organic chemistry has been demonstrated in several contexts, including the efficient synthesis of a variety of biologically important carbohydrate and oligosaccharide-containing natural products, in which carbohydrate-derived cyclic enol ethers (glycals) are valuable electrophilic partners for a variety of glycoside synthesis processes.¹ Our laboratory has developed group VI transition metal-catalyzed processes for the synthesis of structurally diverse carbohydrates arising from non-carbohydrate-derived alkynyl alcohols, which has greatly expanded the scope of available substrates for glycal-based approaches to glycoconjugate synthesis.² The various methods reported by our laboratory for catalytic cycloisomerization have utilized photochemical activation of molybdenum or tungsten carbonyl in the presence of a tertiary amine, with molybdenum catalysis preferred for the formation of five-membered rings³ and tungsten catalysis proving more effective for the synthesis of six⁴- and seven-membered ring cyclic enol ethers⁵ (Figure 1). However, we are aware that the requirement for photochemical activation has limited the exploration and application of this methodology to laboratories that have specialized photochemical equipment. Thus we describe herein a new catalytic system which does not require photochemical activation, based on (methoxymethyl-carbene)pentacarbonyl tungsten (**3**)⁶ as a stable and easily prepared precatalyst for the

Correspondence to: Frank E. McDonald, fmcdona@emory.edu.

Supporting Information **Available**: Experimental procedures, characterization and spectral data on new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

cycloisomerization of various alkynyl alcohols. Our design of the non-photochemical system is predicated on the reactivity of alkyl-substituted Fischer carbenes to undergo deprotonation and in some cases demetallation to vinylic ethers, with amine-metal carbonyl complexes as the byproduct of base-promoted demetallation reactions.⁷

Optimization of conditions was initially conducted primarily with the simple aminoalkynol substrate **4**,⁸ for conversion into the cycloisomeric enol ether **5**. Optimal results were obtained with 25 mol % of tungsten oxacarbene **3** in the presence of 10 equivalents of triethylamine, and warming to 40 °C in THF solvent for 12 hours (Table 1, entry 1), at which time substrate **4** was completely consumed and **5** was produced in 92% isolated yield. The choice of triethylamine as base was critically important to the success of this transformation, as the use of 1,4-diazabicyclo[2.2.2]octane (DABCO)⁹ resulted in low conversion of substrate under non-photochemical conditions. Similar results were obtained with the diastereomeric substrate **6**, as well as several other more complex substrates **8**, **10**, and **12**,¹⁰ with products **9**, **11**, and **13** corresponding to protected glycals of D-acosamine,¹¹ L-vancosamine,¹² and D-saccharosamine,¹³ respectively. Substrate **14**, which differs from **8** only in the protective group pattern of the two oxygen substituents, underwent cycloisomerization under the same conditions to provide the five-membered ring enol ether **15**. With slight increases in temperature and reaction time as well as catalytic loading in some cases, high yield cycloisomerizations of alkynyldiols **16** and **18** to the seven-membered ring glycals **17** and **19** (corresponding to septanose glycals of D-glucal and D-galactal, respectively)^{5,14} were achieved.

Fischer carbene-catalyzed cycloisomerizations of alkynyl alcohols **20** with C3 and C4-oxygen substituents^{9b} required additional optimization relative to the photochemical procedure (Table 2), as substantial amounts of the exocyclic glycal were produced from substrate **20b**. As reported by others,¹⁵ sterically bulky propargylic substituents are required for high *endo*-regioselectivity, but we observed that reduced steric bulk in the O4 methoxymethyl (MOM) protective group required increasing the steric bulk at O3 to the triisopropylsilyl ether **20c** to obtain good *endo*-regioselectivity. The formation of exocyclic glycal **22b** from substrate **20b** was rationalized as the action of tungsten pentacarbonyl as a Lewis acid to promote cyclization to **25** (figure 2), but increasing the steric bulk at O3 in substrate **20c** would increase the steric interaction between η^2 -alkyne-tungsten pentacarbonyl and the O3-protective group, thus disfavoring *exo*-mode cyclization to **25** relative to rearrangement to the vinylidene intermediate **24**, resulting in the formation of glycal **21c** via endocyclic intermediate **26**.¹⁶

The utility of the non-photochemical conditions for tungsten-catalyzed alkynyl alcohol cycloisomerization is highlighted in the synthesis of the (*N,N*-dimethyl)-vancosamine- α -(3'-*O*-methyl)digitoxose disaccharide substructure of altromycin B (**27**, Scheme 1), a glycoconjugate natural product exhibiting *in vivo* activities against P388 leukemia as well as colon, lung, and ovarian tumors.^{17,18} Our synthetic strategy for the altromycin disaccharide relies mainly on iterative application of tungsten-catalyzed alkynol cycloisomerization for each of the carbohydrate moieties.^{2,9,19} The synthesis commenced with NIS-mediated stereoselective *trans*-diaxial glycosylation²⁰ of glycal **21c** with the hydroxy *beta*-lactam

28,^{10,21} followed by radical deiodination to produce *O*-glycoside **29**. The sequence of lactam opening to methyl ketone and exchange of *N*-protective groups to the less basic carbamate of **30**, followed by alkyne desilylation and Felkin-Anh controlled reduction under Luche conditions¹⁰ resulted in stereoselective synthesis of alkynyl alcohol substrate **31**. Cycloisomerization of **31** under the standard conditions developed for the simpler examples in table 1 provided the disaccharide glycal **32** in excellent yield, which was further converted into the *N,N*-dimethylamino-*O*-methyl ether **33**.

While each of the transformations described herein also proceeds by the photochemical method (in some cases with lower catalytic loading), these new conditions now mean that photochemical equipment is no longer required for tungsten-catalyzed cycloisomerization of alkynyl alcohols. In addition, our alkynol cycloisomerization procedure requires neither expensive transition metals nor specialized ligands, and is compatible with a broad range of functional groups.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institutes of Health (R01 CA59703). We thank Mary H. Davidson, Dr. Joseph M. Pletcher, and Matthew A. Boone (Emory University) for providing samples of some of the substrates or advanced synthetic precursors used in this project. We also thank Rui Cao and Dr. Kenneth I. Hardcastle (Emory University X-Ray Crystallography Laboratory) for providing the crystal structure of *beta*-lactam **28**.

References

1. (a) Danishefsky SJ, Roberge JY. *Pure Appl Chem.* 1995; 67:1647.(b) Danishefsky SJ, Bilodeau MT. *Angew Chem Int Ed.* 1996; 35:1380.(c) Seeberger PH, Bilodeau MT, Danishefsky SJ. *Aldrichimica Acta.* 1997; 30:75.(d) Williams, LJ.; Garbaccio, RM.; Danishefsky, SJ. *Carbohydrates in Chemistry and Biology.* Ernst, B.; Hart, GW.; Sinaý, P., editors. Vol. 1. Wiley-VCH; Weinheim: 2000. p. 61(e) Priebe, W.; Grynkiewicz, G. *Glycoscience: Chemistry and Biology.* Fraser-Reid, B.; Tatsuta, K.; Thiem, J., editors. Vol. 1. Springer; Berlin: 2001. p. 749
2. McDonald, FE. *Strategies and Tactics in Organic Synthesis.* Harmata, M., editor. Vol. 5. Elsevier; Amsterdam: 2004. p. 391
3. McDonald FE, Gleason MM. *J Am Chem Soc.* 1996; 118:6648.
4. McDonald FE, Reddy KS, Díaz Y. *J Am Chem Soc.* 2000; 122:4304.
5. Alcázar E, Pletcher JM, McDonald FE. *Org Lett.* 2004; 6:3877. [PubMed: 15469372]
6. Tungsten oxacarbene 3 is stable for several months in the dark Fischer EO, Maasböl A. *Angew Chem Int Ed.* 1964; 3:850.Lam CT, Malkiewicz CD, Senoff CV. *Inorg Synth.* 1977; 17:95.Matsuyama H, Nakamura T, Iyoda M. *J Org Chem.* 2000; 65:4796. [PubMed: 10956454]
7. (a) Kreiter CG. *Angew Chem Int Ed.* 1968; 7:390.(b) Fischer EO, Plabst D. *Chem Ber.* 1974; 107:3326.(c) Casey CP, Anderson RL. *J Chem Soc Chem Commun.* 1975; 895(d) Bernasconi CF. *Chem Soc Rev.* 1997; 26:299.
8. Davidson MH, McDonald FE. *Org Lett.* 2004; 6:1601. [PubMed: 15128246]
9. In contrast, DABCO often gives better results vs. triethylamine in the photoactivated catalytic system McDonald FE, Reddy KS. *J Organometal Chem.* 2001; 617-618:444.McDonald FE, Reddy KS. *Angew Chem Int Ed.* 2001; 40:3653.
10. Cutchins WW, McDonald FE. *Org Lett.* 2002; 4:749. [PubMed: 11869118]

11. (a) Hauser FM, Ellenberger SR. *Chem Rev.* 1986; 86:35.(b) Baer HH, Georges FFZ. *Can J Chem.* 1977; 55:1100.(c) Ginesta X, Pastó M, Pericàs MA, Riera A. *Org Lett.* 2003; 5:3001. [PubMed: 12916966]
12. (a) Smith RM, Johnson AW, Guthrie RD. *J Chem Soc Chem Commun.* 1972; 361(b) Trost BM, Jiang C, Hammer K. *Synthesis.* 2005; 3335(c) Hsu DS, Matsumoto T, Suzuki K. *Synlett.* 2006; 469
13. Kong F, Zhao N, Siegel M, Janota K, Ashcroft JS, Koehn FE, Borders DB, Carter GT. *J Am Chem Soc.* 1998; 120:13301.
14. (a) Snyder NL, Haines HM, Peczu MW. *Tetrahedron.* 2006; 62:9301.(b) Ng CJ, Stevens JD. *Carbohydr Res.* 1996; 284:241.
15. (a) Wipf P, Graham TH. *J Org Chem.* 2003; 68:8798. [PubMed: 14604347] (b) Moilanen SB, Tan DS. *Org Biomol Chem.* 2005; 3:798. [PubMed: 15731865]
16. (a) Sheng Y, Musaev DG, Reddy KS, McDonald FE, Morokuma K. *J Am Chem Soc.* 2002; 124:4149. [PubMed: 11942854] (b) Nowroozi-Isfahani T, Musaev DG, McDonald FE, Morokuma K. *Organometallics.* 2005; 24:2921. [PubMed: 16601823] (c) Sordo T, Campomanes P, Diéguez A, Rodriguez F, Fañanás FJ. *J Am Chem Soc.* 2005; 127:944. [PubMed: 15656633]
17. (a) Jackson M, Karwowski JP, Theriault RJ, Hardy DJ, Swanson SJ, Barlow GJ, Tillis PM, McAlpine JB. *J Antibiot.* 1990; 43:223. [PubMed: 2324007] (b) Brill GM, McAlpine JB, Whittern DN, Buko AM. *J Antibiot.* 1990; 43:229. [PubMed: 2324008]
18. For recent synthetic efforts towards substructures of altromycin B Pasetto P, Franck RW. *J Org Chem.* 2003; 68:8042. [PubMed: 14535781] Fei Z, McDonald FE. *Org Lett.* 2005; 7:3617. [PubMed: 16092833] Koo B, McDonald FE. *Org Lett.* 2005; 7:3621. [PubMed: 16092834]
19. (a) McDonald FE, Zhu HYH. *J Am Chem Soc.* 1998; 120:4246.(b) McDonald FE, Wu M. *Org Lett.* 2002; 4:3979. [PubMed: 12599507] (c) Trost BM, Rhee YH. *J Am Chem Soc.* 2002; 124:2528. [PubMed: 11890802]
20. Thiem J, Ossowski P, Schwentner J. *Chem Ber.* 1980; 113:955.
21. Resolution of racemic beta-lactam 28 is described in the Supporting Information. For different applications of tungsten-catalyzed cycloisomerization methodology applied to aminoglycal synthesis, see: Parker KA, Chang W. *Org Lett.* 2005; 7:1785. [PubMed: 15844906] Parker KA, Chang W. *Org Lett.* 2003; 5:3891. [PubMed: 14535736]

Figure 1. General concepts for generation of $(\text{Et}_3\text{N})\text{W}(\text{CO})_5$ as a catalyst for alkynyl alcohol cycloisomerization

Figure 2. Proposal for substituent effects on regioselectivity

Scheme 1. Synthesis of the altromycin disaccharide

Table 1
Representative examples of alkynyl alcohol cycloisomerizations catalyzed by 3

entry	alkynyl alcohol	enol ether	time	% yield
1			12 h	92
2			12 h	95
3			12 h	95
4			12 h	84
5			12 h	94
6			12 h	77
7			24 h (60 °C)	82 ^{a,b}

entry	alkynyl alcohol	enol ether	time	% yield
8	18	19	24 h (60 °C)	82 ^b

^a with 40 mol % **3**.

^b after treatment with Ac₂O, Et₃N, cat. DMAP.

Table 2
Comparison of non-photochemical procedure catalyzed by 3, vs. photochemical procedures

entry	substrate	conditions ^a	% yield 21	% yield 22
1	20a	A	53 ^b	0 ^b
2	20a	B	92	0
3	20b	A	53	35
4	20b	B	65	20
5	20b	C	75	15
6	20c	A	85	9
7	20c	B	88	7
8	20c	C	88	6

^a A: 40 mol % **3**, 10 equiv Et₃N, THF, 60 °C, 24 h.

B: 25 mol % W(CO)₆, 10 equiv Et₃N, THF, hv (350 nm), 60 °C, 6 h.

C: 25 mol % W(CO)₆, 2 equiv DABCO, THF, hv (350 nm), 60 °C, 6 h

^b 30% of **20a** was recovered.