Published in final edited form as:

Curr Heart Fail Rep. 2015 August ; 12(4): 284–293. doi:10.1007/s11897-015-0263-7.

Autonomic Regulation Therapy in Heart Failure

Una Buckley, MD, Kalyanam Shivkumar, MD PhD, and Jeffrey L. Ardell, PhD

Cardiac Arrhythmia Center & Neurocardiology Research Center of Excellence, UCLA David Geffen School of Medicine, 100 Medical Plaza, Suite 660, Los Angeles, CA 90095

Abstract

Autonomic Regulation Therapy (ART) is a rapidly emerging therapy in the management of congestive heart failure secondary to systolic dysfunction. Modulation of the cardiac neuronal hierarchy can be achieved with bioelectronics modulation of the spinal cord, cervical vagus, baroreceptor, or renal nerve ablation. This review will discuss relevant preclinical and clinical research in ART for systolic heart failure. Understanding mechanistically what is being stimulated within the autonomic nervous system by such device-based therapy and how the system reacts to such stimuli is essential for optimizing stimulation parameters and for the future development of effective ART.

Keywords

Heart failure; sympathetic modulation; vagus nerve stimulation; spinal cord stimulation; baroreceptor stimulation; renal denervation

Introduction

In the United States, 1 in 9 deaths each year is related to heart failure with an estimated 5.1 million people affected ¹. The cost of heart failure is thought to be about 32 billion dollars per year as a result of costs to the health service, medications, and absenteeism in the work force ². Heart failure continues to have a high mortality rate, despite advances in pharmacological and device therapy, with the Framingham study reporting a median 5 year

Corresponding Author: Jeffrey L. Ardell, Ph.D., Department of Medicine (Cardiology), UCLA Health System, 47-126 CHS, Los Angeles, CA 90095, 310-825-0417, jardell@mednet.ucla.edu.

Kalyanam Shivkumar MD PhD Professor of Medicine, Radiology & Bioengineering Director UCLA Cardiac Arrhythmia Center & EP Programs Director & Chief Interventional Cardiovascular Programs UCLA Health System Los Angeles, CA 90095
KShivkumar@mednet.ucla.edu

Una Buckley MD Electrophysiology Research Fellow UCLA Cardiac Arrhythmia Center UCLA Health System Los Angeles, CA 90095
UBuckley@mednet.ucla.edu

Author contributions: All authors contributed equally to researching data for this article, writing the manuscript, discussions of content and approved final version before submission.

Compliance with Ethics Guidelines

Conflict of Interest: Jeffrey L. Ardell serves as a scientific advisor to Cyberonics, Inc Kalyanam Shivkumar and Una Buckley declare that they have no conflict of interest.

Human and Animal Rights and Informed Consent

All procedures performed in studies involving human participants were in accordance with the ethical standards of UCLA, the National Institutes of Health and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards].

All procedures performed in studies involving animals were in accordance with the ethical standards of East Tennessee State University and University of California Los Angeles.

survival of 25% in men and 38% in women³. Early diagnosis and treatment may impact the morbidity and mortality⁴. Given the aging population, increase in obesity, diabetes mellitus, and sedentary lifestyle, it is thought that these statistics largely underestimate the prevalence⁵⁻⁷. While these statistics include all forms of heart failure, this review will specifically focus on autonomic modulating device therapy in the setting of heart failure with reduced ejection fraction.

Structure/function organization of the cardiac neuronal hierarchy

Cardiac control is achieved through a hierarchal network that may be considered in three-levels⁸⁻¹⁰. Level 1: CNS neurons (medullary and spinal cord neurons modulated by higher centers); Level 2: Peripheral: extracardiac-intrathoracic neuronal pool; and Level 3: Peripheral: the intrinsic cardiac nervous (ICN) system (Figure 1).

The peripheral layers (Levels 2 and 3) form cardio-centric loops, while the CNS (Level 1) engages neural mechanisms for cardiac and peripheral vasculature regulation^{9, 10}. Acting together, these hierarchical populations coordinate and regulate regional cardiac electrical and mechanical indices throughout each cardiac cycle to assure that cardiac output matches blood flow demands¹¹⁻¹³. To understand network interactions within and between levels 1-3, one must first understand the characteristics of its constituent parts. It is through such understanding that rational neuromodulation therapies can be devised.

Afferent neurons—Afferent neurons associated with cardiac and major vascular sensory neurites transduce the local mechanical and/or chemical milieu of these tissues¹⁴. The somata of these cardiac and vascular sensory neurons are located in: i) nodose ganglia; ii) dorsal root ganglia; iii) intrathoracic extracardiac ganglia; and iv) intrinsic cardiac ganglia¹⁴. The majority of these somata transduce mechanical distortion, the chemical milieu or both to second order neurons in the CNS and periphery¹⁴⁻¹⁶. Cardiac sensory information is fed into each of the three ‘levels’ of the neuronal hierarchy, these sensory inputs are a primary initiator of responses within and between the different levels of the neuraxis^{10, 17, 18}.

Efferent neurons—Efferent neuronal outflow from the autonomic nervous system to the heart depends on central and peripheral mediated neural reflexes^{10, 19, 20}. Cardiac-related parasympathetic efferent preganglionic somata are localized at medullary sites (primarily the Nucleus Ambiguus [NA]) and project to post-ganglionic neurons within intrinsic cardiac ganglia²¹⁻²⁴. Sympathetic preganglionic efferents for cardiac control are localized at the intermedio-lateral cell column of the spinal cord (T1-T5)²⁴. They project to postganglionic neurons located in stellate, middle cervical, superior cervical, mediastinal and intrinsic cardiac ganglia^{24, 25}. Within each nexus point of the neuronal hierarchy for cardiac control, from the CNS to intrinsic cardiac ganglia, network interactions within and between levels is fundamental to network output function (Fig. 1).

Local circuit neurons (LCN)—LCNs are neurons that are not directly transducing cardiac indices (cardiac afferent neurons) or having direct motor function, but clearly play a role to integrate sensory inputs along with inputs from the central nervous system. These

neurons are located throughout all intrathoracic ganglia, including those distributed on the heart^{10, 12, 26}. They receive sensory feedback from the heart and intrathoracic veins and arteries that is multi-dimensional since most sensory neurons transduce both the regional mechanical and the chemical milieu^{12, 27, 28}. Their activity is likewise altered by autonomic neural inputs¹². This population of neurons represents the dominant sub-class of neurons contained within the intrinsic cardiac nervous system and sub-serves major neuronal processing within that network¹⁰.

Autonomic dysregulation is central to the evolution of cardiac pathology^{17, 18, 29}. Mechanistically, this reflects reactive and adaptive responses of the cardiac neural hierarchy that derive from sensory transduction of the stressed/diseased myocardium^{17, 30, 31}. Such changes in neural processing manifest themselves throughout the neuraxis including at the intrinsic cardiac nervous system, intra-thoracic sympathetic ganglia, spinal cord, brainstem and multiple central regions up to the insular cortex^{10, 15, 19, 32}. This functional reorganization leads to a conflict between central and peripheral aspects of the hierarchy^{18, 30, 33}. Altered neural processing leads to maladaptive responses that ultimately results in excessive sympathetic overdrive;^{18, 31, 34, 35} that in turn contributes to the development of cardiac disease including fatal arrhythmias and heart failure^{17, 29}. It is through the understanding of such hierarchical control and how it adapts that a rationale mechanistic based approach can be devised to effectively target specific neural processing of the cardiac nervous system to therapeutically manage cardiac pathology. Figure 1 illustrates several of these neural nexus points and serves as the focal point around which device based neuromodulation actions/reactions must be considered.

Vagal Nerve Stimulation (VNS)

When considering effects of any bio-electronic approach for neuromodulation one must consider both direct and reactive responses. The vagus can be stimulated in many different ways, at a number of different levels, and for multiple pathologies³⁶⁻³⁸. In each case, one must consider the characteristics of the nerves being stimulated (afferent/efferent) and the potential impact of stimulation parameters (frequency, intensity, pulse width, waveform and duty cycle). Ultimately these factors impact both off-target adverse effects and more importantly the efficacy of the applied therapy. With respect to VNS, emerging technologies for cardiovascular disease involve either direct implant of electrodes onto the cervical vagus or non-invasive stimulation via the auricular branch of the cervical vagus. This is an emerging area and our understanding of vagal nerve stimulation is still rudimentary.

VNS preclinical results

In preclinical studies, VNS has documented efficacy to impact cardiac electrical and mechanical function. In a rabbit model, VNS damped the cardiac electrophysiological restitution curve with a corresponding reduction in potential for ventricular fibrillation^{39, 40}. In a porcine model, VNS applied against the stress imposed by acute ischemia-reperfusion was effective in reducing infarct size, stabilizing cardiac electrical function and in protecting mitochondrial function⁴¹. In an acute canine model, the effects of VNS were evaluated against an elevated sympathetic background as induced by left stellate ganglion (LSG) stimulation⁴². They demonstrated that LSG stimulation resulted in increased ventricular

instability and change in spatial heterogeneity which was reversed with VNS. They also showed that LSG stimulation with VNS resulted in a higher VF threshold compared to LSG alone. Vanoli et al⁴³ likewise demonstrated the efficacy of vagal stimulation to prevent sudden death in a canine model with healed myocardial infarction. Since heart failure is associated with a higher risk for sudden cardiac death, neuromodulation therapies that stabilize cardiac electrical function are of obvious clinical importance.

Pre-clinical studies have demonstrated efficacy for VNS to impact the progression of heart failure. In the rat infarct model with heart failure, VNS improved hemodynamics, left ventricular remodelling, and reduced neurohormonal activation⁴⁴. This study demonstrated a reduction in mortality rate at 140 days from 50% in the sham model to 14% with VNS stimulation⁴⁴. In this rat study, VNS was applied at 20Hz, 16% duty cycle (10 sec on, 50 sec off) and at an intensity that decreased the heart rate by 20-30 beats per minute. In the canine pacing induced heart failure model, VNS was associated with an improvement in left ventricular dimensions and down-regulation of important heart failure-related biomarkers including norepinephrine, angiotensin II, and C-reactive protein⁴⁵. In that study VN was stimulated at 20 Hz, with a 53% duty cycle (14 sec on, 12 sec off) and with an intensity sufficient to produce a ~20 beat/min decrease in heart rate. Importantly, more recent preclinical studies have demonstrated that therapeutic benefits of VNS against heart failure progression can be achieved at levels of VNS that induce minimal changes in heart rate. This includes the guinea pig pressure overload model⁴⁶ with right cervical vagal stimulation (20 Hz, 22% duty cycle [14 sec on, 48 sec off]) and with bilateral non-invasive stimulation of the auricular branch of the vagus nerve (20Hz, 50% duty cycle [5 sec on, 5 sec off])^{47, 48}. In both studies contractile function improved and adverse indices of neurohumoral activation reduced towards control.

VNS impacts multiple levels of the hierarchy for cardiac control. Activation of descending efferent projections can mitigate sympatho-excitation via neural interactions within the intrinsic cardiac nervous system^{49, 50}, modulate cardio-cardiac reflexes¹² and impart cardioprotection via effects on cardiomyocytes⁴¹. Activation of ascending afferents can impact central reflexes including those that involve sympathetic and parasympathetic efferent outflows to the heart^{19, 20, 22, 36}. Future pre-clinical studies should expand upon these concepts and consider additional factors including: 1) the impact of optimum medical therapy on neuromodulation; 2) intermittent versus continuous methodologies; 3) differential effects mediated from sites of VNS activation (e.g. intra-thoracic, cervical or auricular); and 4) efficacy against different cardiac pathologies including HFpEF (heart failure with preserved ejection fraction) and HFrEF (heart failure with reduced ejection fraction).

VNS clinical results

The ultimate endgame for preclinical studies in neuromodulation is transitioning to the clinical setting. While VNS has a long history in the treatment of epilepsy and depression^{36, 51, 52} its clinical application for cardiac disease commenced in 2008. In patients with NYHA class II-III heart failure and left ventricular ejection fraction <35%, stimulation titration on the right cervical vagus commenced at 1-2Hz, synchronized to the

cardiac cycle, and with a target heart rate reduction of 5-10 beats per minute⁵³. Patients were followed for 6 months. Results from this initial study included improved left ventricular systolic volume, NYHA classification and quality of life⁵³. This was followed by an open-label phase II trial in patients with reduced ejection fraction and NYHA class II-IV⁵⁴. At 6 months there was an improvement in left ventricular ejection fraction, volumes and 6 minute walk test which was maintained at one year⁵⁴. INOVATE-HF is a continuation of this approach, with a target of 650 patients and an expected primary completion date of December 2016.

NECTAR-HF (Neural cardiac TherApy foR Heart Failure) was a randomized controlled trial with VNS in patients with ejection fraction <35%, increased LV end diastolic dimensions (>55mm), NYHA class III-IV, excluding patients with CRT devices, or QRS>130ms⁵⁵. All patients had a VNS device implanted (Precision™, Boston Scientific Corporation, St Paul, MN, USA) and were randomized in a 2:1 fashion to VNS on or off for 6 months. The stimulation parameters used were 20Hz with a 12.5% duty cycle (10 sec on, 50 sec off) with an average intensity of 1.42±0.8mA. This study failed to reach its primary end point of improvement in left ventricular systolic dimensions and secondary end points of improvement in other echocardiographic parameters, and circulating biomarkers. The study, however, did show an improvement in quality of life and NYHA classification. It is likely that the stimulation parameters, especially intensity, used in this trial may have contributed to the lack of efficacy.

ANTHEM-HF (Autonomic Neural regulation Therapy to Enhance Myocardial function in Heart Failure; Cyberonics, Houston, TX, USA) investigated VNS of the right or left cervical vagus in 60 patients⁵⁶. The main inclusion criteria was ejection fraction less than 40%, LV end diastolic dimensions 50-80mm, and QRS <150ms. Patients were followed up over a 6 month period with up titrations of VNS over 10 weeks to an average intensity of 2.0 ± 0.6mA at 10Hz stimulation, and with a duty cycle of 17.5% (14 sec on, 66 sec off). There was an improvement in left ventricular ejection fraction by 4.5% but the left ventricular end systolic volume did not decrease significantly. There was again an improvement in quality of life, exercise capacity, and NYHA classification. There was no significant difference between left or right cervical vagus stimulation.

In summary, VNS has proven to be safe and feasible for use in humans in the setting of HFrEF. Future studies on VNS should focus on optimization of parameters of stimulation, patient selection and its transition where indicated into standard of care.

Spinal Cord Stimulation

Spinal Cord stimulation (SCS) has a clinical history of 20 years for the treatment of chronic pain and refractory angina pectoris⁵⁷⁻⁵⁹. While initially put forward based upon the gate control theory of pain⁶⁰, subsequent work has demonstrated that SCS is not a masking phenomenon, but instead fundamentally alters the neural-end organ interface. It is usually implemented by placement of a multi-pole electrode over the dorsal column of the thoracic cord and stimulation at parameter sets of 50 Hz, 200 µs pulse width, and intensities of 90% motor threshold^{59, 61}. Early clinical studies speculated that the anti-anginal effects were a reflection of changes in supply/demand at the heart; however, subsequent preclinical studies

determined that SCS doesn't modulate/alter coronary blood flow or LV dynamics during ischemic stress⁶², at least in the acute setting. Beyond its anti-anginal effects⁶¹, SCS exerts multi-factorial cardio-protective influences including reducing atrial and ventricular arrhythmias⁶³⁻⁶⁵, and the apoptotic potential^{66, 67}, while helping to preserve contractile function^{11, 65}. In the acute setting its efficacy is optimum when applied pre-emptively⁶⁷, but reactive and chronic SCS therapy is also cardioprotective^{63, 65}.

SCS pre-clinical results

Cardiac sympathetic afferents transduce information responding to mechanical and chemical stimuli via the intrathoracic (T1-6) and cervical (C8-9) paravertebral sympathetic ganglia to the dorsal root ganglia and subsequently to the spinal cord and higher centers^{14, 15, 19}. The cell bodies that convey the sympathetic afferent visceral input to the brain stem are found in lamina I, V, VII, and X in the C8-T9 dorsal horn^{14, 15, 19}. Central and peripheral reflex processing of that afferent signal likewise contributes to the underlying sympatho-excitation of cardiac disease and the progression into heart failure^{17, 18, 29}. As demonstrated in Figure 1, the spinal cord is one primary nexus point from which such processes can potentially be regulated with appropriate bioelectric medicine.

SCS impacts autonomic reflexes at multiple levels of the cardiac nervous system to impact basal cardiac function and its response to imposed stress. At the spinal cord itself, SCS induces the release of neuromodulators such as dynorphin, blunts the release of primary afferent related neurotransmitters such as substance P, and alters activity with sympathetic preganglionic neurons contained within the intermediolateral cell column^{68, 69}. As such it thereby alters ascending signals to higher centers^{15, 57, 58} and alters autonomic efferent outflows to peripheral aspects of the cardiac nervous system. Within extracardiac sympathetic ganglia the reflex sympatho-excitation imposed by transient myocardial ischemia is blunted while its basal function remains unaltered¹¹. Within the intrinsic cardiac nervous system, a similar blunting of reflex responses to transient ischemic stress is also present, but its basal activity is reduced by SCS⁷⁰; an effect that is manifest over time⁷¹. Subsequent studies have identified local circuit neurons as a primary target for SCS mediated therapy⁶⁴ and that SCS modifies synaptic function without directly targeting transmembrane properties of individual IC neuronal somata⁶³. Overall, such influences can be best characterized as reflex stabilization across the neuraxis for cardiac control.

SCS modulation/stabilization of autonomic responsiveness is reflected in cardioprotection. SCS reduced aberrant electrophysiological activity within the myocardium in chronic animal models with reduced coronary reserve⁷². SCS in a heart failure canine model was effective at reducing ventricular arrhythmias⁷³, improving left ventricular contractile function, and reducing heart failure⁶⁵. A porcine model of ischemic heart failure showed similar results with improved left ventricular function and myocardial strain with SCS⁷⁴. Together, these studies have demonstrated safety and efficacy for SCS for treatment of cardiac pathology.

SCS clinical results

The Defeat-HF trial (NCT01112579), was a randomized, multicenter single blind study of 66 patients with systolic heart failure. The investigators used a single lead in the T2-T4

epidural region with patients randomized 3:2 ratio of stimulation 'on' or 'off', with the spinal cord stimulation for 12 hours per day. After 6 months the controls crossed over to therapy. The inclusion criteria were left ventricular ejection fraction $\geq 35\%$, NYHA class III, QRS duration less than 120ms, left ventricular end diastolic dimensions of 55mm to 80mm on stable heart failure treatments. The preliminary results failed to show an improvement in left ventricular function and dimensions but the device was found to be safe and SCS feasible to do ⁷⁵. There was no difference in freedom from heart failure or hospitalization between the two groups. The lack of efficacy of this trial, compared to preclinical experience, was likely due to sub-optimum stimulus paradigms, especially leaving patients untreated for 12hr/day.

The Spinal Cord Stimulation Heart study was a multicenter, prospective, pilot trial involving SCS in patients with systolic heart failure ⁷⁶. Inclusion criteria were patients with left ventricular ejection fractions around 20-30%, and NYHA class III. Eligible patients had spinal cord stimulators implanted in the T1-T3 epidural space. The stimulation parameters were 50Hz for 24 hours a day. Of the 15 patients that completed the 24 month follow up, there was an improvement in NYHA classification, quality of life, left ventricular end systolic volume and peak oxygen consumption. The study was also found to be safe in the setting of severe systolic heart failure.

In summary, preclinical and clinical studies both substantiate the safety of SCS for management of both cardiac arrhythmias and progression of heart failure. Future studies on SCS should focus on optimization of parameters of stimulation and patient selection. Additional mechanistic studies are likewise required to delineate the precise mechanisms by which SCS exerts its effects on central and peripheral aspects of the cardiac nervous system. Further studies are also required to determine what intracellular pathways are engaged by SCS to render cardiomyocytes stress resistant to ischemic and non-ischemic cardiac stressors.

Baroreceptor Stimulation

The baroreflex is a negative feedback system that is a primary controller of arterial blood pressure. Its afferent input signal derives from mechanoreceptors located in the carotid sinus and aortic arch. Indices of blood pressure are sensed by changes in vessel stretch as transduced by the sensory neurites enveloping both regions. Soma for the aortic arch mechanoreceptors are localized within the nodose ganglion while soma for the carotid sinus afferents are localized to the petrosal ganglia ¹⁴. These signals are transmitted to the Nucleus Tractus Solitarius from which secondary projections arise for control of sympathetic and parasympathetic outflows ¹⁹. Baroreceptor sensitivity is depressed in heart failure due to persistent enhancement of the sympathetic activity, possibly related to central angiotensin II levels ^{18, 77}. Impairment in baroreceptor sensitivity in heart failure is associated with increased mortality ^{30, 78, 79}. Baroreceptor neurons are adaptive and in the presence of persistent neurohormonal and cardiovascular responses the baroreceptors incompletely reset leading to long term alterations in sympathetic activity and arterial pressure ^{18, 80}.

Avoidance of activation of the carotid body chemoreceptors with baroreceptor stimulation is imperative given their role in progression of heart failure by contributing to respiratory

instability and oscillatory breathing (changes in tidal volume and respiratory frequency)⁸¹. This can further exacerbate tonic and chemo-reflex evoked activation of the sympathetic nervous system by causing changes in pH, circulatory delay, and a decrease in systemic oxygen transport⁸¹.

Baroreceptors can be stimulated at different points but clinically the easiest point of stimulation is at the level of the carotid sinus. While early studies with carotid sinus implants were associated with structural damage to implanted areas⁸², recent advances in biotechnology have overcome such problems. Current devices are implanted with electrodes positioned in the carotid perivascular space around the sinus of the carotid arteries with a lead to a pulse generator positioned in the infraclavicular region^{83, 84}. They are usually placed bilaterally. The premise for such therapy is that stimulation of the peripheral baroreceptors fibers increase afferent activity transduced to the Nucleus Tractus Solitarius, which is interpreted as an increase in blood pressure. In reflex response to that afferent signal efferent outflows (sympathetic down and parasympathetic up) are modified leading to reduction in blood pressure and heart rate.

Baroreceptor stimulation preclinical results

Preclinical studies support proof of concept for utilizing bioelectric approaches as applied to the carotid sinus to treat cardiac disease. Such stimulation was correlated with lower plasma norepinephrine, angiotensin II levels, and reduced mortality in a heart failure induced canine model⁸⁴. Improvements in left ventricular function have been demonstrated in a canine heart failure infarct model with advanced heart failure at 3-months. Bilateral activation of the carotid sinus nerves improved left ventricle systolic and diastolic function and reduced heart rate compared to no-treatment controls⁸³. Adverse structural remodelling was likewise mitigated in the treated group. No major safety issues were identified in preclinical studies utilizing carotid sinus stimulation and these studies laid the preclinical foundation for ongoing clinical trials.

Baroreceptor stimulation clinical results

Clinical trials have been taking place to determine the outcomes of the use of baroreceptor stimulation therapy in both systolic and diastolic heart failure. The use of implantable carotid sinus stimulator device (Rheos System) was initially used in hypertension (DEBuT-HT trial – Device Based Therapy in Hypertension Trial) showed a sustained blood pressure drop up to 4 years out and improvement in cardiac function^{85, 86}. To corroborate this data, the Rheos Pivotal Trial, a double-blind randomized placebo-controlled trial, showed 88% maintenance in blood pressure reduction response at a 12 month period⁸⁷. A recent study looking at baroreflex activation therapy in advanced systolic heart failure, in the setting of a narrow QRS, randomized patients to BAT or standard medical therapy. They showed an improvement in quality of life scores, NYHA classification, and NT-pro-BNP but did not show any change in left ventricular function with therapy⁸⁸.

In summary, utilization of the carotid sinus nerve stimulation to modulate baroreflex control mechanisms has proof-of-concept as a safe therapy for heart disease, but with limited experience in the clinical realm. It is likely that treatment efficacy is critically dependent on

underling pathology, so patient selection is critical. Future studies should focus on electrode interfaces, stimulus paradigms, and the potential for closed-loop feedback.

Renal axonal modulation and heart failure

The role of modulation of afferent renal sympathetic neurons has been investigated in preclinical and clinical studies in the setting of heart failure. Renal modulation was accomplished by catheter ablation of the renal arteries under fluoroscopic and/or electroanatomic mapping guidance, after confirmation of the absence of any baseline renal artery stenosis. Renal sympathetic modulation in a rat model with compensated high output heart failure secondary to atrio-venous fistula formation resulted in attenuated sodium excretion after sodium loading ⁸⁹. In post myocardial infarction rats, renal sympathetic modulation resulted in increased sodium excretion and decreased LV filling pressure with improved left ventricular function ⁹⁰. In a rabbit pacing induced heart failure model, renal sympathetic modulation modified angiotensin II release, and preserved renal flow and reno-vascular resistance ⁹¹. In a canine model, renal sympathetic modulation in the setting of a pacing induced heart failure model reduced circulating angiotensin II, aldosterone, BNP, endothelin-1, and reninase ⁹². Similarly, another canine high rate pacing induced heart failure model demonstrated reduced ventricular substrate remodelling and circulating angiotensin II and TGF- β with renal sympathetic modulation compared to controls ⁹³. They subsequently demonstrated that renal sympathetic modulation resulted in attenuation of substrate and electrical remodelling with less inducibility of ventricular fibrillation ⁹⁴.

Renal axonal modulation clinical results

Small case series have demonstrated a reduction of arrhythmia post renal nerve denervation ^{95, 96}. Although this benefit is for arrhythmia burden reduction and not specifically to provide an improvement in left ventricular function there is a significant overlap. In other words, frequent lethal ventricular arrhythmias occur in the setting of LV dysfunction and congestive cardiomyopathy could result in deterioration in heart failure status.

The outcomes of multiple ongoing trials in relation to renal denervation actually directly benefiting left ventricular function will be very interesting. There is evidence to suggest benefits in preserved ejection fraction, probably as a result of an anti-hypertensive effect resulting in less structural remodelling ⁹⁷. This may not translate to a reduced ejection fraction heart failure model given that the majority of these patients have normal or reduced systolic blood pressure.

The REACH- Pilot study ⁹⁸ was a feasibility study in 7 patients with NYHA class III-IV already on maximum medical therapy. These patients had heart failure with left ventricular ejection fractions (EF) of $43 \pm 15\%$. This study failed to show an improvement in left ventricular function but there was a reported improvement in symptoms. The REACH-Trial (NCT01639378) is an ongoing prospective, double blind, randomized control trial looking at safety and efficacy of renal sympathetic modulation in chronic heart failure. Another ongoing feasibility study is Symplicity-HF (NCT01392196) which aims to recruit about 40 NYHA class II-III patients, with EF < 40%, impaired renal function and the primary end

point is safety. A smaller trial from the Czech Republic (NCT01870310) is a randomized, controlled trial in chronic heart failure with EF<35%, with primary end points of NT-proBNP levels, and secondary endpoints of hospitalization, and death due to cardiovascular causes.

Other novel methods of ANS modulation

Initiation and progression of cardiac disease is at its foundation critically dependent upon changes in afferent signalling. Modulation of such afferent signalling is an emerging target for autonomic regulation therapy. In a preclinical study, Wang et al.³⁵, demonstrated that they can mitigate the cardiac afferent mediated sympatho-exciatory reflex by administering Resiniferatoxin (RTX) to the epicardium of an ischemic heart failure rat model. RTX works by blocking the transient receptor potential vallinoid 1 receptor and appears to reduce the cardiac afferent response. They demonstrated in comparison to sham model that RTX prevented increased left ventricular diastolic pressures, lung edema, cardiac hypertrophy, and partially reduced left ventricular dimensions in the failing heart. They also showed that by removing the afferent input there was attenuation in cardiac fibrosis, apoptosis, and reduced expression of fibrotic markers such as TGF- β in the RTX treated group.

Conclusion

Modulation of the autonomic nervous system is an emerging therapy to treat heart failure. It is predicated on using targeted bioelectric approaches to mitigate the maladaptive and excessive neurohumoral response to cardiac disease that are endogenously engaged to help maintain adequate cardiac output. These neurohumoral imbalances result in structural and functional changes within the various elements of the cardiac nervous system and in the cardiac tissues they innervate. Stabilization of imbalances within select elements of the cardiac neuronal hierarchy can reduce arrhythmogenesis and maintain myocardial viability in the setting of ischemic and non-ischemic heart disease. What is essential for this field to move forward is a mechanistic understanding of the induced changes in the neural hierarchy/ cardiac interface in pathological conditions and from that knowledge to design and implement the optimum interfaces and stimulation paradigms to mitigate such adverse responses.

Acknowledgments

Sources of funding and competing interests: Work supported by Heart, Lung and Blood Institute grants HL71830 (JLA) and HL84261 (KS). Private sector funding for studies related to neuromodulation provided in part by St. Jude Medical (JLA) and Cyberonics Inc (JLA).

References

1. Go AS, Mozaffarian D, Roger VL, et al. Heart disease and stroke statistics--2014 update: a report from the American Heart Association. *Circulation*. Jan 21.2014 129:e28–e292. [PubMed: 24352519]
2. Heidenreich PA, Albert NM, Allen LA, et al. Forecasting the impact of heart failure in the United States: a policy statement from the American Heart Association. *Circulation Heart failure*. May. 2013 6:606–619. [PubMed: 23616602]

3. Ho KK, Pinsky JL, Kannel WB, Levy D. The epidemiology of heart failure: the Framingham Study. *Journal of the American College of Cardiology*. Oct.1993 22:6A–13A. [PubMed: 8509564]
4. Mehta PA, Dubrey SW, McIntyre HF, et al. Improving survival in the 6 months after diagnosis of heart failure in the past decade: population-based data from the UK. *Heart*. Nov.2009 95:1851–1856. [PubMed: 19587390]
5. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *Jama*. Feb 26.2014 311:806–814. [PubMed: 24570244]
6. Wiener JM, Tilly J. Population ageing in the United States of America: implications for public programmes. *International journal of epidemiology*. Aug.2002 31:776–781. [PubMed: 12177018]
7. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes care*. May.2004 27:1047–1053. [PubMed: 15111519]
8. Armour JA. Cardiac neuronal hierarchy in health and disease. *Am J Physiol Regul Integr Comp Physiol*. Aug.2004 287:R262–271. [PubMed: 15271675]
9. Kember G, Armour JA, Zamir M. Neural control of heart rate: the role of neuronal networking. *J Theor Biol*. May 21.2011 277:41–47. [PubMed: 21354183]
10. Armour JA. Potential clinical relevance of the 'little brain' on the mammalian heart. *Exp Physiol*. Feb.2008 93:165–176. [PubMed: 17981929]
11. Ardell JL, Cardinal R, Vermeulen M, Armour JA. Dorsal spinal cord stimulation obtunds the capacity of intrathoracic extracardiac neurons to transduce myocardial ischemia. *Am J Physiol Regul Integr Comp Physiol*. Aug.2009 297:R470–477. [PubMed: 19515981]
12. Beaumont E, Salavatian S, Southerland EM, et al. Network interactions within the canine intrinsic cardiac nervous system: implications for reflex control of regional cardiac function. *J Physiol*. Sep 15.2013 591:4515–4533. [PubMed: 23818689]
13. Oppenheimer SM, Gelb A, Girvin JP, Hachinski VC. Cardiovascular effects of human insular cortex stimulation. *Neurology*. Sep.1992 42:1727–1732. [PubMed: 1513461]
14. Armour, JA.; Kember, G. Cardiac sensory neurons.. In: Armour, JA.; Ardell, JL., editors. *Basic and Clinical Neurocardiology*. Oxford University Press; New York: 2004. p. 79-117.
15. Foreman RD. Mechanisms of cardiac pain. *Annu Rev Physiol*. 1999; 61:143–167. [PubMed: 10099685]
16. Fu LW, Longhurst JC. Regulation of cardiac afferent excitability in ischemia. *Handb Exp Pharmacol*. 2009:185–225. [PubMed: 19655108]
17. Fukada K, Kanazawa H, Aizawa Y, Ardell JL, Shivkumar K. Cardiac innervation and sudden cardiac death. *Circulation research*. 2015 in press.
18. Zucker IH, Patel KP, Schultz HD. Neurohumoral stimulation. *Heart Fail Clin* Jan. 2012; 8:87–99.
19. Andresen, MC.; Kunze, DL.; Mendelowitz, D. Central Nervous System Regulation of the Heart.. In: Armour, JA.; Ardell, JL., editors. *Basic and Clinical Neurocardiology*. Oxford University Press; New York: 2004. p. 187-219.
20. Zucker, IH.; Gilmore, JP. *Reflex Control of the Circulation*. CRC Press; Boca Raton: 1991.
21. Blinder KJ, Johnson TA, John Massari V. Negative inotropic vagal preganglionic neurons in the nucleus ambiguus of the cat: neuroanatomical comparison with negative chronotropic neurons utilizing dual retrograde tracers. *Brain Res*. Sep 7.1998 804:325–330. [PubMed: 9757076]
22. Coote JH. Myths and realities of the cardiac vagus. *J Physiol*. Sep 1.2013 591:4073–4085. [PubMed: 23878363]
23. McAllen RM, Salo LM, Paton JF, Pickering AE. Processing of central and reflex vagal drives by rat cardiac ganglion neurones: an intracellular analysis. *J Physiol*. Dec 1.2011 589:5801–5818. [PubMed: 22005679]
24. Hopkins DA, Armour JA. Localization of sympathetic postganglionic and parasympathetic preganglionic neurons which innervate different regions of the dog heart. *J Comp Neurol*. Oct 20.1984 229:186–198. [PubMed: 6501600]
25. Randall, WC. Efferent Sympathetic Innervation of the Heart.. In: Armour, JA.; Ardell, JL., editors. *Neurocardiology*. Oxford University Press; New York: 1994. p. 77-94.

26. Gray AL, Johnson TA, Ardell JL, Massari VJ. Parasympathetic control of the heart. II. A novel interganglionic intrinsic cardiac circuit mediates neural control of heart rate. *J Appl Physiol* (1985). Jun.2004 96:2273–2278. [PubMed: 14978001]
27. Armour JA, Collier K, Kember G, Ardell JL. Differential selectivity of cardiac neurons in separate intrathoracic autonomic ganglia. *Am J Physiol*. Apr.1998 274:R939–949. [PubMed: 9575954]
28. Waldmann M, Thompson GW, Kember GC, Ardell JL, Armour JA. Stochastic behavior of atrial and ventricular intrinsic cardiac neurons. *J Appl Physiol* (1985). Aug.2006 101:413–419. [PubMed: 16645188]
29. Florea VG, Cohn JN. The autonomic nervous system and heart failure. *Circ Res*. May 23.2014 114:1815–1826. [PubMed: 24855204]
30. Billman GE. A comprehensive review and analysis of 25 years of data from an in vivo canine model of sudden cardiac death: implications for future anti-arrhythmic drug development. *Pharmacol Ther*. Sep.2006 111:808–835. [PubMed: 16483666]
31. Ajjola OA, Yagishita D, Reddy NK, et al. Remodeling of Stellate Ganglion Neurons following Spatially Targeted Myocardial Infarction: Neuropeptide And Morphologic Changes. *Heart rhythm : the official journal of the Heart Rhythm Society*. Jan 29.2015
32. Macey PM, Wu P, Kumar R, et al. Differential responses of the insular cortex gyri to autonomic challenges. *Autonomic neuroscience : basic & clinical*. May 21.2012 168:72–81. [PubMed: 22342370]
33. Kember G, Armour JA, Zamir M. Neural control hierarchy of the heart has not evolved to deal with myocardial ischemia. *Physiol Genomics*. Aug 1.2013 45:638–644. [PubMed: 23695889]
34. Vaseghi M, Gima J, Kanaan C, et al. Cardiac sympathetic denervation in patients with refractory ventricular arrhythmias or electrical storm: intermediate and long-term follow-up. *Heart Rhythm*. Mar.2014 11:360–366. [PubMed: 24291775]
35. Wang HJ, Wang W, Cornish KG, Rozanski GJ, Zucker IH. Cardiac sympathetic afferent denervation attenuates cardiac remodeling and improves cardiovascular dysfunction in rats with heart failure. *Hypertension*. Oct.2014 64:745–755. [PubMed: 24980663]
36. Bonaz B, Picq C, Sinniger V, Mayol JF, Clarencon D. Vagus nerve stimulation: from epilepsy to the cholinergic anti-inflammatory pathway. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society*. Mar.2013 25:208–221. [PubMed: 23360102]
37. De Ferrari GM. Vagal stimulation in heart failure. *J Cardiovasc Transl Res*. Apr.2014 7:310–320. [PubMed: 24500409]
38. Pavlov VA, Tracey KJ. The vagus nerve and the inflammatory reflex--linking immunity and metabolism. *Nature reviews Endocrinology*. Dec.2012 8:743–754.
39. Brack KE, Winter J, Ng GA. Mechanisms underlying the autonomic modulation of ventricular fibrillation initiation--tentative prophylactic properties of vagus nerve stimulation on malignant arrhythmias in heart failure. *Heart Fail Rev*. Jul.2013 18:389–408. [PubMed: 22678767]
40. Ng GA. Vagal modulation of cardiac ventricular arrhythmia. *Exp Physiol*. Feb.2014 99:295–299. [PubMed: 24014808]
41. Shinlapawittayatorn K, Chinda K, Palee S, et al. Low-amplitude, left vagus nerve stimulation significantly attenuates ventricular dysfunction and infarct size through prevention of mitochondrial dysfunction during acute ischemia-reperfusion injury. *Heart Rhythm*. Nov.2013 10:1700–1707. [PubMed: 23933295]
42. Huang J, Qian J, Yao W, et al. Vagus nerve stimulation reverses ventricular electrophysiological changes induced by hypersympathetic nerve activity. *Experimental physiology*. Mar.2015 100:239–248. [PubMed: 25720663]
43. Vanoli E, De Ferrari GM, Stramba-Badiale M, Hull SS Jr, Foreman RD, Schwartz PJ. Vagal stimulation and prevention of sudden death in conscious dogs with a healed myocardial infarction. *Circulation research*. May.1991 68:1471–1481. [PubMed: 2019002]
44. Li M, Zheng C, Sato T, Kawada T, Sugimachi M, Sunagawa K. Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats. *Circulation*. Jan 6.2004 109:120–124. [PubMed: 14662714]

45. Zhang Y, Popovic ZB, Bibeovski S, et al. Chronic vagus nerve stimulation improves autonomic control and attenuates systemic inflammation and heart failure progression in a canine high-rate pacing model. *Circulation Heart failure*. Nov.2009 2:692–699. [PubMed: 19919995]
46. Beaumont, E.; Southerland, EM.; Hardwick, JC.; Ryan, SE.; KenKnight, BH.; Ardell, JL. Chronic autonomic regulation therapy mitigates adverse remodeling induced by pressure overload in the guinea pig heart.. Paper presented at: American College of Cardiology; Washington DC.. 2014;
47. Wang Z, Yu L, Chen M, Wang S, Jiang H. Transcutaneous electrical stimulation of auricular branch of vagus nerve: a noninvasive therapeutic approach for post-ischemic heart failure. *International journal of cardiology*. Dec 15.2014 177:676–677. [PubMed: 25449481]
48. Wang Z, Yu L, Wang S, et al. Chronic intermittent low-level transcutaneous electrical stimulation of auricular branch of vagus nerve improves left ventricular remodeling in conscious dogs with healed myocardial infarction. *Circulation Heart failure*. Nov.2014 7:1014–1021. [PubMed: 25332149]
49. McGuirt AS, Schmach DC, Ardell JL. Autonomic interactions for control of atrial rate are maintained after SA nodal parasympathectomy. *Am J Physiol*. Jun.1997 272:H2525–2533. [PubMed: 9227527]
50. Randall DC, Brown DR, McGuirt AS, Thompson GW, Armour JA, Ardell JL. Interactions within the intrinsic cardiac nervous system contribute to chronotropic regulation. *Am J Physiol Regul Integr Comp Physiol*. Nov.2003 285:R1066–1075. [PubMed: 12842863]
51. Ryzi M, Brazdil M, Novak Z, et al. Long-term outcomes in patients after epilepsy surgery failure. *Epilepsy research*. Feb.2015 110:71–77. [PubMed: 25616458]
52. Tisi G, Franzini A, Messina G, Savino M, Gambini O. Vagus nerve stimulation therapy in treatment-resistant depression: a series report. *Psychiatry and clinical neurosciences*. Aug.2014 68:606–611. [PubMed: 25215365]
53. Schwartz PJ, De Ferrari GM, Sanzo A, et al. Long term vagal stimulation in patients with advanced heart failure: first experience in man. *European journal of heart failure*. Sep.2008 10:884–891. [PubMed: 18760668]
54. De Ferrari GM, Crijns HJ, Borggrefe M, et al. Chronic vagus nerve stimulation: a new and promising therapeutic approach for chronic heart failure. *European heart journal*. Apr.2011 32:847–855. [PubMed: 21030409]
55. Zannad F, De Ferrari GM, Tuinenburg AE, et al. Chronic vagal stimulation for the treatment of low ejection fraction heart failure: results of the NEural Cardiac TherApy foR Heart Failure (NECTAR-HF) randomized controlled trial. *European heart journal*. Feb 14.2015 36:425–433. [PubMed: 25176942]
56. Premchand RK, Sharma K, Mittal S, et al. Autonomic Regulation Therapy via Left or Right Cervical Vagus Nerve Stimulation in Patients with Chronic Heart Failure: Results of the ANTHEM-HF Trial. *J Card Fail*. Aug 22.2014
57. Foreman RD, Linderorth B. Neural mechanisms of spinal cord stimulation. *International review of neurobiology*. 2012; 107:87–119. [PubMed: 23206679]
58. Zhang TC, Janik JJ, Grill WM. Mechanisms and models of spinal cord stimulation for the treatment of neuropathic pain. *Brain research*. Jun 20.2014 1569:19–31. [PubMed: 24802658]
59. Mannheimer C, Camici P, Chester MR, et al. The problem of chronic refractory angina; report from the ESC Joint Study Group on the Treatment of Refractory Angina. *Eur Heart J*. Mar.2002 23:355–370. [PubMed: 11846493]
60. Melzack R, Wall PD. Pain mechanisms: a new theory. *Science*. Nov 19.1965 150:971–979. [PubMed: 5320816]
61. Borjesson M, Andrell P, Lundberg D, Mannheimer C. Spinal cord stimulation in severe angina pectoris--a systematic review based on the Swedish Council on Technology assessment in health care report on long-standing pain. *Pain*. Dec.2008 140:501–508. [PubMed: 19004553]
62. Kingma JG Jr, Linderorth B, Ardell JL, Armour JA, DeJongste MJ, Foreman RD. Neuromodulation therapy does not influence blood flow distribution or left-ventricular dynamics during acute myocardial ischemia. *Auton Neurosci*. Aug 13.2001 91:47–54. [PubMed: 11515801]

63. Ardell JL, Cardinal R, Beaumont E, Vermeulen M, Smith FM, Andrew Armour J. Chronic spinal cord stimulation modifies intrinsic cardiac synaptic efficacy in the suppression of atrial fibrillation. *Autonomic neuroscience : basic & clinical*. Dec.2014 186:38–44. [PubMed: 25301713]
64. Gibbons DD, Southerland EM, Hoover DB, Beaumont E, Armour JA, Ardell JL. Neuromodulation targets intrinsic cardiac neurons to attenuate neuronally mediated atrial arrhythmias. *Am J Physiol Regul Integr Comp Physiol*. Feb 1.2012 302:R357–364. [PubMed: 22088304]
65. Lopshire JC, Zhou X, Dusa C, et al. Spinal cord stimulation improves ventricular function and reduces ventricular arrhythmias in a canine postinfarction heart failure model. *Circulation*. Jul 28.2009 120:286–294. [PubMed: 19597055]
66. Southerland EM, Gibbons DD, Smith SB, et al. Activated cranial cervical cord neurons affect left ventricular infarct size and the potential for sudden cardiac death. *Auton Neurosci*. Jul 2.2012 169:34–42. [PubMed: 22502863]
67. Southerland EM, Milhorn DM, Foreman RD, et al. Preemptive, but not reactive, spinal cord stimulation mitigates transient ischemia-induced myocardial infarction via cardiac adrenergic neurons. *Am J Physiol Heart Circ Physiol*. Jan.2007 292:H311–317. [PubMed: 16920800]
68. Ding X, Ardell JL, Hua F, et al. Modulation of cardiac ischemia-sensitive afferent neuron signaling by preemptive C2 spinal cord stimulation: effect on substance P release from rat spinal cord. *Am J Physiol Regul Integr Comp Physiol*. Jan.2008 294:R93–101. [PubMed: 17989135]
69. Ding X, Hua F, Sutherly K, Ardell JL, Williams CA. C2 spinal cord stimulation induces dynorphin release from rat T4 spinal cord: potential modulation of myocardial ischemia-sensitive neurons. *Am J Physiol Regul Integr Comp Physiol*. Nov.2008 295:R1519–1528. [PubMed: 18753268]
70. Foreman RD, Linderorth B, Ardell JL, et al. Modulation of intrinsic cardiac neurons by spinal cord stimulation: implications for its therapeutic use in angina pectoris. *Cardiovasc Res*. Aug.2000 47:367–375. [PubMed: 10946073]
71. Armour JA, Linderorth B, Arora RC, et al. Long-term modulation of the intrinsic cardiac nervous system by spinal cord neurons in normal and ischaemic hearts. *Auton Neurosci*. Jan 10.2002 95:71–79. [PubMed: 11873770]
72. Cardinal R, Ardell JL, Linderorth B, Vermeulen M, Foreman RD, Armour JA. Spinal cord activation differentially modulates ischaemic electrical responses to different stressors in canine ventricles. *Auton Neurosci*. Mar 31.2004 111:37–47. [PubMed: 15109937]
73. Issa ZF, Zhou X, Ujhelyi MR, et al. Thoracic spinal cord stimulation reduces the risk of ischemic ventricular arrhythmias in a postinfarction heart failure canine model. *Circulation*. Jun 21.2005 111:3217–3220. [PubMed: 15956128]
74. Liu Y, Yue WS, Liao SY, et al. Thoracic spinal cord stimulation improves cardiac contractile function and myocardial oxygen consumption in a porcine model of ischemic heart failure. *Journal of cardiovascular electrophysiology*. May.2012 23:534–540. [PubMed: 22151312]
75. Zipes, DP.; Neuzil, P.; Theres, H., et al. Ventricular functional response to spinal cord stimulation of advanced heart failure: primary results of the randomized DEFEAT-HF trial.. Paper presented at: American Heart Association 2014; Chicago.
76. Tse HF, Turner S, Sanders P, et al. Thoracic Spinal Cord Stimulation for Heart Failure as a Restorative Treatment (SCS HEART study): First-in-man experience. *Heart rhythm : the official journal of the Heart Rhythm Society*. Mar.2015 12:588–595. [PubMed: 25500165]
77. Zhang D, Liu J, Zheng H, Tu H, Muelleman RL, Li YL. Effect of angiotension II on voltage-gated sodium currents in aortic baroreceptor neurons and arterial baroreflex sensitivity in heart failure rats. *Journal of hypertension*. Mar 30.2015
78. Nolan J, Batin PD, Andrews R, et al. Prospective study of heart rate variability and mortality in chronic heart failure: results of the United Kingdom heart failure evaluation and assessment of risk trial (UK-heart). *Circulation*. Oct 13.1998 98:1510–1516. [PubMed: 9769304]
79. Schwartz PJ, La Rovere MT, Vanoli E. Autonomic nervous system and sudden cardiac death. Experimental basis and clinical observations for post-myocardial infarction risk stratification. *Circulation*. Jan.1992 85:177–91. [PubMed: 1728509]
80. Sata Y, Kawada T, Shimizu S, Kamiya A, Akiyama T, Sugimachi M. Predominant role of neural arc in sympathetic baroreflex resetting of spontaneously hypertensive rats. *Circulation journal : official journal of the Japanese Circulation Society*. 2015; 79:592–599. [PubMed: 25746544]

81. Marcus NJ, Del Rio R, Schultz EP, Xia XH, Schultz HD. Carotid body denervation improves autonomic and cardiac function and attenuates disordered breathing in congestive heart failure. *The Journal of physiology*. Jan 15.2014 592:391–408. [PubMed: 24247985]
82. Braunwald NS, Epstein SE, Braunwald E. Carotid sinus nerve stimulation for the treatment of intractable angina pectoris: surgical technic. *Annals of surgery*. Nov.1970 172:870–876. [PubMed: 5477663]
83. Sabbah HN, Gupta RC, Imai M, et al. Chronic electrical stimulation of the carotid sinus baroreflex improves left ventricular function and promotes reversal of ventricular remodeling in dogs with advanced heart failure. *Circulation Heart failure*. Jan.2011 4:65–70. [PubMed: 21097604]
84. Zucker IH, Hackley JF, Cornish KG, et al. Chronic baroreceptor activation enhances survival in dogs with pacing-induced heart failure. *Hypertension*. Nov.2007 50:904–910. [PubMed: 17846349]
85. Scheffers JJ, Kroon AA, Schmidli J, et al. Novel baroreflex activation therapy in resistant hypertension: results of a European multi-center feasibility study. *Journal of the American College of Cardiology*. Oct 5.2010 56:1254–1258. [PubMed: 20883933]
86. Bisognano JD, Kaufman CL, Bach DS, et al. Improved cardiac structure and function with chronic treatment using an implantable device in resistant hypertension: results from European and United States trials of the Rheos system. *Journal of the American College of Cardiology*. Apr 26.2011 57:1787–1788. [PubMed: 21511116]
87. Bisognano JD, Bakris G, Nadim MK, et al. Baroreflex activation therapy lowers blood pressure in patients with resistant hypertension: results from the double-blind, randomized, placebo-controlled rheos pivotal trial. *Journal of the American College of Cardiology*. Aug 9.2011 58:765–773. [PubMed: 21816315]
88. Abraham WT, Zile M, Weaver FA, et al. Baroreflex activation therapy for the treatment of heart failure with a reduced ejection fraction. *JACC Heart failure*. 2015 in press.
89. Villarreal D, Freeman RH, Johnson RA, Simmons JC. Effects of renal denervation on postprandial sodium excretion in experimental heart failure. *The American journal of physiology*. May.1994 266:R1599–1604. [PubMed: 8203638]
90. Nozawa T, Igawa A, Fujii N, et al. Effects of long-term renal sympathetic denervation on heart failure after myocardial infarction in rats. *Heart and vessels*. Jan.2002 16:51–56. [PubMed: 11833842]
91. Clayton SC, Haack KK, Zucker IH. Renal denervation modulates angiotensin receptor expression in the renal cortex of rabbits with chronic heart failure. *American journal of physiology Renal physiology*. Jan.2011 300:F31–39. [PubMed: 20962112]
92. Zhao Q, Huang H, Wang X, et al. Changes of serum neurohormone after renal sympathetic denervation in dogs with pacing-induced heart failure. *International journal of clinical and experimental medicine*. 2014; 7:4024–4030. [PubMed: 25550911]
93. Dai Z, Yu S, Zhao Q, et al. Renal sympathetic denervation suppresses ventricular substrate remodelling in a canine high-rate pacing model. *EuroIntervention : journal of EuroPCR in collaboration with the Working Group on Interventional Cardiology of the European Society of Cardiology*. Jul.2014 10:392–399.
94. Guo Z, Zhao Q, Deng H, et al. Renal sympathetic denervation attenuates the ventricular substrate and electrophysiological remodeling in dogs with pacing-induced heart failure. *International journal of cardiology*. Jul 15.2014 175:185–186. [PubMed: 24801082]
95. Bradfield JS, Vaseghi M, Shivkumar K. Renal denervation for refractory ventricular arrhythmias. *Trends in cardiovascular medicine*. Jul.2014 24:206–213. [PubMed: 25017919]
96. Remo BF, Preminger M, Bradfield J, et al. Safety and efficacy of renal denervation as a novel treatment of ventricular tachycardia storm in patients with cardiomyopathy. *Heart rhythm : the official journal of the Heart Rhythm Society*. Apr.2014 11:541–546. [PubMed: 24389229]
97. Brandt MC, Mahfoud F, Reda S, et al. Renal sympathetic denervation reduces left ventricular hypertrophy and improves cardiac function in patients with resistant hypertension. *Journal of the American College of Cardiology*. Mar 6.2012 59:901–909. [PubMed: 22381425]

98. Davies JE, Manisty CH, Petraco R, et al. First-in-man safety evaluation of renal denervation for chronic systolic heart failure: primary outcome from REACH-Pilot study. *International journal of cardiology*. Jan 20.2013 162:189–192. [PubMed: 23031283]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Hierarchy for cardiac control

Figure 1.

This figure demonstrates the complexity of the neuronal hierarchy for cardiac control. Autonomic regulation therapy (ART) can target different structures in the cardiac neuronal hierarchy with promising results utilizing carotid sinus (CSN), dorsal column spinal cord (SCS), and cervical vagus electrical stimulation (VNS). Autonomic control can likewise be impacted by interrupting aberrant cardiac afferent signalling with Resiniferatoxin (RTX) or by renal denervation. Sympath: sympathetic; Parasymp: parasympathetic; LCN: local circuit neuron; DRG: Dorsal root ganglia; Aff. – afferent; T1-T4: first to 4th level of thoracic cord; Ang: Angiotensin; β: beta adrenergic receptor; M: muscarinic receptor; G_s and G_i: g proteins; AC: adenylate cyclase; ATP: adenosine triphosphate; cAMP: cyclic-adenosine monophosphate; Neurite: sensory endings embedded in myocardium; decent: decentralization.