Published in final edited form as:

Soc Sci Med. 2015 November ; 144: 28–37. doi:10.1016/j.socscimed.2015.09.007.

How reliable are self-reports of HIV status disclosure? Evidence from couples in Malawi

Amy A. Conroy [Post-doctoral Fellow] and

Center for AIDS Prevention Studies, University of California San Francisco, 550 16th Street, 3rd Floor, San Francisco, CA 94158

Lauren H. Wong [Post-doctoral Fellow]

Center for AIDS Prevention Studies, University of California San Francisco, 550 16th Street, 3rd Floor, San Francisco, CA 94158

Lauren H. Wong: lauren.bevand@gmail.com

Abstract

Introduction—The majority of research on human immunodeficiency virus (HIV) disclosure utilizes the perspective from a single individual, which cannot be substantiated in the absence of supporting data such as from a primary partner.

Objectives—The objectives of this study were to evaluate: (1) the extent to which self-reported HIV disclosure was confirmed by a primary partner; (2) individual and relationship-level predictors of self-reported versus confirmed disclosure; and (3) whether confirmed disclosure was a stronger predictor of correctly assessing a partner's HIV status compared to self-reported disclosure.

Methods—As part of an 8-wave longitudinal study from 2009–2011 in southern Malawi, 366 individuals (183 couples) were interviewed about their primary relationship (wave 3), individually tested for HIV (wave 4), and then asked whether they disclosed to their primary partner (wave 5).

Results—While 93% of respondents reported that they disclosed, only 64% of respondents had confirmed reports from their partner. Having communicated with partner about HIV was positively associated with self-reported disclosure; this association remained significant but became more precise in the models for confirmed disclosure. Confirmed disclosure, but not self-report, was a significant predictor of correctly assessing a partner's HIV status. Being male, having lower perceived partner infidelity, having higher relationship unity, and testing HIV-negative were positively and significantly associated with correct assessment. Dyadic data from two partners provide an improved measure of disclosure as compared to a single individual's self-report and could be used to identify behavioral and biomedical opportunities to prevent HIV transmission within couples.

Corresponding author: Amy Conroy, Center for AIDS Prevention Studies, University of California San Francisco, 550 16th Street, 3rd Floor, San Francisco, CA 94158. amy.conroy@ucsf.edu. Phone: (303) 522-9439. Fax: (630) 522-9439.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

HIV status disclosure; couples; relationship dynamics; reliability; sub-Saharan Africa; Malawi

Introduction

In sub-Saharan Africa, the success of human immunodeficiency virus (HIV) prevention and treatment depends upon high levels of disclosure to sexual partners. HIV disclosure facilitates the adoption of safer sex behaviors, partner HIV testing, social support, and engagement in HIV care and treatment programs (Brou et al., 2007; Loubiere et al., 2009; Simoni & Pantalone, 2004; Stirratt et al., 2006; Wong et al., 2009). Disclosure to primary partners may encourage disclosure to other relatives and friends, who can provide emotional and economic support necessary for coping and accessing health services (Suzan-Monti et al., 2013). Even for those who test negative—the majority of HIV testers in sub-Saharan Africa—disclosure provides an important opportunity for couples to communicate about risk reduction approaches that might not occur otherwise (Crepaz & Marks, 2004; Desgrees-Du-Lou et al., 2007). Disclosure of negative results has the potential to increase trust and intimacy, and strengthen relationship quality, which is associated with many positive health outcomes in couples (Lewis et al., 2006). Despite these benefits, disclosure of positive results can invite stigma, abandonment, discrimination, violence, and emotional distress (Obermeyer et al., 2011). Thus, decisions on when, how, and to whom to disclose are often described as a complex calculus weighing the perceived risks and benefits (Black & Miles, 2002).

Research on individuals who disclose to their sexual partners suggests that socio-economic and relationship factors play an important role. Demographic factors related to non-disclosure include lower socio-economic status, being unmarried, and younger age (Anglewicz & Chintsanya, 2011; King et al., 2008; Wong et al., 2009). Relationship dynamics such as couple communication around HIV may facilitate disclosure, while other factors related to power imbalances may interfere with disclosure (Katz et al., 2009; Maman et al., 2001; Maman et al., 2003). Indeed, women who worried about relationship dissolution, loss of economic support, and intimate partner violence were less likely to disclose to sexual partners (Maman et al., 2001; Maman et al., 2003; Medley et al., 2004). Decisions to disclose may also depend on beliefs about whether a partner's HIV status is similar or different to one's own. In seroconcordant positive partnerships, disclosure may be higher if partners expect to rely on each other for social support and coping with HIV/AIDS. Conversely, in serodiscordant partnerships, disclosure may be lower as these relationships are more prone to violence and dissolution following disclosure, especially for HIV-positive women (Maman et al., 2003; Maman et al., 2002; Porter et al., 2004). While studies have shown that HIV-negative individuals are more likely to disclose to partners than HIV-positive individuals (Anglewicz et al., 2011), little research has examined whether couples' discrepancy in HIV status itself predicts disclosure.

To date, most research on HIV disclosure has relied upon self-reported data from the perspective of a single individual (e.g., see Medley et al., 2004; Obermeyer et al., 2011;

Simoni et al., 2004), and thus cannot be substantiated in the absence of other data (Oley et al., 2004; Venkatesh et al., 2011). First, the reliability of disclosure reports may be affected by differences in perceptions around what is 'disclosure', which can occur through verbal and non-verbal gestures or behaviors (Wong et al., 2011). Second, as with other sensitive measures around sexuality and HIV/AIDS, disclosure reports are subject to social desirability bias. In sub-Saharan Africa, intentions to disclose are conflicted between the perceived ethical responsibility of protecting a partner and concerns of abandonment and violence (King et al., 2008; WHO, 2003). Thus, those who believe they will be morally judged for placing their partner at risk for HIV may be more likely to over-report disclosure. Finally, disclosure is a bi-directional process involving both a transmitter and receiver and thus, a single partner's account cannot capture both sides of the interaction.

Without an objective measure of disclosure such as direct observation, couples studies offer a novel opportunity to assess the reliability of self-reports by using partners' confirmatory accounts. Indeed, two couples studies from sub-Saharan Africa provide evidence that respondents may not disclose even when they say they do. Among married couples from Malawi, 85% of HIV-positive men and 93% of HIV-positive women who reported disclosing had their spouse's confirmation (Anglewicz et al., 2011). In a study from Kenya, men were also more likely to confirm their partner's reports of disclosure (93%) as compared to women (71%) (Katz et al., 2009). However, these two studies did not evaluate respondents' reports of their partner's HIV status following disclosure, which would provide stronger evidence that disclosure occurred.

If confirmed reports of disclosure are less prone to error than an individual's self-report, they should result in better assessments of a partner's HIV status. At the same time, knowledge of test results may not completely explain perceptions of a partner's HIV status. Research from sub-Saharan Africa suggests that people use locally adaptive strategies to socially diagnose HIV/AIDS in their communities. For example, in rural Malawi, Watkins (2004) found that people rely on the physical signs of AIDS and information from their social networks to assess who is HIV-positive. Also in Malawi, spouses were more likely to incorrectly perceive their partner was HIV-positive if they suspected partner infidelity even after adjusting for previous HIV tests and couple communication around HIV (Anglewicz et al., 2008). Furthermore, there are widespread beliefs regarding the idea of a shared HIV status among married couples, which may not reflect a partner's actual HIV status (Lingappa et al., 2008; Njau et al., 2011). As more people test for HIV, these diagnostic techniques will evolve to blend beliefs about HIV with knowledge of partner's HIV status obtained through disclosure (Reniers & Helleringer, 2011).

In this study, we utilized a longitudinal couples dataset of women and their male partners who were separately tested for HIV and then asked four months later whether they disclosed to their primary partner. This design allowed us to circumvent three common methodological limitations observed in previous studies: (1) the inability to control for the timing of testing and when the disclosure occurred; (2) the reliance on self-reported HIV status in the absence of serostatus biomarkers; and (3) the inability to assess the reliability of self-reported disclosure. Our objectives were three-fold. First, we examined the extent to which self-reported disclosure was confirmed by a partner and whether rates of disclosure

differed by gender (objective 1). In accordance with classical test theory (Lord & Novick, 1968), we expected that a multi-item measure of disclosure (i.e., confirmed disclosure) would be more reliable than a single-item measure from one partner (i.e., self-reported disclosure). Second, using separate models for the two disclosure outcomes, we compared the patterns of associations between individual and relationship factors with self-reported versus confirmed disclosure (objective 2). Third, we investigated whether there were differences in correct assessment of a partner's HIV serostatus using both self-reported disclosure and confirmed disclosure (objective 3). We hypothesized that if confirmed disclosure was a better measure of disclosure, our confidence intervals would be smaller and more precise (objective 2) and respondents would more accurately report on their partner's HIV serostatus (objective 3).

Methods

Participants and procedures

The data were from *Tsogolo la Thanzi* (TLT; “Healthy Futures” in Chichewa), a prospective cohort study that took place from 2009–2011 in the Balaka district of southern Malawi (details available online at: <http://sites.psu.edu/tltc/>). TLT was designed to study how young women and their male partners who have grown up during the HIV/AIDS epidemic make decisions about marriage, childbearing, and sexual behavior. The TLT cohort consisted of approximately 1500 women aged 15–25 who were randomly selected from a household listing of all residents living within a 7-kilometer radius of the Balaka town center. The women were given enumerated tokens and could recruit up to three romantic male partners (e.g., husbands and boyfriends) to participate in the study. Recruiting up to three partners allowed women to enroll secondary partners; however, it was not expected that all women would enroll multiple partners. A romantic partner was defined for respondents as “any relationship that was sexual as well as any relationship where you felt affectionate towards someone even if it was not a sexual relationship”. Male partners who were interested in participating in the study could present the token at the TLT research center to enroll. The use of enumerated tokens allowed for the verification of each male participant as a named partner and the ability to identify his female partner in the sample, which has been a successful approach in other studies (Helleringer & Kohler, 2007).

The TLT baseline sample consisted of 1505 randomly selected women and 616 of their male partners (2121 individuals)—as shown in Figure 1. As noted earlier, several steps were required for male partners to make it into this sample: (1) the women needed to report on the male partner in the survey, (2) give the token to the named male partner, and (3) the male partner needed to present the token at the research center and be matched to a woman in the study. Of the 1505 women at baseline, 17.5% reported on zero partners, 36.2% reported on one partner, 28.3% reported on two partners, and 18% reported on three partners. However, for those who reported on more than one partner, only one male partner per woman was enrolled in the study. It is possible that only more serious male partners felt comfortable enrolling in the study. To provide a brief background on the 1505 women, 42.2% were married, the mean age was 21, and the mean years of education was 7.3. The retention rate

for the cohort of 1505 women was relatively high: 91.1% at wave 3, 89.2% at wave 4, and 82.2% at wave 5.

Women and their male partners were interviewed at 4-month intervals over a period of approximately three years (for a total of 8 waves). At each wave, respondents were administered two separate questionnaires by trained TLT research assistants (local Malawians): (1) a general questionnaire covering life-events during the inter-survey period, including: marriage, divorce, pregnancy, self-reported health, and health-seeking behaviors; and (2) a partnership questionnaire covering questions about each named partner, including: the partner type, relationship status, sexual behavior, and beliefs about the partner's sexual behavior, health behavior, and likelihood of HIV infection.

To address our research questions, we constructed a sub-sample from the baseline set of women and male partners (see Figure 1). At wave 3, a module on relationship power was added to the partnership questionnaire. The relationship power module was designed to understand power dynamics within sexual relationships, as opposed to other relationship types, given the accompanying risks for HIV infection and other negative health outcomes that occur within such relationships (Blanc, 2001; Pulerwitz et al., 2000). Thus, respondents were asked the power statements if they reported a current sexual partner including a spouse, live-in partner, or steady boyfriend/girlfriend. If a respondent reported more than one ongoing sexual relationship, the relationship power questions were asked with regards to the most serious relationship. If the respondent was married, their spouse automatically served as the reference partner. At wave 3, 1338 individuals out of 2121 individuals completed the relationship power module. A couples' dataset was created for respondents who answered the power questions about each other at wave 3. Of the 1338 individuals, 406 did not have participating partner in the study and were dropped. This process resulted in 932 individuals (466 couples) at wave 3 (see Figure 1).

One objective of the TLT study was to assess how changes in knowledge of HIV status influence reproductive and prevention goals. At baseline, women and their male partners were randomly assigned to three HIV testing groups. Group 1 was tested for HIV every four months at each wave. Group 2 was tested for HIV at the end of the first year (wave 4) and at the end of the study (wave 8). Group 3 was only tested for HIV test at the end of the study (wave 8). Thus, at wave 4, approximately two-thirds of the entire TLT sample received an HIV test. Respondents who tested HIV-positive at an earlier wave were not re-tested at wave 4. We used an intention-to-treat approach (Hulley et al., 2013) and included these respondents in our analysis since they were randomly assigned to test at wave 4. Trained TLT counselors tested respondents separately from their partners in private rooms at the TLT research center following the serial algorithm employed by the Malawi Ministry of Health: 1) Determine assay (Abbott Laboratories, USA); 2) Unigold (Trinity Biotech, Ireland) for reactive respondents; and 3) SD Bioline (Standard Diagnostics Inc., Korea) to confirm discordant tests. Of the 466 couples at wave 3, we dropped 28 couples who did not both participate in wave 4 and 232 couples who were not assigned to test for HIV at wave 4. This resulted in 206 couples tested at wave 4 (see Figure 1).

At wave 5, all TLT respondents were asked if they disclosed the results of their last HIV test to their spouse/primary partner. For those tested via TLT's wave 4, their last HIV test most likely occurred at wave 4. Respondents were not informed that their partner was tested at wave 4, allowing for the process of disclosure to occur naturally. We merged the 206 couples who tested at wave 4 with their disclosure data from wave 5 (see Figure 1). After removing couples who had missing data or dissolved relationships, the final sample consisted of 366 individuals (183 couples) (see Figure 1). Of the 183 women in the final sample, 99% reported on a single male partner and for the one woman who reported on two partners, the secondary partner was not enrolled in the TLT. The retention rate was 83% over the period from wave 3 to 5. To check for any bias introduced by attrition, we compared the final sample of 183 couples to the 40 couples who were considered lost (see Figure 1; $28+10+2=40$ couples). The 40 couples were similar to the 183 couples on key demographic and relationship characteristics with the exception of marital status. The 40 couples were slightly less likely to be married (90% versus 96%; $p=0.02$), which is expected as unmarried partners would be less likely to participate in the study than spouses.

Measures

Dependent variables

Self-reported disclosure: At wave 5, respondents were asked, "The last time you tested, to whom if anyone did you tell your results?" Respondents were asked whether they told a spouse/long-term partner, other sexual partner, a relative, friend, doctor/traditional healer, or other person (multiple people could be selected). A binary variable was created to measure whether a respondent disclosed to their spouse/long-term partner (*yes/no*). Regarding disclosure to an "other sexual partner", 100% of respondents reported *no* to this question. Given that the disclosure question was not asked specifically in reference to the participating partner, we performed additional checks. We found that in 97% of couples, both partners listed each other as spouses, which provides assurance that these respondents reported disclosure in reference to their participating partner. We also evaluated the remaining discrepant couples on a case-by-case basis. These couples had been together for over 2.5 years and had participated in multiple waves, and thus we included them in the sample because we were confident that disclosure was in reference to the participating partner.

Confirmed disclosure: At wave 5, respondents were asked whether their partner had tested for HIV in the last 4 months. If yes, respondents were asked, "the last time [your partner] was tested for HIV, did he/she tell you his/her result?" We assumed that those who did not know their partners tested in the last 4 months were not disclosed to. Confirmed disclosure was coded as *yes* only if a respondent reported that he/she disclosed and their partner confirmed it, and as *no* if a respondent reported that he/she did not disclose to their partner.

Correct assessment of partner's HIV status: At wave 5, respondents were asked, "What is the likelihood that your partner is currently infected with HIV?" Response options included *no likelihood* (1), *low* (2), *medium* (3), *high* (4), *I know she/he is infected with HIV* (5), or *Don't know* (88). We used their partner's HIV serostatus from wave 4 to create a binary variable indicating whether or not a respondent correctly assessed partner HIV status at wave 5. If a respondent indicated that their partner had a certain likelihood of HIV infection

(i.e., “I know he/she is infected”) and their partner tested HIV-positive, they were coded as *correct*. If a respondent indicated that their partner had no or low likelihood of HIV infection and their partner tested HIV-negative, they were also coded as *correct*. We included “low” responses in the *correct* category because beliefs about the window period and counseling messages about the need to re-test after three months (e.g., see MacPhail et al., 2008) could prompt respondents to report a “low” (versus “no”) likelihood of HIV infection—which would be correct. Respondents who either incorrectly assessed their partner's HIV status (e.g., reported *no likelihood* but their partner tested positive) or were uncertain of their partner's status (reported a *medium* or *high* likelihood) were coded as *incorrect*.

Independent variables

Perceived HIV status concordance: We used respondents' own HIV status and their perception of a partner's likelihood of HIV infection at wave 4 to create a perceived HIV status concordance variable with three categories: (1) belief that partners are seroconcordant; (2) belief that partners are serodiscordant; and (3) uncertain about a partner's HIV status. Respondents who tested HIV-negative and reported that their partner had no/low likelihood of HIV infection were coded as *seroconcordant*. Similarly, respondents who tested HIV-positive and reported that their partner had a certain likelihood of a HIV infection were coded as *seroconcordant*. Conversely, respondents who tested HIV-positive, but reported that their partner had no/low likelihood of HIV infection, were coded as *serodiscordant*. Respondents who tested HIV-negative, but reported that their partner had a certain likelihood of HIV infection, were also coded as *serodiscordant*. Respondents who reported their partner had a medium or high likelihood of HIV infection or didn't know their partner's likelihood of HIV infection were coded as *uncertain*.

Relationship unity: At wave 3, respondents were asked their level of agreement with three statements: “My partner shows he/she cares about me”; “When I need my partner's assistance, he/she is there to help me”; and “My partner and I discuss important matters together”. Response options ranged from 1 (*strongly disagree*) to 4 (*strongly agree*). We computed the mean score for the three unity items ($\alpha=0.65$). Higher scores indicated greater unity (refer to Conroy, 2014 for details).

Relationship dominance: At wave 3, respondents were asked, “In your relationship, who would you say is generally in charge?” with answer choices of *respondent*, *equal control*, or *partner*. Given the distribution of responses, a binary variable was created to examine *egalitarian/female-dominated* versus *male-dominated* relationships.

Physical violence: At wave 3, respondents were asked, “Has your partner ever hurt you by beating you?” (*yes/no*) (adapted from Pulerwitz et al. 2000). While multiple forms of physical abuse are possible, the question was restricted to “beating” to reflect the predominant local term used to describe physical abuse in Malawi.

Perceived partner infidelity: At wave 5, perceived partner infidelity was captured with the statement: “My partner is probably having sex with someone else”. Response options ranged

from 1 (*strongly disagree*) to 4 (*strongly agree*), which were then collapsed into *disagreement* and *agreement*.

Couple communication around HIV: At baseline, respondents were asked, “have you ever talked with [partner's name] about the risk of HIV?” (*yes/no*).

Disclosure to other relatives/friends: Using the self-reported disclosure question described earlier, we created a binary variable to indicate whether the respondent disclosed the results of their most recent HIV test result to another relative or friend (*yes/no*).

Couple-level characteristics: We included several couple-level characteristics (i.e., values were constant within the couple) as explanatory variables. For relationship type, respondents were asked whether their participating partner was a spouse or other type of partner (e.g., steady boyfriend/girlfriend, infrequent sexual partner, etc.) We created a binary variable for relationship type (0=*non-married*; 1=*married or cohabitating*). For the six couples who were discrepant in their reports of relationship type, we manually recoded the variable based on the response patterns present across both partners and multiple waves of data. Relationship duration was computed by subtracting the date of the survey from the date the couple first started spending time together as more than friends. Using both partners' responses, we computed the mean relationship duration for the couple. For shared children, a binary variable was created to indicate whether a couple had at least one child together. For partners who were discrepant (5.5% of couples), we used the female partner's response, which we believed would be more accurate.

Demographic variables—We included several demographics, namely age and two measures of socio-economic status (years of education and household economic status), as continuous variables. For developing countries such as Malawi, it is common to use asset indices as a proxy measure of household wealth. Using the principal component analysis approach recommended by Filmer and Pritchett (2001), we computed a weighted household wealth index based on the ownership of 9 durable consumer goods (e.g., bicycle, television, radio, bed with mattress) and 21 indicators for characteristics of the household dwelling (e.g., type of roof, type of flooring, electricity). We then computed a couple-level mean using both partners' responses.

Data Analysis

We assessed gender differences in all sample characteristics using either ANOVA (for continuous variables) or chi-square tests (for categorical variables). For objective 1, we conducted Fisher's exact tests to evaluate differences in levels of self-reported and confirmed disclosure by gender and HIV status. For objective 2, we fit a single multilevel structural equation model (SEM) for self-reported and confirmed disclosure to simultaneously compare the patterns of associations for the main predictor variables. An SEM framework allowed us to model the correlated residuals between self-reported and confirmed disclosure. This correlation represents the effects of un-modeled variables that could jointly affect both disclosure outcomes, which if not controlled for, could bias our estimates. Indeed, the SEM results showed that the covariance between the error terms for

the two disclosure outcomes (between-couples) was 2.99 ($p=0.01$), indicating that both outcomes were correlated after controlling for observed characteristics.

Our SEM incorporated a multilevel approach, which allowed us to solve an inherent problem of non-independence with dyadic data (Kenny et al., 2006). We theorized that because dyad members interact and influence each other's behavior, individuals who disclosed their HIV status would be more likely to have a partner who also disclosed as compared to two unpaired men and women in the sample. Failure to account for this non-independence, even small levels (0.05 or less), can lead to overinflated alpha levels and type I errors (Hox & Kreft, 1994). Others have argued that if there is a nested hierarchy in the data, analysts should always use a multilevel (versus single-level) model irrespective of the level of non-independence (Nezlak, 2008). Thus, given the hierarchical structure of the dyadic data, our intention to model the odds of disclosure for the average couple, and the desire to err on the side of caution with non-independence, we conducted a multilevel analysis. As such, we treated both dyad members' estimates of disclosure as the dependent variables while accounting for the higher-level couple unit.

We evaluated the level of non-independence in disclosure by computing the intraclass correlation coefficient (ICC) for each of the disclosure variables. We fit an empty model with no predictors and then used the post-estimation command in Stata 13 to compute the ICC for logistic regression. The ICC was calculated using the variance of the random intercept and the assumed variance of the error term in a logistic regression model ($\pi^2/3$) (Rodriguez & Elo, 2003). For self-reported disclosure, the variance of the random intercept was 1.17 (95% CI: 0.41-3.30). The ICC for self-reported disclosure showed that approximately 29.3% of the variance was explained by the dyad (95% CI: 0.05-0.77). For confirmed disclosure, the variance of the random intercept was 0.45 (95% CI: 0.06-3.22). The ICC for confirmed disclosure showed that 12.1% of the variance was explained by the dyad (95% CI: 0.02-0.49).

The final multilevel SEM was fit using the maximum likelihood estimator in Mplus 6.11. At the individual level (within-couples), we configured a model with two arrows leading from each predictor to self-reported disclosure (first arrow) and confirmed disclosure (second arrow). At the second level (between-couples), we included a random intercept for self-reported disclosure and confirmed disclosure, and allowed the residual errors between the two to co-vary. We computed crude odds ratios (ORs) by fitting the model with each predictor one at a time and then computed adjusted odds ratios (aORs) by including all predictors in the model simultaneously. To confirm that a multilevel model was necessary, we tested the significance of the variance components (between-couples) for the full regression model via a Wald test. The results indicated significant couple-level variability ($\chi^2(3)=29.99$; $p<.05$) even after accounting for all other predictor's variance, supporting the need for a multilevel model.

For objective 3, we hypothesized that if confirmed disclosure was a better gauge of disclosure than self-reported, respondents would be more likely to correctly assess their partner's HIV status. To test this hypothesis, we ran separate chi-squared tests for each disclosure outcome to assess any differences in a respondent's correct assessment of

partner's status. We then conducted a multivariate analysis using the multilevel approach described above to evaluate whether the two disclosure measures were associated with correct assessment. The ICC for correct assessment showed that 29.3% of the variance was explained by the dyad (95% CI: 0.11-0.58). Based on the background literature, we included additional predictors in the multivariate models including: gender, perceptions of a partner's infidelity, HIV status of respondent, and number of partner's past HIV tests. To fit these models, we used the `melogit` command in Stata 13 for multilevel mixed-effects logistic regression with binary outcomes.

Several quality control checks were performed. We examined the level of missing data in the sample, which was minimal. We checked for multi-collinearity among all independent variables by examining the variance inflation factor (VIF), which were all well below the recommended cutoff of 10 (Agresti & Finlay, 2009). We checked for non-normality of all continuous variables. Because relationship unity was slightly skewed to the left, we transformed the mean score by computing the exponential value.

Results

Table 1 shows the demographic characteristics for the analytic sample of the 366 individuals (183 couples). The mean age was 25 and the mean education level reflected a primary school education level. Couples were primarily married/cohabitating (96.2%), together for an average of 5 years, and had at least one child together (77.6%). Over 90% of respondents had communicated with their partner about HIV. On average, respondents reported high relationship unity ($M=3.8$, $SD=0.4$). Women were more likely to report being physically abused (9.3% vs. 0.6%; $p<.05$) and to suspect partners of infidelity (25.7% vs. 6.6%; $p<.05$) as compared to men. Almost 86% of respondents believed that the couple was seroconcordant, 4.1% believed the couple was serodiscordant, and 10.4% were uncertain. By wave 4, 7.4% of the sample tested HIV-positive. Of the 27 HIV-positive respondents, 12 had tested positive at a previous wave. Most couples were seroconcordant negative (89.6%), 4.4% were seroconcordant positive, and 6.0% were serodiscordant.

For objective 1, we compared the rates of self-reported and confirmed disclosure (Figure 2). Over 93% of respondents reported that they had disclosed their HIV status to their partner. However, only 64.3% of respondents had a partner who confirmed their disclosure. There were 12 individuals (3%) who reported that they did not disclose to their partners, but their partner reported they did. For self-reported disclosure, 87.4% of couples had mutually disclosed. However, using the confirmed disclosure measure, only 42.6% of couples had mutually disclosed. Women were significantly more likely to have their disclosure confirmed by a partner as compared to men (72.0% vs. 56.6%; $p<.01$). We also examined whether rates of self-reported disclosure and confirmed disclosure differed by respondent's HIV status but found no significant differences.

For objective 2, we compared the patterns of associations between predictor variables and both self-reported and confirmed disclosure (Table 2). Because we used a conservative model that controlled for the correlated residuals between self-reported and confirmed disclosure, we also report marginally significant findings ($p<.10$) below. For self-reported

disclosure, the multivariate results showed that the odds of self-reported disclosure for the average couple were approximately 27 times higher when respondents communicated with partners about HIV ($aOR=26.69$; $p<.001$). We also found that the odds of self-reported disclosure were 1.05 times higher for each one-unit increase in the transformed (exponential value) relationship unity variable ($aOR=1.05$; $p=.073$). In addition, the odds of self-reported disclosure were 89% lower when respondents were uncertain about their partner's HIV status ($aOR=0.11$; $p<.05$). Finally, the odds of self-reported disclosure were 91% lower for respondents who had disclosed recent HIV test results to another relative or friend ($p<.001$).

In general, patterns of predictors for confirmed disclosure were similar to self-reported disclosure with a few exceptions for demographic variables (Table 2). For confirmed disclosure, the multivariate results showed that the odds of confirmed disclosure for the average couple were 55% lower for men as compared to women ($aOR=0.44$; $p<.05$). The multivariate results also showed that the odds of confirmed disclosure decreased by 8% for each one-year increase in age ($aOR=0.92$; $p<.05$). In addition, the odds of confirmed disclosure increased by 19% for each additional year of education ($aOR=1.19$; $p<.05$). Consistent with self-reported disclosure, the odds of confirmed disclosure were 2.81 times higher when respondents had communicated with partners about HIV ($p<.05$). Similarly, the odds of confirmed disclosure were 59% lower when respondents were uncertain about their partner's HIV status ($aOR=0.41$; $p=.062$). Regardless of statistical significance of each predictor, all odds ratios were attenuated and confidence intervals became smaller in the model for confirmed disclosure as compared to self-report—suggesting that the results for confirmed disclosure are more precise.

For objective 3, we examined the association between the two HIV disclosure measures, other explanatory variables, and correct assessment of partner's HIV status at wave 5 (Table 3). Approximately 80% of respondents correctly assessed their partner's HIV status. For self-reported disclosure, a respondent's accuracy in assessing their partner's HIV status was higher when a partner disclosed than when they did not (79.5% vs. 64.0%), although this difference was not statistically significant. For confirmed disclosure, a respondent's accuracy in assessing their partner's HIV status was significantly higher with confirmed disclosure as compared to non-disclosure (85.0% vs. 66.9%; $p<.001$). The multivariate results showed that confirmed disclosure (Model 2), but not self-reported disclosure (Model 1), was predictive of correctly assessing a partner's HIV status (Table 3). The odds of correctly assessing a partner's HIV status were 2.76 times higher with confirmed disclosure as compared to non-disclosure ($aOR=2.76$; $p<.01$). We also found that correct assessment of a partner's HIV status was significantly and positively associated with being male ($aOR=2.91$; $p<.01$), lower perceived partner infidelity ($aOR=0.28$, $p<.01$), higher relationship unity ($aOR=1.03$; $p<.05$), and if the respondent tested HIV-negative at wave 4 ($aOR=0.11$; $p<.001$).

Discussion

Given that partners were tested for HIV as part of a research study rather than on their own, we expected to find very high rates of confirmed disclosure (objective 1). Instead, we found that while over 93% of respondents reported that they disclosed, there were major

discrepancies when partners' confirmed reports were taken into account; only 64% of respondents had disclosed according to the confirmed disclosure measure. We also found significant gender differences in levels of confirmed disclosure, with men more likely to confirm a female partner's report than women to confirm men's. One explanation for these differences may be that social desirability bias is higher in men's self-reports. Two other studies from Malawi and Kenya also found that women were less likely to confirm their partner's reported disclosure as compared to men (Anglewicz et al., 2011; Katz et al., 2009). Our findings are also consistent with other spousal reports on HIV-related behavior. In Kenya, Miller and colleagues (2009) found that when husbands and wives were discordant about whether they communicated about AIDS, men were generally more likely to say "yes" while women were more likely to say "no". Together, these results highlight the role that gender plays in reporting of HIV disclosure, suggesting that future efforts focus on addressing the issue of social desirability among men.

In our primary analysis (objective 2), we found that communication about HIV risk was associated with disclosure regardless of whether a partner confirmed. Open communication regarding the risks of HIV may provide a strong foundation for couples to later discuss HIV test results with each other, as other studies have demonstrated (King et al., 2008; Maman et al., 2001). In contrast, relationship unity was not associated with confirmed disclosure and only marginally associated with self-reported disclosure in the positive direction. Together, these findings suggest that efforts to improve HIV disclosure may yield greater returns by focusing more on communication skills than aspects of unity. We also found that uncertainty about a partner's HIV status was negatively associated with disclosure regardless of whether a partner confirmed (although only marginally significant for confirmed disclosure). This pattern provides evidence of the role that perceptions play in partners' decisions to disclose, which could be circumvented if couples jointly tested for HIV and disclosed together. Finally, disclosure to a friend/relative was negatively associated with self-reported disclosure to a primary partner, which may suggest that the provision of social support from others may obviate the need to inform a spouse. However, this measure was assessed at wave 5 when the disclosure outcome was captured (cross-sectional), which does not allow us to make statements about causality. The same results did not hold for confirmed disclosure.

Beyond these associations, there is evidence that a partner's confirmed report may improve the reliability of self-reported disclosure. The odds ratios and confidence intervals for HIV communication and uncertainty about a partner's HIV status became more precise in the model for confirmed disclosure than for self-reported disclosure. According to classical test theory (Lord et al., 1968), adding items to a scale increases the reliability of the measure by reducing error score variance and increasing the true score variance. With less error in the measure, the 95% confidence intervals should theoretically become smaller—as we found for confirmed disclosure.

In our final analysis, we hypothesized that if confirmed disclosure was a better measure, respondents should be more accurate at assessing their partner's HIV status (objective 3). Indeed, we found that confirmed disclosure, but not self-reported disclosure, was significantly associated with correctly assessing a partner's HIV status. In addition, we found

that respondents were more accurate about their partner's HIV status if they were male, had higher relationship unity, had lower perceived partner infidelity, and were HIV-negative. Although we did not examine sexual risk behavior as an outcome of disclosure, another study from Malawi among married couples found that perceived partner HIV status had greater influence than actual partner HIV status on the acceptability of and use of condoms (Anglewicz & Clark, 2013). Thus, while efforts aimed at increasing knowledge of one's own HIV status through uptake of HIV testing are vitally important for HIV prevention, the powerful role of individual beliefs and relationship factors should not be underestimated.

Limitations

Two limitations of this study are noteworthy. First, to answer novel research questions related to HIV disclosure within couples, we selected a sub-sample of 183 couples from the full TLT sample of 1500 young women and their romantic male partners. The selection of the couples sub-sample limits our ability to draw conclusions about the larger TLT study population. While the sub-sample was significantly smaller than the full sample, most of the exclusions related to being a couple (e.g., still together, both enrolled in the study) rather than to factors like attrition. The largest drop in sample size occurred for respondents who did not meet the criteria of having an ongoing sexual relationship with a partner in enrolled in the study (see Figure 1). While TLT enrolled male partners using a looser definition of “romantic” partner, our analysis was restricted to sexually active couples. There are likely to be differences in the frequency and correlates of disclosure between sexual versus non-sexual partnerships. For example, disclosure to non-sexual partners may be a precursor to sex and driven more by desires to advance a relationship or establish commitment. While such differences are important and need to be better understood, our focus on couples who are already engaging in sex with the potential to transmit the HIV virus calls for more immediate attention. Because our goal was to study these couples, the couples sub-sample was more likely to be married (97.8% vs. 42.2%; $p<.001$) with lower years of education ($M=6.5$ vs. 7.3; $p<.001$) than the full sample of 1505 women. Thus, we cannot generalize to the sample of 1505 women who are inherently different. Future research is needed to study HIV disclosure in representative samples of couples at the population level and within other types of couples such as pre-marital partnerships, non-sexual partnerships, extra-marital partnerships, and older couples with different relationship dynamics. Our research provides a starting point for this area of research.

Second, because we used a subset of a population-based sample (rather than HIV-positive individuals or a sample of clinic patients who may be sicker and more likely to be HIV positive), the majority of respondents were HIV-negative. Thus, low numbers of HIV-positive individuals in the sample limited our ability to understand how patterns of disclosure may differ by HIV status. This limitation extends to our ability to adequately test for associations between couple HIV status concordance and disclosure. Third, given our relatively small sample size, we did not have enough statistical power to adequately assess statistical interactions by gender. We did test for statistical interactions in initial analyses, which were non-significant—potentially due to a lack of power. Future disclosure studies should replicate our findings using larger, more diverse samples of couples with mixed HIV status. Fourth, we found that couple communication was a significant predictor of

disclosure, however, this variable was only collected at baseline. Therefore, we could not account for any changes in couple communication over time. Finally, while we can infer that confirmed disclosure is a superior measure in comparison to a single partner's self-report, we did not have the ability to validate the measure against a gold standard. Our research makes an incremental advance in the field, but does not solve all problems related to self-reported survey data.

Study Implications

Despite these limitations, this study is one of the first in sub-Saharan Africa to evaluate whether having dyadic data from two partners improves the measurement of disclosure based on a single partner's account. Indeed, the use of confirmatory reports from a partner may be even more important in samples of HIV-positive individuals who might be more inclined to provide socially-desirable responses due to stigma and fear. Thus, we recommend that researchers and policymakers take caution when interpreting and applying findings from analyses on self-reported disclosure from a single individual, as these reports may be more prone to Type I errors.

Our findings highlight the need for couple-focused interventions in sub-Saharan Africa, which have shown to be effective for HIV prevention (Burton et al., 2010). We found that even under highly favorable conditions, couples may not discuss their HIV test results with each other. There are several implications for couple-focused interventions. First, efforts could center on improving HIV-related communication and other relationship dynamics to provide couples with the skills necessary to test for HIV and disclose together. For example, in South Africa, research is underway to test a couples-based intervention that aims to build positive relationship dynamics as a pathway to increased uptake of couples HIV testing and counseling (CHTC) (Darbes et al., 2014). However, our recommendation for such interventions is based on this study's findings among mostly married, seroconcordant couples. More research is needed to understand how to effectively intervene with serodiscordant couples and those with different relationship dynamics, who may require more intensive efforts. Second, CHTC itself is an effective strategy to reduce the risk of HIV transmission in serodiscordant couples (Allen et al., 2003) and could alleviate the burden of HIV communication and disclosure on partners by shifting responsibility to trained healthcare providers. Lessons could be drawn from HIV testing interventions using peer or community-based counselors (Baiden et al., 2007; Peltzer et al., 2010), who may be familiar with the challenges faced by couples in their communities and thus well positioned to encourage couples to seek CHTC.

Conclusions

This study illustrates the persistent challenges faced by couples around HIV disclosure, which are likely to be more pronounced in serodiscordant or other types of couples. Mutual HIV disclosure within committed relationships is required to improve accuracy of risk perceptions and the adoption of healthier behaviors—regardless of HIV status. Non-disclosure of HIV status to spouses and partners has been noted as a key concern for initiation of antiretroviral therapy (ART), retention in care, and ART adherence, prevention of mother-to-child transmission, and treatment as prevention in serodiscordant couples (e.g.,

see Medley et al., 2004; Tenthani et al., 2014). This is a particularly timely issue in sub-Saharan Africa as more people qualify for ART at higher CD4 counts through policies such as “Option B+” (Tenthani et al., 2014). As treatment options evolve rapidly in sub-Saharan Africa, it remains unclear how these healthcare changes intersect with disclosure in couples. Future studies utilizing dyadic data are needed to more fully understand the complex array of psychosocial, socio-economic, and relationship factors affecting HIV disclosure in couples to identify appropriate opportunities to intervene at the couple-level.

Acknowledgments

This research uses data from *Tsogolo La Thanzi* (TLT), a research project designed by Jenny Trinitapoli and Sara Yeatman, and funded by the National Institute of Child Health and Human Development [R01-HD058366]. Refer to <http://projects.pop.psu.edu/tlt> for more information about the TLT dataset (including response rates), to request data access, and for replication files. This research was also supported by two training grants [F31-MH093260 and T32-MH19105] and a center grant [P30-MH062246] from the National Institute of Mental Health. The authors thank Dr. Torsten Neilands and Dr. Steven Gregorich for statistical guidance.

References

- Agresti, A.; Finlay, B. *Statistical Methods for the Social Sciences*. 4th. Upper Saddle River, New Jersey: Pearson Prentice Hall; 2009.
- Allen S, Meizen-Derr J, Kautzman M, et al. Sexual behavior of HIV discordant couples after HIV counseling and testing. *AIDS*. 2003; 17:733–740. [PubMed: 12646797]
- Anglewicz P, Chintsanya J. Disclosure of HIV status between spouses in rural Malawi. *AIDS Care*. 2011; 23(8):998–1005. [PubMed: 21390889]
- Anglewicz P, Clark S. The effect of marriage and HIV risks on condom use acceptability in rural Malawi. *Social Science & Medicine*. 2013; 97:29–40. [PubMed: 24161086]
- Anglewicz PA, Bignami-Van Assche S, Clark S, Mkandawire J. HIV risk among currently married couples in rural Malawi: What do spouses know about each other? *AIDS and Behavior*. 2008; 14(1):103–112. [PubMed: 19051003]
- Baiden F, Akanlu G, Hodgson A, Akweongo P, Debpuur C, Binka F. Using lay counsellors to promote community-based voluntary counselling and HIV testing in rural northern Ghana: A baseline survey on community acceptance and stigma. *J Biosocial Science*. 2007; 39:721–733. [PubMed: 17207292]
- Black BP, Miles MS. Calculating the risks and benefits of disclosure in African American women who have HIV. *Journal of Obstetric and Gynecological Neonatal Nursing*. 2002; 31:688–697.
- Blanc A. The effect of power in sexual relationships on sexual and reproductive health: An examination of the evidence. *Studies in Family Planning*. 2001; 32(3):189–213. [PubMed: 11677692]
- Brou H, Djohan G, Becquet R, et al. When Do HIV-Infected Women Disclose Their HIV Status to Their Male Partner and Why? A Study in a PMTCT Programme, Abidjan. *PLoS Medicine*. 2007; 4(12):e342. [PubMed: 18052603]
- Burton J, Darbes LA, Operario D. Couples-focused behavioral interventions for prevention of HIV: Systematic review of the state of evidence. *AIDS Behavior*. 2010; 14:1–10. [PubMed: 18843530]
- Conroy AA. Gender, power, and intimate partner violence: A study on couples from rural Malawi. *Journal of Interpersonal Violence*. 2014; 29(5):866–888. [PubMed: 24227592]
- Crepaz N, Marks G. Serostatus disclosure, sexual communication and safer sex in HIV-positive men. *AIDS Care*. 2004; 15:379–87. [PubMed: 12745398]
- Darbes LA, Van Rooyen H, Hosegood V, Ngubane T, Johnson MO, Fritz K, McGrath N. Uthando Lwethu (‘our love’): a protocol for a couples-based intervention to increase testing for HIV: a randomized controlled trial in rural KwaZulu-Natal, South Africa. *Trials*. 2014; 15(64)
- Desgrees-Du-Lou A, Brou H, Djohan G, Becquet R, Ekouevi DK, Zanou B. Beneficial effects of offering prenatal HIV counselling and testing on developing a HIV preventive attitude among couples, Abidjan, 2002–2005. *AIDS and Behavior*. 2007; 13(2):348–355. [PubMed: 17985231]

- Filmer D, Pritchett LH. Estimating wealth effects without expenditure data--or tears: An application to educational enrollments in states of India. *Demography*. 2001; 38(1):115–132. [PubMed: 11227840]
- Helleringer S, Kohler HP. Sexual network structure and the spread of HIV in Africa: evidence from Likoma Island, Malawi. *AIDS*. 2007; 21(17):2323–2332. [PubMed: 18090281]
- Hox JJ, Kreft IGG. Multilevel analysis methods. *Sociological Methods and Research*. 1994; 22:283–299.
- Hulley, SB.; Cummings, SR.; Browner, WG.; Grady, DG.; Newman, TB. *Designing Clinical Research*. 4th. Lippincott Williams & Wilkins; 2013.
- Katz DA, Kiarie JN, John-Stewart GC, Richardson BA, John FN, Farquhar C. HIV testing men in the antenatal setting: understanding male non-disclosure. *International Journal of STD and AIDS*. 2009; 20(11):765–767. [PubMed: 19833691]
- Kenny, DA.; Kashy, DA.; Cook, WL. *Dyadic data analysis*. New York: The Guilford Press; 2006.
- King R, Katuntu D, Lifshay J, et al. Processes and Outcomes of HIV Serostatus Disclosure to Sexual Partners among People Living with HIV in Uganda. *AIDS and Behavior*. 2008; 12:232–243. [PubMed: 17828450]
- Lewis MA, McBride CM, Pollak KI, Puleo E, Butterfield RM, Emmons KM. Understanding health behavior change among couples: An interdependence and communal coping approach. *Social Science & Medicine*. 2006; 62:1369–1380. [PubMed: 16146666]
- Lingappa JR, Lambdin B, Bukusi EA, Ngure K, Kavuma L, Inambao M, et al. Regional differences in prevalence of HIV-1 discordance in Africa and enrollment of HIV-1 discordant couples into an HIV-1 prevention trial. *PLOS One*. 2008; 3(1):e1411. [PubMed: 18183292]
- Lord, FM.; Novick, MR. *Statistical theories of mental test scores*. Reading, MA: Addison-Wesley; 1968.
- Louibiere S, Peretti-Watal P, Boyer S, Blanche J, Abega SC, Spire B. HIV disclosure and unsafe sex among HIV-infected women in Cameroon: Results from the ANRS-EVAL study. *Social Science & Medicine*. 2009; 69:885–891. [PubMed: 19560244]
- MacPhail CL, Pettifor A, Coates T, Rees H. You must do the test to know your status: Attitudes to HIV voluntary counseling and testing for adolescents among South African youth and parents. *Health Education and Behavior*. 2008; 35(1):87–104. [PubMed: 16870815]
- Maman S, Hogan NM, Kilonza GP. Women's barriers to HIV-1 testing and disclosure: challenges for HIV-1 voluntary counselling and testing. *AIDS Care*. 2001; 13(5):595–603. [PubMed: 11571006]
- Maman S, Mbwambo J, Hogan NM, Weiss E, Kilonzo GP, Sweat MD. High rates and positive outcomes of HIV serostatus disclosure to sexual partners: Reasons for cautious optimism from a voluntary counseling and testing clinic in Dar es Salaam, Tanzania. *AIDS and Behavior*. 2003; 7:373–382. [PubMed: 14707534]
- Maman S, Mbwambo JK, Hogan NM, Kilonza GP, Campbell JC, Weiss E, Sweat MD. HIV-positive women report more life-time partner violence: Findings from a voluntary counselling and testing clinic in Dar es Salaam, Tanzania. *American Journal of Public Health*. 2002; 92:1331–1337. [PubMed: 12144993]
- Medley A, Garcia-Moreno C, McGill S, Maman S. Rates, barriers and outcomes of HIV serostatus disclosure among women in developing countries: implications for prevention of mother-to-child transmission programmes. *Bulletin of the World Health Organization*. 2004; 82:299–307. [PubMed: 15259260]
- Miller A, Golding L, Ngula KW, et al. Couples' communication on sexual and relational issues among the Akamba in Kenya. *African Journal of AIDS Research*. 2009; 8(1):51–60. [PubMed: 25864476]
- Nezlek JB. An Introduction to Multilevel Modeling for Social and Personality Psychology. *Social and Personality Psychology Compass*. 2008; 2(2):842–860.
- Njau B, Watt MH, Ostermann J, Manongi R, Sikkema KJ. Perceived acceptability of home-based couples voluntary HIV counseling and testing in Northern Tanzania. *AIDS Care*. 2011; 24(4):413–419. [PubMed: 21939369]
- Obermeyer CM, Bajjal P, Pegurri E. Facilitating HIV Disclosure Across Diverse Settings: A Review. *American Journal of Public Health*. 2011; 101(6):1011–1023. [PubMed: 21493947]

- Olley BO, Seedat S, Stein DJ. Self-Disclosure of HIV Serostatus in Recently Diagnosed Patients with HIV in South Africa. *African Journal of Reproductive Health*. 2004; 8(2):71–76. [PubMed: 15623121]
- Peltzer K, Tabane C, Matseke G, Simbayi L. Lay counsellor-based risk reduction intervention with HIV positive diagnosed patients at public HIV counselling and testing sites in Mpumalanga, South Africa. *Evaluation and Program Planning*. 2010; 33:379–385. [PubMed: 20416948]
- Porter L, Hao L, Bishai D, Serwadda D, Wawer MJ, Lutalo T, Gray R. HIV status and union dissolution in sub-Saharan Africa: The case of Rakai, Uganda. *Demography*. 2004; 41(3):465–482. [PubMed: 15461010]
- Pulerwitz J, Gortmaker SL, DeJong W. Measuring relationship power in HIV/STD research. *Sex Roles*. 2000; 42(7/8):637–660.
- Reniers G, HELLERINGER S. Serosorting and the Evaluation of HIV Testing and Counseling for HIV Prevention in Generalized Epidemics. *AIDS and Behavior*. 2011; 15:1–8. [PubMed: 20683650]
- Rodriguez G, Elo I. Intra-class correlation in random-effects models for binary data. *The Stata Journal*. 2003; 3(1):32–46.
- Simoni JM, Pantalone DW. Secrets and Safety in the Age of AIDS: Does HIV Disclosure Lead to Safer Sex? *Topics in HIV Medicine*. 2004; 12(4):110–118.
- Stirratt MJ, Remien RH, Smith A, Copeland OQ, Dolezal C, Krieger D. The role of HIV serostatus disclosure in antiretroviral medication adherence. *AIDS and Behavior*. 2006; 10(5):483–493. [PubMed: 16721505]
- Suzan-Monti M, Kouanfack C, Boyer S, et al. Impact of HIV Comprehensive Care and Treatment on Serostatus Disclosure among Cameroonian Patients in Rural District Hospitals. *PLOS One*. 2013; 8(1)
- Tenthani L, Haas AD, Tweya H, et al. Retention in care under universal antiretroviral therapy for HIV-infected pregnant and breastfeeding women ('Option B+') in Malawi. *AIDS*. 2014; 28:589–598. [PubMed: 24468999]
- Venkatesh KK, Madiba P, De Bruyn G, Lurie MN, Coates TJ, Gray GE. Who Gets Tested for HIV in a South African Urban Township? Implications for Test and Treat and Gender-Based Prevention Interventions. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2011; 56(2):151–165. [PubMed: 21084993]
- Watkins SC. Navigating the AIDS epidemic in rural Malawi. *Population and Development Review*. 2004; 30(4):673–705.10.1111/j.1728-4457.2004.00037.x
- WHO. Gender dimensions of HIV status disclosure to sexual partners: Rates, barriers, and outcomes; A review paper. Geneva, Switzerland: World Health Organization; 2003.
- Wong LH, Van Rooyen H, Modiba P, et al. Test and Tell: Correlates and Consequences of Testing and Disclosure of HIV Status in South Africa (HPTN 043 Project Accept). *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2009; 50(2):215–222. [PubMed: 19131885]
- Wong, RK.; Higgins, J.; Nakyanjo, N. Relationship Dynamics and HIV Disclosure: A Qualitative Study in Rakai District, Uganda; Paper Presented at the Annual Meeting of the American Public Health Association; Washington, D.C.. 2011.

Research Highlights

- Self-reported HIV disclosure and a partner's confirmatory account are often discrepant.
- Confirmatory reports of a partner's disclosure may improve the reliability of self-reports.
- Confirmatory reports of a partner's disclosure predict correct assessment of HIV status.

Figure 1. Flow chart for the creation of the analytic sample of couples, Tsogolo La Thanzi study

Figure 2. Discrepancies between self-reported disclosure and confirmed disclosure for the entire couples full sample ($N=366$ individuals; 183 couples) and by gender. *Chi-square test, $p < .05$.

Table 1

Descriptive characteristics for the couples sample, Tsogolo La Thanzi ($N=183$ couples).

Characteristics	Wave	Total (N=366)		Women (N=183)		Men (N=183)	
		%	Mean (SD)	%	Mean (SD)	%	Mean (SD)
Individual-level variables							
Age in years (range: 16-42)	3		24.7 (4.6)*		22.1 (2.7)		27.3 (4.6)
Years of education (range: 0-13)	3		7.1 (2.8)*		6.5 (2.5)		7.7 (3.1)
Couple communication around HIV/AIDS	1	90.8		88.6		92.9	
Relationship unity (mean score; range 1-4)	3		3.8 (0.4)		3.8 (0.4)		3.8 (0.4)
Ever physically abused by partner	3	4.9*		9.3		0.6	
Relationship dominated by male partner	3	86.6		85.3		88.0	
Having sex with another partner in past 4 months	4	2.4*		0.0		4.8	
Consistent condom use with partner in the study	4	1.6		1.1		2.2	
Perceived partner infidelity	5	16.1*		25.7		6.6	
Perceived HIV status concordance	4						
Belief that partners are seroconcordant		85.5*		80.3		90.7	
Belief that partners are serodiscordant		4.1		4.4		3.9	
Unsure about partner's status		10.4		15.3		5.5	
Disclosure of last HIV test result to other relative/friend	5	24.9		25.7		24	
HIV-positive (serostatus)	4	7.4		8.2		6.6	
Couple-level variables							
Relationship type (married)	4	96.2		96.2		96.2	
Household wealth index (range: -2.5 to 6.5)	3		-1.1 (1.5)		-1.1 (1.5)		-1.1 (1.5)
Relationship duration in years (range: 0.5-12)	3		5.0 (2.7)		5.0 (2.7)		5.0 (2.7)
Shared child(ren)	3	77.6		77.6		77.6	

Note.* ANOVA or chi-square tests indicates a significant gender difference at $p<0.05$. **A.bbr.** AIDS=Acquired immunodeficiency syndrome. HIV= Human immunodeficiency virus. SD=Standard deviation.

Table 2

Odds ratios from a multilevel structural equation model predicting HIV status disclosure, for Wave 5: Tsogolo La Thanzi ($N=366$ individuals; 183 couples).

Variable	Self-Reported Disclosure		Confirmed Disclosure	
	Crude OR (95% CIs)	Adjusted OR (95% CIs)	Crude OR (95% CIs)	Adjusted OR (95% CIs)
<i>Individual-level variables</i>				
Gender (Reference: Female)	0.70 (0.27, 1.83)	0.55 (0.09, 3.24)	0.41 (0.25, 0.69)***	0.44 (0.22, 0.91)*
Age (years)	0.95 (0.86, 1.06)	0.95 (0.79, 1.15)	0.91 (0.86, 0.97)**	0.92 (0.84, 0.99)*
Years of education	1.06 (0.87, 1.29)	1.26 (0.95, 1.68) [†]	1.10 (1.00, 1.19)*	1.19 (1.07, 1.34)**
Couple communication around HIV/AIDS	10.80 (2.36, 49.46)**	26.69 (3.31, 214.35)***	2.42 (1.01, 5.81)*	2.81 (1.09, 7.27)*
Relationship unity (exponential)	1.07 (1.02, 1.12)**	1.05 (1.00, 1.11) [†]	1.01 (0.99, 1.03)	1.00 (0.97, 1.02)
Ever physically abused by partner	1.46 (0.12, 18.09)	6.99 (0.22, 226.83)	1.36 (0.41, 4.52)	0.99 (0.27, 3.64)
Relationship dominated by male partner (Reference: Egalitarian/Male-dominated)	1.64 (0.39, 6.87)	1.24 (0.14, 11.08)	0.79 (0.36, 1.75)	0.69 (0.28, 1.66)
Perceived HIV status concordance Belief that partners are seroconcordant (Reference)	1.00	1.00	1.00	1.00
Belief that partners are serodiscordant	0.73 (0.04, 14.89)	0.18 (0.01, 5.85)	1.12 (0.28, 4.43)	1.05 (0.22, 4.97)
Unsure about partner's status	0.10 (0.02, 0.47)**	0.11 (0.02, 0.79)*	0.56 (0.24, 1.29)	0.41 (0.16, 1.05) [†]
Disclosure to relative or friend	0.14 (0.04, 0.45)***	0.09 (0.02, 0.46)***	0.75 (0.41, 1.37)	0.74 (0.38, 1.42)
<i>Couple-level variables</i>				
Relationship type (Reference: Unmarried)	3.96 (0.28, 56.10)	0.58 (0.00, 105.00)	1.48 (0.33, 6.64)	2.08 (0.35, 12.44)
Household wealth index	0.79 (0.55, 1.07)	0.72 (0.43, 1.23)	1.00 (0.82, 1.18)	1.00 (0.80, 1.24)
Relationship duration (years)	1.08 (0.85, 1.36)	0.91 (0.62, 1.34)	1.04 (0.93, 1.17)	1.07 (0.92, 1.25)
Shared child(ren)	2.05 (0.50, 8.35)	1.73 (0.15, 20.47)	1.49 (0.74, 2.99)	1.33 (0.52, 3.35)

Note.

[†] $p < .10$;

* $p < .05$;

** $p < .01$;

*** $p < .001$. **Abbr.** AIDS=Acquired immunodeficiency syndrome. CIs=Confidence intervals. HIV= Human immunodeficiency virus. OR=Odds ratio.

Table 3

Odds ratios from multi-level mixed effects logistic regression models predicting a respondent's correct assessment of their partner's HIV status using self-reported disclosure (Model 1) and confirmed disclosure (Model 2 for Wave 5: Tsogolo La Thanzi ($N=366$ individuals; 183 couples)).

Variable	Model 1		Model 1	
	Self-Reported Disclosure (as the primary predictor)	Adjusted OR (95% CIs)	Confirmed Disclosure (as the primary predictor)	Adjusted OR (95% CIs)
<i>Primary predictors</i>				
Partner's self-reported disclosure	2.87 (0.94, 8.81)			
Partner's confirmed disclosure			2.76 (1.35, 5.64)	**
<i>Secondary predictors</i>				
Gender (Reference: Female)	3.34 (1.67, 6.66)	**	2.91 (1.43, 5.94)	**
Perceived partner infidelity	0.33 (0.15, 0.77)	*	0.28 (0.11, 0.67)	**
Relationship unity (exponential)	1.04 (1.01, 1.06)	**	1.03 (1.00, 1.06)	*
Communication with partner about HIV/AIDS	1.23 (0.44, 3.41)		1.17 (0.41, 3.36)	
HIV-positive test result at wave 4 (respondent)	0.10 (0.04, 0.30)	***	0.11 (0.04, 0.35)	***
Number of previous HIV tests at TLT (partner)	1.31 (1.01, 1.70)	*	1.20 (0.92, 1.57)	

Note.

* $p<.05$;

** $p<.01$;

*** $p<.001$.

Abbr. AIDS=Acquired immunodeficiency syndrome. CIs=Confidence intervals. HIV= Human immunodeficiency virus. OR=Odds ratio. TLT=Tsogolo La Thanzi.