Published in final edited form as:

Front Biol (Beijing). 2015 June ; 10(3): 221–229. doi:10.1007/s11515-015-1359-x.

Role of upstream stimulatory factor 2 in diabetic nephropathy

Shuxia Wang

Department of Pharmacology and Nutritional Sciences, University of Kentucky, Lexington, KY 40536, USA

Shuxia Wang: swang7@uky.edu

Abstract

Diabetic nephropathy (DN) is the most common cause of end-stage renal disease (ESRD). About 20%–30% of people with type 1 and type 2 diabetes develop DN. DN is characterized by both glomerulosclerosis with thickening of the glomerular basement membrane and mesangial matrix expansion, and tubulointerstitial fibrosis. Hyperglycemia and the activation of the intra-renal renin-angiotensin system (RAS) in diabetes have been suggested to play a critical role in the pathogenesis of DN. However, the mechanisms are not well known.

Studies from our laboratory demonstrated that the transcription factor—upstream stimulatory factor 2 (USF2) is an important regulator of DN. Moreover, the renin gene is a downstream target of USF2. Importantly, USF2 transgenic (Tg) mice demonstrate a specific increase in renal renin expression and angiotensin II (AngII) levels in kidney and exhibit increased urinary albumin excretion and extracellular matrix deposition in glomeruli, supporting a role for USF2 in the development of diabetic nephropathy. In this review, we summarize our findings of the mechanisms by which diabetes regulates USF2 in kidney cells and its role in regulation of renal renin-angiotensin system and the development of diabetic nephropathy.

Keywords

USF2; renin-angiotensin system; TGF- β ; renal fibrosis

Diabetic nephropathy

Diabetic nephropathy (DN) is one of the most prevalent diabetic microvascular complications and characterized by both glomerulosclerosis with thickening of the glomerular basement membrane and mesangial matrix expansion, and tubulointerstitial fibrosis (Lim, 2014). A relationship between hyperglycemia and DN has been established in both animal and clinical studies. DN starts with microalbuminuria through proteinuria, azotaemia, and culminates in ESRF. Before the onset of overt proteinuria, renal function changes include renal hyperfiltration, hyperperfusion, and increasing capillary permeability to macromolecules. During this early stage, the hypertrophic glomeruli exhibit normal

Compliance with ethics guidelines

Shuxia Wang declares that there is no conflict of interest. This article does not contain studies with human subjects. It contains studies with animal subjects, which conformed to the National Institutes of Health Guide for the Care and Use of Laboratory Animals and were approved by the University of Kentucky Institutional Animal Care and Use Committee.

structure. After several years of diabetes, the increased thickness of the glomerular basement membrane, as well as mesangial expansion with the accumulation of matrix are the fundamental changes. These changes correlate strongly with the clinical onset of proteinuria, hypertension and kidney failure. Production of the extracellular matrix by mesangial cells contributes to progressive glomerulosclerosis.

The mechanisms of DN are complicated. Both metabolic and hemodynamic factors are involved in the development of DN (reviewed by Cooper(2001) and Forbes et al. (2007)). Metabolic factors include advanced glycated end products (AGE) (Petrica et al., 2015), oxidative stress (Dabhi and Mistry, 2015), aldose reductase/polyol pathway flux and hexosamine flux (Giacco and Brownlee, 2010). Hemodynamic factors include the renin-angiotensin system (RAS), endothelin and nitric oxide etc. These factors act independently or coordinately to activate intracellular second messengers such as PKC, NF- κ B, and MAPK or various cytokines including TGF- β , connective tissue growth factor (CTGF), vascular endothelial growth factor (VEGF), insulin-like growth factor (IGF), and inflammatory cytokines (IL-1, IL-6 or TNF- α) (Navarro-González and Mora-Fernández, 2008) leading to renal functional and structural changes as described above. Among the many mediators of DN, TGF- β has been considered to be an important cytokine in mediating renal fibrosis in diabetes.

The renin angiotensin system (RAS) and DN

Systemic RAS (classic RAS) and its role in DN

The RAS is a multi-step enzymatic cascade to produce a family of angiotensin peptides that control blood pressure and fluid homeostasis. The RAS has been acknowledged as an endocrine, paracrine, autocrine, and intracrine system (Navar et al., 2002; Kobori et al., 2007; Re, 2007). The components of the RAS, including angiotensinogen, renin, and ACE are present systemically for endocrine production of AngII, termed the classic RAS. Angiotensinogen, the only known precursor to AngII, is constitutively secreted from liver hepatocytes as well as other cell types and generally considered to be non-rate limiting in the production of systemic AngII (Peach, 1977). Angiotensinogen is cleaved by the aspartyl protease renin to generate the decapeptide angiotensin I (AngI). For systemic production of AngII, renin synthesis and release from the kidney juxtaglomerular (JG) cells of the afferent arteriole is the rate limiting step. Hydrolysis of the C-terminal amino acids of AngI by the dicarboxypeptidase angiotensin converting enzyme (ACE) results in the formation of AngII, the primary peptide of the RAS with biological activity. A homolog of ACE, termed ACE-2, has been recently discovered, which hydrolyzes AngI to Ang(1–9) and AngII to Ang(1–7) (Donoghue et al., 2000). ACE2 has been suggested as the catabolic arm of the RAS (Ferrario et al., 2005), blunting the actions of AngII and favoring effects of the vasodilator Ang(1–7). AngII exhibits the majority of its well-known physiological effects via the angiotensin type 1 receptor (AT1 receptor).

Accumulating evidence suggests that the RAS plays an important role in the pathogenesis of diabetic renal disease (Carey and Siragy, 2003; Gurley and Coffman, 2007; Yamout et al., 2014). Studies shows that the renin inhibitor (aliskiren) ameliorates renal damage in diabetic TG (mREN-2) rats (Kelly et al., 2007; Feldman et al., 2008) or in type 1 or 2 diabetics

(Parving et al., 2008; Wang et al., 2014). ACE inhibitors (ACEI) or AT1 receptor antagonists reduce mesangial sclerosis and proteinuria in animal models of both type 1 and type 2 diabetes (Andersen et al., 2000; Chan et al., 2000; Lewis et al., 2001; Brenner et al., 2001; Lewis, 2002; Carey and Siragy, 2003). A large scale clinical trial (RENNAL) demonstrated that losartan, an AT1 receptor antagonist, delayed the progression to proteinuria and end stage renal disease in type 2 diabetics (Brenner et al., 2001; Zhang et al., 2005). These beneficial effects of RAS inhibitors in the prevention of diabetic renal disease suggest that AngII, acting through the AT1 receptor, is a major mediator of progressive renal injury. AngII has many actions that might contribute to DN (Carey and Siragy, 2003), most prominently stimulating extracellular matrix protein synthesis in the kidney through induction of TGF- β expression (Kagami et al., 1994; Sharma et al., 1999; Singh et al., 1999; Erman et al., 2004). In addition to the contribution of AngII to DN, recent studies have suggested that ACE2 (Wong et al., 2007) or the newly described prorenin/renin receptor (Ichihara et al., 2006) plays a role in DN.

Intrarenal RAS and its role in DN

In addition to the systemic RAS, the kidney contains a complete intrarenal RAS (Carey and Siragy, 2003). Each RAS component has been localized to kidney (Navar et al., 1996; Miyata et al., 1999; Carey and Siragy, 2003). The intrarenal RAS is regulated independently from the systemic RAS. In general, circulating components of the RAS are normal or suppressed in diabetic patients (Burden and Thurston, 1979; Price et al., 1999), suggesting that a local RAS in the kidney is the primary contributor to AngII effects in the kidney with diabetes (Anderson et al., 1993; Border and Noble, 1998b; Ruiz-Ortega et al., 1998; Leehey et al., 2000; Zhang et al., 2002; Tamura et al., 2005). Accumulating evidence suggests that the intrarenal RAS is activated under diabetic conditions. Studies have shown that STZ-induced diabetes in rats activates the intrarenal RAS (Anderson et al., 1993), including activation of a glomerular RAS (Singh et al., 2005; Leehey et al., 2008). Diabetes also alters the distribution of RAS components within the various kidney compartments. In experimental diabetes, redistribution of ACE to vascular and glomeruli sites has been demonstrated by Anderson et al. (1993).

It has been demonstrated that glomerular mesangial cells express mRNA for renin, AGT, and ACE (Andrade et al., 2002). Moreover, mesangial cells in culture are able to synthesize, store, and secrete both renin and prorenin (Andrade et al., 2002) as well as AngII, and contain both the AT1 receptor (Miyata et al., 1999) and prorenin/renin receptor (Nguyen et al., 1996; Nguyen et al., 1998). Previous studies demonstrated that high glucose stimulates intracellular renin activity and AngII generation in mesangial cells (Vidotti et al., 2004; Wang et al., 2013). In addition to mesangial cells, podocytes and proximal tubular cells have been shown to express all components of the RAS, and interestingly RAS components in these cells are also stimulated by high glucose exposure (Zhang et al., 2002; Velez et al., 2007; Yoo et al., 2007; Durvasula and Shankland, 2008; Sonneveld et al., 2014), and have been suggested to contribute to the development of diabetic nephropathy (Phillips et al., 1999; Hong et al., 2001; Hsieh et al., 2003; White, 2006; Phillips, 2007; Kanwar et al., 2008; Liu et al., 2008; Niranjana et al., 2008; Márquez et al., 2015).

TGF- β and DN

TGF- β s are a family of cytokines that influence cell growth, differentiation, apoptosis, inflammatory processes, and gene expression. TGF- β is synthesized and secreted in a latent complex (latent TGF- β) (Gleizes et al., 1997). It must be converted to the active state before binding to its receptors and eliciting cellular functions. Latent TGF- β can be activated by a number of factors, including heat, extreme pH, plasmin, integrin binding, reactive oxygen species, and thrombospondin 1 (Lyons et al., 1990; Rifkin et al., 1993; Schultz-Cherry and Murphy-Ullrich, 1993; Schultz-Cherry et al., 1994; Taipale et al., 1995; Daniel et al., 2007).

The importance of TGF- β in the development of DN is well established. Increased expression of both active and total TGF- β in glomeruli occurs in both patients and in animal models of diabetes (Park et al., 2014; Shankland et al., 1994; Sharma and Ziyadeh, 1994; Young et al., 1995; Bertolucci et al., 1996; Sharma et al., 1997; Border and Noble, 1998a; Hong et al., 2001; Vasanthakumar et al., 2015). TGF- β is a major mediator of extracellular matrix deposition in the diabetic kidney (Border et al., 1995; Chen et al., 2001; Ziyadeh, 2004). The fibrogenic effects of TGF- β are due to its upregulation of extracellular matrix components, including collagen, fibronectin, osteopontin, and the downregulation of matrix degrading enzymes (Ziyadeh et al., 1994; Oh et al., 1998; Isono et al., 2002). TGF- β 1-specific neutralizing antibodies diminish overproduction of extracellular matrix proteins secreted by mesangial cells incubated with high glucose (Sharma et al., 1996).

Upstream stimulatory factors (USFs) in DN

USFs, including USF1 and USF2, were initially characterized as transcription factors implicated in the regulation of the adenovirus major late promoter. In mammals, USF1 and USF2 are ubiquitously expressed with a molecular weight of 43 kDa and 44 kDa, respectively. They predominantly localize in nuclei (Qyang et al., 1999) and belong to the Myc family of transcription factors characterized by a basic/helix loop helix/leucine zipper domain responsible for dimerization and DNA binding. These ubiquitous factors form homo- and hetero-dimers and recognize *in vitro* a CACGTG core sequence termed E box. Through binding to E boxes of target genes, USF factors regulate the gene expression (Rippe et al., 1997; Vallet et al., 1997; Qian et al., 1999; Kingsley-Kallesen et al., 2001; Nicolas et al., 2001; Bidder et al., 2002; Zhu et al., 2005; Chen et al., 2006). The USF2 structure and its regulated genes are illustrated in Fig. 1. USF1 and USF2 null mice have been generated and their phenotypes have been described previously. An embryonic lethal phenotype was observed with the double null mouse mutants. For USF1 deficient mice, no physical abnormalities have been reported in adult mice, except for the occasional occurrence of epileptic seizures in female USF1 null mice. USF2 null mice, however, show a severe phenotype including growth defects, abnormalities in fertility, mammary gland malfunction, an impaired transcriptional response to glucose in liver, and multivisceral iron overload. Although the pups display an obvious growth defect, and have an increased prenatal mortality rate (40%–50%), the surviving pups subsequently develop in an apparently normal fashion. USF2 null mice have a decreased lifespan (2.5 to 4.5 months in males; 10 months in females).

High glucose or glycated albumin upregulates USF2 expression in mesangial cells at the transcriptional level

Studies from our laboratory demonstrated that USFs are transcription factors involved in glucose mediated upregulation of thrombospondin 1 (TSP1) gene expression and TGF- β activity in glomerular mesangial cells and that these effects of USF2 contribute to diabetic renal complications (Wang et al., 2004). We also showed that treatment of rat mesangial cells (RMCs) with high glucose (30 mM) upregulates USF2 but not USF1 protein accumulation in mesangial cells through the activation of PKC, ERK, and p38 MAPK pathways (Wang et al., 2004). Furthermore, high glucose exposure stimulated USF2 gene transcription. Using the luciferase-promoter deletion assay, site-directed mutagenesis, and transactivation assay, we identified a glucose-responsive element in the USF2 gene promoter (– 1740 to – 1620, relative to the transcription start site) and demonstrated that glucose-induced USF2 expression is mediated through a cAMP-response element binding protein (CREB)-dependent transactivation of the USF2 promoter.

In addition to hyperglycemia, glycated proteins have been shown to accumulate in the kidneys of diabetic patients and contribute to DN. We found that glycated albumin upregulated USF2 expression (mRNA and protein) in a dose- and time-dependent manner. We also demonstrated that glycated albumin stimulated USF2 gene expression at the transcriptional level. By using the luciferase-promoter deletion assay, site-directed mutagenesis, and transactivation assay, we identified a glycated albumin-responsive region in the USF2 gene promoter (– 837 to – 430, relative to the transcription start site) and demonstrated that glycated albumin-induced USF2 expression was mediated through NF- κ B-dependent transactivation of the USF2 promoter. Furthermore, glycated albumin increased nuclear NF- κ B subunit-p65 protein levels. siRNA-mediated p65 knockdown prevented glycated albumin-induced USF2 gene expression (promoter activity, mRNA, and protein levels). Taken together, these data suggest that glycated albumin upregulated USF2 gene transcription in MCs through NF- κ B-dependent transactivation of the USF2 promoter (Li and Wang, 2010).

High glucose upregulates USF2 in renal proximal tubular cells through angiotensin II-dependent activation of CREB

In addition to glomerular mesangial cells, our studies found that high glucose upregulated USF2 expression and increased extracellular matrix accumulation in human renal proximal tubular cells (HK-2 cells); both were inhibited by siRNA-mediated USF2 knockdown. In addition, high glucose stimulated angiotensinogen and renin expression, increased renin activity, and resulted in increased angiotensin II formation. Treatment of HK-2 cells with an angiotensin II receptor 1 (AT1) blocker- losartan- prevented high-glucose-induced USF2 expression and high-glucose-enhanced phos-phorylation of CREB (cAMP response element binding protein). These data established that high glucose stimulated USF2 expression in HK-2 cells, at least in part, through angiotensin II-AT1-dependent activation of CREB, which can contribute to diabetic tubulointerstitial fibrosis (Visavadiya et al., 2011).

USF2 regulates glucose-induced TGF- β levels in mesangial cells

It has been shown that USF2 can bind to the TGF- β gene promoter and regulates its expression (Riccio et al., 1992; Kingsley-Kallesen et al., 2001; Weigert et al., 2004; Zhu et al., 2005). To determine the role of USF2 in glucose-mediated stimulation of TGF- β production, we utilized USF2-siRNA (from Santa Cruz) to knock down USF2 gene in mouse mesangial cells and then examined the effect of high glucose (30 mM glucose) on TGF- β levels in these cells using the method as described previously (Wang et al., 2004). As shown in Fig. 2, siRNA-USF2 transfection significantly inhibited USF2 expression under both normal and high glucose conditions (around 80% inhibition). Importantly, high glucose stimulated increases in active and total TGF- β levels were inhibited by siRNA-USF2 transfection. siRNA-USF2 transfection had no effect on basal TGF- β production (under 5mM glucose conditions). These data are consistent with that from Weigert et al. (2004), indicating that USF2 regulates glucose-induced TGF- β production in mesangial cells.

Overexpression of USF2 accelerates diabetic kidney injury

To define the functional role of USF2 in the development of diabetic nephropathy (DN) *in vivo*, we generated the global USF2 transgenic mice (Liu et al., 2007). Type 1 diabetes was induced in USF2 (Tg) and their wild-type littermates (WT) by injection of streptozotocin. Control USF2 (Tg) mice (~6 months old) exhibited increased urinary albumin excretion. These mice also exhibited glomerular hypertrophy, accompanied by increased TSP1, active TGF- β , fibronectin accumulation in the glomeruli compared with control WT littermates. Type 1 diabetes onset further augmented glomerular USF2 protein levels, the urinary albumin excretion and glomerular hypertrophy and fibrosis in the USF2 (Tg) mice. Together, these data indicate that functional changes (albuminuria) and structural changes (mesangial matrix expansion) exhibited in kidneys from USF2 (Tg) mice under basal conditions, which were significantly augmented by diabetes onset, suggesting that overexpression of USF2 accelerates diabetic kidney injury.

Activation of renal RAS in USF2 transgenic mice

It has been shown that USF1/2 binds to the mouse renin promoter and regulates renin gene expression in a kidney-derived renin expressing cell line (As4.1) (Pan et al., 2001). Therefore, the regulation of the RAS in USF2 (Tg) mice were determined. Our data demonstrated that renal renin expression (mRNA and protein levels), renin activity, and glomerular AngII levels were upregulated in USF2 transgenic mice, but plasma renin concentration was not altered. Systolic blood pressure was not altered in USF2 (Tg) mice as compared to littermates control. Moreover, renin receptor protein levels in kidney homogenates or the mRNA levels of angiotensinogen, ACE, ACE2, AT1a and AT2 receptors (determined by real-time PCR) were not changed in kidneys from USF2 transgenic mice as compared to control littermates (Shi et al., 2009). Collectively, these results suggest that USF2 represents an important mechanism of intra-renal RAS activation and local AngII formation.

Model of the role of USF2 in the development of diabetic nephropathy

Summary

Underdiabetic conditions USF2 expression is increased in the kidneys. Increased glomerular USF2 levels correlates with the glomerular mesangial matrix expansion. Furthermore, over-expression of USF2 accelerates the development of diabetic nephropathy in a type 1 diabetic mouse model, supporting the role of this transcription factor in the development of diabetic nephropathy. Our studies describe a novel effect of USF2 to promote renin gene expression and AngII formation in mesangial cells. This effect of USF2 was linked to glucose-induced regulation of TGF- β production, suggesting that this pathway may contribute to the development of diabetic nephropathy (Fig. 3).

Future directions

Our studies demonstrated that USF2 transgenic mice exhibited increased renal renin gene expression, renin activity and AngII levels without alteration of other components of the renin-angiotensin system (RAS) including angiotensinogen, ACE, ACE2, renin receptors, AT1 or AT2 receptors (Liu et al., 2007). Moreover, plasma renin concentration was not altered in USF2 transgenic mice. Together, these results suggest that increased renin enhances AngII synthesis, leading to the activation of the renal RAS in USF2 (Tg) mice, and contributing to the development of nephropathy. Therefore, the mechanistic role of RAS in USF2-mediated acceleration of diabetic nephropathy warrants future investigation.

Acknowledgments

This work was supported by National Institute of Diabetes and Digestive and Kidney Disease Grant DK081555 (to S. Wang).

References

- Andersen S, Tarnow L, Rossing P, Hansen BV, Parving HH. Renoprotective effects of angiotensin II receptor blockade in type 1 diabetic patients with diabetic nephropathy. *Kidney Int.* 2000; 57(2): 601–606. [PubMed: 10652037]
- Anderson S, Jung FF, Ingelfinger JR. Renal renin-angiotensin system in diabetes: functional, immunohistochemical, and molecular biological correlations. *Am J Physiol.* 1993; 265(4 Pt 2):F477–F486. [PubMed: 8238377]
- Andrade AQ, Casarini DE, Schor N, Boim MA. Characterization of renin mRNA expression and enzyme activity in rat and mouse mesangial cells. *Braz J Med Biol Res.* 2002; 35(1):17–24. [PubMed: 11743610]
- Bertoluci MC, Schmid H, Lachat JJ, Coimbra TM. Transforming growth factor-beta in the development of rat diabetic nephropathy. A 10-month study with insulin-treated rats. *Nephron.* 1996; 74(1):189–196. [PubMed: 8883039]
- Bidder M, Shao JS, Charlton-Kachigian N, Loewy AP, Semenkovich CF, Towler DA. Osteopontin transcription in aortic vascular smooth muscle cells is controlled by glucose-regulated upstream stimulatory factor and activator protein-1 activities. *J Biol Chem.* 2002; 277(46):44485–44496. [PubMed: 12200434]
- Borch-Johnsen K, Andersen PK, Deckert T. The effect of proteinuria on relative mortality in type 1 (insulin-dependent) diabetes mellitus. *Diabetologia.* 1985; 28(8):590–596. [PubMed: 4054448]
- Border WA, Noble NA. Evidence that TGF-beta should be a therapeutic target in diabetic nephropathy. *Kidney Int.* 1998a; 54(4):1390–1391. [PubMed: 9773681]

- Border WA, Noble NA. Interactions of transforming growth factor-beta and angiotensin II in renal fibrosis. *Hypertension*. 1998b; 31(1 Pt 2):181–188. [PubMed: 9453300]
- Border WA, Noble NA, Ketteler M. TGF-beta: a cytokine mediator of glomerulosclerosis and a target for therapeutic intervention. *Kidney Int Suppl*. 1995; 49(61):S59–S61. [PubMed: 7674597]
- Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S, the RENAAL Study Investigators. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med*. 2001; 345(12):861–869. [PubMed: 11565518]
- Burden AC, Thurston H. Plasma renin activity in diabetes mellitus. *Clin Sci (Lond)*. 1979; 56(3):255–259. [PubMed: 477208]
- Carey RM, Siragy HM. The intrarenal renin-angiotensin system and diabetic nephropathy. *Trends Endocrinol Metab*. 2003; 14(6):274–281. [PubMed: 12890592]
- Chan JC, Ko GT, Leung DH, Cheung RC, Cheung MY, So WY, Swaminathan R, Nicholls MG, Critchley JA, Cockram CS. Long-term effects of angiotensin-converting enzyme inhibition and metabolic control in hypertensive type 2 diabetic patients. *Kidney Int*. 2000; 57(2):590–600. [PubMed: 10652036]
- Chen L, Shen YH, Wang X, Wang J, Gan Y, Chen N, Wang J, LeMaire SA, Coselli JS, Wang XL. Human prolyl-4-hydroxylase alpha(I) transcription is mediated by upstream stimulatory factors. *J Biol Chem*. 2006; 281(16):10849–10855. [PubMed: 16488890]
- Chen S, Hong SW, Iglesias-de la Cruz MC, Isono M, Casaretto A, Ziyadeh FN. The key role of the transforming growth factor-beta system in the pathogenesis of diabetic nephropathy. *Ren Fail*. 2001; 23(3–4):471–481. [PubMed: 11499562]
- Cooper ME. Interaction of metabolic and haemodynamic factors in mediating experimental diabetic nephropathy. *Diabetologia*. 2001; 44(11):1957–1972. [PubMed: 11719827]
- Corre S, Galibert MD. Upstream stimulating factors: highly versatile stress-responsive transcription factors. *Pigment Cell Res*. 2005; 18(5):337–348. [PubMed: 16162174]
- Dabhi B, Mistry KN. Oxidative stress and its association with TNF-alpha-308 G/C and IL-1alpha-889 C/T gene polymorphisms in patients with diabetes and diabetic nephropathy. *Gene*. 2015; 56(2):197–202. [PubMed: 25732517]
- Daniel C, Schaub K, Amann K, Lawler J, Hugo C. Thrombospondin-1 is an endogenous activator of TGF-beta in experimental diabetic nephropathy *in vivo*. *Diabetes*. 2007; 56(12):2982–2989. [PubMed: 17878288]
- Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, Donovan M, Woolf B, Robison K, Jeyaseelan R, Breitbart RE, Acton S. A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1–9. *Circ Res*. 2000; 87(5):E1–E9. [PubMed: 10969042]
- Durvasula RV, Shankland SJ. Activation of a local renin angiotensin system in podocytes by glucose. *Am J Physiol Renal Physiol*. 2008; 294(4):F830–F839. [PubMed: 18216149]
- Erman A, Veksler S, Gafter U, Boner G, Wittenberg C, van Dijk DJ. Renin-angiotensin system blockade prevents the increase in plasma transforming growth factor beta 1, and reduces proteinuria and kidney hypertrophy in the streptozotocin-diabetic rat. *J Renin Angiotensin Aldosterone Syst*. 2004; 5(3):146–151. [PubMed: 15526251]
- Feener EP, King GL. Vascular dysfunction in diabetes mellitus. *Lancet*. 1997; 350(Suppl 1):SI9–SI13. [PubMed: 9250277]
- Feldman DL, Jin L, Xuan H, Contrepas A, Zhou Y, Webb RL, Mueller DN, Feldt S, Cumin F, Maniara W, Persohn E, Schuetz H, Jan Danser AH, Nguyen G. Effects of aliskiren on blood pressure, albuminuria, and (pro)renin receptor expression in diabetic TG (mRen-2)27 rats. *Hypertension*. 2008; 52(1):130–136. [PubMed: 18490518]
- Ferrario CM, Trask AJ, Jessup JA. Advances in biochemical and functional roles of angiotensin-converting enzyme 2 and angiotensin-(1–7) in regulation of cardiovascular function. *Am J Physiol Heart Circ Physiol*. 2005; 289(6):H2281–H2290. [PubMed: 16055515]
- Fogo AB. Mesangial matrix modulation and glomerulosclerosis. *Exp Nephrol*. 1999; 7(2):147–159. [PubMed: 10213868]

- Forbes JM, Fukami K, Cooper ME. Diabetic nephropathy: where hemodynamics meets metabolism. *Exp Clin Endocrinol Diabetes*. 2007; 115(2):69–84. [PubMed: 17318765]
- Giacco F, Brownlee M. Oxidative stress and diabetic complications. *Circ Res*. 2010; 107(9):1058–1070. [PubMed: 21030723]
- Gleizes PE, Munger JS, Nunes I, Harpel JG, Mazzieri R, Noguera I, Rifkin DB. TGF-beta latency: biological significance and mechanisms of activation. *Stem Cells*. 1997; 15(3):190–197. [PubMed: 9170210]
- Gurley SB, Coffman TM. The renin-angiotensin system and diabetic nephropathy. *Semin Nephrol*. 2007; 27(2):144–152. [PubMed: 17418683]
- Hadsell DL, Bonnette S, George J, Torres D, Klimentidis Y, Gao S, Haney PM, Summy-Long J, Soloff MS, Parlow AF, Sirito M, Sawadogo M. Diminished milk synthesis in upstream stimulatory factor 2 null mice is associated with decreased circulating oxytocin and decreased mammary gland expression of eukaryotic initiation factors 4E and 4G. *Mol Endocrinol*. 2003; 17(11):2251–2267. [PubMed: 12907752]
- Hong SW, Isono M, Chen S, Iglesias-De La Cruz MC, Han DC, Ziyadeh FN. Increased glomerular and tubular expression of transforming growth factor-beta1, its type II receptor, and activation of the Smad signaling pathway in the *db/db* mouse. *Am J Pathol*. 2001; 158(5):1653–1663. [PubMed: 11337363]
- Hsieh TJ, Fustier P, Zhang SL, Filep JG, Tang SS, Ingelfinger JR, Fantus IG, Hamet P, Chan JS. High glucose stimulates angiotensinogen gene expression and cell hypertrophy via activation of the hexosamine biosynthesis pathway in rat kidney proximal tubular cells. *Endocrinology*. 2003; 144(10):4338–4349. [PubMed: 12960040]
- Ichihara A, Suzuki F, Nakagawa T, Kaneshiro Y, Takemitsu T, Sakoda M, Nabi AH, Nishiyama A, Sugaya T, Hayashi M, Inagami T. Prorenin receptor blockade inhibits development of glomerulosclerosis in diabetic angiotensin II type 1a receptor-deficient mice. *J Am Soc Nephrol*. 2006; 17(7):1950–1961. [PubMed: 16738017]
- Ishimura E, Sterzel RB, Morii H, Kashgarian M. Extracellular matrix protein: gene expression and synthesis in cultured rat mesangial cells. *Nihon Jinzo Gakkai Shi*. 1992; 34(1):9–17. [PubMed: 1593802]
- Isono M, Chen S, Hong SW, Iglesias-de la Cruz MC, Ziyadeh FN. Smad pathway is activated in the diabetic mouse kidney and Smad3 mediates TGF-beta-induced fibronectin in mesangial cells. *Biochem Biophys Res Commun*. 2002; 296(5):1356–1365. [PubMed: 12207925]
- Kagami S, Border WA, Miller DE, Noble NA. Angiotensin II stimulates extracellular matrix protein synthesis through induction of transforming growth factor-beta expression in rat glomerular mesangial cells. *J Clin Invest*. 1994; 93(6):2431–2437. [PubMed: 8200978]
- Kanwar YS, Wada J, Sun L, Xie P, Wallner EI, Chen S, Chugh S, Danesh FR. Diabetic nephropathy: mechanisms of renal disease progression. *Exp Biol Med (Maywood)*. 2008; 233(1):4–11. [PubMed: 18156300]
- Kelly DJ, Zhang Y, Moe G, Naik G, Gilbert RE. Aliskiren, a novel renin inhibitor, is renoprotective in a model of advanced diabetic nephropathy in rats. *Diabetologia*. 2007; 50(11):2398–2404. [PubMed: 17828524]
- Kingsley-Kallesen M, Luster TA, Rizzino A. Transcriptional regulation of the transforming growth factor-beta 2 gene in glioblastoma cells. *In Vitro Cell Dev Biol Anim*. 2001; 37(10):684–690. [PubMed: 11776974]
- Kobori H, Nangaku M, Navar LG, Nishiyama A. The intrarenal renin-angiotensin system: from physiology to the pathobiology of hypertension and kidney disease. *Pharmacol Rev*. 2007; 59(3):251–287. [PubMed: 17878513]
- Lawrence DA. Transforming growth factor-beta: a general review. *Eur Cytokine Netw*. 1996; 7(3):363–374. [PubMed: 8954178]
- Leehey DJ, Singh AK, Alavi N, Singh R. Role of angiotensin II in diabetic nephropathy. *Kidney Int Suppl*. 2000; 77(8):S93–S98. [PubMed: 10997697]
- Leehey DJ, Singh AK, Bast JP, Sethupathi P, Singh R. Glomerular renin angiotensin system in streptozotocin diabetic and Zucker diabetic fatty rats. *Transl Res*. 2008; 151(4):208–216. [PubMed: 18355768]

- Lewis EJ. The role of angiotensin II receptor blockers in preventing the progression of renal disease in patients with type 2 diabetes. *Am J Hypertens.* 2002; 15(10 Pt 2):123S–128S. [PubMed: 12383593]
- Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I, the Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med.* 2001; 345(12):851–860. [PubMed: 11565517]
- Li Y, Wang S. Glycated albumin upregulates upstream stimulatory factor 2 gene transcription in mesangial cells. *Am J Physiol Renal Physiol.* 2010; 299(1):F121–F127. [PubMed: 20410211]
- Lim A, Kh. Diabetic nephropathy- complications and treatment. *Int J Nephrol Renovasc Dis.* 2014; 7:361–381. [PubMed: 25342915]
- Liu F, Brezniceanu ML, Wei CC, Chénier I, Sachetelli S, Zhang SL, Filep JG, Ingelfinger JR, Chan JS. Overexpression of angiotensinogen increases tubular apoptosis in diabetes. *J Am Soc Nephrol.* 2008; 19(2):269–280. [PubMed: 18057217]
- Liu S, Shi L, Wang S. Overexpression of upstream stimulatory factor 2 accelerates diabetic kidney injury. *Am J Physiol Renal Physiol.* 2007; 293(5):F1727–F1735. [PubMed: 17881461]
- Lyons RM, Gentry LE, Purchio AF, Moses HL. Mechanism of activation of latent recombinant transforming growth factor beta 1 by plasmin. *J Cell Biol.* 1990; 110(4):1361–1367. [PubMed: 2139036]
- Márquez E, Riera M, Pascual J, Soler MJ. Renin-angiotensin system within the diabetic podocyte. *Am J Physiol Renal Physiol.* 2015; 308(1):F1–F10. [PubMed: 25339703]
- Miyata N, Park F, Li XF, Cowley AW Jr. Distribution of angiotensin AT1 and AT2 receptor subtypes in the rat kidney. *Am J Physiol.* 1999; 277(3 Pt 2):F437–F446. [PubMed: 10484527]
- Navar LG, Harrison-Bernard LM, Nishiyama A, Kobori H. Regulation of intrarenal angiotensin II in hypertension. *Hypertension.* 2002; 39(2 Pt 2):316–322. [PubMed: 11882566]
- Navar LG, Inscho EW, Majid SA, Imig JD, Harrison-Bernard LM, Mitchell KD. Paracrine regulation of the renal microcirculation. *Physiol Rev.* 1996; 76(2):425–536. [PubMed: 8618962]
- Navarro-González JF, Mora-Fernández C. The role of inflammatory cytokines in diabetic nephropathy. *J Am Soc Nephrol.* 2008; 19(3):433–442. [PubMed: 18256353]
- Nguyen G, Bouzhir L, Delarue F, Rondeau E, Sraer JD. Evidence of a renin receptor on human mesangial cells: effects on PAI1 and cGMP. *Nephrologie.* 1998; 19(7):411–416. [PubMed: 9857376]
- Nguyen G, Delarue F, Berrou J, Rondeau E, Sraer JD. Specific receptor binding of renin on human mesangial cells in culture increases plasminogen activator inhibitor-1 antigen. *Kidney Int.* 1996; 50(6):1897–1903. [PubMed: 8943472]
- Nicolas G, Bennoun M, Devaux I, Beaumont C, Grandchamp B, Kahn A, Vaulont S. Lack of hepcidin gene expression and severe tissue iron overload in upstream stimulatory factor 2 (USF2) knockout mice. *Proc Natl Acad Sci USA.* 2001; 98(15):8780–8785. [PubMed: 11447267]
- Niranjan T, Bielez B, Gruenwald A, Ponda MP, Kopp JB, Thomas DB, Susztak K. The Notch pathway in podocytes plays a role in the development of glomerular disease. *Nat Med.* 2008; 14(3):290–298. [PubMed: 18311147]
- Oh JH, Ha H, Yu MR, Lee HB. Sequential effects of high glucose on mesangial cell transforming growth factor-beta 1 and fibronectin synthesis. *Kidney Int.* 1998; 54(6):1872–1878. [PubMed: 9853252]
- Pan L, Black TA, Shi Q, Jones CA, Petrovic N, Loudon J, Kane C, Sigmund CD, Gross KW. Critical roles of a cyclic AMP responsive element and an E-box in regulation of mouse renin gene expression. *J Biol Chem.* 2001; 276(49):45530–45538. [PubMed: 11564732]
- Park JT, Kato M, Lanting L, Castro N, Nam BY, Wang M, Kang SW, Natarajan R. Repression of let-7 by transforming growth factor- β 1-induced Lin28 upregulates collagen expression in glomerular mesangial cells under diabetic conditions. *Am J Physiol Renal Physiol.* 2014; 307(12):F1390–F1403. [PubMed: 25354942]
- Parving HH, Persson F, Lewis JB, Lewis EJ, Hollenberg NK, the AVOID Study Investigators. Aliskiren combined with losartan in type 2 diabetes and nephropathy. *N Engl J Med.* 2008; 358(23):2433–2446. [PubMed: 18525041]

- Peach MJ. Renin-angiotensin system: biochemistry and mechanisms of action. *Physiol Rev.* 1977; 57(2):313–370. [PubMed: 191856]
- Petrica L, Vlad A, Gluhovschi G, Gadalean F, Dumitrascu V, Vlad D, Popescu R, Velciov S, Gluhovschi C, Bob F, Ursoniu S, Petrica M, Jianu DC. Glycated peptides are associated with the variability of endothelial dysfunction in the cerebral vessels and the kidney in type 2 diabetes mellitus patients: a cross-sectional study. *J Diabetes Complications.* 2015; 29(2):230–237. [PubMed: 25511877]
- Phillips A. The role of proximal tubular cells in interstitial fibrosis: understanding TGF-beta1. *Chang Gung Med J.* 2007; 30(1):2–6. [PubMed: 17477023]
- Phillips A, Janssen U, Floege J. Progression of diabetic nephropathy. Insights from cell culture studies and animal models. *Kidney Blood Press Res.* 1999; 22(1–2):81–97. [PubMed: 10352411]
- Price DA, Porter LE, Gordon M, Fisher ND, De'Oliveira JM, Laffel LM, Passan DR, Williams GH, Hollenberg NK. The paradox of the low-renin state in diabetic nephropathy. *J Am Soc Nephrol.* 1999; 10(11):2382–2391. [PubMed: 10541298]
- Qian J, Kaytor EN, Towle HC, Olson LK. Upstream stimulatory factor regulates Pdx-1 gene expression in differentiated pancreatic beta-cells. *Biochem J.* 1999; 341(Pt 2):315–322. [PubMed: 10393088]
- Qyang Y, Luo X, Lu T, Ismail PM, Krylov D, Vinson C, Sawadogo M. Cell-type-dependent activity of the ubiquitous transcription factor USF in cellular proliferation and transcriptional activation. *Mol Cell Biol.* 1999; 19(2):1508–1517. [PubMed: 9891084]
- Re R. Intracellular renin-angiotensin system: the tip of the intracrine physiology iceberg. *Am J Physiol Heart Circ Physiol.* 2007; 293(2):H905–H906. [PubMed: 17526648]
- Riccio A, Pedone PV, Lund LR, Olesen T, Olsen HS, Andreassen PA. Transforming growth factor beta 1-responsive element: closely associated binding sites for USF and CCAAT-binding transcription factor-nuclear factor I in the type 1 plasminogen activator inhibitor gene. *Mol Cell Biol.* 1992; 12(4):1846–1855. [PubMed: 1549130]
- Rifkin DB, Kojima S, Abe M, Harpel JG. TGF-beta: structure, function, and formation. *Thromb Haemost.* 1993; 70(1):177–179. [PubMed: 8236098]
- Rippe RA, Umezawa A, Kimball JP, Breindl M, Brenner DA. Binding of upstream stimulatory factor to an E-box in the 3'-flanking region stimulates alpha1(I) collagen gene transcription. *J Biol Chem.* 1997; 272(3):1753–1760. [PubMed: 8999857]
- Ruiz-Ortega M, Lorenzo O, Egido J. Angiotensin III up-regulates genes involved in kidney damage in mesangial cells and renal interstitial fibroblasts. *Kidney Int Suppl.* 1998; 68(5):S41–S45. [PubMed: 9839282]
- Sawadogo M, Roeder RG. Interaction of a gene-specific transcription factor with the adenovirus major late promoter upstream of the TATA box region. *Cell.* 1985; 43(1):165–175. [PubMed: 4075392]
- Schultz-Cherry S, Lawler J, Murphy-Ullrich JE. The type 1 repeats of thrombospondin 1 activate latent transforming growth factor-beta. *J Biol Chem.* 1994; 269(43):26783–26788. [PubMed: 7929414]
- Schultz-Cherry S, Murphy-Ullrich JE. Thrombospondin causes activation of latent transforming growth factor-beta secreted by endothelial cells by a novel mechanism. *J Cell Biol.* 1993; 122(4):923–932. [PubMed: 8349738]
- Shankland SJ, Scholey JW, Ly H, Thai K. Expression of transforming growth factor-beta 1 during diabetic renal hypertrophy. *Kidney Int.* 1994; 46(2):430–442. [PubMed: 7967355]
- Sharma K, Eltayeb BO, McGowan TA, Dunn SR, Alzahabi B, Rohde R, Ziyadeh FN, Lewis EJ. Captopril-induced reduction of serum levels of transforming growth factor-beta1 correlates with long-term renoprotection in insulin-dependent diabetic patients. *Am J Kidney Dis.* 1999; 34(5):818–823. [PubMed: 10561136]
- Sharma K, Jin Y, Guo J, Ziyadeh FN. Neutralization of TGF-beta by anti-TGF-beta antibody attenuates kidney hypertrophy and the enhanced extracellular matrix gene expression in STZ-induced diabetic mice. *Diabetes.* 1996; 45(4):522–530. [PubMed: 8603776]
- Sharma K, Ziyadeh FN. Renal hypertrophy is associated with upregulation of TGF-beta 1 gene expression in diabetic BB rat and NOD mouse. *Am J Physiol.* 1994; 267(6 Pt 2):F1094–F01. [PubMed: 7810696]

- Sharma K, Ziyadeh FN, Alzahabi B, McGowan TA, Kapoor S, Kurnik BR, Kurnik PB, Weisberg LS. Increased renal production of transforming growth factor-beta1 in patients with type II diabetes. *Diabetes*. 1997; 46(5):854–859. [PubMed: 9133555]
- Shi L, Nikolic D, Liu S, Lu H, Wang S. Activation of renal renin-angiotensin system in upstream stimulatory factor 2 transgenic mice. *Am J Physiol Renal Physiol*. 2009; 296(2):F257–F265. [PubMed: 19004931]
- Singh R, Alavi N, Singh AK, Leehey DJ. Role of angiotensin II in glucose-induced inhibition of mesangial matrix degradation. *Diabetes*. 1999; 48(10):2066–2073. [PubMed: 10512375]
- Singh R, Singh AK, Leehey DJ. A novel mechanism for angiotensin II formation in streptozotocin-diabetic rat glomeruli. *Am J Physiol Renal Physiol*. 2005; 288(6):F1183–F1190. [PubMed: 15701818]
- Sirito M, Lin Q, Deng JM, Behringer RR, Sawadogo M. Overlapping roles and asymmetrical cross-regulation of the USF proteins in mice. *Proc Natl Acad Sci USA*. 1998; 95(7):3758–3763. [PubMed: 9520440]
- Sirito M, Lin Q, Maity T, Sawadogo M. Ubiquitous expression of the 43- and 44-kDa forms of transcription factor USF in mammalian cells. *Nucleic Acids Res*. 1994; 22(3):427–433. [PubMed: 8127680]
- Skyler JS. Diabetic complications. The importance of glucose control. *Endocrinol Metab Clin North Am*. 1996; 25(2):243–254. [PubMed: 8799699]
- Sonneveld R, van der Vlag J, Baltissen MP, Verkaart SA, Wetzels JF, Berden JH, Hoenderop JG, Nijenhuis T. Glucose specifically regulates TRPC6 expression in the podocyte in an AngII-dependent manner. *Am J Pathol*. 2014; 184(6):1715–1726. [PubMed: 24731445]
- Taipale J, Lohi J, Saarinen J, Kovanen PT, Keski-Oja J. Human mast cell chymase and leukocyte elastase release latent transforming growth factor-beta 1 from the extracellular matrix of cultured human epithelial and endothelial cells. *J Biol Chem*. 1995; 270(9):4689–4696. [PubMed: 7876240]
- Tamura J, Konno A, Hashimoto Y, Kon Y. Upregulation of renal renin-angiotensin system in mouse diabetic nephropathy. *Jpn J Vet Res*. 2005; 53(1–2):13–26. [PubMed: 16190318]
- Vallet VS, Henrion AA, Bucchini D, Casado M, Raymondjean M, Kahn A, Vaulont S. Glucose-dependent liver gene expression in upstream stimulatory factor 2^{-/-} mice. *J Biol Chem*. 1997; 272(35):21944–21949. [PubMed: 9268329]
- Vasanthakumar R, Mohan V, Anand G, Deepa M, Babu S, Aravindhan V. Serum IL-9, IL-17, and TGF- β levels in subjects with diabetic kidney disease (CURES-134). *Cytokine*. 2015; 72(1):109–112. [PubMed: 25542095]
- Velez JC, Bland AM, Arthur JM, Raymond JR, Janecz MG. Characterization of renin-angiotensin system enzyme activities in cultured mouse podocytes. *Am J Physiol Renal Physiol*. 2007; 293(1):F398–F407. [PubMed: 17429035]
- Vidotti DB, Casarini DE, Cristovam PC, Leite CA, Schor N, Boim MA. High glucose concentration stimulates intracellular renin activity and angiotensin II generation in rat mesangial cells. *Am J Physiol Renal Physiol*. 2004; 286(6):F1039–F1045. [PubMed: 14722017]
- Visavadiya NP, Li Y, Wang S. High glucose upregulates upstream stimulatory factor 2 in human renal proximal tubular cells through angiotensin II-dependent activation of CREB. *Nephron, Exp Nephrol*. 2011; 117(3):e62–e70. [PubMed: 20814220]
- Wang S, Skorzewski J, Feng X, Mei L, Murphy-Ullrich JE. Glucose up-regulates thrombospondin 1 gene transcription and transforming growth factor-beta activity through antagonism of cGMP-dependent protein kinase repression via upstream stimulatory factor 2. *J Biol Chem*. 2004; 279(33):34311–34322. [PubMed: 15184388]
- Wang W, Qiu L, Howard A, Solis N, Li C, Wang X, Kopp JB, Levi M. Protective effects of aliskiren and valsartan in mice with diabetic nephropathy. *J Renin Angiotensin Aldosterone Syst*. 2014; 15(4):384–395. [PubMed: 25031296]
- Wang Z, Ni J, Shao D, Liu J, Shen Y, Zhou L, Huang Y, Yu C, Wang J, Xue H, Lu L. Elevated transcriptional co-activator p102 mediates angiotensin II type 1 receptor up-regulation and extracellular matrix overproduction in the high glucose-treated rat glomerular mesangial cells and isolated glomeruli. *Eur J Pharmacol*. 2013; 702(1–3):208–217. [PubMed: 23376562]

- Weigert C, Brodbeck K, Sawadogo M, Haring HU, Schleicher ED. USF proteins induce human TGF-beta1 gene activation via the glucose response element – 1013/ – 1002 in mesangial cells—upregulation of USF activity by the hexosamine biosynthetic pathway. *J Biol Chem.* 2004; 279:2161–2168. [PubMed: 15001118]
- White KE. Research into the glomerular podocyte—is it relevant to diabetic nephropathy? *Diabet Med.* 2006; 23(7):715–719. [PubMed: 16842474]
- Wong DW, Oudit GY, Reich H, Kassiri Z, Zhou J, Liu QC, Backx PH, Penninger JM, Herzenberg AM, Scholey JW. Loss of angiotensin-converting enzyme-2 (Ace2) accelerates diabetic kidney injury. *Am J Pathol.* 2007; 171(2):438–451. [PubMed: 17600118]
- Yamout H, Lazich I, Bakris GL. Blood pressure, hypertension, RAAS blockade, and drug therapy in diabetic kidney disease. *Adv Chronic Kidney Dis.* 2014; 21(3):281–286. [PubMed: 24780456]
- Yoo TH, Li JJ, Kim JJ, Jung DS, Kwak SJ, Ryu DR, Choi HY, Kim JS, Kim HJ, Han SH, Lee JE, Han DS, Kang SW. Activation of the renin-angiotensin system within podocytes in diabetes. *Kidney Int.* 2007; 71(10):1019–1027. [PubMed: 17361112]
- Young BA, Johnson RJ, Alpers CE, Eng E, Gordon K, Floege J, Couser WG, Seidel K. Cellular events in the evolution of experimental diabetic nephropathy. *Kidney Int.* 1995; 47(3):935–944. [PubMed: 7752595]
- Zhang SL, To C, Chen X, Filep JG, Tang SS, Ingelfinger JR, Chan JS. Essential role(s) of the intrarenal renin-angiotensin system in transforming growth factor-beta1 gene expression and induction of hypertrophy of rat kidney proximal tubular cells in high glucose. *J Am Soc Nephrol.* 2002; 13(2):302–312. [PubMed: 11805157]
- Zhang Z, Shahinfar S, Keane WF, Ramjit D, Dickson TZ, Gleim GW, Mogensen CE, de Zeeuw D, Brenner BM, Snapinn SM. Importance of baseline distribution of proteinuria in renal outcomes trials: lessons from the reduction of endpoints in NIDDM with the angiotensin II antagonist losartan (RENAAL) study. *J Am Soc Nephrol.* 2005; 16(6):1775–1780. [PubMed: 15872078]
- Zhu Y, Casado M, Vaulont S, Sharma K. Role of upstream stimulatory factors in regulation of renal transforming growth factor-beta1. *Diabetes.* 2005; 54(7):1976–1984. [PubMed: 15983197]
- Ziyadeh FN. Mediators of diabetic renal disease: the case for tgf-Beta as the major mediator. *J Am Soc Nephrol.* 2004; 15(Suppl 1):S55–57. [PubMed: 14684674]
- Ziyadeh FN, Sharma K, Ericksen M, Wolf G. Stimulation of collagen gene expression and protein synthesis in murine mesangial cells by high glucose is mediated by autocrine activation of transforming growth factor-beta. *J Clin Invest.* 1994; 93(2):536–542. [PubMed: 8113392]

Figure 1.

Schematic illustration of USF2 structure and its regulated genes. USR: USF-specific region; B: basic region; HLH: helix–loop–helix domain; LZ: leucine zipper domain.

Figure 2.

siRNA-USF2 transfection inhibits high glucose levels-induced TGF- β production. Mouse mesangial cells were transfected with siRNA-USF2 or control siRNA for 48 h. Then cells were treated with normal or high glucose media for 24 h. Conditioned media were collected and used for measuring active and total TGF- β levels by PAI-1/lucifersae assay. Cells were harvested to determine the USF2 protein levels in nuclear extracts by immunoblotting. TBP (anti-TATA binding protein) was used as internal nuclear loading control. Data are represented as mean of 3 replicates $\pm$ S.D. *, $p < 0.05$ vs. NG. &, $p < 0.05$ vs. NG of control siRNA. # $p < 0.05$ vs. HG of control siRNA.

Figure 3.

Model of the role of USF2 in DN. Diabetic conditions upregulate kidney USF2 levels. USF2 binds to renin promoter and increases renin gene expression and stimulates angiotensin II production in kidney cells. Increased angiotensin II increases TGF- β levels and then extracellular matrix production in kidney cells, which leads to renal fibrosis.