CASE REPORT

Multifocal chorioretinitis with progressive subretinal fibrosis in a young child

Laura R Steeples,¹ Jane Ashworth,¹ Nicholas Jones²¹Department of Ophthalmology, Manchester Royal Eye Hospital, Manchester, UK²Manchester Royal Eye Hospital, Manchester, UK**Correspondence to** Professor Nicholas Jones, nicholas.jones@cmft.nhs.uk

Accepted 2 October 2015

SUMMARY

Progressive subretinal fibrosis with uveitis (PSFU) syndrome is a rare clinical entity presenting with multifocal granulomatous chorioretinitis and subsequently developing extensive subretinal fibrosis over weeks or months. Subretinal fibrosis affecting the macula may cause profound and permanent vision loss. We report the presentation and management of this condition in a child, to our knowledge, the youngest reported. The condition caused substantial visual loss and required high-dose systemic corticosteroid and immunosuppression with methotrexate and ciclosporin to control aggressive intraocular inflammation.

BACKGROUND

The syndrome of progressive subretinal fibrosis with uveitis (PSFU) is very rare even in a specialised uveitis practice. The patient in this case demonstrated a severe acute presentation with aggressive, progressive subretinal fibrosis. The development of PSFU in a young child is unique, to date, with almost all previous reports being in adults, and a very small number in teenagers. We describe the substantial management challenges of this complex syndrome in a young child.

CASE PRESENTATION

A 3-year-old Caucasian boy presented to our paediatric uveitis clinic with right mydriasis and photophobia. He had no ophthalmic history. Six weeks before presentation, he had received oral antibiotics for an episode of tonsillitis. He had no other significant medical or family history.

At presentation, right (R) visual acuity (VA) was 6/19 and the left (L), 6/19. A right relative afferent pupillary defect was detected. Mild non-granulomatous anterior uveitis and moderate vitritis were seen in both eyes. A large number of pale choroidal lesions, each with a very distinct central punctate dot, was seen throughout both fundi, extending from the posterior pole to the far periphery (figure 1). There was also patchy periphlebitis with calibre changes.

Both optic nerve heads were swollen, with clustered peri-papillary lesions with oedema, particularly in the right eye. Submacular fluid was evident in both eyes and there was deep intra-retinal haemorrhage at the posterior poles. The retinal pigment epithelium showed signs of fragmentation.

INVESTIGATIONS

The patient was extensively investigated for infectious and inflammatory aetiology by a

multidisciplinary team including paediatric rheumatology and immunology. He had a normal full blood count, no inflammatory markers and normal serum angiotensin enzyme; serology for syphilis, toxoplasmosis, borreliosis, brucellosis, Epstein-Barr virus and herpes simplex virus was negative. Antistreptolysin (ASO) and antistreptodornase titres were within normal limits. He was IgG positive for both measles and Varicella zoster virus but with no evidence of recent infection. Mantoux and gamma-interferon tests for tuberculosis were both negative. Subsequent brain MRI with angiography was normal, lumbar puncture showed no pleocytosis and cerebrospinal fluid (CSF) was negative for enteroviruses and herpesviruses. Immune profile testing was normal. Autoantibody testing was negative for antinuclear antibodies, antidouble-stranded DNA, myeloperoxidase, proteinase 3, rheumatoid factor, extractable nuclear antigens and anticardiolipin. Aqueous sampling was PCR negative for herpesviruses and fungi. Radiography showed a degree of sinusitis; ENT (ear, nose, and throat) investigation demonstrated no evidence of fungal or other infection.

Serial retinal ultra-widefield imaging, optical coherence tomography (OCT) scans and examination under anaesthesia, including fundus fluorescein angiography (FFA), were performed and were critical for assessing disease activity, and for demonstrating the extent and progress of retinal and choroidal lesions. FFA was also used to diagnose development of secondary choroidal neovascular membrane (CNVM). Retinal photography was a particularly helpful adjunct to clinical examination in such a young patient, where cooperation for assessment was limited.

Figure 1 Retinal images of right and left eyes at presentation (Optomap images, Optos) demonstrating a large number of white–yellow choroidal inflammatory lesions with central punctate dots. The lesions extended from the posterior pole to the far periphery. Both optic nerves are swollen with peri-papillary lesions evident. Optical coherence tomography images are superimposed and demonstrate subretinal fluid in the left eye and peri-papillary retinal thickening in the right eye.

CrossMark

To cite: Steeples LR, Ashworth J, Jones N. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-212526

Figure 2 Right and left posterior poles 7 weeks after presentation (top images) showing incipient subretinal fibrosis and lesions becoming confluent. Eleven weeks after presentation (bottom images) the fibrosis is spreading and becoming contiguous, with individual lesions coalescing and remaining active.

DIFFERENTIAL DIAGNOSIS

The initial differential diagnosis included post-streptococcal uveitis following recent tonsillitis, atypical punctate inner choroidopathy (PIC) and immune reaction to occult infection. However, no evidence of infection was found and post-streptococcal disease was excluded. The clinical behaviour of the condition, with aggressive multifocal choroiditis and, subsequently, progressive subretinal fibrosis, became consistent with clinical descriptions of PSFU.

TREATMENT

Empirical treatment for post-streptococcal uveitis was started immediately, with oral penicillin V 125 mg four times a day and oral prednisolone 25 mg (1.25 mg/kg) daily. Penicillin was discontinued on receipt of normal ASO titres, and oral steroid was slowly tapered to 10 mg/day. The VA improved to R 6/12 and L 6/6 and the uveitis improved. The L subretinal fluid cleared and was reduced significantly in the R. However, the chorioretinal lesions progressively increased in size and began to coalesce in both eyes, at which time (7 weeks after presentation) the

diagnosis of PSFU became apparent (figure 2). Confluent progressive subretinal fibrosis emerged in the R papillomacular region and in the L inferior macula. The VA deteriorated to R 3/60 and L 6/9 (figure 2). Oral methotrexate 10 mg weekly was started alongside oral steroid 10 mg/day. However, 5 months after presentation, increasing right intra-retinal fluid was noted and FFA under general anaesthesia (figure 3) demonstrated an active peri-papillary CNVM in that eye, which therefore underwent four successive injections of intravitreal bevacizumab at monthly intervals; stability on OCT and FFA was achieved.

OUTCOME AND FOLLOW-UP

Seven months after presentation, VA was R 6/60 and L 6/24. Increasing bilateral intra-retinal fluid was treated with increased oral prednisolone (20 mg/day; 1 mg/kg) and, subsequently, bilateral intravitreal bevacizumab injections. Subretinal fibrosis progressed in both eyes (figure 4), encircling the left macula with VA of R 6/60 and L 6/30. Oral ciclosporin, initially 25 mg two times a day rising to 50 mg two times a day (5 mg/kg) was therefore added.

Figure 3 Hand-held fluorescein angiography performed under general anaesthesia 5 months after presentation demonstrating late staining of lesions, disc leakage and a right peri-papillary choroidal neovascular membrane.

Figure 4 Retinal images of right and left eyes 9 months later showing expansion of lesions with development of extensive subretinal fibrosis in the papillomacular region in the right eye and encircling the macular region in the left eye. Optical coherence tomography images showing fibrosis and intra-retinal fluid in the right eye and fibrosis around the fovea in the left eye with central macular retinal atrophy.

Currently, 15 months after presentation, the condition appears to have stabilised, with no further progression of subretinal fibrosis and VA of R 6/38 and L 6/15. The R macula is permanently compromised by subretinal fibrosis and chronic intraretinal fluid. The L fovea is currently spared fibrosis but remains compromised by chronic intraretinal fluid, and threatened by any potential extension of fibrosis (figure 5). Local visual impairment teams are supporting the patient and his family, and a formal visual impairment registration has been completed.

DISCUSSION

The syndrome of PSFU was first described by Palestine¹ in 1984, and there have been fewer than 50 reports of affected patients in the literature in the subsequent 31 years. A degree of subretinal fibrosis is intrinsic to many patients with PIC,² but the degree of involvement and inexorable progression experienced in PSFU is qualitatively different. Prior to the appearance of this patient, this tertiary referral centre had diagnosed 83 patients with PIC but no patients with PSFU, from 3000 new presentations over 24 years.³ The condition usually affects adults, and the youngest patient previously reported was 14 years of age.^{4 5} Confluent subretinal fibrosis in PSFU is usually described as being concentrated at the posterior pole, and this is reflected in our patient, with ultra-widefield imaging showing no peripheral fibrosis despite previous pan-fundal multifocal chorioretinitis. Understanding of the pathogenesis of this condition is limited; histopathological studies (inevitably

available only in the late stages of the disease) demonstrate that the disease is predominantly CD20 B-cell driven with stimulation of fibroblasts.⁶ The underlying trigger is unknown, with infection and auto-antigen exposure hypothesised by Lim *et al.*⁶ Recently, subclinical fungal infection was raised as a potential trigger,⁷ but no evidence of this was found in our patient, who was investigated extensively with no evidence of infection or autoimmunity.

The optimal treatment of PSFU is unknown, with various immunomodulatory strategies described in the literature, including high-dose oral steroids,¹ immunosuppression including cyclophosphamide, ciclosporin and azathioprine,^{4 8} and, recently, biological therapy with infliximab⁹ and rituximab.⁷ Despite potent immunosuppression, PSFU can blind.¹⁰ The aggressive behaviour of the condition in our patient demonstrates this; systemic steroid with methotrexate failed to stop progression; the current stability on ciclosporin may indicate effective therapy or spontaneous resolution.

Our patient underwent sequential treatment in a logical progression: initial empirical penicillin treatment was justified until negative ASO titres were demonstrated; use of a high-dose oral steroid reflected the degree of intraocular inflammation; immunosuppression with methotrexate and, subsequently, ciclosporin, was eventually followed by stabilisation. However, substantial permanent visual loss occurred, and the risk of further fibrosis or secondary CNVM in the left eye remains a significant concern to the clinicians as well as to the child's parents; the use of rituximab in such a situation has been discussed.

The management of complex inflammation in a young child can be very challenging; difficulties in examination and imaging give potential for delayed diagnosis; repeated examination or intervention under general anaesthesia may be necessary and systemic immunosuppression requires careful monitoring. The lack of an evidence base in this situation created difficulties with the choice of immunosuppressive agent, appropriate dosage and duration of therapy.

Figure 5 The left posterior pole 15 months after presentation. There is almost a complete circle of subretinal fibrosis around the fovea, with macular atrophy and chronic intra-retinal oedema.

Learning points

- ▶ Progressive subretinal fibrosis with uveitis may present in a wide age range from young children to the elderly, and may cause substantial visual loss.
- ▶ Ultra-widefield retinal imaging and optical coherence tomography are helpful in documenting extent and progression of chorioretinal lesions, subretinal fibrosis and secondary choroidal neovascular membranes.
- ▶ Aggressive systemic immunosuppression may be necessary to prevent permanent visual loss.

Acknowledgements This research was facilitated by the Greater Manchester Local Clinical Research Network.

Contributors All the authors have been involved in the management of this patient from the outset. All contributed to the writing of the paper. LRS and NJ performed the literature search.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Palestine AG, Nussenblatt RB, Parver LM, *et al.* Progressive subretinal fibrosis and uveitis. *Br J Ophthalmol* 1984;68:667–73.
- 2 Gerstenblith AT, Thorne JE, Sobrin L, *et al.* Punctate inner choroidopathy: a survey analysis of 77 persons. *Ophthalmology* 2007;114:1201–4.
- 3 Jones NP. Manchester uveitis clinic: the first 3000 patients—epidemiology and casemix. *Ocul Immunol Inflamm* 2015;23:118–26.
- 4 Cantrill HL, Folk JC. Multifocal choroiditis associated with progressive subretinal fibrosis. *Am J Ophthalmol* 1986;101:170–80.
- 5 Onoda S, Shibuya K, Miyasaka H, *et al.* Multifocal choroiditis with subretinal fibrosis. *Nihon Ganka Gakkai Zasshi* 1997;101:711–17.
- 6 Lim WK, Chee SP, Sng I, *et al.* Immunopathology of progressive subretinal fibrosis: a variant of sympathetic ophthalmia. *Am J Ophthalmol* 2004;138:475–7.
- 7 Cornish KS, Kuffova L, Forrester JV. Treatment of diffuse subretinal fibrosis uveitis with rituximab. *Br J Ophthalmol* 2015;99:153–4.
- 8 Amaro MH, Muccioli C, Motta MM. Progressive subretinal fibrosis and multifocal granulomatous chorioretinitis. *Arq Bras Oftalmol* 2006;69:413–15.
- 9 Adán A, Sanmartí R, Burés A, *et al.* Successful treatment with infliximab in a patient with diffuse subretinal fibrosis syndrome. *Am J Ophthalmol* 2007;143:533–4.
- 10 Brown J Jr, Folk JC, Reddy CV, *et al.* Visual prognosis of multifocal choroiditis, punctate inner choroidopathy, and the diffuse subretinal fibrosis syndrome. *Ophthalmology* 1996;103:1100–5.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow