Published in final edited form as:

Tetrahedron Lett. 2015 June 3; 56(23): 3507–3510. doi:10.1016/j.tetlet.2015.02.064.

A Practical, Multi-gram Synthesis of (±)-Herbindole A, (±)-Herbindole B, and (±)-Herbindole C from a Common Intermediate via 6,7-Indole Aryne Cycloaddition and Pd(0)-Catalyzed Cross-Coupling Reactions

Nalin Chandrasoma¹, Sivadarshini Pathmanathan², and Keith R. Buszek^{1,2}

Keith R. Buszek: buszekk@umkc.edu

¹Department of Chemistry, University of Missouri, 205 Kenneth A. Spencer Chemical Laboratories, 5100 Rockhill Road, Kansas City, MO 64110

²Center of Excellence in Chemical Methodologies and Library Development (KU-CMLD), University of Kansas, Delbert M. Shankel Structural Biology Center, 2034 Becker Drive, Lawrence, KS 66047

Abstract

A practical, multi-gram 10-step synthesis of racemic herbindole A, B, and C from a common intermediate is described. The key step features a remarkably regioselective C-7 metal-halogen exchange and elimination from a Bartoli-generated *N*-*t*-butyldimethylsilyl-4,6,7-tribromo-5-methylindole scaffold to afford the 6,7-indole aryne. Cycloaddition with cyclopentadiene, oxidative cleavage, and Fujimoto reduction gave a common intermediate from which all three herbindoles were readily derived. A final Pd(0)-catalyzed Negishi and Stille cross-coupling reaction at the C-4 bromide afforded each of the herbindoles on a multigram scale.

Graphical Abstract

Keywords

herbindole; trikentrin; indole; indolyne; aryne; hetaryne; cycloaddition; cross-coupling; Bartoli; Negishi

Correspondence to: Keith R. Buszek, buszekk@umkc.edu.

Supplementary data

Supplementary data (¹H and ¹³C NMR data for all new compounds reported and experimental details for their preparation can be found under supplementary material as a PDF document) associated with this article can be found, in the online version, at <http://dx.doi.org/>.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The herbindoies **1-3** are three analogous 6,7-benzannulated indole natural products that were isolated from the Western Australian sponge *Axinella sp.* and reported by Scheuer in 1990 (Figure 1).¹ They are close structural relatives of the trikentrins that were discovered four years earlier² and are members of a larger though still uncommon class of sinistrally annulated or substituted indoles.

The other benzannulated members include the teleocidins (6,7-benzannulation),³ the nodulisporic acids (5,6-benzannulation),⁴ the penetrems (4,5-benzannulation),⁵ and the notoamides and related stephacidins (6,7-heterobenzannulation).⁶ The biosynthetic origins of the herbindoies (and the trikentrins) remain unknown, though they are probably not derived from tryptophan due to the lack of substitution at the C-3 position.

The herbindoies were reported to exhibit cytotoxic and antifeedant properties, although the levels of these activities were not given in the original citation or to our knowledge in any subsequent report. We recently synthesized the first library of benzannulated indoles inspired by the herbindoies and the trikentrins in an effort to develop new chemotypes in support of drug discovery initiatives currently underway by us and our collaborators.⁷ Several members of this library exhibited provocative levels of antiproliferative activity that mimicked the effect of vincristine, but not that of taxol, on tubulin polymerization.^{7a} Other members of these annulated indole libraries remarkably increased the rate and level of actin polymerization in a manner similar to jasplakinolide, suggesting, *inter alia*, that they might also stabilize the cleavage furrow to block cytokinesis.^{7b} Since these compelling biological results have already validated the significance of such libraries, the development of additional small molecule collections based on these systems clearly has much merit.

The synthesis of this library relied on a key strategy of regioselective C-7 metal-halogen exchange of a Bartoli-generated⁸ 4,6,7-tribromoindole scaffold **4** that permitted the facile generation of the 6,7-indole aryne, **6**,^{9,10} followed by cycloaddition with cyclopentadiene and furan (Scheme 1).

The remaining C-4 aryl bromide was subjected to various Pd(0)-catalyzed cross-coupling reactions to create a diverse library of compounds **8a** and **8b** that were predicted to show *a priori* favorable pharmacokinetic properties.^{7a} During the course of these investigations, it occurred to us that a practical, multi-gram synthesis of each of the herbindoies should be possible by a simple modification of the library tactic. Practical and efficient routes to important natural products have been recognized as major contemporary objectives of synthetic organic chemists.¹¹ We now report in this Letter an efficient, practical and economical route to the herbindoies from a common intermediate *via* our indole aryne cycloaddition and cross-coupling methodology,^{7a} and on a scale that is sufficient to meet the demand for these fascinating compounds.

The herbindoies and the trikentrins have been characterized as “deceptively simple”¹² natural products “whose relatively small size belies their complexity”.^{13a} Indeed, several novel approaches to address these challenging systems have been reported by other laboratories. The first successful total synthesis efforts by Natsume in 1992 were ostensibly designed to confirm the structure of the herbindoies and to establish their absolute

configuration.¹⁴ Subsequent total syntheses by Kerr relied on a novel quinone imine cycloaddition as the key step,¹³ while a more recent report by Sato used a Rh(I)-catalyzed [2+2+2] cyclization process to access an optically pure late-stage 2,3-dihydro indole intermediate.¹⁵ None of these approaches, notwithstanding their importance and synthetic elegance, appeared to be amenable to a scalable synthesis of the herbindoles.

Our retrosynthetic analysis is shown in Scheme 2, and was influenced in part by our own previous experience with the total synthesis of herbindole A and B.¹⁶ The aim was to effect a regioselective metal-halogen exchange from the 4,6,7-tribromo-5-methylindole system **14** at C-7, thereby generating the requisite indole aryne **12**.

Although we previously demonstrated that similar regioselective exchanges occurred at C-7 in the 4,6,7-tribromo- and 5,6,7-tribromoindole cases,^{17,18} it was not at all obvious that the presence of an electron-donating alkyl group at C-5 would favorably impact the proposed regioselectivity. Presumably the alternative exchange at C-6 would still afford the indole aryne; however, preferential or exclusive exchange at C-4 would render the plan moot.

The synthesis of the 4,6,7-tribromo-5-methylindole began with commercially available and economically feasible 5-methyl-2-nitroaniline **16** (Scheme 3). Bromination (HBr, 3 eq; H₂O₂, 2 eq; MeOH, rt, 6 h) afforded the 2,4-dibromo-3-methyl-6-nitroaniline, **17** in quantitative yield on a 20 g scale. Diazotization (CuBr₂, 1.3 eq; t-BuONO, 1.6 eq; MeCN, 65 °C, 1 h) consistently gave the 1,3,4-tribromo-2-methyl-5-nitrobenzene **15** in 91% yield, again on a 20 g scale.

Application of the Bartoli reaction (CH₂=CHMgBr, 3.5 eq; THF, -40 °C, 1 h) to this intermediate on a 15 g scale gave the desired indole **18** (40%) in only three steps. While we have successfully employed the Bartoli reaction previously in natural products total synthesis and library efforts, our experience has been that the yields with polyhalogenated systems are significantly diminished at scales above about 5 g.^{7a} The current example is therefore a fortuitous exception. Finally, the indole nitrogen was protected in 88% yield with a TBS group (NaH, 2.0 eq; Et₃N, 2.0 eq; TBSOTf, 2.0 eq; rt, 0.5 h) which we found to be quite robust.¹⁹

With the desired scaffold in hand, we tested the concept of regioselective C-7 metal-halogen exchange by treating the indole with n-BuLi (2.0 eq; PhMe, -78 °C) and quenching with water (Scheme 4). Gratifyingly, only the C-7 protonated product **13b** was isolated, although the reasons for this selectivity remain a mystery.²⁰ Repeating this process in the presence of cyclopentadiene (10 eq) and slowing warming to room temperature gave the desired Diels-Alder cycloadduct **11** in 90% yield on a 10+ g scale.

Alternatively, the generation of the 6,7-indole aryne could be generated at 0 °C in toluene with only a slight diminution of yield (83–85%). Pd(0)-catalyzed Negishi cross-coupling [Pd₂(dba)₃, 8 mol%; t-Bu₃P•HBF₄, 20 mol%; R₂Zn, 3.0 eq; THF, 60 °C, 2 h] at the remaining C-4 aryl bromide with Me₂Zn and Et₂Zn gave the corresponding and previously reported late-stage intermediates **19** and **20** in 70% and 68% yields, respectively, thus comprising a formal total synthesis of herbindole A and B.¹⁶

Ozonolysis of **11** followed by reduction of the ozonide with sodium borohydride (12 eq) gave the diol **10** in 85% yield (Scheme 5). Adapting the Fujimoto reduction²¹ used by Kerr^{13a} to our intermediate (MsCl, 2.2 eq; Et₃N, 4.0 eq; CH₂Cl₂, 0 °C to rt; then NaI (15 eq); Zn (60 eq); glyme, 80 °C, 12 h) afforded the late-stage common intermediate **9** that was used for all three racemic herbindoles.

The intermediate **9** was routinely prepared on a 10-gram scale and was subjected to the same Negishi cross-coupling conditions as described earlier. The yields with the respective dialkylzinc reactants improved to 86% (R = methyl) and 84% (R = ethyl) with this scaffold compared to the cycloadduct **11**. This improved cross-coupling reaction has been ascribed to the lack of a potentially interfering strained olefin, an observation also made during our benzannulated indole library synthesis.^{7a} Desilylation with tetrabutylammonium fluoride in THF was unexceptional and delivered in each case herbindole A **1** and herbindole B **2** in 96% yield.

To complete the series with herbindole C, we elected, after much trial and error, to adapt the same Stille cross-coupling tactic that was used in the final step of our recent total synthesis of (±)-*cis*-trikentrin B.¹⁸ As before, the use of the triphenylarsine ligand and microwave heating (150 °C, 23 min) was necessary to achieve acceptable yields of protected herbindole C **24**. Deprotection afforded the desired product in 98% yield.

The present work provides yet another important and instructive example of a regioselective metal-halogen exchange reaction and indole aryne cycloaddition in a tribromoindole scaffold. This finding, combined with the scalability of the reactions, affords for the first time the synthesis of each of the herbindoles in the multi-gram quantities that are needed for further study and research. To put this achievement into some perspective, the Scheuer report of the isolation of the herbindoles in 1990 resulted in just 230 mg of all three herbindoles combined, from about 0.75 kg dry weight of the sponge.

Given that benzannulated indole libraries inspired by the herbindoles and the structurally similar trikentrins show promising preliminary results (e.g., vincristine surrogates),^{7a,b,c} the need for a practical synthesis of the herbindoles is clear and compelling. With sufficient quantities in hand, modification of the herbindoles, primarily around the pyrrole ring, is presently underway. A report of this activity, along with the biological annotation of the resulting libraries, will be disclosed as developments warrant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We acknowledge support of this work by the National Institutes of Health, NIGMS, Grant R01 GM069711 (KRB). We also acknowledge additional support of this work by the National Institutes of Health, The University of Kansas Chemical Methodologies and Library Development Center of Excellence (KU-CMLD), NIGMS, Grant P50 GM069663. We further acknowledge Mr. Alok Nerurkar for his assistance and expertise with the preparation of this manuscript. We are grateful to Dr. Chris Sakai (Kibo Biosciences, LLC) for his valuable advice and technical assistance with this project.

References and notes

1. Herb R, Carroll AR, Yoshida WY, Scheuer PJ. *Tetrahedron*. 1990; 46:3089–3092.
2. Capon RJ, MacLeod JK, Scammells PJ. *Tetrahedron*. 1986; 42:6545–6550.
3. Nakatsuka S, Matsuda T, Goto T. *Tetrahedron Lett*. 1987; 38:3671–3674.
4. Singh SB, Ondeyka JG, Jayasuriya H, Zink DL, Ha SN, Dahl-Roshak A, Greene J, Kim JA, Smith MM, Shoop W, Tkacz JS. *J Nat Prod*. 2004; 67:1496–1506. [PubMed: 15387649]
5. Smith AB III, Kanoh N, Ishiyama H, Hartz RA. *J Am Chem Soc*. 2000; 122:11254–11255.
6. Kato H, Yoshida T, Tokue T, Nojiri Y, Hirota H, Ohta T, Williams RM, Tsukamoto S. *Angew Chem Int Ed*. 2007; 46:2254–2256.
7. (a) Thornton PD, Brown N, Hill D, Neuenswander B, Lushington GH, Santini C, Buszek KR. *ACS Comb Sci*. 2011; 13:443–448. [PubMed: 21668016] (b) Perchellet JPH, Waters AM, Perchellet EM, Thornton PD, Brown N, Hill D, Neuenswander B, Lushington GH, Santini C, Chandrasoma N, Buszek KR. *Anticancer Res*. 2012; 32:4671–4684. [PubMed: 23155229] (c) Perchellet JPH, Perchellet EM, Singh CR, Monnet MT, Studer E, Thornton PD, Brown N, Hill D, Neuenswander B, Lushington GH, Santini C, Buszek KR. *Anticancer Res*. 2014; 34:1643–1655. [PubMed: 24692693] (d) Brown N, Xie B, Markina N, VanderVelde D, Perchellet JPH, Perchellet EM, Crow KR, Buszek KR. *Bioorg Med Chem Lett*. 2008; 18:4876–4879. [PubMed: 18703334] (e) Brown N, Gao G, Minatoya M, Xie B, VanderVelde D, Lushington GH, Perchellet JPH, Perchellet EM, Crow KR, Buszek KR. *J Comb Chem*. 2008; 10:628–631. [PubMed: 18693764]
8. Dalpozzo R, Bartoli G. *Curr Org Chem*. 2005; 9:163–178.
9. (a) Buszek KR, Luo D, Kondrashov M, Brown N, VanderVelde D. *Org Lett*. 2007; 9:4135–4137. [PubMed: 17880092] (b) Brown N, Luo D, VanderVelde D, Yang S, Brassfield A, Buszek KR. *Tetrahedron Lett*. 2009; 50:63–65. [PubMed: 21057588] (c) Garr AN, Luo D, Brown N, Cramer CJ, Buszek KR, VanderVelde D. *Org Lett*. 2010; 12:96–99. [PubMed: 19961152] (d) Nerurkar A, Chandrasoma N, Maina L, Brassfield A, Luo D, Brown N, Buszek KR. *Synthesis*. 2013; 45:1843–1852.
10. For recent reviews of arynes and hetarynes in organic synthesis, see: Gampe CM, Carreira EM. *Angew Chem Int Ed*. 2012; 51:3766–3778. Tadross PM, Stoltz BM. *Chem Rev*. 2012; 112:3550–3577. [PubMed: 22443517] Goetz AE, Shah TK, Garg NK. *Chem Commun*. 2014 Ahead of print.
11. Wender PA, Miller BL. *Nature*. 2009; 460:197–201. [PubMed: 19587760]
12. Liu W, Lim HJ, RajanBabu TV. *J Am Chem Soc*. 2012; 134:5496–5499. [PubMed: 22394308]
13. (a) Jackson SK, Banfield SC, Kerr MA. *Org Lett*. 2005; 7:1215–1218. [PubMed: 15787470] (b) Jackson SK, Kerr MA. *J Org Chem*. 2007; 72:1405–1411. [PubMed: 17256909]
14. Muratake H, Mikawa A, Natsume M. *Tetrahedron Lett*. 1992; 33:4595–4598.
15. Saito N, Ichimaru T, Sato Y. *Org Lett*. 2012; 14:1914–1917. [PubMed: 22452396]
16. Buszek KR, Brown N, Luo D. *Org Lett*. 2009; 11:201–204. [PubMed: 19055375]
17. Brown N, Luo D, Decapo JA, Buszek KR. *Tetrahedron Lett*. 2009; 50:7113–7115. [PubMed: 20877442]
18. Chandrasoma N, Brown N, Brassfield A, Nerurkar A, Suarez S, Buszek KR. *Tetrahedron Lett*. 2013; 54:913–917. [PubMed: 25278636]
19. Indoles protected with electron-withdrawing groups such as sulfonyl do not undergo metal-halogen exchange. Interestingly, unprotected indoles do participate in metal-halogen exchange, but at the C-6 position instead (see reference 9d).
20. *Ab initio* calculations with DFT methods (reference 9c) performed on the indole arylene, benzofuran, and benzothiophene systems revealed the 6,7-aryne bond to be highly polarized, with a nucleophilic C-7 carbon. See also: Brown N, Buszek KR. *Tetrahedron Lett*. 2012; 53:4022–4025. [PubMed: 25278635]
21. Fujimoto Y, Tatsuno T. *Tetrahedron Lett*. 1976; 37:3325–3326.

Figure 1.
The structures of the herbindoles.

Scheme 1.

Formation of 6,7-indole aryne and cycloaddition via regioselective metal-halogen exchange and elimination.

Scheme 2.
Retrosynthetic analysis of the herbindoles.

Scheme 3.

Synthesis of the 4,6,7-tribromo-5-methylindole scaffold.

Scheme 4.

Regioselective C-7 metal-halogen exchange/elimination and cycloaddition.

Scheme 5.

Fujimoto reduction and Pd(0)-catalyzed Negishi and Stille cross-coupling reactions for herbindole A, B, and C.