Published in final edited form as:

J Comp Psychol. 2015 August ; 129(3): 291–303. doi:10.1037/a0039435.

Social conditioned place preference in the captive ground squirrel *Ictidomys tridecemlineatus*: Social reward as a natural phenotype

Garet P. Lahvis^{a,b,*}, Jules B. Panksepp^{a,c,*}, Bruce C. Kennedy^{a,b}, Clarinda R. Wilson^b, and Dana K. Merriman^d

^aDepartment of Behavioral Neuroscience, Oregon Health and Science University, Portland, Oregon

^bDepartment of Surgery, University of Wisconsin, Madison, Wisconsin

^cNeuroscience Training Program, University of Wisconsin, Madison, Wisconsin

^dDepartment of Biology and Microbiology, University of Wisconsin, Oshkosh, Wisconsin

Abstract

Social behaviors of wild animals are often considered within an ultimate framework of adaptive benefits versus survival risks. By contrast, studies of laboratory animals more typically focus on affective aspects of behavioral decisions, whether a rodent derives a rewarding experience from social encounter and how this experience might be initiated and maintained by neural circuits. Artificial selection and inbreeding have rendered laboratory animals more affiliative and less aggressive than their wild conspecifics, leaving open the possibility that social reward is an artifact of domestication. We compared social behaviors of wild and captive population of juvenile 13-lined ground squirrels (*Ictidomys tridecemlineatus*), the latter 2nd and 3rd generation descendents of wild individuals. At an age corresponding to emergence from the burrow, postnatal day 38, captive squirrels engaged in vigorous social approach and play; these juvenile behaviors declined significantly by postnatal day 56. Similarly, young wild squirrels expressed social proximity and play, affiliative interactions declined with summer's progression and were replaced by agonistic chasing behaviors. Social conditioned place preference testing (conditioned postnatal days 40–50) indicated that adolescent squirrels derived a rewarding experience from social reunion. Our results support the contention that undomesticated rodents have the capacity for social reward and more generally suggest the possibility that positive affective experiences may support group cohesion and social cooperation in the wild.

Keywords

social reward; classical conditioning; pro-social behavior; ground squirrel; play-fighting

Correspondence to Garet P. Lahvis, PhD, Department of Behavioral Neuroscience, Oregon Health and Science University, 3181 SW Sam Jackson Park Road, Mail Code L470, Portland, Oregon 97239-3011, USA, lahvisg@ohsu.edu.

*Authors contributed equally to this work.

Introduction

Mammals express an ontogeny of social behaviors, beginning with mother-infant interactions, continuing through playful interactions with peers, and maturing into the more stereotypical, sex-dependent patterns characteristic of adults (Wolff & Sherman, 2008). Though sharing commonalities, the sociality of ground squirrel species varies in both duration and degree (Armitage, 1981; Hare & Murie, 2008; Rayor & Armitage, 1991). Juvenile ground squirrels engage in social play soon after they emerge from their natal burrows (Steiner, 1971; Yeaton, 1972). As summer progresses, play behaviors diminish (Betts, 1976; Nunes, Muecke, Anthony, & Batterbee, 1999), with adolescents often wandering beyond their natal range, an action that can lead to emigration (Barash, 1974; Holekamp, 1984; McCarley, 1966; Michener & Michener, 1977; Rayor & Armitage, 1991). Even among the more social Columbian ground squirrels, *S. columbianus*, adult social interactions rarely seem affiliative, typically consisting of dyads and triads (Manno, 2008). Among less social species, including *I. tridecemlineatus*, arctic ground squirrels, *S. parryii plesius*, and Richardson's ground squirrels, *S. richardsonii*, social interactions are expressed rather as conflicts for mating opportunities and territory (Lacey, Wieczorek, & Tucker, 1997; Schwagmeyer & Woontner, 1986; Schwagmeyer & Parker, 1987; Luna & Baird, 2004).

Mammalian social behaviors are often considered within the ultimate framework of adaptive benefits versus survival risks (Silk, 2007). For instance, juvenile play is thought to improve physical and emotional responsiveness to unexpected events (Nunes et al., 1999; Spinka, Newberry, & Bekoff, 2001), familiarize juveniles with self handicap and fair behavior (Bekoff, 2004), improve abilities to cope with social challenges (van den Berg et al., 1999), establish dominance relationships (Blumstein, Chung, & Smith, 2013), and help refine abilities to respond to subtle and ambiguous social signals (Pellis, Pellis, & Reinhart, 2010). Such adaptive benefits are difficult to test (Burghardt, 2005; Pellis & Pellis, 2006) but are suggested by evidence that social play promotes normal brain development (Gordon, Burke, Akil, Watson, & Panksepp, 2003; Pellis & Pellis, 2007).

Like juvenile play, dispersal also has adaptive benefits that likely vary among squirrel species (Bowler & Benton, 2005; Lawson Handley & Perrin, 2007; Wolff, 1994). Benefits include competitive mating advantages (Dobson, 1982), access to higher quality resources (Dobson, 1979), avoidance of inbreeding (Wauters & Dhondt, 1993) and evasion from parasitism (Hoogland, 1979); these benefits can offset dispersal's enormous survival risks (Byrom & Krebs, 1999). Proximate explanations for variations in adolescent dispersal include differences in environmental harshness (Barash, 1974; Ritchie & Belovsky, 1990), adult aggression toward maturing adolescents (Steiner, 1972; Wolff, 1993), and ontogenetic changes promoting excursion behavior (Holekamp, 1984; Holekamp, 1984). Proximate explanations might also include psychological changes. Exploratory activity of laboratory rodents can be greater among adolescents than either younger juveniles or adults (Macri, Adriani, Chiarotti, & Laviola, 2002). Affiliative social behaviors also change with maturation (Panksepp et al., 2007; Terranova, Laviola, & Alleva, 1993). Exploratory and social behaviors utilize neurocircuits (Laviola, Macri, Morley-Fletcher, & Adriani, 2003; Liu & Wang, 2003; Resendez, Kuhnmuensch M., Krzywosinski T., & Aragona, 2012).

activated by natural and drug rewards (Spear, 2000; Kelley & Berridge, 2002; Dölen et al. 2013).

We can infer an animal subject finds a stimulus rewarding *if* the subject returns to an environment associated with that stimulus (Glickman & Schiff, 1967; Schneirla, 1959). Measures of conditioned place preference (CPP) allow us to formally assess this possibility. In the conditioning phase, the subject is alternated between one environment paired with the presence of a stimulus and a second environment associated with its absence. During the test phase, both environments are available to the subject, though no stimulus is present, and the subject is allowed to amble back and forth between the two conditioned environments (Bardo & Bevins, 2000; Tzschentke, 2007). CPP has demonstrated that laboratory rodents typically prefer environments associated with social access, a behavior driven by anticipation of a social reward (Calcagnetti & Schechter, 1992; Douglas, Varlinskaya, & Spear, 2004) and by avoidance of environments paired with social isolation (Panksepp & Lahvis, 2007).

Though laboratory rodents can express social reward-related phenotypes during a CPP test, similar processes may not of necessity regulate the social behaviors of wild rodents. Confinement of laboratory rodents for generations to small cages fosters a selection bias for individuals with relatively low levels of territoriality and activity concomitant with tolerance for high housing densities (Boice, 1981; Chalfin et al., 2014; Kunzl, Kaiser, Meier, & Sachser, 2003; Kunzl & Sachser, 1999). Thus, conceivably, multi-generational confinement might cultivate phenotypes expressing strong responses to social CPP tests. Many laboratory rodents were selected for ‘desirable’ traits (Beck et al., 2000; Wade & Daly, 2005) and then inbred for over 200 generations for medical research (see <http://jaxmice.jax.org/findmice/index.html>). Though social CPP has been designed primarily with face and construct validity relative to psychological concepts, this test has not been utilized for rodents lacking an extensive history with domestication and genetic inbreeding. This experimental gap leads to an obvious question regarding the applicability and relevance of such testing scenarios to wild or captive rodents.

In this paper, we present findings from a novel experimental opportunity that employs laboratory tests of rodent social interaction (Panksepp et al., 2007) and social reward (Panksepp & Lahvis, 2007) to study captive 13-lined ground squirrels, *Ictidomys tridecemlineatus*, 2nd and 3rd generation descendents of wild individuals (Merriman et al., 2012). Our choice of generation for using captive born ground squirrels is supported by studies showing that wild Norway rats robustly express wild behaviors within the first 2–5 generations of captivity in the absence of any direct selection for tameness (Stryjek & Pisula, 2008). Litters of 13-lined ground squirrel are born in late May to early June (McCarley, 1966; Merriman, Lahvis, Jooss, Gesicki, & Schill, 2012), emerging from their burrows at about five weeks of age, forage, engage in play, then disperse within a month of their emergence (McCarley, 1966; Rayor & Armitage, 1991; Wistrand, 1974). Young females more commonly remain within the colony (Schwagmeyer, 1980), whereas males typically emigrate (Armitage, 1981; McCarley, 1966).

We compared social interaction (SI) in captive, same- and mixed-sex dyads, with contemporaneous observations of wild squirrels at times approximating emergence from the burrow and dispersal. We found that captive and wild ground squirrels expressed analogous declines in affiliative behaviors with the progression of summer. Employing an adapted social CPP paradigm (Panksepp & Lahvis, 2007), we found that adolescent squirrels expressed a robust place preference for environments associated with social interactions, indicating social reward from reunion with their peers. Overall, squirrel social behaviors resemble patterns expressed by highly domesticated and inbred rodents, strengthening the value of using psychological approaches for understanding the natural developmental transitions in rodent social affiliation.

Method

Capture

Detailed descriptions of the capture, housing and husbandry of 13-lined ground squirrels have been previously published (Vaughan, Gruber, Michalski, Seidling, & Schlink, 2006b). A Scientific Collector's Permit (NER SCP 144) was obtained from the Wisconsin Department of Natural Resources prior to capture. The colony of 13-lined ground squirrels used here was derived from animals trapped at cemeteries located in Oshkosh, WI (44.02°N 88.53°W) and Neenah, WI (44.19°N 88.46°W).

Husbandry

Thirteen-lined ground squirrels were bred for 2 generations in the University of Wisconsin-Oshkosh colony located at the Halsey Science Center (Oshkosh, WI, USA). Subjects used for social behavior and CPP testing (see below) were either 2nd or 3rd generation descendents of individuals captured during the initial trapping excursions. Except for purposes of breeding (see below), adult squirrels were housed individually in standard polypropylene cages (412x210x200 mm) that contained shaved aspen bedding (Teklad Aspen Sani-Chips, Harlan, Indianapolis, IN, USA) and a plastic, rain gutter tube (200x65x65 mm) available for use as a burrow. Squirrels received *ad libitum* access to water and commercial dog chow (IAMS Chunks) supplemented with sunflower seeds. Ambient temperature was maintained at $68 \pm 2^\circ\text{F}$ and the light-dark cycle was periodically adjusted to correspond with sunrise and sunset in the location captured until squirrels entered hibernation in the fall. During hibernation, squirrels were housed at $38 \pm 3^\circ\text{F}$ in constant darkness. After arousal and recovery from hibernation, sexually mature male and female squirrels were housed together for up to 4 weeks before pregnant females were moved to single housing for birth and pup upbringing. Pups born in the enclosures received a combination of dog chow, cat chow (Purina Indoor Formula) and sunflower seeds until weaning.

Juveniles used in this study were not handled from birth or otherwise tamed, so their handling was thus no different from that of animals recently caught from the wild (Vaughan, Gruber, Michalski, Seidling, & Schlink, 2006a). Individuals were thus highly capable of escape and expressed some skittish behaviors. All cage changes were performed with the home cage placed in a deep plastic "tote" bin. Soft cloth nets identical to those used for field

captures were used for the occasional escapee. Whenever it was necessary to transfer or restrain them, leather welder's gloves were used to protect from inevitable bites.

On postnatal day (PD) 35/36, squirrels were weaned into mixed-sex social groups (2 males and 2 females) from either a single litter (6 groups) or a combination of multiple litters (8 groups), and these groupings served as the subsequent housing arrangement between all of the behavioral procedures. Statistical differences between squirrels from full-sibling vs. partial-sibling groups were not detected and are not considered further. All cages of weaned juveniles were moved to a separate room outside the main colony that received natural lighting through room windows.

Social interaction test

Social interaction (SI) tests were conducted on PD 38/39 and again on PD 56/57 (Fig. 1). To engender social motivation, each individual within a social group was isolated into a clean cage 24 h prior to testing (see Panksepp, Wong, Kennedy, & Lahvis (2008) for rationale). One male and one female individual from each social group were randomly designated as test squirrels and the remaining two as stimulus squirrels. All test/stimulus combinations were evaluated (i.e., F-F, F-M, M-F and M-M), and these designations were maintained for the tests on PD 38/39 and PD 56/57. To identify individuals, the back of each squirrel was marked with hair dye. On test day, squirrels were habituated to a windowless experimental room (approximately 4x4m) illuminated with dim fluorescent lighting (adjacent to the colony room) for at least 90 min before testing. Five min prior to testing, the cage top was replaced with clear Plexiglas®. The SI test consisted of placing a stimulus squirrel into the home cage of a test squirrel from the same social group. Behaviors of the test and stimulus squirrels were video-recorded with an overhead camcorder (Sony, DCR-VX2100) for a 5-min period. Following each test, squirrels were returned to their original social group in a fresh cage. Testing was conducted in the afternoon from 1200–1900 h. All laboratory procedures were conducted from June 20, 2007 (sunrise 0511 h, sunset 2040 h) to July 7, 2007 (sunrise 0518 h, sunset 2040 h). Following completion of the first SI test, squirrels were returned to housing in their respective social groups.

Three categories of behavior were assessed during the SI test; social play (Nunes et al., 1999; Pasztor, Smith, MacDonald, Michener, & Pellis, 2001; Pellis, MacDonald, & Michener, 1996), social investigation (Grubitz, 1963; Michener & Sheppard, 1972; Panksepp et al., 2007) and vigilant behavior (Arenz & Leger, 1999). See Table 1 for descriptions of each behavioral category. Behaviors were analyzed with computer-assisted analysis software (ButtonBox v5.0, Behavioral Research Solutions) by two independent observers, and all presented data and statistical analyses are based on an average of these two measurements (inter-rater reliability, Pearson's correlation coefficient for social investigation – $r = 0.90$, play behavior – $r = 0.98$ and vigilance – $r = 0.80$). A total of 44 squirrels were tested for SI.

Social conditioned place preference test (SCPP)

After 24 h in their home cage, on PD 39/40, social groups were housed together in a fresh cage containing one of two novel conditioning environments (see below). The conditioning

environments were composed of either ‘paper’ (soft paper bedding; Cellu-Dri Soft, Shepherd Specialty Papers) and two smooth 1 ½” PVC couplers or ‘corncob’ (¼ inch grain size corncob bedding; Harlan, Teklad) and two threaded 1 ½” PVC couplers (see Panksepp & Lahvis (2007) for additional background on SCPP experiments). After the first 24 h social conditioning period, squirrels were separated and isolated into a fresh cage containing the alternate conditioning environment. Social and isolate housing contexts and their respective conditioning environments were alternated every 24 h over a 10-day period (see Fig. 1). In both environments, squirrels were provided with *ad libitum* dog chow and water. To assess whether squirrels had a natural preference for either of the 2 conditioning environments, a control group (n=15) was alternated daily as a social unit between the two environments (i.e., no conditioning).

Habituation to the testing situation entailed placing squirrels individually into a CPP testing apparatus after conditioning on days 8 and 9, and they were allowed to move freely about without the conditioning environments present for 20 min under dim fluorescent lighting. The CPP apparatus (constructed of ABS Plastic) consisted of three adjacent compartments (each 480mm x 247mm x 200mm) accessible to the test squirrel via square openings (76x76mm). On the test day (PD 49/50), the ‘paper’ and ‘corncob’ environments were assembled in the peripheral compartments. An individual test squirrel was placed in the central compartment (‘lexan’ plastic floor) and the apparatus was covered with a transparent Plexiglas® top. Movement between compartments was recorded from an overhead-video camera for 30 min under dim lighting. Following the SCPP test, squirrels were returned to their original social groups in a fresh cage. Two independent observers analyzed 40% (n=20) of the SCPP tests (inter-rater reliability, Pearson’s correlation coefficient, $r = 0.99$). Data from a small subset of ground squirrels (n=7) was excluded from statistical analyses because these squirrels failed to explore the conditioning environments during testing, defined as remaining motionless for 15min of the 30 min test or repeatedly attempting to escape from the testing arena. Six additional squirrels were excluded due to technician error during the conditioning procedure. A total of 41 squirrels were evaluated for SCPP. The UW-Oshkosh IACUC approved all experimental procedures (protocol number: 0-03-0026-000188-5-18-07).

Field observations

The University of Wisconsin-Madison permitted us to observe and video-record 13-lined ground squirrels at a field site (43.06° N, 89.52° W) maintained by the Agriculture Research Station (ARS), approximately 90 miles from Oshkosh, WI, where the laboratory studies were conducted. The site consisted of over 60% perennial ryegrass (*Lolium perenne*), along with tall fescue (*Festuca arundinacea*) and common meadow grass (*Poa pratensis*). Field observations were conducted at various time points between 1530 and 1900 h over a 5-week period between late June and early August of 2007 (6/26-8/02). The length of each observation period varied with levels of squirrel activity, weather conditions and use of the field by ARS staff. Observation periods occurred 3–4 days/week and each period lasted 107 ± 28 min/day (mean $\pm$ s.d.). ARS staff avoided the field site during observation periods. Ground squirrel activity was monitored for the occurrence of interactions between 2 or more squirrels within an area of approximately 400 m² and ranging from approximately 10 to 30

meters from the observer. Social interactions observed on the site were video-recorded with a professional video camera under 20X optical zoom (XL2 20x IS zoom, Canon). Once a social interaction began, recording continued until the interacting ground squirrels dispersed, entered a burrow or disappeared into taller grass.

To quantify field observations of wild squirrels, an observer first evaluated behaviors while watching video recordings in real time. Beginning anytime two or more squirrels entered the camera's field of view, the observer noted the time stamp and then classified behaviors according to an ethogram (Table 2). In subsequent analysis, the observer returned to each time stamp and conducted frame-by-frame analysis of the video recordings at 30-fold slower speed via iMovie (Apple). In cases where there were questions about a particular behavior, a second rater was consulted. A second independent observer analyzed 50% (n=122) of the field behaviors to assess reliability between observers (inter-rater agreement, Cohen's kappa coefficient, $\kappa = 0.72$). Social behaviors were classified according to an ethogram similar to the one used for captive squirrel behavior that included the affiliative behaviors observed in captivity along with spatial proximity and chasing behaviors observed in the wild (Betts, 1976; Michener & Sheppard, 1972). See Table 2 for descriptions of each behavioral category. Frequencies of each behavior were tallied for each week of observation.

Statistical analyses

A 2x4 analysis of variance (ANOVA) was used to evaluate effects of age and the sex pairing of squirrels on play behavior, social investigation and vigilance, respectively, during the SI tests. For SCPP testing, a 2x3 ANOVA was utilized, with sex and experimental group as between-subjects factors. Specific comparisons between groups were conducted with orthogonal contrasts (which include averaged error variance across all experimental groups). Data from the field are presented as descriptive statistics for weekly totals of individual types of social behavior (e.g., approach, play, etc.). 'Expected' weekly counts for the combined social behavior measure were generated assuming an equal distribution over the 5-week observation period from late-June to early August and are represented in Figure 4. Expected values for each specific type of social behavior were adjusted for observed changes in social behavior (combined) across the 5-week period. All analyses were performed using JMP v8.0.1 (SAS Institute Inc.). Statistical significance was set at $P < 0.05$.

Results

Social interaction test

Adolescent squirrels engaged in more play behavior during the first SI test (on PD 38/39) than the second test (PD 56/57) occurring 20 days later (Fig. 2A, main effect of age, $F[1,42] = 5.2$, $P = 0.03$, $R^2=0.08$). Play behavior was also sensitive to the sex of the interacting squirrels (Fig. 2A', main effect of sex, $F[3,40] = 5.6$, $P = 0.003$, $R^2=0.27$), with the highest level of play observed during interactions between two males (orthogonal contrast, male-male pairs vs. all other groups, $F[1,36] = 15.9$, $P = 0.0003$). The influence of sex on social play did not differ between the first and second SI tests (age x sex interaction, $F[3,40] = 1.7$, $P = 0.19$).

Social investigation declined between PD 38 and PD 56 (Fig. 2B, main effect of age, $F[1,42] = 4.4$, $P = 0.04$, $R^2=0.08$) and was dependent on the sex of both the test and stimulus animal (Fig. 2B', main effect of sex, $F[3,40] = 5.2$, $P = 0.004$, $R^2=0.26$). Social investigation was highest during female-female interactions compared to all other sex pairings (orthogonal contrast, female-female pairs vs. all other groups, $F[1,36] = 15.1$, $P = 0.0004$). The influence of sex on social investigation did not vary with age (age x sex interaction, $F[3,40] = 0.9$, $P = 0.44$).

The expression of vigilant behavior was greater during the second SI test than during the first test (Fig. 2C, main effect of age, $F[1,42] = 13$, $P = 0.0009$, $R^2=0.24$). Vigilant behavior was not dependent on the sex of the interacting squirrels (Fig. 2C', main effect of sex, $F[3,40] = 0.4$, $P = 0.73$). Moreover, vigilance during social interactions was negatively correlated with social play (Pearson's correlation coefficient, $r = -0.38$, d.f. = 43, $P = 0.009$) and with social investigation ($r = -0.24$, d.f. = 43, $P = 0.12$), although the latter relationship failed to reach statistical significance. There was no relationship between play behavior and social investigation ($r = -0.05$, d.f. = 43, $P = 0.76$). During SI testing, the time that elapsed while squirrels were engaged in social investigation, play and vigilance accounted for less than half of the 5-min testing session (Fig. 2D), indicating that the associations between these behaviors were not impacted by the duration of the testing period.

Social conditioned place preference test

Social conditioning altered the time spent in each of the peripheral compartments of the CPP arena (Fig. 3, main effect of conditioning, $F[2,38] = 11.8$, $P < 0.0001$, $R^2=0.38$). Although unconditioned squirrels expressed a modest preference for the paper bedding ($t = -1.44$, d.f. = 15, $P = 0.08$), association of the paper environment with social housing resulted in a CPP relative to this control group (orthogonal contrast, $F[1,34] = 4.9$, $P = 0.03$). When social exposure was paired with the corncob bedding, social conditioning resulted in an increased duration spent in the corncob bedding (orthogonal contrast, $F[1,34] = 10.3$, $P = 0.003$). There was no effect of the sex of test animals on social conditioning (main effect of sex, $F[1,39] = 1.4$, $P = 0.25$) or an interaction (sex x conditioning interaction, $F[2,38] = 1.5$, $P = 0.24$).

Field observations

Wild 13-lined ground squirrels expressed both social play and social investigation behaviors. In all, 228 instances of social behavior were observed during 30 h of video recordings collected in the field over a 5-week period when ground squirrels were visible. In the field, proximity (two or more ground squirrels within 2 body lengths of each other) accounted for 60% of all social behaviors. Chasing accounted for 15% of the total social behaviors observed, followed by approach, play and investigation, which each contributed 7–8%.

The total number of social behaviors observed each week varied substantially (Fig. 4A), but generally decreased over the 5-week period (95 social behaviors were observed in Week 1 vs. 12 social behaviors during Week 5). Affiliative social behaviors were more typical during the first few weeks of observation, whereas agonistic social behaviors were common

at later time points. Behaviors categorized as approach, investigation, play and proximity (Fig. 4B–E) were predominantly observed in late June – mid July (Weeks 1–3), but rarely from late July to early August (Weeks 4–5). By contrast, chasing behavior (Fig. 4F) occurred most frequently during mid July and August (Weeks 3–5).

We observed 228 distinct bouts of ground squirrel social behavior. We did not capture and tag squirrels so were unable to identify individuals. Over our 5 weeks of field observations, we video recorded four or more individuals in close proximity on 3 separate occasions. Considering the low frequency of interactions known to occur between adults, these observations suggest an absolute lower boundary on the number of juveniles sampled in the colony. However, we estimate a considerably larger number of juveniles. For thirteen-lined ground squirrels, 100 burrow entrances correspond to approximately 14 nesting burrows (Desha, 1966). Our estimate of 50 to 100 burrow entrances at the colony indicates 7 to 14 nesting burrows. Each nesting burrow supports on average 8 pups per litter (Rongstad, 1965; Streubel & Fitzgerald, 1978), so we estimate 56–112 juveniles in our colony. The video camera did not provide sufficient field-of-view to record all social interactions observed, so some of the residing juveniles may not have been sampled.

Discussion

At a juvenile age corresponding to emergence from the burrow (PD 38/39), captive 13-lined ground squirrels engaged in vigorous social interactions that differed between sexes: male dyads more typically in bouts of play fighting and females in social investigation. These sex-dependent social patterns were maintained, albeit at lower overall levels, when squirrels were tested as older adolescents (PD 56/57). Our results mirror an early study of captive *S. richardsonii*, showing that non-sibling females preferentially interacted with ‘recognitive’ behaviors (Sheppard & Yoshida, 1971), akin to social investigation, whereas male dyads engaged in ‘agonistic’ behaviors, which included rough-and-tumble behaviors not deemed to be playful. These contrasting interpretations may reflect differences in squirrel rearing environment; most juveniles in the Sheppard and Yoshida study were trapped in wild environments that afforded social refuge so subsequent confinement may have fostered aggression. A different explanation is that Sheppard and Yoshida may have misinterpreted play fighting as aggression. The first report of play-fighting among ground squirrels was published in the same year (Steiner, 1971) and only much later systematically dissociated from agonistic behavior (Pellis & Pellis, 1987; Pellis, 1988). More recently, free-living male juveniles of *S. richardsonii* and *S. beldingi* were shown to engage in more play fighting than females (Nunes et al., 1999; Pasztor et al., 2001). This sexually dimorphic pattern is also expressed by juvenile laboratory rats (Olloff & Stewart, 1978; Pellis, 2002) and play-fighting was found to be rewarding, as revealed by CPP experiments (Calcagnetti & Schechter, 1992; Sivi, 1998). Physiological studies show that play fighting is influenced by activation of perinatal steroid receptors (Meaney & Stewart, 1981; Olesen, Jessen, Auger, & Auger, 2005) and supported by juvenile reward circuits (Burgdorf, Panksepp, Beinfeld, Kroes, & Moskal, 2006; Sivi, Fleischhauer, Kerrigan, & Kuhlman, 1996; Trezza & Vanderschuren, 2008). Environmental harshness may influence the timing of play and dispersal via its effects on calorie intake, a factor that moderates play behavior in laboratory

rats (Siviy & Panksepp, 1985) and free living squirrels (Nunes et al., 1999) and can influence reward circuits (Alamy & Bengelloun, 2012; Vucetic et al., 2010).

Captive and wild squirrels expressed analogous declines in affiliative social behaviors over the summer season despite profound differences in their rearing environments. Unlike their wild counterparts, captive squirrels were provisioned with predictable food availability and quality. To maintain defined levels of social motivation during SI testing (Panksepp et al., 2008), we exposed captive squirrels to specific durations of social isolation. To eliminate possible influences of changing adult sociality on adolescent behavior, we restricted the social environment of captive juveniles to interactions their peers. Were adults to become aggressive in response to more fully-grown adolescents, these behaviors might, in turn, have promoted a shift from affiliative to agnostic behaviors among juveniles. Confining captive squirrels to 'adult-free' housing, we imposed a less variable, more unnatural, social environment. In light of these considerable differences, the similar seasonal decline in affiliative social behaviors of captive and wild squirrels underscores common developmental changes. Taken together, this comparison supports the idea that factors intrinsic to the maturing 13-line ground squirrel moderate sociality, akin to the idea that an 'ontogenetic switch' within the adolescent's 'internal milieu' influences social development (Holekamp, 1984).

We also urge caution in relating social behaviors expressed in nature with those expressed in the lab. Comparative field studies indicate 13-lined ground squirrels are among the least social of the ground squirrel species (Rayor & Armitage, 1991; Wistrand, 1974). Some studies consider *I. tridecemlineatus* to be 'asocial' (McCarley, 1966) and see also (Luna & Baird, 2004), forming colonies not out of social attraction, rather because they share a preference for complex habitats that *reduce* social contact (McCarley, 1966). By contrast, we were impressed by the high levels of social investigation expressed by captive squirrels when contrasted with our own experience with laboratory mice. Captive squirrels spent 25–40% of a 5-min behavioral test interacting socially (see Figure 2d). This level of social investigation lies between levels expressed by a gregarious inbred laboratory mouse strain (~60%) and those of a less social strain (~20%) (Fairless et al., 2008; Panksepp et al., 2007; Sankoorikal, Kaercher, Boon, Lee, & Brodtkin, 2006), a comparison that highlights the confusing species- and context- specific connotations of the term 'sociality.' The labels 'social' and 'asocial' reference the overall social nature of a species but are, in fact, based upon single social phenotypes under specific environmental conditions, classifications applied to monogamous versus polygamous voles (McGraw & Young, 2010), to mouse strains expressing different levels of social approach (Moy et al., 2007), and to ground squirrels living in overlapping versus discontinuous territories. Such labels become confusing when considered in juxtaposition. Our finding of a robust expression of social reward by *I. tridecemlineatus* suggests that a comprehensive study of social CPP responses across ground squirrel species would improve our understanding of the putative relationship between social motivation and patterns of play and dispersal.

Field and laboratory studies also impose their own limitations to how we measure sociality. For instance, ultrasonic vocalizations are difficult to record in the field. By contrast, laboratory studies offer limited options for rodents to make complex social decisions and

confinement to a cage can alter some (Holmes & Mateo, 2007), but not all (Mateo & Holmes, 1999) social phenotypes. To integrate field and laboratory nomenclature for rodent social behavior (Blumstein et al., 2010; Cacioppo & Berntson, 2005), general terms like ‘social and ‘asocial’ might be replaced by ethological terms that describe social behaviors within specific experimental, environmental and social contexts.

This study provides the first demonstration of social CPP in an undomesticated rodent. When laboratory rodents prefer an environment paired with social housing, a common interpretation is that they derive a pleasurable experience from social interaction (and/or discomfort from social deprivation (Panksepp et al., 2007). Use of the CPP test has demonstrated that positive affective experiences occur during juvenile social interactions (Calcagnetti & Schechter, 1992; Douglas et al., 2004), mating opportunities (Camacho, Sandoval, & Paredes, 2004; Jenkins & Becker, 2003), access to offspring (Mattson, Williams, Rosenblatt, & Morrell, 2001) and even aggressive social interactions (Martinez, Guillen-Salazar, Salvador, & Simon, 1995; Tzschenke, 2007). Importantly, domestication removes some sources of natural selection faced by wild ancestors and can alter the social phenotypes of rodents (Nelson & Chiavegatto, 2001). Tame animals tend to be less aggressive toward conspecifics (Boreman & Price, 1972; Ebert, 1976) and more readily engage in mating opportunities without requiring mate choice (Drickamer, Gowaty, & Holmes, 2000; Manning, Potts, Wakeland, & Dewsbury, 1992; Penn & Potts, 1999). Thus, it is conceivable that responses of laboratory mice and rats to social CPP tests reflect the artificial selection and/or genetic fixation associated with domestication. Our present finding of social CPP in captive 13-lined ground squirrels argues against this, demonstrating social interaction is similarly rewarding to wild-derived animals with minimal domestication – for a species not thought to be particularly social in the wild.

The possibility that positive emotions accompany social interactions in the wild may be central to answering questions about the selective advantages of animal play (Allen & Bekoff, 2005) and altruistic social behavior (Hui & Deacon, 2009; Roberts, 2005). Social isolation impairs brain development (Black & Greenough, 1998; Champagne & Curley, 2005; Wiedenmayer, 2009), immune reactivity (Boissy et al., 2007; Shanks, Renton, Zalcman, & Anisman, 1994; Tuchscherer, Kanitz, Puppe, & Tuchscherer, 2010), burn and wound healing (Detillion, Craft, Glasper, Prendergast, & DeVries, 2004; eri et al., 2010), response to ischemia (Norman et al., 2010), recovery from social defeat (Ruis et al., 1999), resiliency to metastasis (Wu et al., 2000), and competence in social hierarchies (van den Berg et al., 1999). These studies suggest that psychological responses to social context can moderate systemic physiology (e.g. circulating corticosteroids and pro-inflammatory cytokines) and influence survivability. The experience of social reward may thereby serve as substrate for cooperative animal societies, as sufficient interdependence or a ‘stake’ among its individuals (Roberts, 2005), one that exists irrespective of the classical requirements for the evolution of altruism (i.e., repeated interaction, recognition, and ‘active’ reciprocity). Positive affective experiences associated with social reunion could be supportive mechanisms in this regard (Hui & Deacon, 2009) and our present study demonstrates that social reward is operational in an outbred, minimally domesticated rodent. Thus, our findings suggest that the psychological concept of social reward is deserving of ample consideration in interpreting the behavioral features of animal societies in the wild.

Acknowledgments

Support

This work was supported by NIH research grants DA022543 and P30HD03.

References

- Alamy M, Bengelloun WA. Malnutrition and brain development: An analysis of the effects of inadequate diet during different stages of life in rat. *Neuroscience & Biobehavioral Reviews*. 2012; 36(6):1463–1480. [PubMed: 22487135]
- Allen C, Bekoff M. Animal play and the evolution of morality: an ethological approach. *Topoi*. 2005; 24(2):125–135.
- Arenz CL, Leger DW. Thirteen-lined ground squirrel (Sciuridae: *Spermophilus tridecemlineatus*) antipredator vigilance. *Animal Behaviour*. 1999; 57(1):97. [PubMed: 10053076]
- Armitage KB. Sociality as a life-history tactic of ground squirrels. *Oecologia*. 1981; 48(1):36–49.
- Barash DP. The evolution of marmot societies: a general theory. *Science*. 1974; 185(4149):415–420. [PubMed: 17743072]
- Bardo MT, Bevins RA. Conditioned place preference: what does it add to our preclinical understanding of drug reward? *Psychopharmacology*. 2000; 153(1):31–43. [PubMed: 11255927]
- Beck JA, Lloyd S, Hafezparast M, Lennon-Pierce M, Eppig JT, Festing MF, et al. Genealogies of mouse inbred strains. *Nature Genetics*. 2000; 24(1):23–25. [PubMed: 10615122]
- Bekoff M. Wild justice and fair play: cooperation, forgiveness, and morality in animals. *Biology and Philosophy*. 2004; 19(4):489–520.
- Betts BJ. Behaviour in a population of Columbian ground squirrels, *Spermophilus columbianus columbianus*. *Animal Behaviour*. 1976; 24(3):652–680.
- Black, JE.; Greenough, WT. Developmental approaches to the memory process. In: Martinez, JL., Jr; Kesner, RP., editors. *Neurobiology of learning and memory*. San Diego: Academic Press; 1998. p. 55–88.
- Blumstein DT, Chung LK, Smith JE. Early play may predict later dominance relationships in yellow-bellied marmots (*Marmota flaviventris*). *Proceedings of the Royal Society B: Biological Sciences*. 2013; 280(1759):20130485. [PubMed: 23536602]
- Blumstein DT, Ebensperger LA, Hayes LD, Vásquez RA, Ahern TH, Burger JR, et al. Toward an integrative understanding of social behavior: new models and new opportunities. *Frontiers in behavioral neuroscience*. 2010; 4
- Boice R. Behavioral comparability of wild and domesticated rats. *Behavior Genetics*. 1981; 11(5): 545–553. [PubMed: 7325955]
- Boissy A, Manteuffel G, Jensen MB, Moe RO, Spruijt B, Keeling LJ, et al. Assessment of positive emotions in animals to improve their welfare. *Physiology & Behavior*. 2007; 92(3):375–397. [PubMed: 17428510]
- Boreman J, Price E. Social dominance in wild and domestic Norway rats (*Rattus norvegicus*). *Animal Behaviour*. 1972; 20(3):534–542.
- Bowler DE, Benton TG. Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biological Review*. 2005; 80:205–225.
- Burgdorf J, Panksepp J, Beinfeld MC, Kroes RA, Moskal JR. Regional brain cholecystokinin changes as a function of rough-and-tumble play behavior in adolescent rats. *Peptides*. 2006; 27(1):172–177. [PubMed: 16143427]
- Burghardt, GM. *The genesis of animal play: Testing the limits*. Cambridge, MA: MIT Press; 2005.
- Byrom AE, Krebs CJ. Natal dispersal of juvenile arctic ground squirrels in the boreal forest. *Canadian Journal of Zoology*. 1999; 77(7):1048–1059.
- Cacioppo, JT.; Berntson, GG. *Social neuroscience: Key readings*. New York, NY: Psychology Press; 2005.

- Calcagnetti DJ, Schechter MD. Place conditioning reveals the rewarding aspect of social interaction in juvenile rats. *Physiology & Behavior*. 1992; 51(4):667–672. [PubMed: 1594664]
- Camacho F, Sandoval C, Paredes R. Sexual experience and conditioned place preference in male rats. *Pharmacology, Biochemistry and Behavior*. 2004; 78(3):419–425.
- Chalfin L, Dayan M, Levy DR, Austad SN, Miller RA, Iraqi FA, et al. Mapping ecologically relevant social behaviours by gene knockout in wild mice. *Nature Communication*. 2014; 5
- Champagne FA, Curley JP. How social experiences influence the brain. *Current opinion in neurobiology*. 2005; 15(6):704–709. [PubMed: 16260130]
- Desha PG. Observations on the burrow utilization of the thirteen-lined ground squirrel. *The Southwestern Naturalist*. 1966:408–410.
- Detillion CE, Craft TKS, Glasper ER, Prendergast BJ, DeVries AC. Social facilitation of wound healing. *Psychoneuroendocrinology*. 2004; 29(8):1004–1011. [PubMed: 15219651]
- Dobson FS. An experimental study of dispersal in the California ground squirrel. *Ecology*. 1979:1103–1109.
- Dobson SF. Competition for mates and predominant juvenile male dispersal in mammals. *Animal Behaviour*. 1982; 30(4):1183–1192.
- Douglas LA, Varlinskaya EI, Spear LP. Rewarding properties of social interactions in adolescent and adult male and female rats: Impact of social versus isolate housing of subjects and partners. *Developmental Psychobiology*. 2004; 45(3):153–162. [PubMed: 15505797]
- Drickamer LC, Gowaty PA, Holmes CM. Free female mate choice in house mice affects reproductive success and offspring viability and performance. *Animal Behaviour*. 2000; 59:371–378. [PubMed: 10675259]
- Ebert PD. Agonistic behavior in wild and inbred *Mus musculus*. *Behavioral Biology*. 1976; 18(2):291–294. [PubMed: 1033758]
- Fairless AH, Dow HC, Toledo MM, Malkus KA, Edelman M, Li H, et al. Low sociability is associated with reduced size of the corpus callosum in the BALB/cJ inbred mouse strain. *Brain Research*. 2008:16.
- Glickman SE, Schiff BB. A biological theory of reinforcement. *Psychological Review*. 1967; 74(2): 81–109. [PubMed: 5342347]
- Gordon NS, Burke S, Akil H, Watson SJ, Panksepp J. Socially-induced brain ‘fertilization’: play promotes brain derived neurotrophic factor transcription in the amygdala and dorsolateral frontal cortex in juvenile rats. *Neuroscience letters*. 2003; 341(1):17–20. [PubMed: 12676333]
- Grubitz, G. The social behavior of the thirteen-lined ground squirrel (*Citellus tridecemlineatus*). University of Oklahoma; Norman, Oklahoma: 1963.
- Hare, JF.; Murie, JO. Ecology, Kinship, and Ground Squirrel Sociality: Insights from Comparative Analyses. In: Wolff, JO.; Sherman, PW., editors. *Rodent societies: an ecological and evolutionary perspective*. University of Chicago Press; 2008. p. 345–355.
- Holekamp, K. Dispersal in ground-dwelling sciurids. In: Murie, JO.; Michener, GR., editors. *Biology of ground-dwelling squirrels: annual cycles, behavioral ecology, and sociality*. Lincoln, Nebraska, USA: University of Nebraska Press; 1984. p. 297–320.
- Holekamp KE. Natal Dispersal in Belding’s Ground Squirrels (*Spermophilus beldingi*). *Behavioral Ecology and Sociobiology*. 1984; 16(1):21–30.
- Holmes WG, Mateo JM. Kin recognition in rodents: issues and evidence. *Rodent Societies: an Ecological and Evolutionary Perspective*. 2007:216–228.
- Hoogland JL. Aggression, ectoparasitism, and other possible costs of prairie dog (*sciuridae*, *cynomys* spp.) coloniality *Behaviour*. 1979; 69(1/2):1–35.
- Hui J, Deacon T. The evolution of altruism via social addiction. *Proceedings of the British Academy*. 2009; 158:181–203.
- eri SÖ, Dü ünceli F, Erzik C, Uslu B, Arbak S, Ye en BÇ. Oxytocin or social housing alleviates local burn injury in rats. *Journal of Surgical Research*. 2010; 162(1):122–131. [PubMed: 19628229]
- Jenkins WJ, Becker JB. Female rats develop conditioned place preferences for sex at their preferred interval. *Hormones and Behavior*. 2003; 43(4):503–507. [PubMed: 12788296]

- Kelley AE, Berridge KC. The neuroscience of natural rewards: relevance to addictive drugs. *The Journal of Neuroscience*. 2002; 22(9):3306–3311. [PubMed: 11978804]
- Kunzl C, Kaiser S, Meier E, Sachser N. Is a wild mammal kept and reared in captivity still a wild animal? *Hormones and Behavior*. 2003; 43(1):187–196. [PubMed: 12614649]
- Kunzl C, Sachser N. The behavioral endocrinology of domestication: A comparison between the domestic guinea pig (*Cavia aperea*, *porcellus*) and its wild ancestor, the cavy (*Cavia aperea*). *Hormones and Behavior*. 1999; 35(1):28–37. [PubMed: 10049600]
- Lacey EA, Wieczorek JR, Tucker PK. Male mating behaviour and patterns of sperm precedence in Arctic ground squirrels. *Animal Behaviour*. 1997; 53(4):767–779.
- Laviola G, Macri S, Morley-Fletcher S, Adriani W. Risk-taking behavior in adolescent mice: psychobiological determinants and early epigenetic influence. *Neuroscience & Biobehavioral Reviews*. 2003; 27(1–2):19–31. [PubMed: 12732220]
- Lawson Handley LJ, Perrin N. Advances in our understanding of mammalian sex-biased dispersal. *Molecular Ecology*. 2007; 16(8):1559–1578. [PubMed: 17402974]
- Liu Y, Wang ZX. Nucleus accumbens oxytocin and dopamine interact to regulate pair bond formation in female prairie voles. *Neuroscience*. 2003; 121(3):537–544. [PubMed: 14568015]
- Luna LD, Baird TA. Influence of density on the spatial behavior of female thirteen-lined ground squirrels, *Spermophilus tridecemlineatus*. *The Southwestern Naturalist*. 2004; 49(3):350–358.
- Macri S, Adriani W, Chiarotti F, Laviola G. Risk taking during exploration of a plus-maze is greater in adolescent than in juvenile or adult mice. *Animal Behaviour*. 2002; 64(4):541–546.
- Manning, CJ.; Potts, WK.; Wakeland, EK.; Dewsbury, DA. What's wrong with MHC mate choice experiments?. In: Doty, RL.; Muller-Schwarze, D., editors. *Chemical signals in vertebrates*. New York: Plenum; 1992. p. 229–235.
- Manno TG. Social networking in the Columbian ground squirrel, *Spermophilus columbianus*. *Animal Behaviour*. 2008; 75(4):1221–1228.
- Martinez M, Guillen-Salazar F, Salvador A, Simon VM. Successful intermale aggression and conditioned place preference in mice. *Physiology & Behavior*. 1995; 58(2):323–328. [PubMed: 7568436]
- Mateo JM, Holmes WG. How rearing history affects alarm-call responses of Belding's ground squirrels (*Spermophilus beldingi*, Sciuridae). *Ethology*. 1999; 105:207–222.
- Mateo JM, Holmes WG. Plasticity of alarm-call response development in Belding's ground squirrels (*Spermophilus beldingi*, Sciuridae). *Ethology*. 1999; 105:193–206.
- Mattson BJ, Williams S, Rosenblatt JS, Morrell JI. Comparison of two positive reinforcing stimuli: pups and cocaine throughout the postpartum period. *Behavioral Neuroscience*. 2001; 115(3):683–694. [PubMed: 11439457]
- McCarley H. Annual cycle, population dynamics and adaptive behavior of *Citellus tridecemlineatus*. *Journal of Mammalogy*. 1966; 47:294–316.
- McGraw LA, Young LJ. The prairie vole: an emerging model organism for understanding the social brain. *Trends in Neurosciences*. 2010; 33(2):103–109. [PubMed: 20005580]
- Meaney MJ, Stewart J. Neonatal-androgens influence the social play of prepubescent rats. *Hormones & Behavior*. 1981; 15(2):197–213. [PubMed: 7250910]
- Merriman DK, Lahvis GP, Jooss M, Gesicki JA, Schill K. Current practices in a captive breeding colony of 13-lined ground squirrels. *Lab animal*. 2012; 41(11):315–325. [PubMed: 23079915]
- Michener GR, Michener DR. Population structure and dispersal in Richardson's ground squirrels. *Ecology*. 1977:359–368.
- Michener GR, Sheppard DH. Social behavior between adult female Richardson's ground squirrels (*Spermophilus richardsonii*) and their own and alien young. *Canadian Journal of Zoology*. 1972; 50:1343–1349.
- Moy SS, Nadler JJ, Young NB, Perez A, Holloway LP, Barbaro RP, et al. Mouse behavioral tasks relevant to autism: Phenotypes of 10 inbred strains. *Behavioural Brain Research*. 2007; 176(1):4–20. [PubMed: 16971002]
- Nelson RJ, Chiavegatto S. Molecular basis of aggression. *Trends in Neurosciences*. 2001; 24(12):713–719. [PubMed: 11718876]

- Norman GJ, Zhang N, Morris JS, Karelina K, Berntson GG, DeVries AC. Social interaction modulates autonomic, inflammatory, and depressive-like responses to cardiac arrest and cardiopulmonary resuscitation. *Proceedings of the National Academy of Sciences*. 2010; 107(37):16342–16347.
- Nunes S, Muecke EM, Anthony JA, Batterbee AS. Endocrine and Energetic mediation of play behavior in free-living Belding's ground squirrels. *Hormones and Behavior*. 1999; 36(2):153–165. [PubMed: 10506539]
- Olesen KM, Jessen HM, Auger CJ, Auger AP. Dopaminergic activation of estrogen receptors in neonatal brain alters progesterin receptor expression and juvenile social play behavior. *Endocrinology*. 2005; 146(9):3705–3712. [PubMed: 15919740]
- Olioff M, Stewart J. Sex differences in the play behavior of prepubescent rats. *Physiology & Behavior*. 1978; 20(2):113–115. [PubMed: 662934]
- Panksepp JB, Jochman K, Kim JU, Koy JJ, Wilson ED, Chen Q, et al. Affiliative behavior, ultrasonic communication and social reward are influenced by genetic variation in adolescent mice. *PLoS ONE [Electronic Resource]*. 2007; 2:e351.10.1371/journal.pone.0000351
- Panksepp JB, Lahvis GP. Social reward among juvenile mice. *Genes, Brain, & Behavior*. 2007; 6(7): 661–671.
- Panksepp JB, Wong JC, Kennedy BC, Lahvis GP. Differential entrainment of a social rhythm in adolescent mice. *Behavioral Brain Reseach*. 2008; 195(2):239–245.
- Passtor TJ, Smith LK, MacDonald NK, Michener GR, Pellis SM. Sexual and aggressive play fighting of sibling Richardson's ground squirrels. *Aggressive Behavior*. 2001; 27:323–337.
- Pellis S. Sex differences in play fighting revisited: Traditional and nontraditional mechanisms of sexual differentiation in rats. *Archives of Sexual Behavior*. 2002; 31(1):17–26. [PubMed: 11910788]
- Pellis SM. Agonistic versus amicable targets of attack and defense: Consequences for the origin, function, and descriptive classification of play-fighting. *Aggressive Behavior*. 1988; 14(2):85–104.
- Pellis SM, MacDonald NL, Michener GR. Lateral display as a combat tactic in richardson's ground squirrel *Spermophilus richardsonii*. *Aggressive Behavior*. 1996; 22(2):119–134.
- Pellis SM, Pellis VC. Play-fighting differs from serious fighting in both target of attack and tactics of fighting in the laboratory rat *Rattus norvegicus*. *Aggressive Behavior*. 1987; 13(4):227–242.
- Pellis SM, Pellis VC. Play and the development of social engagement: A comparative perspective. *The Development of Social Engagement : Neurobiological Perspectives: Neurobiological Perspectives*. 2006:247.
- Pellis SM, Pellis VC. Rough-and-Tumble Play and the Development of the Social Brain. *Current Directions in Psychological Science*. 2007; 16(2):95–98.
- Pellis, SM.; Pellis, VC.; Reinhart, CJ. The evolution of social play. In: Wothman, C.; Plotsky, P.; Schecter, D.; CCA, editors. *Formative experiences: The interaction of caregiving, culture, and developmental psychobiology*. Cambridge University Press; 2010. p. 404-431.
- Penn DJ, Potts WK. The evolution of mating preferences and major histocompatibility complex genes. *The American Naturalist*. 1999; 153(2):145–164.
- Rayor LS, Armitage KB. Social behavior and space-use of young of ground-dwelling squirrel species with different levels of sociality. *Ethology Ecology & Evolution*. 1991; 3(3):185–205.
- Resendez SL, Kuhnmuench M, Krzywosinski T, Aragona BJ. kappa-Opioid receptors within the nucleus accumbens shell mediate pair bond maintenance. *The Journal of Neuroscience*. 2012; 32(20):6771–6784. [PubMed: 22593047]
- Ritchie ME, Belovsky GE. Sociality of Columbian ground squirrels in relation to their seasonal energy intake. *Oecologia*. 1990; 83:495–503.
- Roberts G. Cooperation through interdependence. *Animal Behavior*. 2005; 70:901–908.
- Rongstad OJ. A life history study of thirteen-lined ground squirrels in southern Wisconsin. *Journal of Mammalogy*. 1965; 46(1):76–87.
- Ruis M, Te Brake J, Buwalda B, De Boer S, Meerlo P, Korte S, et al. Housing familiar male wildtype rats together reduces the long-term adverse behavioural and physiological effects of social defeat. *Psychoneuroendocrinology*. 1999; 24(3):285–300. [PubMed: 10101734]

- Sankoorikal GM, Kaercher KA, Boon CJ, Lee JK, Brodtkin ES. A mouse model system for genetic analysis of sociability: C57BL/6J versus BALB/cJ inbred mouse strains. *Biological Psychiatry*. 2006; 59(5):415–423. [PubMed: 16199013]
- Schneirla, TC., editor. An evolutionary and developmental theory of biphasic processes underlying approach and withdrawal. University of Nebraska Press; 1959.
- Schochet TL, Kelley AE, Landry CF. Differential behavioral effects of nicotine exposure in adolescent and adult rats. *Psychopharmacology (Berl)*. 2004; 175(3):265–273. [PubMed: 15098085]
- Schochet TL, Kelley AE, Landry CF. Differential expression of arc mRNA and other plasticity-related genes induced by nicotine in adolescent rat forebrain. *Neuroscience*. 2005; 135(1):285–297. [PubMed: 16084664]
- Schwagmeyer P, Woontner S. Scramble competition polygyny in thirteen-lined ground squirrels: the relative contributions of overt conflict and competitive mate searching. *Behavioral Ecology and Sociobiology*. 1986; 19(5):359–364.
- Schwagmeyer PL. Alarm calling behavior of the thirteen-lined ground squirrel, *Spermophilus tridecemlineatus*. *Behavioral Ecology and Sociobiology*. 1980; 7(3):195–200.
- Schwagmeyer PL, Parker GA. Queuing for mates in thirteen-lined ground squirrels. *Animal Behaviour*. 1987; 35(4):1015–1025.
- Shanks N, Renton C, Zalcman S, Anisman H. Influence of change from grouped to individual housing on a T-cell-dependent immune response in mice: Antagonism by diazepam. *Pharmacology Biochemistry and Behavior*. 1994; 47(3):497–502.
- Sheppard DH, Yoshida SM. Social behavior in captive Richardson's ground squirrels. *Journal of Mammalogy*. 1971:793–799.
- Silk JB. The adaptive value of sociality in mammalian groups. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2007; 362(1480):539–559.
- Siviy SM, Fleischhauer AE, Kerrigan LA, Kuhlman SJ. D dopamine receptor involvement in the rough-and-tumble play behavior of juvenile rats. *Behavioral Neuroscience*. 1996; 110(5):1168. [PubMed: 8919019]
- Siviy SM, Panksepp J. Energy balance and play in juvenile rats. *Physiology & Behavior*. 1985; 35(3):435–441. [PubMed: 2999842]
- Spear LP. Neurobehavioral changes in adolescence. *Current Directions in Psychological Science*. 2000; 9(4):111–114.
- Spinka M, Newberry RC, Bekoff M. Mammalian play: training for the unexpected. *The Quarterly Review of Biology*. 2001; 76(2):143–168.
- Steiner AL. Play Activity of Columbian Ground Squirrels. *Zeitschrift fur Tierpsychologie*. 1971; 28(3):247–261.
- Steiner AL. Mortality Resulting from Intraspecific Fighting in Some Ground Squirrel Populations. *Journal of Mammalogy*. 1972; 53(3):601–603.
- Streubel DP, Fitzgerald JP. *Spermophilus tridecemlineatus*. *Mammalian Species*. 1978:1–5.
- Stryjek R, Pisula W. Warsaw wild captive Pisula Stryjek rats (WWCPS)-establishing a breeding colony of Norway Rat in captivity. *Polish Psychological Bulletin*. 2008; 39(2):67–70.
- Terranova ML, Laviola G, Alleva E. Ontogeny of amicable social behavior in the mouse: gender differences and ongoing isolation outcomes. *Developmental Psychobiology*. 1993; 26(8):467–481. [PubMed: 8293893]
- Trezza V, Vanderschuren LJMJ. Cannabinoid and opioid modulation of social play behavior in adolescent rats: Differential behavioral mechanisms. *European Neuropsychopharmacology*. 2008; 18(7):519–530. [PubMed: 18434104]
- Tuchscherer M, Kanitz E, Puppe B, Tuchscherer A. Altered immunomodulation by glucocorticoids in neonatal pigs exposed to a psychosocial stressor. *Pediatric Research*. 2010; 68(6):473–478. [PubMed: 20717071]
- Tzschentke TM. Measuring reward with the conditioned place preference (CPP) paradigm: Update of the last decade. *Addiction Biology*. 2007; 12(3–4):227–462. [PubMed: 17678505]

- van den Berg CL, Hol T, Van Ree JM, Spruijt BM, Everts H, Koolhaas JM. Play is indispensable for an adequate development of coping with social challenges in the rat. *Developmental Psychobiology*. 1999; 34(2):129–138. [PubMed: 10086231]
- Vaughan DK, Gruber AR, Michalski ML, Seidling J, Schlink S. Capture, care, and captive breeding of 13-lined ground squirrels, *Spermophilus tridecemlineatus*. *Lab Animal*. 2006b; 35(4):33–40. [PubMed: 16582898]
- Vucetic Z, Totoki K, Schoch H, Whitaker KW, Hill-Smith T, Lucki I, et al. Early life protein restriction alters dopamine circuitry. *Neuroscience*. 2010; 168(2):359–370. [PubMed: 20394806]
- Wade CM, Daly MJ. Genetic variation in laboratory mice. *Nature Genetics*. 2005; 37(11):1175–1180. [PubMed: 16254563]
- Wauters L, Dhondt A. Immigration pattern and success in red squirrels. *Behavioral Ecology and Sociobiology*. 1993; 33(3):159–167.
- Wiedenmayer CP. Plasticity of defensive behavior and fear in early development. *Neuroscience & Biobehavioral Reviews*. 2009; 33(3):432–441. [PubMed: 19073211]
- Wistrand H. Individual, social, and seasonal behavior of the thirteen-lined ground squirrel (*Spermophilus tridecemlineatus*). *Journal of Mammalogy*. 1974:329–347.
- Wolff JO. What is the role of adults in mammalian juvenile dispersal? *Oikos*. 1993:173–176.
- Wolff JO. More on juvenile dispersal in mammals. *Oikos*. 1994:349–352.
- Wolff, JO.; Sherman, PW., editors. *Rodent societies: an ecological and evolutionary perspective*. University of Chicago Press; 2008.
- Wu W, Yamaura T, Murakami K, Murata J, Matsumoto K, Watanabe H, et al. Social isolation stress enhanced liver metastasis of murine colon 26-L5 carcinoma cells by suppressing immune responses in mice. *Life Sciences*. 2000; 66(19):1827–1838. [PubMed: 10809180]
- Yeaton RI. Social Behavior and Social Organization in Richardson's Ground Squirrel (*Spermophilus richardsonii*) in Saskatchewan. *Journal of Mammalogy*. 1972; 53(1):139–147.

Figure 1. Developmental timeline of experimental procedures and illustrated representation of social conditioned place preference

The top of the figure depicts the ages when post-weanling squirrels were tested for SI and conditioned and tested for social CPP, respectively. Grouped squirrels on the left side, in the bedding represented by gray dots, are undergoing social conditioning in a ‘corncob’ environment with tubes threaded within the interior. The isolated squirrel on the right side, in the bedding represented by diagonal lines, is undergoing conditioning alone in a ‘paper’ environment with tubes containing a smooth interior. The social CPP testing arena is illustrated at the bottom.

Figure 2. Behavioral differences between early and late adolescent ground squirrels in the laboratory

Labels on the abscissa indicate the age at testing and the sex of the interacting squirrels (test-stimulus combinations). (A) Play behavior of test squirrels diminished across adolescent development from PD38 to PD56 and (A') was most highly expressed in male-male pairs. (B) Social investigation was greater at PD38 versus PD56 and (B') was expressed more within female-female pairs. (C) Vigilance behavior was higher on PD56 versus PD38 and (C') did not differ between the sex-pairings of squirrels. (D) 'Time budgets' for squirrels during the 5-min test period demonstrating that a majority of time was spent engaged in behaviors that were not social in nature. N's = 4–7 squirrels per sex combination. All data are presented as the mean $\pm$ standard error. (* $P < 0.05$ ** $P < 0.001$).

Figure 3. Social conditioned place preference in adolescent ground squirrels

Preference scores were generated by subtracting the amount of time each squirrel spent in the ‘paper’ environment from the amount of time in the ‘corncob’ environment. A positive preference score therefore indicates an individual spent more time in the corncob environment during the 30-min trial whereas a negative score indicates more time spent in the paper environment. N's = 6–18 squirrels per group. All data are presented as the mean $\pm$ standard error. (* $P < 0.05$ ** $P < 0.001$).

Figure 4. Social behaviors of adolescent ground squirrels in the field

Expected weekly counts of social behaviors were generated assuming an equal distribution of the total social behaviors ($n=228$) over the 5-week observation period from late June and early August. (A) The number of social behaviors observed each week deviated from the expected values and generally decreased over the 5-week period. Social behaviors were categorized as approach, play, investigation, maintaining proximity or chasing (see Table 2), and total counts were tallied for each week. Observed behaviors in the field varied significantly from the expected values. (B) Approach, (C) play, (D) investigation and (E) proximity were predominantly observed during weeks 1–3 while (F) agonistic chasing became more frequent in weeks 3–5. Total minutes of observation: week 1 = 291 min, week 2 = 357 min, week 3 = 551 min, week 4 = 298 min, week 5 = 321 min. All data are presented as frequencies of occurrence.

Table 1

Ethogram of social interaction test in the laboratory.

Category	Behavior	Description
Social investigation	grasping	placing both forepaws on back of other juvenile without lower body 'mounting'
	allo-grooming	grooming or licking any area of other juvenile
	sniffing and pursuit	sniffing or snout contact of head, flank or anogenital areas, pursuit within one body-length
Social play	wrestling	facing other juvenile, often in a face-to-face clench or embrace, pecking at juvenile's neck, throat, cheeks, chest or abdomen, without inflicting bite wounds
	boxing	batting other juvenile with forepaws, usually reciprocated
	mounting	climbing on other juvenile's back, wrapping forepaws around chest or abdomen, then attempting intromission
Vigilance	stand/freeze	rearing onto hind feet and maintaining upright posture

Table 2

Ethogram of social behaviors in the field.

Category	Behavior	Description
Affiliative	approach	approaching a stationary squirrel or following a moving squirrel without threat posture characteristic of chase behaviors
	social investigation	See 'Social investigation' category in Table 1
	play	two squirrels engaging in any non-snout directed physical contact, including mounting, boxing and wrestling
	proximity	within two body-lengths of another squirrel but not engaging in other social behavior
Agonistic	chasing	aggressive pursuit of another squirrel, typically preceded or followed by a threatening posture involving an upright stance and tail flicking