Published in final edited form as:

Alcohol Clin Exp Res. 2015 November ; 39(11): 2085–2094. doi:10.1111/acer.12887.

Clinical Impact of Alcohol Related Cirrhosis in the Next Decade: Estimates Based on Current Epidemiological Trends in the United States

John Guirguis, MD¹, Jagpreet Chhatwal, PhD³, Jaividhya Dasarathy, MD², John Rivas, MD¹, David McMichael, MS, Laura E. Nagy, Ph.D⁴, Arthur J McCullough, MD¹, and Srinivasan Dasarathy, MD¹

¹Departments of Gastroenterology, Hepatology and Transplant Surgery, Cleveland Clinic, Cleveland OHIO.

²Department of Family Medicine, Metro Health Medical Center, Cleveland, OHIO.

³Department of Health Services Research, Massachusetts General Hospital, Harvard Medical School, Boston.

⁴Department of Pathobiology Lerner Research Institute The Cleveland Clinic Foundation.

Abstract

Background—Identifying changes in the epidemiology of liver disease is critical for establishing health-care priorities and allocating resources to develop therapies. The projected contribution of different etiologies towards development of cirrhosis in the United States was estimated based on current publications on epidemiological data and advances in therapy. Given the heterogeneity of published reports and the different perceptions, that are not always reconcilable, a critical overview rather than a formal meta-analysis of the existing data and projections for the next decade were performed.

Methods—Data from the World Health Organization (WHO) Global Status Report on Alcohol and Health of 2014, Scientific Registry of Transplant Recipients (SRTR) from 1999 to 2012, National Institute on Alcohol Abuse and Alcoholism (NIAAA) and the Centers for Disease Control and Prevention were inquired to determine future changes in the epidemiology of liver disease.

Results—Alcohol consumption has increased over the past 60 years. In 2010, transplant related costs for liver recipients were the highest for hepatitis C (~\$124 million) followed by alcohol related cirrhosis (~\$86 million). We anticipate a significant reduction in incidence cirrhosis due to causes other than alcohol because of the availability of high efficiency antiviral agents for hepatitis C, universal and effective vaccination for hepatitis B, relative stabilization of the obesity trends in the United States and novel, potentially effective therapies for non-alcoholic

---

Address for correspondence. Srinivasan Dasarathy MD, NE4 208, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44195, dasaras@ccf.org, Tel 2164442980, Fax 216445388.

Conflict of interest: Authors declared there is no financial conflict of interest.

steatohepatitis (NASH). The proportion of alcohol related liver disease (ALD) is therefore likely to increase in both the population as a whole and in the liver transplant wait-list.

Conclusions—

Keywords

Alcohol; liver disease; cirrhosis; hepatitis C; health care costs

INTRODUCTION

Cirrhosis is the twelfth leading cause of death in the United States, and is a consequence of a number of etiologies that result in necroinflammation and fibrosis of the liver (Hoyert DL, 2012). Cirrhosis in 2011 resulted in about 34,869 deaths in the United States of which alcohol related cirrhosis accounted to over half of these deaths (Chen C. M, 2014d). Hepatitis C virus (HCV) and alcohol related cirrhosis are the most common causes of chronic liver disease in the United States (Mueller et al., 2009, Scaglione et al., 2014). Hepatitis B leads to chronic infection in about 5%-10 % of infected people and only about 2.1%-6% of patients with hepatitis B infection will develop cirrhosis (Chu and Liaw, 2006). Non alcoholic fatty liver disease is now recognized to the most common form of chronic liver disease with a population prevalence of 20-25%. The population prevalence of non alcoholic steatohepatitis, the more severe form of NAFLD, is estimated to be 3-5% and between 15-20% of patients with NASH progress to cirrhosis. Given the large number of patients affected with NAFLD and NASH, the incidence and prevalence of cirrhosis related to fatty liver are expected to increase if no effective therapies are developed (Fargion et al., 2014).

Several recent changes will affect the epidemiology of alcohol related cirrhosis in the United States, and potentially the rest of the world. The availability of highly effective antiviral therapies of HCV and updates in the screening policy by the Centers for Disease Control and Prevention (CDC) are projected to make hepatitis C a rare disease by 2036 in the United States (Kabiri et al., 2014). Programs for universal vaccination have decreased the prevalence of hepatitis B virus (HBV) in the United States (Goyal and Murray, 2014). In addition, the relative stabilization of obesity epidemic could eventually affect the prevalence of non-alcoholic steatohepatitis (NASH) (Ogden et al., 2014).

In contrast, the number of patients with alcohol related cirrhosis has remained relatively unchanged in last 12 years, between 60,000 and 65,000 patients (Chen C. M, 2014b). Despite the fact that only 10%–16% of the alcohol drinkers develop cirrhosis (Robert E. Mann and Richard, 2004), recent trends in the epidemiology of liver disease are likely to result in alcohol to be the most common etiology of cirrhosis in the United States. There have been no advances in therapies that have had a clinically significant impact on the prevalence of alcohol use disorders.

The focus of policy makers towards allocation of resources is based on a number of factors, including the prevalence, complications related to cirrhosis, availability of effective therapies to combat these complications and the potential economic impact of the disease

(Brand et al., 2004). Despite recognizing that alcohol abuse is a major cause of liver disease, most epidemiological studies have focused on the prevalence of viral hepatitis (El-Serag, 2012) and non-alcohol related fatty liver disease (NAFLD) (Zezos and Renner, 2014) and not on alcohol related cirrhosis.

We explored trends in the etiology and epidemiology of liver disease and focused specifically on alcohol related cirrhosis because the prevalence of liver disease not due to alcohol is expected to decrease due to a variety of factors as mentioned earlier. We therefore hypothesized that the prevalence and consequently costs of alcohol related cirrhosis would be higher in the next decade compared to that of non-alcohol related cases of cirrhosis. Even though the spectrum of alcohol related liver disease extends from steatosis to cirrhosis, we have elected to examine the epidemiology of alcohol related cirrhosis because the long-term health care impact of alcohol use disorders is related primarily to cirrhosis. We examined our hypothesis by analyzing published data and public domain databases including the scientific registry of transplant recipients, National Institute on Alcohol Abuse and Alcoholism, Medicare costs, and reports by the CDC and the World Health Organization (WHO).

MATERIAL AND METHODS

We searched the following databases: PubMed, Ovid, and Google Scholar. The search terms included alcohol, liver disease, cirrhosis, epidemiology, prevalence, incidence, and costs. Published data from the National Institute of Alcohol Abuse and Alcoholism, WHO, CDC, Scientific Registry of Transplant Recipients were also evaluated. Following the search, only publications from 2000-2015 that specially addressed the prevalence of cirrhosis were included in the analysis because these were believed to permit the best assessment of current trends in epidemiology of liver disease and allow us to estimate the projections for the next decade. Details of results obtained from the search results are described for each specific database or publication.

Published studies

NIAAA Surveillance reports—Data from National Institutes of Health (NIAAA) surveillance reports # 99 and # 100 published in December 2014 were analyzed to determine the extent of alcohol related disorders from the number of discharges of patients with a primary diagnosis of alcohol related disorder from 2000 to 2012, mortality rates in alcohol related cirrhosis compared to other causes of cirrhosis and the percentage of alcohol related discharges in the cirrhotic population were also obtained from these reports (Chen C. M, 2014d, Chen C. M, 2014b). We also determined the median duration of stay per hospitalization, the cost per hospitalization to examine if the costs of managing patients with alcoholic liver disease was different from that of other causes of cirrhosis (Chen C. M, 2014b, Chen C. M, 2014a). The NIAAA surveillance report # 98 was used to determine the per capita alcohol consumption in gallons per year in the United States from 1950 to 2012 (Chen C. M, 2014d).

WHO global status report on alcohol and health—The per capita alcohol consumption calculated in liters per year from the WHO global status report on alcohol and

health was converted to gallons per year in order to match the data from the NIAAA surveillance report # 98. The NIAAA surveillance report #98 and WHO global status report in alcohol and health were then used to examine the current consumption and projections were made using a best fit (least squares) linear regression to predict future per capita alcohol consumption in the United States (Montgomery and Vining, 2012).

SRTR report—The details of patients with cirrhosis due to different etiologies who are listed for liver transplantation were obtained from the 2014 SRTR report (OPTN/SRTR 2012 Annual Data Report. Rockville, 2014). The number of patients who underwent transplantation, waiting time for transplantation and death on the wait list and after transplantation were analyzed to project future trends for alcoholic cirrhosis. The SRTR data was however not used to estimate the actual prevalence of alcohol related liver disease because only 5% or less of patients with alcoholic cirrhosis are considered for liver transplantation (O'Grady, 2006). This estimate is supported by our analysis of the data from UNOS and the NIAAA surveillance report #100. The number of deaths in the United States from alcoholic cirrhosis in 2011 was ~16,500 and the number with alcoholic cirrhosis being transplanted in 2011 was 1004. The proportion of transplanted to the number of alcoholic cirrhotic patients needing transplant (transplanted + death from alcoholic cirrhosis) during 2011 showed that only 5% of patients with alcoholic cirrhosis were considered for transplant (OPTN/SRTR 2012 Annual Data Report. Rockville, 2014, Chen C. M, 2014c)

Published data review about recent changes in HCV epidemiology—The current prevalence of HCV was determined from a recent NIH-sponsored study that projected the changing prevalence of HCV in the United States, considering updates in the CDC screening policy and availability of highly-effective direct-acting antiviral agents (Kabiri et al., 2014).

CDC Behavioral risk factor Surveillance report—Obesity data was extracted from annual CDC Behavioral Risk Factor surveillance reports for the last 13 years (Centers for Disease Control and Prevention, 2015).

Analysis—This study seeks to extrapolate future trends in the burden of alcohol related liver disease over the next decade in the United States. Projections are based on a number of published reports that take different, and often not reconcilable, perspectives. Since the primary focus of the present studies was to estimate the future clinical impact of alcohol related liver disease, a formal meta-analysis of the published data could not be performed. A descriptive analysis of high quality data with reproducible and reliable outcomes was conducted followed by projections on the trends in the epidemiology of cirrhosis in the next decade.

We presented data on historic and future trends. Trends in alcohol related discharges, per capita consumption, alcohol related death, transplant and listed numbers in alcoholic population were computed based on historic data from published SRTR and NIAAA data. We used the best-fit (least squares) linear regression on the last 12 years data to project current year's estimate using FORECAST function of Microsoft Excel. For example, the projections in year 2013 were based on the best-fit regression on data from years 2001-2012, and the projections for year 2014 were based on the data from years 2002-2013.

RESULTS

The future of alcohol related liver disease should be discussed in the context of liver disease due to other etiologies. Analysis of the anticipated changes in the epidemiology of both HCV and NAFLD disease, two other major causes of liver disease currently is necessary to estimate the future epidemiology of liver disease in the United States.

The overlooked risk of alcohol

Analysis of hospital discharge rates showed that ~50% of all discharges related to cirrhosis are linked to the diagnosis of alcohol related cirrhosis as shown in **Figure 1**. A small but sustained increase in the number of patients discharged from US hospitals between 2000 and 2012 with diagnoses related to alcohol was observed as shown in **Figure 2a**. These numbers are anticipated to increase in the next 10 years (**Figure 2b**).

The mortality due to cirrhosis based on different etiologies shows that crude death rate from cirrhosis is about 11.2 deaths per 100,000, of which 5.4 deaths were related to alcohol related cirrhosis constituting more than 45% of the total mortality of cirrhosis (**Figure 3**). It was also observed that from 2007 to 2011, cirrhosis related mortality rate was highest in patients with alcohol related cirrhosis. Projections from these numbers show that mortality related to alcohol related cirrhosis and ALD will continue to increase in the next 10 years if efforts are not undertaken to halt the prevalence and progression (**Figure 3b**).

The annual cost of admissions for alcohol related cirrhosis was ~\$848 million while that of other specified causes of cirrhosis was ~\$200 million (Chen C. M, 2014b). Data from the same report showed that the length of stay was ~5.9 days for each group, the median cost per hospital stay was ~8,000 for each group. If the length of stay and the cost of each hospitalization were similar, then these data would suggest that the readmission rate for alcoholic cirrhotic population is nearly 4 times higher than that of other causes of cirrhosis based on the total annual cost. This is consistent with a previous report that suggested significantly higher readmission rates in alcohol related cirrhosis population (Ganesh et al., 2013).

The per capita alcohol consumption in the United States from 1950 to 2012 is shown in **Figure 4**. The WHO global status report on alcohol and health of 2014 (Organization, 2014) and NIAAA surveillance report #98 (Chen C. M, 2014a) showed that the rate of alcohol consumption in the US steadily increased from 1992 to 2012. Projections made by the WHO suggest that population consumption of alcohol will increase in the next decade before it may stabilize, but is not likely to decrease. Our estimates based on the data of the past decade and even those prior to 2000 suggest that the consumption of alcohol will increase in the next decade with consequent adverse impact on liver disease.

Hepatitis B and C

Hepatitis B was a major cause of cirrhosis in the United States when a universal effective vaccination policy was introduced in 1991 (Wasley et al., 2010) that significantly decreased the prevalence of HBV. Specifically, among children (6-19 years), the prevalence decreased from 0.24% during the period 1988-1994 to 0.05% during the period 1999-2006 (Wasley et

al., 2010) and the incidence of HBV declined from 3 in 100,000 to 0.19 per 100,000 population in the period 1990–2004 (Shepard et al., 2006). Currently, HCV is the major cause of cirrhosis and most frequent indication for liver transplantation in the United States (Tsoulfas et al., 2009, Scaglione et al., 2014). In 2012, the CDC and U.S. Preventive Services Task Force updated the hepatitis C screening guidelines to include a one-time screening of people born between 1945 and 1965 (Moyer, 2013). In 2014, new oral direct-acting antiviral agents become available that are highly effective in patients with chronic HCV (Beinhardt et al., 2015). The burden of HCV disease has therefore been estimated to decrease substantially in the next decade (**Figure 5a**), and HCV is projected to become a rare disease in the United States by 2036 (Kabiri et al., 2014). Lower HCV burden is predicted to result in a reduction in the incidence of HCV-associated decompensated cirrhosis and hepatocellular carcinoma by 26% and 17%, respectively, in the next 10 years (**Figure 5b**) (Kabiri et al., 2014).

Liver Transplantation

HCV and alcohol related liver diseases are currently the major indications for liver transplantation (Tsoulfas et al., 2009, Scaglione et al., 2014). Medicare cost for liver transplantation in 2010 for HCV and alcohol related cirrhosis was reported to be 124.2 million and 83.2 million dollars respectively (OPTN/SRTR 2012 Annual Data Report. Rockville, 2014). With the availability of the highly effective new antiviral medications for HCV, the number of patients being transplanted for HCV will decrease based on the modeling study on the changing burden of hepatitis C (Kabiri et al., 2014). The authors estimated that the number of liver transplants related to HCV will peak in year 2016 and decrease by 18% and 70% in the next 10 and 20 years, respectively (**Figure 5C**).

Despite an increase in the number of patients with alcohol related cirrhosis being listed for liver transplantation, the number of patients with alcohol related cirrhosis that are transplanted is decreasing (OPTN/SRTR 2012 Annual Data Report. Rockville, 2014). Continued increase in numbers of non-transplanted alcohol related cirrhosis, as predicted from our projections based on current data are shown in **Figure 6**, will result in an increase in health care costs related to the management of complications related to alcohol related cirrhosis.

The number of patients stratified by diagnosis who are listed for liver transplantation is shown in **Figure 7a** (OPTN/SRTR 2012 Annual Data Report. Rockville, 2014). The most rapid increase in patients listed for as well as those who underwent liver transplantation over the past 10 years is due to HCC while the number of patients being listed or transplanted with diagnoses other than HCC is relatively unchanged. We calculated the ratio of the number of patients who underwent transplantation to those who were listed, to compare the proportion of listed patients who were transplanted. We found that the *listed to transplant ratio* in patients with alcohol related cirrhosis was relatively stable over 5 years although in the last 2 years it started to increase and this can be attributed to the widening gap between the patients listed to those transplanted as shown in **Figure 7b**.

Non-alcohol related fatty liver disease and obesity

The current trends in the prevalence of obesity are shown in **Figure 8**. Consistent with the obesity epidemic, NASH-related cirrhosis is projected to be the major indication for liver transplantation by the end of this decade (Zezos and Renner, 2014). However, recent epidemiological data show that the rate of increase in prevalence of obesity in the United States is decreasing as observed from the change in the slope of the prevalence (**Figure 8**). In addition, although the number of patients with NAFLD has tripled since 2004, it is believed that patients with NAFLD are less likely to undergo liver transplantation and are less likely survive on the transplant waiting list than patients with alcohol related liver disease (Wong et al., 2015). This may be related to cardiovascular and cerebrovascular disease being the major causes of death in NAFLD patients (Rafiq et al., 2009). Furthermore, a number of effective therapies for the management of NASH are being evaluated and may contribute to a reduction in prevalence of NASH (Neuschwander-Tetri et al., 2015). In contrast to NAFLD, the major cause of death in alcohol abuse disorders and alcohol related liver disease is related to liver disease and its complications (Domanski et al., 2012, Morgan, 1994). The focus of therapies in NAFLD therefore is on the management of the metabolic components and systemic disorders; while in alcohol use disorders, the primary goal is to prevent and manage the consequences of liver disease.

DISCUSSION

The present analysis is the first of its kind to critically evaluate multiple large data sources and examine the potential clinical significance of alcohol related liver disease relative to other causes of liver injury and cirrhosis. Our analysis showed that alcohol abuse and related liver injury are of high significance in the United States. However, a recent analysis showed that ALD is the most prevalent cause of liver disease in Europe also (Mathurin and Bataller, 2015). We have focused on the current patterns and predicted the future trends in the epidemiology of alcohol related liver disease in the United States because novel and effective therapies for other forms of liver disease are currently available or are likely to become available for use in clinical practice while there are no effective therapies to treat alcohol use disorders. We demonstrate that alcohol related liver disease is not only one of the most common causes of liver disease at present (Iruzubieta et al., 2013) but is also likely to be the most prevalent liver disease in the next decade.

Alcohol abuse has been a significant clinical problem in the United States and a number of medical problems are linked to alcohol abuse like dependence, psychosis and cirrhosis (Frazier et al., 2011). Over the last 15 years, alcohol consumption (measured as per capita consumption) has increased that is also reflected in the numbers of patients with alcohol related disorders (Chen C. M, 2014a). We identified three relatively recent reviews, which focused on the risk factors of alcohol related liver disease, overall trends in mortality, drinking patterns, and outcomes of liver transplantation (Basra and Anand, 2011, Robert E. Mann and Richard, 2004, Bhupinderjit, 2013). These reviews were published before the recent publications in the epidemiology of cirrhosis, availability of highly effective therapies for HCV and did not address future trends in prevalence of cirrhosis due to alcohol and other causes. Previous publications on the epidemiology of alcohol also did not take into

consideration the rapid advances in the management of non-alcohol related liver disease because the highly effective novel therapies for HCV and potential therapies for NASH were not available at that time. A critical analysis of the SRTR data also reiterated that alcohol will be the major cause of cirrhosis in the United States. Our interpretation however, needs to be tempered by the fact that despite data in the SRTR being robust, transplant related data do not necessarily reflect the prevalence of alcohol related liver disease accurately because patients with ALD do not get listed or transplanted in the same proportion as those with non-alcohol related liver disease (OPTN/SRTR 2012 Annual Data Report. Rockville, 2014). Therefore, SRTR based data are likely to underestimate the actual prevalence of alcohol related liver disease.

The present study includes a critical analysis of epidemiological trends and health care costs as well as predictions based on modeling for the prevalence of HCV, HBV and obesity in the next decades (Beste and Ioannou, 2015, Goyal and Murray, 2014). The new highly effective anti-HCV therapies are orally administered, have low rates of adverse effects and are effective in patients who have failed previous treatments (Moreno et al., 2015) in contrast to the last 3 decades of interferon based therapies. However, long-term viral clearance, development of resistance and possible decline in efficacy with progressive disease are considerations that will become known over time. Finally, progression to cirrhosis can only be determined in large populations who are followed over extended periods of time. The current high cost of these medications (Petta and Craxi, 2015) is also a factor that may limit use of these agents and affect the current models of the epidemiology of HCV that predict a rapid decline in the proportion of HCV cirrhosis by 2036 (Kabiri et al., 2014). Counterarguments, however, include the consistent reduction in cost of any novel therapy, the potential for even more effective therapies directed against HCV - and possibly HBV - that may decrease the prevalence of HCV even more rapidly than has been currently estimated (Hatzakis et al., 2015). On balance, one may argue that if the current trends in response to antiviral agents continue, HCV is likely to be of much lower clinical significance than it was in the previous decade.

Similarly, the effect of nearly universal HBV vaccination globally will decrease the existing pool of HBV carriers combined with the high efficacy of therapy for HBV acquired in adulthood (McMahon and Brian, 2014), is expected to reduce the prevalence of HBV rapidly. Finally, since HBV and HCV are major causes of hepatocellular carcinoma (HCC) (Sarvari et al., 2013), the most rapidly increasing indication for liver transplantation, a reduction in prevalence of the underlying diseases will result in a lower number of patients due to viral hepatitis and HCC who are either listed or undergo liver transplantation.

We have used data from multicenter, double blind, randomized controlled treatment trials for NASH, as well as, a recent data that the epidemic of obesity in the United States is stabilizing (Yanovski and Yanovski, 2011, Centers for Disease Control and Prevention, 2015), to predict that the clinical consequences of NASH will decrease over time. Compared to the relatively short period of time that effective treatments for NASH have been developed, the success rate of treatments for alcohol abuse and acute alcohol related hepatitis has been very low (Iruzubieta et al., 2013, Litten et al., 2012). Furthermore, since the major cause of death in patients with NAFLD is related to cardiovascular and

cerebrovascular disease (Neuschwander-Tetri et al., 2015), it is possible that even if NASH is a significant cause of cirrhosis, the major focus will continue to be the management of components of the metabolic syndrome.

Changes in the prevalence and outcomes in non-alcoholic liver disease will by necessity, increase the relative prevalence of alcohol related liver disease (ALD) because attempts to reduce alcohol consumption using anti addiction medications, nonpharmacological therapies, and attempts to develop molecular targeted therapies to prevent and reverse ALD have been unsuccessful to date especially when compared with the highly effective therapies for other liver diseases (Beste and Ioannou, 2015, Goyal and Murray, 2014). Health care costs related to alcohol are also predicted to increase over the next decades. Published data demonstrate that currently HCV has the highest cost of management (Basu, 2015). However, the economic consequences of ALD will be significantly higher in the next decade based on the predicted increase in per capita alcohol consumption (**Figure 4**), relatively low transplant/listing ratio resulting in a larger number patients with alcoholic liver disease, and more severe malnutrition and sarcopenia that adversely affect outcomes in cirrhosis (Dasarathy, 2014).

A formal statistical modeling, meta-analysis or forecasting were not performed because of the heterogeneity of the studies, the effect of time on behavioral patterns, and the very recent advances in therapeutics that have not been used for sufficiently long period of time to be allow precise and confident projections. We, however, made several simplifying assumptions about the future projections. We assumed that the trend in disease burden is linear; however, in practice such relationship could be nonlinear. For further and better future analysis, we need a comprehensive simulation model that takes into account multiple factors such as patient demographics, changing alcohol consumption patterns, and natural history of disease. Finally, the validity of the underlying data is the major factor that determines our analysis and interpretations. By selecting publications from agencies tasked with collecting the data used in this study we have used the best available data for the present study.

The projections made in this paper were based on the preceding 12 years of data. Since we are dealing a system that is affected by a number of biological, psychological, social and economic factors, the overall trend is not expected to be linear and some cases the projections increased at a higher rate. For instance, the projection of year 2015 is based on data from 2003–2014, which has a very different trend than from 2000–2003 (**Figure 2b**, **Figure 3b** and **Figure 4**). We also did not compare the cost benefit of liver transplantation for alcohol related liver disease with the costs of readmission to the hospital and non-transplant management therapies. The cost of liver transplantation could potentially be counterbalanced by the economic benefit and the reduction in health care costs after successful transplantation. Studying this is beyond the scope of this paper and will need further analysis.

Despite these limitations, the major strengths of the study are the consistent interpretation from a number of very disparate data sources and being the first study to have made projections regarding liver disease in the coming decades. We conclude that alcohol related

cirrhosis and ALD will be a major clinical challenge for not only hepatologists and transplant surgeons but for the health care system given the high costs of managing these patients. We based our conclusions on current trends that show a progressive increase in alcohol consumption, costs of management of alcohol related liver disease, relatively low transplant/listing ratio, and lack of targeted and effective therapies in combination with the rapid availability and/or the clinical success of novel and highly effective therapies for other causes of cirrhosis. To avoid the health care crises related to alcohol related liver disease, there is an urgent need to identify the molecular mechanisms of not only alcohol related liver disease but also its attendant complications that worsen clinical outcomes.

Acknowledgments

Partial support for this work was provided by:

NIDDK ROI DK 083414 (SD)

NIAAA UOI AA 021890 (SD, AJM)

NIAAA R21 AA022742 (SD)

NIAAA UOI AA 021890 (LEN)

REFERENCES

- Basra S, Anand BS. Definition, epidemiology and magnitude of alcoholic hepatitis. *World journal of hepatology*. 2011; 3:108–113. [PubMed: 21731902]
- Basu A. Financing cures in the United States. Expert review of pharmacoeconomics & outcomes research. 2015; 15:1–4. [PubMed: 25482146]
- Beinhardt S, Peck-Radosavljevic M, Hofer H, Ferenci P. Interferon-free antiviral treatment of chronic hepatitis C in the transplant setting. *Transpl Int*. 2015
- Beste LA, Ioannou GN. Prevalence and Treatment of Chronic Hepatitis C Virus Infection in the US Department of Veterans Affairs. *Epidemiologic reviews*. 2015
- Bhupinderjit, AKSa. Recent Trends in the Epidemiology of Alcoholic Liver Disease. *Clinical Liver Disease*. 2013; 2:53–56.
- Brand DA, Viola D, Rampersaud P, Patrick PA, Rosenthal WS, Wolf DC. Waiting for a liver--hidden costs of the organ shortage. *Liver Transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2004; 10:1001–1010. [PubMed: 15390326]
- Centers for Disease Control and Prevention NCfCDPaHP, Division of Population Health. BRFSS Prevalence and Trends Data [online] 2015. 2015. URL:<http://wwwdev.cdc.gov/brfss/brfssprevalence/>. in Series BRFSS Prevalence and Trends Data [online] 2015. URL:<http://wwwdev.cdc.gov/brfss/brfssprevalence/>
- Chen C, M YHY. Surveillance Report #98: APPARENT PER CAPITA ALCOHOL CONSUMPTION: NATIONAL, STATE, AND REGIONAL TRENDS, 1977–2012. 2014aNational Institute on Alcohol Abuse and Alcoholism (NIAAA)Bethesda, MD in Series Surveillance Report #98: APPARENT PER CAPITA ALCOHOL CONSUMPTION: NATIONAL, STATE, AND REGIONAL TRENDS, 1977–2012
- Chen C, M YHY. Surveillance Report #99: Trends in Alcohol-Related Morbidity among Community Hospital Discharges, United States 2000-2012. 2014bNational Institute on Alcohol Abuse and Alcoholism (NIAAA)Bethesda, MD in Series Surveillance Report #99: Trends in Alcohol-Related Morbidity among Community Hospital Discharges, United States 2000-2012
- Chen C, M YHY. Surveillance Report #100: LIVER CIRRHOSIS MORTALITY IN THE UNITED STATES: NATIONAL, STATE, AND REGIONAL TRENDS, 2000–2011. 2014c:2000–2011. in

Series Surveillance Report #100: LIVER CIRRHOSIS MORTALITY IN THE UNITED STATES: NATIONAL, STATE, AND REGIONAL TRENDS

- Chen C, M YHY. Surveillance Report #100: LIVER CIRRHOSIS MORTALITY IN THE UNITED STATES: NATIONAL, STATE, AND REGIONAL TRENDS, 2000–2011. 2014dNational Institute on Alcohol Abuse and Alcoholism (NIAAA)Bethesda, MD in Series Surveillance Report #100: LIVER CIRRHOSIS MORTALITY IN THE UNITED STATES: NATIONAL, STATE, AND REGIONAL TRENDS, 2000–2011. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism (NIAAA)
- Chu CM, Liaw YF. Hepatitis B virus-related cirrhosis: natural history and treatment. *Seminars in liver disease*. 2006; 26:142–152. [PubMed: 16673292]
- Dasarathy S. Treatment to improve nutrition and functional capacity evaluation in liver transplant candidates. *Current treatment options in gastroenterology*. 2014; 12:242–255. [PubMed: 24691782]
- Domanski JP, Park SJ, Harrison SA. Cardiovascular disease and nonalcoholic fatty liver disease: does histologic severity matter? *Journal of clinical gastroenterology*. 2012; 46:427–430. [PubMed: 22469639]
- El-Serag HB. Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology*. 2012; 142:1264–1273. e1261. [PubMed: 22537432]
- Fargion S, Porzio M, Fracanzani AL. Nonalcoholic fatty liver disease and vascular disease: state-of-the-art. *World J Gastroenterol*. 2014; 20:13306–13324. [PubMed: 25309067]
- Frazier TH, Stocker AM, Kershner NA, Marsano LS, McClain CJ. Treatment of alcoholic liver disease. *Therapeutic advances in gastroenterology*. 2011; 4:63–81. [PubMed: 21317995]
- Ganesh S, Rogal SS, Yadav D, Humar A, Behari J. Risk factors for frequent readmissions and barriers to transplantation in patients with cirrhosis. *PLoS One*. 2013; 8:e55140. [PubMed: 23383085]
- Goyal A, Murray JM. The impact of vaccination and antiviral therapy on hepatitis B and hepatitis D epidemiology. *PloS one*. 2014; 9:e110143. [PubMed: 25313681]
- Hatzakis A, Chulanov V, Gadano AC, Bergin C, Ben-Ari Z, Mossong J, Schreter I, Baatarkhuu O, Acharya S, Aho I, Anand AC, Andersson MI, Arendt V, Arkkila P, Barclay K, Bessone F, Blach S, Blokhina N, Brunton CR, Choudhuri G, Cisneros L, Croes EA, Dahgwahdorj YA, Dalgard O, Daruich JR, Dashdorj NR, Davaadorj D, de Knecht RJ, de Vree M, Estes CA, Flisiak R, Gane E, Gower E, Halota W, Henderson C, Hoffmann P, Hornell J, Houlihan D, Hrusovsky S, Jarcuska P, Kershenobich D, Kostrzevska K, Kristian P, Leshno M, Lurie Y, Mahomed A, Mamonova N, Mendez-Sanchez N, Norris S, Nurmukhametova E, Nymadawa P, Oltman M, Oyunbileg J, Oyunsuren T, Papatheodoridis G, Pimenov N, Prabdial-Sing N, Prins M, Radke S, Rakhmanova A, Razavi-Shearer K, Reesink HW, Ridruejo E, Safadi R, Sagalova O, Sanchez Avila JF, Sanduijav R, Saraswat V, Seguin-Devaux C, Shah SR, Shestakova I, Shevaldin A, Shibolet O, Silva MO, Sokolov S, Sonderup M, Souliotis K, Spearman CW, Staub T, Stedman C, Strebkova EA, Struck D, Sypsa V, Tomasiewicz K, Undram L, van der Meer AJ, van Santen D, Veldhuijzen I, Villamil FG, Willemse S, Zuckerman E, Zuure FR, Puri P, Razavi H. The present and future disease burden of hepatitis C virus (HCV) infections with today's treatment paradigm - volume 2. *Journal of viral hepatitis*. 2015; 22(Suppl 1):26–45. [PubMed: 25560840]
- Hoyert, DLXJD. Preliminary data for 2011. *National vital statistics reports*. Vol. 61. National Center for Health Statistics; Hyattsville, MD.: 2012.
- Iruzubieta P, Crespo J, Fabrega E. Long-term survival after liver transplantation for alcoholic liver disease. *World journal of gastroenterology : WJG*. 2013; 19:9198–9208. [PubMed: 24409048]
- Kabiri M, Jazwinski AB, Roberts MS, Schaefer AJ, Chhatwal J. The changing burden of hepatitis C virus infection in the United States: model-based predictions. *Annals of Internal Medicine*. 2014; 161:170–180. [PubMed: 25089861]
- Litten RZ, Egli M, Heilig M, Cui C, Fertig JB, Ryan ML, Falk DE, Moss H, Huebner R, Noronha A. Medications development to treat alcohol dependence: a vision for the next decade. *Addiction Biology*. 2012; 17:513–527. [PubMed: 22458728]
- Mathurin P, Bataller R. Trends in the management and burden of alcoholic liver disease. *J Hepatol*. 2015; 62:S38–S46. [PubMed: 25920088]

- McMahon, ASFL.; Brian, J. Chronic Hepatitis B: Update 2009. AASLD guidelines; 2014. in Series Chronic Hepatitis B: Update 2009
- Montgomery, DCPEA.; Vining, GG. Introduction to linear regression analysis. John Wiley & Sons; 2012. p. 821
- Moreno C, Hezode C, Marcellin P, Bourgeois S, Francque S, Samuel D, Zoulim F, Grange JD, Shukla U, Lenz O, Ouwerkerk-Mahadevan S, Fevery B, Peeters M, Beumont M, Jessner W. Efficacy and safety of simeprevir with PegIFN/ribavirin in naive or experienced patients infected with chronic HCV genotype 4. *Journal of hepatology*. 2015; 62(5):1047–55. [PubMed: 25596313]
- Morgan MY. The prognosis and outcome of alcoholic liver disease. *Alcohol and alcoholism*. 1994; 2:335–343. [PubMed: 8974353]
- Moyer VA. Summaries for patients: screening for hepatitis C virus infection in adults: U.S. Preventive Services Task Force recommendation statement. *Annals of Internal Medicine*. 2013; 159:I–32.
- Mueller S, Millonig G, Seitz HK. Alcoholic liver disease and hepatitis C: a frequently underestimated combination. *World journal of gastroenterology : WJG*. 2009; 15:3462–3471. [PubMed: 19630099]
- Neuschwander-Tetri BA, Loomba R, Sanyal AJ, Lavine JE, Van Natta ML, Abdelmalek MF, Chalasani N, Dasarathy S, Diehl AM, Hameed B, Kowdley KV, McCullough A, Terrault N, Clark JM, Tonascia J, Brunt EM, Kleiner DE, Doo E, for the NCRN. Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet* 2015 Mar 14. 2015; 385(9972):946.
- O'Grady JG. Liver transplantation alcohol related liver disease: (deliberately) stirring a hornet's nest! *Gut*. 2006; 55:1529–1531. [PubMed: 17047102]
- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *Jama*. 2014; 311:806–814. [PubMed: 24570244]
- Rockville, M. Annual Data Report. Department of Health and Human Services, Health Resources and Services Administration; 2014. p. 201OPTN/SRTR 2012 Annual Data Report in Series Annual Data Report
- Organization WHO. Global status report on alcohol and health. 2014 in Series Global status report on alcohol and health.
- Petta S, Craxi A. Current and future HCV therapy: do we still need other anti-HCV drugs? *Liver international : official journal of the International Association for the Study of the Liver*. 2015; 35(Suppl 1):4–10. [PubMed: 25529081]
- Rafiq N, Bai C, Fang Y, Srishord M, McCullough A, Gramlich T, Younossi ZM. Long-term follow-up of patients with nonalcoholic fatty liver. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association*. 2009; 7:234–238. [PubMed: 19049831]
- Robert, E.; Mann, RGS.; Richard, G. The Epidemiology of Alcoholic Liver Disease. NIAAA; 2004.
- Sarvari J, Mojtahedi Z, Taghavi SA, Kuramitsu Y, Shamsi Shahrabadi M, Ghaderi A, Nakamura K. Differentially Expressed Proteins in Chronic Active Hepatitis, Cirrhosis, and HCC Related to HCV Infection in Comparison With HBV Infection: A proteomics study. *Hepatitis monthly*. 2013; 13:e8351. [PubMed: 24066001]
- Scaglione S, Kliethermes S, Cao G, Shoham D, Durazo R, Luke A, Volk ML. The Epidemiology of Cirrhosis in the United States: A Population-based Study. *J Clin Gastroenterol*. 2014
- Shepard CW, Simard EP, Finelli L, Fiore AE, Bell BP. Hepatitis B virus infection: epidemiology and vaccination. *Epidemiologic reviews*. 2006; 28:112–125. [PubMed: 16754644]
- Tsoufas G, Goulis I, Giakoustidis D, Akriviadis E, Agorastou P, Imvrios G, Papanikolaou V. Hepatitis C and liver transplantation. *Hippokratia*. 2009; 13:211–215. [PubMed: 20011084]
- Wasley A, Kruszon-Moran D, Kuhnert W, Simard EP, Finelli L, McQuillan G, Bell B. The prevalence of hepatitis B virus infection in the United States in the era of vaccination. *The Journal of infectious diseases*. 2010; 202:192–201. [PubMed: 20533878]
- Wong RJ, Aguilar M, Cheung R, Perumpail RB, Harrison SA, Younossi ZM, Ahmed A. Nonalcoholic steatohepatitis is the second leading etiology of liver disease among adults awaiting liver transplantation in the United States. *Gastroenterology*. 2015; 148:547–555. [PubMed: 25461851]

- Yanovski SZ, Yanovski JA. Obesity prevalence in the United States--up, down, or sideways? *N Engl J Med*. 2011; 364:987–989. [PubMed: 21410367]
- Zezos P, Renner EL. Liver transplantation and non-alcoholic fatty liver disease. *World journal of gastroenterology : WJG*. 2014; 20:15532–15538. [PubMed: 25400437]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Number of discharged patients from the US hospitals with alcohol related diagnosis: Hospital discharges in the United States related to cirrhosis show that over half are due to alcohol related cirrhosis. NIAAA Data report #99 (December 2014).

Figure 2.

Panel A: Number of patient discharged from US hospitals who carried a diagnosis related to alcohol abuse: Patients discharged from hospitals in the United States with a diagnosis of alcohol disorder. Numbers are in the 1000's. Data derived from NIAAA Surveillance Report #99 (December 2014). Panel B: Projection on future numbers of discharged patients from US hospitals with Alcohol related diagnosis: Alcohol related diagnosis is predicted to continue increasing in the next 10 years as shown in the projection made from the current hospital discharges

Figure 3.

Panel A: Death rate in cirrhotic population: High yearly death rate from alcohol related cirrhosis compared with other specific and non-specific causes of cirrhosis. Data from NIAAA surveillance report # 100 (December 2014). Panel B: Projection using linear regression model on death in cirrhotic population rate till 2025: Projection made on the death rate in the cirrhotic population with different etiologies.

Figure 4.

Per capita alcohol consumption from 1950 till 2010 and projection till 2025: Per capita alcohol consumption during the last 6 decades in the United States data was driven from NIAAA surveillance report # 98 (April 2014). Projection was made based on the past as well on the current rates of alcohol consumptions in the United States.

Figure 5.

Panel A: The changing burden of hepatitis C virus infection in the United States: model-based predictions: The prevalence of HCV is decreasing in the United States because of improved HCV screening and availability of new antiviral agents (Kabiri et al. 2014). Panel B: The changing prevalence of HCV-associated decompensated cirrhosis and hepatocellular carcinoma in the United States: The prevalence of HCV-associated decompensated cirrhosis and hepatocellular carcinoma in the United States is expected to decrease due to the advances in management of HCV infection (Kabiri et al. 2014). Panel C: Anticipated numbers of HCV cirrhosis associated transplant till 2050. Due to the expected decrease in HCV prevalence, the number of patients with HCV cirrhosis undergoing liver transplantation will also decrease over time.

Figure 6.

Projection using linear regression model on the numbers of alcohol related cirrhosis on the transplant list till 2025: An increase in the number of patients with alcohol related cirrhosis on the transplant waiting list and transplanted patients compared to other etiologies of cirrhosis is anticipated. Data from SRTR report 2014.

Figure 7.

Panel A; Number of patients who underwent liver transplantation for cirrhosis due to different etiologies: Number of patients with alcohol related cirrhosis who underwent liver transplantation has been nearly unchanged during the past 4 years. Panel B: Percentage of listed to transplanted patients. Ratio of listed to transplanted patient numbers with a diagnosis of alcohol related cirrhosis is decreasing. Data from the SRTR report 2014.

Figure 8.

Prevalence of obesity and overweight people in the United States from 2000 to 2013.

Prevalence of obesity has increased over the last 13 years but the rate of increase is lower since 2007. Data from the CDC behavioral risk factor surveillance report 2015.