Published in final edited form as:

Chem Commun (Camb). 2008 May 28; (20): 2322–2324. doi:10.1039/b717747j.

Spectral observation of conversion between ionized vs. non-ionized proton-coupled electron transfer interfaces†

Elizabeth R. Young, Joel Rosenthal, and Daniel G. Nocera

Department of Chemistry, 6-335, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA 02139-4307, USA

Abstract

Two-point hydrogen bonding between acid and base functionalities provides a convenient method for the modular assembly of proton-coupled electron transfer (PCET) networks, especially when that interface comprises an amidinium and two-point anionic partner; a system is presented that permits the proton configuration within the interface to be determined when pK_a values of the conjugate acids are known.

Proton-coupled electron transfer (PCET) may be established within donor–acceptor dyads assembled by a hydrogen bonded interface.^{1,2} PCET networks assembled from Watson–Crick base pairs³ and dicarboxylic acid dimers⁴ do not support proton transfer within the interface thus limiting the extent to which PCET may be investigated. For this reason, the asymmetric proton interface of an amidinium bound to two-point, anionic partners, especially carboxylate, has come to the fore.^{5–10} The two tautomers shown in Chart 1 are possible: an ionized amidinium–carboxylate or a non-ionized amidine–carboxylic acid, both of which can support proton transfer along a PCET coordinate.

The amidinium–carboxylate interface combines the dipole of an electrostatic ion pair interaction, a two-point hydrogen bond, and secondary electrostatic interactions to produce a highly stable proton interface.^{11–15} These factors along with the relative pK_a values of amidinium and carboxylate, conspire to determine the extent to which the interface configuration is ionized.¹⁶ The electrostatic interactions within the interface stabilize charge accumulation, leading in some cases to an ionized interface even when a simple pK_a calculation predicts formation of a non-ionized interface in a solution of a given dielectric constant.¹⁰ We now show the pK_a value necessary to achieve the ionized vs. the non-ionized interface by stepping through a series of binding moieties (BMs) with different acidity constants. The transition from a non-ionized to ionized interface occurs as the acidity of the BM increases.

Efficient binding of an amidinium (amH^+) to carboxylate and sulfonate anions occurs in aprotic solvents of low dielectric constant. A series of BMs of varying pK_a values in ACN was chosen as summarized in Table 1.^{17,18} The trend in the dependence of the $\text{amH}^+:\text{BM}^-$

†Electronic supplementary information (ESI) available: Experimental procedures, additional spectra and charts. See DOI: 10.1039/b717747j

Daniel G. Nocera: nocera@mit.edu; Fax: 01 617 253 7670; Tel: 01 617 253 5537.

association and interface configuration on the dielectric constant of the solvent can be probed along the series tetrahydrofuran (THF, $\epsilon = 7.58$), dichloromethane (DCM, $\epsilon = 8.93$) and acetonitrile (ACN, $\epsilon = 37.5$).¹⁹

Interrogation of the proton configuration of the amidinium interface was accomplished using purpurin 1, in which conjugation is maintained between the amidinium and porphyrin *via* an unsaturated five-membered ring. We have shown that the absorption spectrum of 1 is sensitive to its protonation state.¹⁰ Reduction of the five-membered isocyclic ring of 1 yields the corresponding chlorin homologue. Unlike 1, the absorption spectrum of the chlorin–amidinium is invariant with pH (see Fig. S1, ESI[†]).

Fig. 1 shows the absorption spectrum of 1 before (**1-amH**⁺) and after (**1-am**) addition of the base 4-dimethylaminopyridine (DMAP). Deprotonation of the amidinium functionality in ACN is indicated by a shift in the absorption maximum of the Soret band from 428 to 423 nm (Fig. 1(c)). Similar spectral trends are observed in THF and DCM (Fig. 1(a) and (b), respectively). Absorption spectra (Fig. S2–S4, ESI[†]) demonstrate analogous spectral shifts with each BM and exhibit well-anchored isosbestic points during the titration of the **BM** with **1-amH**⁺. A 1:1 binding motif is signified from a Benesi–Hildebrand analysis of the titration data ($1/A$ vs. $1/[BM]$) (Fig. S2–S4, ESI[†]).²⁰ The binding constant, K_{assoc} , for 1:2 is found to be $\sim 4.5 \times 10^4 \text{ M}^{-1}$ and for 1:3 is $\sim 1.5 \times 10^2 \text{ M}^{-1}$ in ACN. The K_{assoc} for **4**, **5** and **6** are all the same within error and are measured to be $\sim 9.9 \times 10^3 \text{ M}^{-1}$.

The endpoint spectrum of each titration reveals the configuration of the proton interface as indicated by the final position of the Soret ($S_2 \leftarrow S_0$ transition at 430 nm) or Q-band ($S_1 \leftarrow S_0$ transition at ~ 650 nm) maximum. Conversion of **1-amH**⁺ to **1-am** in the non-ionized interface is indicated by a shift of the peak maxima to wavelengths nearly coincident with that observed for **1-am** upon deprotonation of **1-amH**⁺ with DMAP. We interpret the small 1–2 nm shifts of the Soret band of **1-amH**⁺ in the presence of the BM to the establishment of a non-ionized amidine–carboxylic acid interface.¹⁰

Fig. 1 (inset) shows the final Soret peak position of 1 in each solvent for all BMs studied. The solid data points at each end represent the peak position of **1-amH**⁺ (highest wavelength) and **1-am** (lowest wavelength). The dotted line in each plot shows an approximate boundary for the transition between the ionized to non-ionized interface as determined by the final Soret position, and augmented by spectral analysis of the initial and final spectra (see ESI[†] for details of spectral analysis). The unavailability of additional BMs with precise pK_a values limits the observation of a clearly identifiable pK_a transition between the two tautomers.

Experimental observation of the proton interface configuration for each BM and determination of K_{assoc} for a given interface enables the construction of the thermodynamic cycle shown in Scheme 2. Since the interface under consideration is formed from amH^+ and $^-\text{O}_2\text{C}$ as the reactants (Scheme 2 top, left), either an $\text{amH}^+ \dots ^-\text{O}_2\text{C}$ or an $\text{am} \dots \text{HO}_2\text{C}$

[†]Electronic supplementary information (ESI) available: Experimental procedures, additional spectra and charts. See DOI: 10.1039/b717747j

interface is produced. The driving force for formation of the former (G_1) is measured directly from the measurement of K_{assoc} for the ionized **1-amH⁺·O₂C** complex, while the driving force for the latter (G_3) is determined for the non-ionized **1-am...HO₂C** complex (Scheme 2, inset). The driving force for formation of am...HO₂C from the neutral analogues (G_2) can be determined by the difference between the G_3 and $\Delta G_{\text{p}K_a}$ as shown by the inset in Scheme 2,

$$\Delta G_2 = \Delta G_3 - \Delta G_{\text{p}K_a} \quad (1)$$

Accordingly, the stabilization imparted by proton transfer within the interface ($G_{\text{interface}}$) can be determined from the free energies of the square scheme and the $\text{p}K_a$ s of the constituent moieties ($\Delta G_{\text{p}K_a}$).

Although the binding of the various BMs of Table 1 to **1-amH⁺** was carried out in ACN, DCM and THF, the thermodynamic description and stabilization energy of the interface outlined above can only be rigorously determined for ACN since $\text{p}K_a$ values for the BMs used in this study have been determined in this solvent. Inspection of Fig. 1(c) reveals that conversion between the tautomeric forms of the interface occurs between BMs **4** and **5**. Thus, Scheme 2 is analyzed for these two BMs as they most closely define the razor's edge between formation of the ionized and neutral hydrogen bonding interface. Accordingly, these two points allow for the determination of the minimum interface stabilization necessary to obtain an ionized interface as per eqn (2):

$$\Delta G_{\text{interface}} = (\Delta G_2 + \Delta G_{\text{p}K_a}) - \Delta G_1 \quad (2)$$

The am...HO₂C tautomer is formed upon association of **1-amH⁺** with **4**, allowing for the determination of G_2 via eqn (1). G_1 is obtained from $K_{\text{assoc}}(\mathbf{1}:\mathbf{5})$, since association of **1-amH⁺** with **5** generates the ionized **amH⁺·O₂C** complex. That the ionized interface for **1:5** persists when the $\text{p}K_a$ difference dictates formation of the am...HO₂C interface reveals the importance of electrostatic contributions to the interface configuration. The electrostatic stabilization of the interface imparted by formation of the ionized **amH⁺·O₂C** complex ($G_{\text{interface}}$) can be determined using eqn (2); it is minimally 0.18 eV in ACN. This electrostatic interaction energy can be calculated for point charges, q_1 and q_2 , according to $V = (q_1 q_2) / (4\pi\epsilon r)$ where ϵ is the dielectric constant of the medium. An electrostatic stabilization energy of 0.10 eV is calculated for the amidinium-carboxylate salt bridge at a distance of $r = 3.9 \text{ \AA}$ in THF. This distance is measured between the central carbon atoms of the amidinium and carboxylate groups from the energy optimized structure.⁸ The experimental results reported here agree with a simple ion solvation model. The discrepancy likely arises from the approximation that the electrostatic charge within the interface is confined to a point-dipole (Table 2).

While the dearth of known $\text{p}K_a$ values in DCM and THF preclude a similar thermodynamic analysis in those solvents, a qualitative understanding of interface behaviour is inferred from experimental results. In the dielectric environment of DCM, the proton interface maintains the ionized configuration at the same $\text{p}K_a$ difference as ACN, however in the lower

dielectric environment of THF conversion to the ionized tautomer shifts to lower pK_a . THF does not tolerate charge buildup to the extent that ACN and DCM do, and thereby favours a non-ionized interface configuration at a lower free energy.

The energetics of the amidinium-carboxylate interface are essential to assigning a general interface stabilization energy with the goal of accurately projecting the tautomeric state of the proton interface to PCET systems. For example, the hydrogen bonded donor-acceptor, system D-[H⁺]-A (D = amidinium appended Zn(II)TMP, A = naphthalene diimide carboxylic acid) reveals a PCET rate of $9.50 \times 10 \text{ s}^{-1}$ with a kinetic isotope effect (KIE) of 1.06 at room temperature. In this case, the interface is the non-ionized amidine-carboxylic acid tautomer. When this D-[H⁺]-A system is subtly altered to produce an ionized amidinium-sulfonate interface (A = naphthalene diimide sulfonic acid) of similar driving force, (Table 1, represented by the **1:3** interface), the change in interface configuration has a clear effect on the PCET rate and solvent dependent ET parameters ($|V|$, λ , E_{act}). The PCET rate roughly doubles to $19 \times 10 \text{ s}^{-1}$ with a KIE of 1.17 and more strikingly ET parameters show a clear solvent dependence through the ionized interface, which is absent in the first system. The enhancement of PCET rate and solvent dependent ET parameters for the same D-A pair bearing different tautomeric forms of the interface, illustrates the pronounced effect that proton position exerts on photoinduced charge transfer.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Notes and references

1. Chang, C.J.; Brown, J.D.; Chang, M.C.Y.; Baker, E.A.; Nocera, D.G. *Electron Transfer in Chemistry*. Balzani, V., editor. Vol. 3.2.4. Wiley-VCH; Weinheim, Germany: 2001. p. 409-461.
2. Cukier RI, Nocera DG. *Annu Rev Phys Chem*. 1998; 49:337-369. [PubMed: 9933908]
3. Sessler, J.L.; Wang, B.; Springs, S.L.; Brown, C.T. *Comprehensive Supramolecular Chemistry*. Murakami, Y., editor. Vol. 4. Pergamon Press; Oxford: 1996. p. 311-336. and references therein
4. Turró C, Chang CK, Leroi GE, Cukier RI, Nocera DG. *J Am Chem Soc*. 1992; 114:4013-4015.
5. Roberts JA, Kirby JP, Nocera DG. *J Am Chem Soc*. 1995; 117:8051-8052.
6. Roberts JA, Kirby JP, Wall ST, Nocera DG. *Inorg Chim Acta*. 1997; 263:395-405.
7. Roberts JA, Kirby JP, Nocera DG. *J Am Chem Soc*. 1997; 119:9230-9236.
8. Damrauer NH, Hodgkiss JM, Rosenthal J, Nocera DG. *J Phys Chem B*. 2004; 108:6315-6321. [PubMed: 18950117]
9. Hodgkiss JM, Damrauer NH, Presse S, Rosenthal J, Nocera DG. *J Phys Chem B*. 2006; 110:18853-8. [PubMed: 16986876]
10. Rosenthal J, Hodgkiss JM, Young ER, Nocera DG. *J Am Chem Soc*. 2006; 128:10474-10483. [PubMed: 16895413]
11. Pranata J, Wierschke SG, Jorgensen WL. *J Am Chem Soc*. 1991; 113:2810-2819.
12. Uchimaru T, Korchowiec J, Tsuzuki S, Matsumura K, Kawahara S. *Chem Phys Lett*. 2000; 318:203-209.
13. Lukin O, Leszczynski J. *J Phys Chem A*. 2002; 106:6775-6782.
14. Lukin O, Leszczynski J. *J Phys Chem A*. 2003; 107:9251-9252.
15. Schlund S, Schmuck C, Engels B. *J Am Chem Soc*. 2005; 127:11115-11124. [PubMed: 16076219]

16. Hodgkiss, JM.; Rosenthal, J.; Nocera, DG. Handbook of Hydrogen Transfer Physical and Chemical Aspects of Hydrogen Transfer. Hynes, JT.; Klinman, JP.; Limbach, H-H.; Schowen, RL., editors. Vol. 2.4.17. Wiley-VCH; Weinheim, Germany: 2006. p. 503-562.
17. Izutsu, K. Acid–Base Dissociation Constants in Dipolar Aprotic Solvents. Blackwell Scientific; Cambridge, USA: 1990.
18. The pK_a range reported for each BM are as follows: acetate (4) $pK_a = 20\text{--}22$; chloroacetate (5) $pK_a = 15\text{--}19$; and, dichloroacetate (6) $pK_a = 13\text{--}15$. The pK_a of phenylsulfonate (3) is estimated to be 8 from values for p-toluenesulfonic acid, $pK_a = 8$ and 2,5-dichlorobenzenesulfonic acid, $pK_a = 6.2\text{--}6.4$.
19. Marcus, Y. Ion Solvation. Wiley; Chichester: 1985. p. 136-137.
20. Connors, KA. Binding Constants: A Measurement of Molecular Complex Stability. Wiley; New York: 1987.
21. Marcus, Y. Ion Solvation. John Wiley; Chichester: 1985. p. 136
22. Young ER, Rosenthal JR, Hodgkiss JM, Nocera DG. manuscript in preparation.

Fig. 1.

UV-visible absorption spectra of $\mathbf{1-amH}^+$, solid line; $\mathbf{1-am}$ formed by deprotonated of $\mathbf{1-amH}^+$ with DMAP, dotted line. (a) $\lambda_{\text{max}}(\mathbf{1-amH}^+) = 430$ nm shifts to $\lambda_{\text{max}}(\mathbf{1-am}) = 426$ nm in THF, 2000 eq. DMAP, (b) $\lambda_{\text{max}}(\mathbf{1-amH}^+) = 436$ nm shifts to $\lambda_{\text{max}}(\mathbf{1-am}) = 426$ nm in DCM, 550 eq. DMAP and (c) $\lambda_{\text{max}}(\mathbf{1-amH}^+) = 428$ nm shifts to $\lambda_{\text{max}}(\mathbf{1-am}) = 423$ nm in ACN, 950 eq. DMAP. Inset: Soret peak position of the purpurin *vs.* $\Delta G_{\text{p}K_a}$ for each of the associated $\mathbf{1:BM}$ complexes. The highest and lowest peak maxima (solid circles) represent $\mathbf{1-amH}^+$, solid line; $\mathbf{1-am}$, respectively.

Scheme 1.

Scheme 2.

Square scheme representation of free energies involved in determining the tautomeric form of the interface.

Chart 1.

Table 1

ΔG_{pK_a} and interface configuration for **1:BM** complexes

Binding moiety	pK_a^a	$\Delta G_{pK_a}/eV$ (1:BM) ^b	Ionized interface?		
			THF	DCM	ACN
1 Purpurin-amH ⁺	9.55				
2 3,5-Dinitrobenzoate	17	0.44	✓	✓	✓
3 Phenylsulfonate	7	-0.15	✓	✓	✓
4 Acetate	21	0.68			
5 Chloroacetate	18	0.50	✓		
6 Dichloroacetate	14	0.26	✓	✓	✓

^a In CH₃CN, from ref. 17.

^b At 298 K, calculated from pK_a values recorded in ACN, $\Delta G_{pK_a} = -RT[pK_a(BM)/pK_a(1-am^+)]$.

Table 2

Calculated free energy values for thermodynamic considerations for assignment of DGInterface in CAN.

	Binding moiety (BM)	pK_a^a	G_1^b/eV	G_2/eV	G_3/eV^b	$\Delta G_{pK_a}/\text{eV}$	$G_{\text{interface}}/\text{eV}$
1	Purpurin-amH ⁺	9.55	–	–	–	–	–
2	3,5-Dinitrobenzoate	17	–0.28	–	–	–0.44	–
3	Phenylsulfonate	7	–0.13	–	–	0.15	–
4	Acetate	21	–	0.44	–0.24	–0.68	0.18
5	Chloroacetate	18	–0.24	–	–	–0.50	–
6	Dichloroacetate	14	–0.24	–	–	–0.26	–

^a In CH₃CN, from ref. 17.

^b $G = -RT \ln K_{\text{assoc}}(\mathbf{1}:\mathbf{BM})$.

^c At 298 K, calculated from pK_a values recorded in ACN, $\Delta G_p K_a = -RT \ln K_a(\mathbf{BM})/pK_a(\mathbf{1-amH}^+)$.