Published in final edited form as:

Synlett. 2015 October ; 26(16): 2293–2295. doi:10.1055/s-0035-1560186.

A Scalable Protocol for the Regioselective Alkylation of 2-Methylcyclohexane-1,3-dione with Unactivated sp^3 Electrophiles

Robert J. Sharpe, Maribel Portillo, Robert A. Vélez, and Jeffrey S. Johnson

Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3290, USA

Jeffrey S. Johnson: jsj@unc.edu

Abstract

A method for the C-selective alkylation of 2-methylcyclohexane-1,3-dione with unactivated sp^3 electrophiles is accomplished via alkylation and subsequent deprotection of the derived ketodimethyl hydrazones. The present method provides a high-yielding entry to dialkyl cycloalkanones that cannot be accessed via direct alkylation of 2-methylcyclohexane-1,3-dione. The title reaction may be useful in the scalable preparation of terpene and steroidal building blocks in the arena of natural product synthesis.

Keywords

alkylation; regioselectivity; hydrazones; ketones; alkyl halides; steroids; terpenoids

The 2,2-dialkylcyclohexane-1,3-dione subunit has seen widespread use as a building block in the preparation of terpene and steroidal natural products.¹ The preparation of these compounds is often accomplished via the alkylation of 2-methylcyclohexane-1,3-dione (**1**, Scheme 1) with activated sp^3 electrophiles (e.g., MeI, $H_2C=CHCH_2Br$, BnBr, etc.) or π -electrophiles (e.g. α,β -unsaturated carbonyls). While alkylation with these electrophiles is efficient, the alkylation of **1** with unactivated sp^3 electrophiles is a significant challenge due to preferential O-alkylation in these cases, limiting the effectiveness of this method in target-oriented synthesis.² While a number of methods directed toward C-selective alkylation of **1** have been described,³ these methods have been primarily limited to activated alkyl electrophiles. Piers and others have described preparation of 2,2-dialkylcyclohexane-1,3-diones **3** via alkylation and hydrolysis of 1,5-dimethoxy-1,4-cyclohexadienes **2**,⁴ however,

Correspondence to: Jeffrey S. Johnson, jsj@unc.edu.

Supporting Information

Supporting information for this article is available online at <http://dx.doi.org/10.1055/s-0035-1560186>.

the stoichiometric requirements of *t*-BuLi and HMPA limit scalability and practicality of this protocol.

As part of a research problem in our previously described total synthesis of the indole diterpenoid paspaline,⁵ we required an efficient and scalable alkylation of **1** or its derivatives. We were encouraged by the work of Enders and co-workers,^{3d,6} which demonstrated C-selective alkylation of cyclic ketohydrazones **4** with activated electrophiles (MeI, HC≡CCH₂Br, BnBr, EtO₂CCH₂Br) and EtI. We found that ketohydrazone **4** gave exclusively C-alkylation even with more hindered and functionalized electrophiles; subsequent hydrolysis gave the desired diones **5**. Herein, we describe this extension as a scalable protocol for the preparation of 2,2-dialkylcyclohexane-1,3-diones for use in the arena of natural product synthesis.

A screen of reaction conditions revealed the optimal protocol: Deprotonation of ketohydrazone **4** with KH in THF followed by addition of the electrophile provided access to the intermediate dialkyl ketohydrazones in good yields (Table 1). Hydrolysis of the intermediate alkylation products was completed via Corey's method [Cu(OAc)₂, aq THF]⁷ or upon treatment with Oxone in aqueous acetone.⁸ Employing these conditions, alkylation of **4** with 1-iodooctane (entry 1) gave exclusively C-alkylation in 92% yield, which upon hydrolysis, gave diketone **5a** in 97% yield (89%, two steps). Extending the carbon chain in the alkyl iodide (entries 2 and 3) gave the same C-alkylation selectivity in slightly decreased yields. Alkenyl halides (entries 4 and 5) also performed well (78% and 92%, respectively), which upon hydrolysis provided diketones **5d,e** bearing valuable functional handles. Alkylation of **4** with an iodide bearing a protected hydroxyl group (entry 6) also underwent efficient alkylation (73%, two steps), although the analogous protected aminoiodide (entry 7) failed to react with **4** under any conditions examined. Secondary alkyl halides (entry 8) also underwent alkylation with complete C-regioselectivity, albeit in decreased yields (42%, two steps) to give diketone **5h**. Furthermore, alkyl bromides (entries 9–11) also underwent C-selective alkylation. The alkyl bromide bearing an electrophilic epoxide functionality (entry 11) underwent alkylation (and not the undesired epoxide opening) to give the corresponding diketone upon hydrolysis (52%, two steps).

To demonstrate the scalability of this protocol, the reaction of ketohydrazone **4** with 5-iodo-2-methylpent-2-ene was carried out on an 11.00 gram (65 mmol) scale with complete C-alkylation selectivity (Scheme 2); the sequence provided >10 grams (76% yield) of diketone **5d** over two steps. Notably, purification of the intermediate alkylated hydrazone was omitted, enabling preparation of **5d** requiring only a single purification. Compound **5d** is a critical intermediate in our previously described total synthesis of paspaline.⁵

In summary, a practical and scalable alkylation of 2-methyl-1,3-cyclohexanediones using unactivated electrophiles has been enabled via alkylation⁹ and deprotection^{10,11} of the corresponding ketohydrazones. The scalability and applicability of this protocol has been demonstrated via the 10 gram scale synthesis of diketone **5d**. This methodology may be useful to synthetic chemists in the preparation of 'quaternized' cyclohexane building blocks for natural product synthesis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The project described was supported by Award No. R01 GM084927 from the National Institute of General Medical Sciences. R.J.S. acknowledges an NSF Graduate Research Fellowship and ACS Division of Organic Chemistry Graduate Fellowship. M.P. acknowledges a fellowship from the McNair Undergraduate Scholars Program. R.A.V. acknowledges support from the NSF REU program (CHE-1460874).

References and Notes

1. For illustrative examples of the use of 2,2-dialkylcyclohexanediones in target-oriented synthesis, see: Corey EJ, Virgil SC. *J Am Chem Soc.* 1990; 112:6429. Nicolaou KC, Yang Z, Liu JJ, Ueno H, Nantermet PG, Guy RK, Caliborne CF, Renaud J, Couladouros EA, Paulvannan K, Sorenson EJ. *Nature* (London, UK). 1994; 367:630. [PubMed: 7906395] Solé D, Bonjoch J, García-Rubio S, Peidró E, Bosch J. *Angew Chem Int Ed.* 1999; 38:395. Nicolaou KC, Pfefferkorn JA, Kim S, Wei HX. *J Am Chem Soc.* 1999; 121:4724. Phillips EM, Roberts JM, Scheidt KA. *Org Lett.* 2010; 12:2830. [PubMed: 20469924] Chin ZH, Chen ZM, Zhang YQ, Tu YQ, Zhang FM, Bian Z, Marvin CC, Pettersson M, Martin SF. *J Am Chem Soc.* 2014; 136:14184. [PubMed: 25211501]
2. (a) House, HO. *Modern Synthetic Reactions*. Benjamin; Menlo Park: 1972. p. 520-532. (b) Stowell, JC. *Carbanions in Organic Synthesis*. Wiley; New York: 1979. p. 205-210. (c) Caine, D. *Comprehensive Organic Synthesis*. Trost, BM.; Fleming, I., editors. Vol. 3. Pergamon Press; Oxford: 1991. p. 54-58. (d) Johnson WS, Lunn WH, Fitz K. *J Am Chem Soc.* 1964; 86:1972.
3. (a) Rajamannar T, Palani N, Balasubramanian KK. *Synth Commun.* 1993; 23:3095. (b) Fujiwara M, Hitomi K, Baba A, Matsuda H. *Chem Lett.* 1994; 875. (c) Bedekar AV, Watanabe T, Tanaka K, Fuji K. *Synthesis.* 1995; 1069. (d) Demir AS, Enders D. *J Prakt Chem.* 1997; 339:553.
4. (a) Piers E, Grierson JR. *J Org Chem.* 1977; 42:3755. (b) Tamai Y, Hagiwara H, Uda H. *J Chem Soc, Chem Commun.* 1982:502. (c) Watanabe H, Iwamoto M, Nakada M. *J Org Chem.* 2005; 70:4652. [PubMed: 15932301] (d) Sparling BA, Moebius DC, Shair MD. *J Am Chem Soc.* 2013; 135:644. [PubMed: 23270309]
5. Sharpe RJ, Johnson JS. *J Am Chem Soc.* 2015; 137:4968. [PubMed: 25856767]
6. (a) Corey EJ, Enders D. *Chem Ber.* 1978; 111:1337. (b) Corey EJ, Enders D. *Chem Ber.* 1978; 111:1362.
7. Corey EJ, Knapp S. *Tetrahedron Lett.* 1976; 17:3667.
8. Hajipour AR, Mahboubghah N. *Org Prep Proced Int.* 1999; 31:112.
9. **Typical Procedure for Alkylation of 4KH** (30% dispersion in oil, 0.20 g, 1.50 mmol, 1.50 equiv) was washed free of oil with PE and suspended in anhydrous THF (3.0 mL). The suspension was transferred to a flame-dried, 10 mL round-bottomed flask under an atmosphere of N₂, and the suspension was cooled to -78 °C. Hydrazone **4** (0.25 g, 1.50 mmol, 1.50 equiv) was dissolved in THF (1.0 mL) and added dropwise to the KH suspension. The resulting solution was warmed to 0 °C and stirred 4.5 h. The reaction was recooled to -78 °C, and 1-iodooctane (0.18 mL, 1.00 mmol, 1.00 equiv) was added drop-wise. The mixture was allowed to stir overnight, allowing the reaction to slowly warm to r.t. as the dry ice bath evaporated. After 14 h, the reaction was quenched via addition of sat. NH₄Cl aq (4.0 mL), and the mixture was partitioned in a separatory funnel. The organic layer was separated, and the aqueous layer was extracted with Et₂O (3 × 10 mL). The combined organic extracts were washed with brine (10 mL), dried with MgSO₄ and concentrated in vacuo. The product was purified via flash chromatography (hexanes–EtOAc, 80:20 to 70:30) to afford the product ketohydrazone (0.25 g, 91% yield) as a yellow oil.
10. **Typical Procedure for Hydrazone Hydrolysis (Method 1)** A 10 mL round-bottomed flask was charged with Cu(OAc)₂·H₂O (0.20 g, 1.00 mmol, 2.00 equiv) and H₂O (3.3 mL), and the solution was allowed to stir until complete dissolution of Cu(OAc)₂·H₂O was observed, typically 2 min. The intermediate ketohydrazone derived from alkylation of **4** with 1-iodooctane (0.14 g, 0.50 mmol, 1.00 equiv) was dissolved in THF (3.3 mL) and added to the Cu(OAc)₂·H₂O solution. The

resulting reaction mixture was allowed to stir until complete conversion of the starting material was observed by TLC analysis, typically 12 h. The reaction mixture was concentrated on a rotary evaporator to remove THF, and the remaining solution was quenched with sat. NH_4Cl aq (5 mL) and diluted with CH_2Cl_2 (5 mL). The mixture was partitioned in a separatory funnel, the organic layer was separated, and the aqueous layer was extracted with CH_2Cl_2 (3×10 mL). The combined organic extracts were dried with Na_2SO_4 and concentrated in vacuo. The product was purified via flash chromatography (hexanes–EtOAc, 90:10 to 80:20) to afford diketone **5a** (0.12 g, 99% yield) as a yellow oil.

11. **Typical Procedure for Hydrazone Hydrolysis (Method 2)** A 20 mL scintillation vial was charged with acetone (2.0 mL) and Oxone (1.20 g, 2.00 mmol, 4.00 equiv) with magnetic stirring. The intermediate ketohydrazone derived from alkylation of **4** with 2-iodopropane (0.11 g, 0.50 mmol, 1.00 equiv) was dissolved in acetone (0.7 mL) and transferred to the Oxone solution at r.t., producing a white suspension. The resulting mixture was stirred 2 h, whereupon the solution was concentrated on a rotary evaporator to remove the acetone. The remaining solution was diluted with H_2O (5 mL) and transferred to a separatory funnel. The aqueous layer was extracted with EtOAc (2×7 mL) and CH_2Cl_2 (2×7 mL), and the combined organic extracts were dried with MgSO_4 and concentrated in vacuo. The product was purified via flash chromatography (hexanes–EtOAc, 90:10 to 80:20) to afford diketone **5h** (0.051 g, 65% yield) as a yellow oil.

(a) alkylation of **1** with unactivated sp^3 electrophiles

(b) alkylation of 1,3-cyclohexadienes (ref. 4)

(c) this work

Scheme 1.

C-Selective alkylations of 2-alkylcyclohexane-1,3-diones and their derivatives

Scheme 2. Large-scale synthesis of dialkyldiketone **5d** (10 g scale)

Table 1Substrate Scope of the Alkylation/Hydrolysis of Ketohydrazone **4**^a

Entry	Alkyl halide	Product	Alkylation yield (%) ^b	Hydrolysis method (yield, %) ^c
1		5a	92	1 ^d (97)
2		5b	81	1 (81)
3		5c	90	1 (81)
4		5d	78	1 (91)
5		5e	92	1 (80)
6		5f	90	1 (81)
7		5g	0	—
8		5h	67	2 ^e (63)
9		5i	68	1 (78)
10		5j	94	2 (80)
11		5k	67	1 (77)

^a Yields refer to isolated analytically pure products. Yields are reported as averages of at least two experiments.^b Reactions were carried out with 1.0 mmol alkyl halide, 1.5 mmol **4**, and 1.5 mmol of KH.^c Reactions were carried out with 0.5 mmol of intermediate ketohydrazone.

^d Method 1: Cu(OAc)₂ (2.0 equiv), THF–H₂O (1:1), r.t.

^e Method 2: Oxone (4.0 equiv), acetone–H₂O (1:3), r.t.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript