Published in final edited form as:

Angew Chem Int Ed Engl. 2015 September 1; 54(36): 10600–10603. doi:10.1002/anie.201503773.

Styrene Aziridination by Iron(IV) Nitrides

Dr. Salvador B. Muñoz III^{[a],[+]}, Dr. Wei-Tsung Lee^{[a],[+]}, Dr. Diane A. Dickie^[b], Dr. Jeremiah J. Scepianiak^[c], Deepak Subedi^[c], Dr. Maren Pink^[a], Prof. Michael D. Johnson^[c], and Prof. Jeremy M. Smith^[a]

Jeremy M. Smith: smith962@indiana.edu

^[a]Department of Chemistry, Indiana University, 800 E. Kirkwood Avenue, Bloomington, IN 47403, USA

^[b]Department of Chemistry and Chemical Biology, The University of New Mexico, Albuquerque NM 87131, USA

^[c]Department of Chemistry and Biochemistry, New Mexico State University, Las Cruces NM 88003, USA

Abstract

Thermolysis of the iron(IV) nitride complex $\text{PhB}(\text{tBuIm})_3\text{Fe}\equiv\text{N}$ with styrene leads to formation of the high spin iron(II) aziridino complex $\text{PhB}(\text{tBuIm})_3\text{Fe-N}(\text{CH}_2\text{CHPh})$. Similar aziridination occurs with both electron-rich and electron-poor styrenes, while bulky styrenes hinder the reaction. The aziridino complex $\text{PhB}(\text{tBuIm})_3\text{Fe-N}(\text{CH}_2\text{CHPh})$ acts as a nitride synthon, reacting with electron poor styrenes to generate their corresponding aziridino complexes, i.e. aziridine cross-metathesis. Reaction of $\text{PhB}(\text{tBuIm})_3\text{Fe-N}(\text{CH}_2\text{CHPh})$ with Me_3SiCl releases the *N*-functionalized aziridine $\text{Me}_3\text{SiN}(\text{CH}_2\text{CHPh})$ while simultaneously generating $\text{PhB}(\text{tBuIm})_3\text{FeCl}$. This closes a synthetic cycle for styrene aziridination by a nitride complex. While the less hindered iron(IV) nitride complex $\text{PhB}(\text{MesIm})_3\text{Fe}\equiv\text{N}$ reacts with styrenes below room temperature, only bulky styrenes lead to tractable aziridino products.

Keywords

iron; nitrides; alkenes

The facile reactivity of aziridines, coupled with their ability to undergo highly regio- and stereoselective transformations, makes them important and versatile synthons in the preparation of small molecules (e.g. pharmaceuticals) and materials (e.g. medical devices).^[1] The aziridine moiety is also found in some natural products and synthetic compounds that have interesting or useful biological properties, making them of interest as synthetic targets in their own right.^[2]

Correspondence to: Jeremy M. Smith, smith962@indiana.edu.

[+]⁺These two authors contributed equally to this paper.

Supporting information for this article is given via a link at the end of the document.

Given their importance, it is not surprising that numerous schemes for the synthesis of aziridines have been devised.^[3] A particularly appealing strategy involves the catalytic transfer of nitrenes to alkenes, which has been extensively investigated and a wide range of metal-based catalysts have been reported.^[4] These catalysts include many examples of iron-based complexes that are catalysts for aziridine synthesis by sulfilimine transfer.^[5] More recently, the family of iron-based catalysts has been expanded to include complexes that facilitate alkyl- and aryl nitrene transfer to styrenes and aliphatic alkenes.^[6]

Transition metal nitrides might also be expected to show similar reactivity, in which a two electron nitrogen atom transfer reaction from the nitride to the alkene leads to formation of a coordinated aziridino ligand. A conceptually attractive feature of such a reaction is that the newly-formed aziridino ligand provides a platform for accessing a wider range of *N*-functionalized aziridines than is possible by current nitrene transfer strategies. However, the majority of nitride ligands are notoriously unreactive, and examples of nitride ligands reacting with hydrocarbons are notable for their rarity.^[7] To the best of our knowledge, there is only one example of alkene aziridination by a nitride complex, in which addition of pyridine to the Ru^{VI} nitride complex [(salchda)Ru≡N]⁺ (salchda = *N,N'*-bis(salicylidene)-*o*-cyclohexyldiamine dianion) activates the nitride ligand towards alkene aziridination.^[8] A handful of other nitride complexes have been reported to react with alkenes although aziridino ligand formation does not occur, e.g. *cis*-[(terpy)Os(N)Cl₂]⁺ inserts the nitrogen atom into the C=C bond of stilbene and conjugated dienes, generating azallenium products.^[9]

In contrast to the low reactivity of most transition metal nitride complexes, the iron(IV) complexes PhB(RIm)₃Fe≡N (R = *t*Bu **1**, Mes **2**) which are supported by bulky tris(carbene)borate ligands (Fig. 1),^{[10],[11]} are able to access diverse reaction pathways. Thus, we have chronicled two electron nitrogen atom transfer reactions to substrates such as C≡O, C≡NR and PR₃,^{[10],[11a],[12],[13]} one electron reactions with hydrogen atom donors and organic radicals,^{[11b],[14]} and cycloadditions with conjugated dienes.^[14]

A detailed experimental and computational investigation into the reaction of **2** with phosphines has provided evidence for the ambiphilic nature of the nitride ligand.^[13] An important observation from this study is that the transition state for nitrogen atom transfer involves a σ -symmetry interaction between the nitride LUMO and phosphine HOMO as well as a π -symmetry interaction between the nitride HOMO and the phosphine LUMO. Since this orbital interaction is reminiscent of bonding between low valent metals and unsaturated hydrocarbons, we were stimulated to investigate the reactivity of iron(IV) nitrides towards alkenes. In this communication we report the reactions of **1** and **2** with a range of styrenes, which leads to formation of the corresponding iron(II) aziridino complexes. Evidence for the reversibility of aziridino ligand formation as well as the ability to release an *N*-functionalized aziridine from the metal is also demonstrated.

Heating a solution of complex **1** with excess styrene leads to formation of the high spin (*S* = 2) iron(II) aziridino complex PhB(*t*BuIm)₃Fe-N(CH₂CHPh) **3** in high yield (Scheme 1). The molecular structure of **3** has been determined by single crystal X-ray diffraction (Scheme 1, inset). To the best of our knowledge, this the first structurally characterized complex

featuring an anionic aziridino, rather than neutral aziridine, ligand. The aziridino C(20)-N(5) and C(21)-N(5) bonds of **3** are shorter and the C(20)-N(5)-C(21) bond angle larger than is observed for neutral aziridine ligands, where the corresponding metrics average 1.486(1) Å and 59.5(1.0)°, respectively.^[15] As expected from the anionic nature of the aziridino ligand, the Fe(1)-N(5) bond length is shorter than observed in iron aziridine complexes^[16] but similar to high spin iron(II) amido complexes.^[17] As with the latter complexes, the nitrogen atom of the aziridino ligand in **3** is planar, with the sum of angles around N(5) = 359.9(6)°. The other metrical parameters are unexceptional for a high spin iron(II) tris(carbene)borate complex.^{[11],[12],[13],[18]}

Complex **1** reacts similarly with other styrenes, although the reaction is sensitive to the size of the styrene substrate. While all *para*-substituted styrenes investigated provide the corresponding aziridino complexes, as do *o*-(CH₃)C₆H₄C(H)=CH₂ and C₁₀H₇CH=CH₂, there is no reaction with the bulkier styrene 2,4,6-(CH₃)₂C₆H₃C(H)=CH₂. In addition, no reaction between **1** and β -substituted styrenes, e.g. C₆H₅C(H)=C(H)CH₃, is observed.

While the reactions of nitride complex **1** generally require heat, the reactions of nitride complex **2** occur under ambient conditions. Thus, **2** reacts at room temperature with 3,5-(CF₃)₂C₆H₃CH=CH₂ and C₁₀H₇C(CH₃)=CH₂ to afford the corresponding high spin iron(II) aziridino complexes **4** and **5** as the sole reaction products (Scheme 2). These complexes have also been structurally characterized, showing similar metrical parameters to **3**, except for significant pyramidalization of the aziridino nitrogen atom (sum of angles around N(7) = 336.2(9)° and 348.2(4)° for **4** and **5**, respectively). While **2** reacts with these bulky styrenes under very mild conditions, even below room temperature, the reaction scope of this complex is generally more limited than that of **1**, with less bulky styrene substrates leading to the formation of intractable products.

These synthetic results therefore establish an unusual transformation for a nitride ligand, in which two-electron nitrogen atom transfer to styrenes provides the corresponding aziridino complexes. As mentioned above, there are a handful of transition metal nitride complexes that react with alkenes,^[9] and in only one instance does a two-electron nitrogen atom transfer reaction occur to generate the three-membered aziridine ring.^[8]

Intriguingly, the aziridino complex **3** can be used as an iron nitride synthon. Specifically, heating **3** with (*p*-NO₂C₆H₄)CH=CH₂ leads to formation of the aziridino complex PhB(*t*BuIm)₃Fe-N(CH₂CH(*p*-(NO₂)C₆H₄)) and styrene, i.e. aziridine cross-metathesis (Scheme 3). Similar aziridine cross-metathesis occurs when **3** is treated with electron-poor styrenes such as (*p*-ClC₆H₄)CH=CH₂ and (*p*-CF₃C₆H₄)CH=CH₂, although no reaction occurs with more electron rich styrenes. While detailed studies aimed at elucidating the aziridination mechanism are ongoing, this cross metathesis reactivity strongly suggests the aziridination reaction to be reversible.

In contrast to the reactions with styrenes, which lead to aziridino ligand formation, neither nitride complex reacts with aliphatic alkenes. However, it is possible to independently prepare an iron(II) aziridino complex that would result from such a reaction. Treating PhB(*t*BuIm)₃FeCl with LiN(CH₂CHCH₃) yields the corresponding aziridino complex

PhB(*t*BuIm)₃Fe-N(CH₂CHCH₃) **6** in high yield (Scheme 4). This complex, which has also been structurally characterized,^[15] shows no propensity for releasing propene under thermal or photochemical conditions, nor does it undergo aziridine cross-metathesis reactions with added styrenes. These results therefore suggest that a kinetic barrier prevents the iron nitrides from accessing similar aziridination reactivity with aliphatic alkenes.

As suggested at the outset, the iron(II) aziridino complexes have potential application as synthons for a range of *N*-functionalized aziridines. In a proof-of-concept experiment, treating the aziridino complex **3** with one equivalent Me₃SiCl leads to quantitative formation of the iron(II) complex PhB(*t*BuIm)₃FeCl and the corresponding *N*-silylated aziridine Me₃SiN(CH₂CHPh). To the best of our knowledge, *N*-silylated aziridines are not accessible by current nitrene transfer strategies.^[19] In addition, since the iron(II) chloride complex is readily converted to **1**,^[18] this reaction establishes a synthetic cycle for styrene aziridination using an iron(IV) nitride (Scheme 5). Preliminary results suggest that access to other *N*-functionalized aziridines will be similarly facile, as illustrated by the synthesis of the *N*-methylated aziridine MeN(CH₂CHPh) by reaction of **3** with MeI.

In summary, the reactivity of a suitable iron(IV) nitrides allows a cycle for the synthesis of *N*-functionalized aziridines to be developed. Work to investigate the mechanism of the aziridination reaction is underway, as are efforts to develop catalytic cycles for the synthesis of range of *N*-functionalized aziridines using iron(IV) nitride complexes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding from the DOE-BES (DE-FG02-08ER15996) and Indiana University is gratefully acknowledged. JMS is a Dreyfus Teacher-Scholar. SBM thanks NSF-AMP (HRD 0929343) and NIH-RISE (R25 GM061222-11) for financial support. The Bruker X8 X-ray diffractometer was purchased via an NSF CRIF:MU award to The University of New Mexico, CHE-0443580.

References

1. Gianolio, DA.; Johnston, EA.; Miller, RJ. US Patent. 7,387,836. Jun 17. 2008
2. a) Sweeny J. Chem Soc Rev. 2002; 31:247–258. [PubMed: 12357722] b) Yudin, AK., editor. Aziridines and Epoxides in Organic Synthesis. Wiley; Weinheim: 2006.
3. a) Müller P, Fruit C. Chem Rev. 2003; 103:2905–2920. [PubMed: 12914485] b) Degennaro L, Trinchera P, Luisi R. Chem Rev. 2014; 114:7881–7929. [PubMed: 24823261]
4. Recent reviews: Karila D, Dodd RH. Curr Org Chem. 2011; 15:1507–1523. Chang JWW, Ton TMU, Chan PWH. Chem Rec. 2011; 11:331–357. [PubMed: 22121122] Chow TWS, Qchen G, Liu Y, Zhou CY, Che CM. Pure Appl Chem. 2012; 84:1685–1704. Uchida T, Katsuki T. Chem Rec. 2014; 14:117–129. [PubMed: 24449500]
5. Selected examples: Simonato JC, Pécaut J, Scheidt RW, Marchon JC. Chem Commun. 1999:989–990. Simkhovich L, Gross Z. Tetrahedron Lett. 2001; 42:8089–8092. Heuss BD, Mayer MF, Dennis S, Hossain MM. Inorg Chim Acta. 2003; 342:301–304. Vyas R, Gao GY, Harden JD, Zhang XP. Org Lett. 2004; 6:1907–1910. [PubMed: 15176780] Avenier F, Latour JM. Chem Commun. 2004:1544–1545. Yan S-Y, Wang Y, Shu Y-J, Liu H-H, Zhou X-G. J Mol Cat A. 2006; 248:148–151. Klotz KL, Slominski LM, Hull AV, Gottsacker VM, Mas-Ballesté R, Que L Jr, Halfen JA. Chem Commun. 2007:2063–2065. Nakanishi M, Salit A-F, Bolm C. Adv Synth Cat. 2008;

- 350:1835–1840. Liu P, Wong ELM, Yuen AWH, Che CM. *Org Lett*. 2008; 10:3275–3278. [PubMed: 18582067] Klotz KL, Slominski LM, Riemer ME, Phillips JA, Halfen JA. *Inorg Chem*. 2009; 48:801–803. [PubMed: 19102690] Liang L, Lv H, Yu Y, Wang P, Zhang JL. *Dalton Trans*. 2012; 41:1457–1460. [PubMed: 22187065] Liang S, Jensen MP. *Organometallics*. 2012; 31:8055–8058.
6. a) Liu Y, Che CM. *Chem Eur J*. 2010; 16:10494–10501. [PubMed: 20648488] b) King ER, Hennessy ET, Betley TA. *J Am Chem Soc*. 2011; 133:4917–4923. [PubMed: 21405138] c) Cramer SA, Jenkins DM. *J Am Chem Soc*. 2011; 133:19342–19345. [PubMed: 22081884] d) Hennessy ET, Liu RY, Iovan DA, Duncan RA, Betley TA. *Chem Sci*. 2014; 5:1526–1532. e) Cramer SA, Sánchez RH, Brakhage DF, Jenkins DM. *Chem Commun*. 2014; 50:13967–13970.
7. Smith JM. *Progress Inorg Chem*. 2014; 58:417–470.
8. Man W-L, Lam WWY, Yiu S-M, Lau T-C, Peng S-M. *J Am Chem Soc*. 2004; 126:15336–15337. [PubMed: 15563133]
9. a) Brown SN. *J Am Chem Soc*. 1999; 121:9752–9753. b) Maestri AG, Taylor SD, Schuck SM, Brown SN. *Organometallics*. 2004; 23:1932–1946.
10. Smith JM, Subedi D. *Dalton Trans*. 2012; 41:1423–1429. [PubMed: 22113554]
11. Scepianiak JJ, Young JA, Bontchev RP, Smith JM. *Angew Chem Int Ed*. 2009; 48:3158–3160.
12. Scepianiak JJ, Bontchev RP, Johnson DL, Smith JM. *Angew Chem Int Ed*. 2011; 50:6630–6633.
13. Scepianiak JJ, Margarit CG, Harvey JN, Smith JM. *Inorg Chem*. 2011; 50:9508–9511. [PubMed: 21902179]
14. Lee W-T, Juarez RA, Scepianiak JJ, Muñoz SB III, Dickie DA, Wang H, Smith JM. *Inorg Chem*. 2014; 53:8425–8430. [PubMed: 25068927]
15. See SI for additional details.
16. a) Timken MD, Strouse CE, Soltis SM, Daverio SA, Hendrickson DN, Abdel-Mawgoud AM, Wilson SR. *J Am Chem Soc*. 1986; 108:395–402. [PubMed: 22175454] b) Conti AJ, Chadha RK, Sena KM, Rheingold AL, Hendrickson DN. *Inorg Chem*. 1993; 32:2670–2680.
17. Recent examples of structurally characterized Fe(II) amidos: Mankad NP, Muller P, Peters JC. *J Am Chem Soc*. 2010; 132:4083–4085. [PubMed: 20199026] Cowley RE, DeYonker NJ, Eckert NA, Cundari TR, DeBeer S, Bill E, Ottenwaelde X, Flaschenriem C, Holland PL. *Inorg Chem*. 2010; 49:6172–6187. [PubMed: 20524625] King ER, Hennessy ET, Betley TA. *J Am Chem Soc*. 2011; 133:4917–4923. [PubMed: 21405138] Zadrozny JM, Atanasov M, Bryan AM, Lin CY, Rekker BD, Power PP, Neese F, Long JR. *Chem Sci*. 2013; 4:125–138. Lee WT, Jeon IR, Xu S, Dickie DA, Smith JM. *Organometallics*. 2014; 33:5654–5659. A full listing of these complexes is provided in the SI.
18. a) Scepianiak JJ, Fulton MD, Bontchev RP, Duesler EN, Kirk ML, Smith JM. *J Am Chem Soc*. 2008; 130:10515–10517. [PubMed: 18630913] b) Scepianiak JJ, Harris TD, Vogel CS, Sutter JJ, Meyer K, Smith JM. *J Am Chem Soc*. 2011; 133:3824–3827. [PubMed: 21366252] c) Smith JM, Mayberry DE, Margarit CG, Sutter JJ, Wang H, Meyer K, Bontchev RP. *J Am Chem Soc*. 2012; 134:6516–6519. [PubMed: 22452612] d) Lin HJ, Siretanu D, Dickie DA, Subedi D, Scepianiak JJ, Mitcov D, Clérac R, Smith JM. *J Am Chem Soc*. 2014; 136:13326–13332. [PubMed: 25157642]
19. Strategies for the assembly of N-silylated aziridines: Komatsu M, Ohno M, Tsuno S, Ohshiro Y. *Chem Lett*. 1990:575–576. Rasmussen KG, Jørgensen KA. *J Chem Soc, Chem Commun*. 1995:1401–1402. Arbuzov PV, Voronkov MG, Mirskov RG, Bakhareva EV, Rakhlin VI. *Russ Chem Bull*. 1996; 45:759–760. Vedejs E, Piotrowski DW, Tucci FC. *J Org Chem*. 2000; 65:5498–5505. [PubMed: 10970287]

Figure 1.

The iron(IV) nitrides $\text{PhB(RIm)}_3\text{Fe}\equiv\text{N}$ ($\text{R} = t\text{Bu}$ **1**, Mes **2**).

Scheme 1.

Reaction of **1** with a range of styrenes, leading to formation of the corresponding aziridino complexes. Inset: X-ray crystal structure of **3**, where Ar = C₆H₅. Thermal ellipsoids at 50%, hydrogen atoms and most of the tris(carbene)borate ligand omitted for clarity. Fe(1) – N(5) 1.935(4); Fe(1) – C(1) 2.092(4); Fe(1) – C(8) 2.109(6); N(5) – C(20) 1.417(11); N(5) – C(21) 1.317(14); C(20) – C(21) 1.476; C(1) – Fe(1) – C(8) 90.58(16); Fe(1) – N(5) – C(20) 142.0(5); Fe(1) – N(5) – C(21) 152.5(6); C(20) – N(5) – C(21) 65.3(6).

Scheme 2.

Reaction of **2** with bulky styrenes to afford the corresponding aziridino complexes **4** and **5**. Inset: X-ray crystal structures of **4** (left) and **5** (right). Thermal ellipsoids at 50%, hydrogen atoms and most of the tris(carbene)borate ligand omitted for clarity.

Scheme 3.
Aziridine cross-metathesis.

Scheme 4.

Synthesis of $\text{PhB}(t\text{BuIm})_3\text{Fe-N}(\text{CH}_2\text{CHCH}_3)$ **6**.

Scheme 5.

Synthetic cycle for styrene aziridination by an iron(IV) nitride complex.