Published in final edited form as:

J Med Econ. 2013 July ; 16(7): 845–858. doi:10.3111/13696998.2013.800523.

Cost-Effectiveness of Dabigatran (150 mg Twice Daily) and Warfarin in Patients 65 Years with Non-Valvular Atrial Fibrillation

Brian M Salata, MD, MS^a, David W Hutton, PhD^b, Deborah A Levine, MD, MPH^c, James B Froehlich, MD, MPH^d, and Geoffrey D Barnes, MD, MSc^d

^aNew York Presbyterian/Weill Cornell Medical Center, New York, NY

^bUniversity of Michigan School of Public Health, Ann Arbor, MI; Institute for Healthcare Policy and Innovation, University of Michigan, Ann Arbor, MI

^cDivision of General Medicine, Dept of Internal Medicine, University of Michigan, Ann Arbor, MI; Veterans Affairs Center for Clinical Management Research, Ann Arbor, MI; Institute for Healthcare Policy and Innovation, University of Michigan, Ann Arbor, MI; Department of Neurology and Stroke Program, University of Michigan, Ann Arbor, MI

^dUniversity of Michigan Frankel Cardiovascular Center, Ann Arbor, MI ; Institute for Healthcare Policy and Innovation, University of Michigan, Ann Arbor, MI

Abstract

Dabigatran has been shown to be superior to warfarin for stroke prevention in non-valvular atrial fibrillation (NVAF) but with higher out-of-pocket costs for patients. Although dabigatran has been shown to be cost-effective from a societal perspective, cost implications for individual patients and insurers are not well-described. We aimed to assess cost perspectives of each payer (Medicare and patient) in relation to administration, monitoring and adverse outcomes for dabigatran and warfarin in patients with and without prescription drug coverage. Using a Markov model, we performed a decision analysis comparing 2 treatment strategies (adjusted-dose warfarin and dabigatran 150 mg twice daily) in patients 65 years old with NVAF, CHADS₂ scores 1 and Medicare insurance. Patients have a quality-adjusted life expectancy of 8.998 quality adjusted life years (QALYs) with warfarin and 9.39 QALYs with dabigatran 150 mg twice daily. From Medicare's perspective, the incremental cost-effectiveness ratio (ICER) comparing dabigatran with warfarin was \$35,311 for patients with Part D coverage and cost-saving for patients without coverage. From the patient's perspective, the ICER comparing dabigatran with warfarin was cost-saving for patients with Part D coverage and \$63,884 for patients without coverage. In patients 65 years or older with NVAF and prescription insurance coverage, dabigatran 150 mg twice daily is both cost-effective (Medicare's perspective) and cost-saving (patient perspective) when compared

Corresponding Author: Geoffrey D Barnes, MD, MSc, University of Michigan Frankel Cardiovascular Center, 2800 Plymouth Rd, Building 14 – Room G100, Ann Arbor, MI 48109-2800, gbarnes@umich.edu, 734-998-7955 (p), 734-998-8018 (f).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

to warfarin, at a willingness-to-pay threshold of \$100,000. However, patients without prescription drug coverage have a high out-of-pocket cost burden with dabigatran therapy, leading to a reduction in its cost-effectiveness when compared to warfarin therapy. In conclusion, this Markov model suggests that Medicare Part D coverage influences the cost-effectiveness of dabigatran 150 mg daily compared to adjusted-dose warfarin from multiple payer perspectives.

Keywords

anticoagulant; cost-effectiveness; cost-benefit analysis; stroke

For decades, standard therapy for non-valvular atrial fibrillation has been dose-adjusted warfarin. However, due to a narrow therapeutic window, inter-individual variability, drug-drug interactions and drug-food interactions, alternative oral agents were developed with the potential to minimize those limitations. One alternative, dabigatran 150 mg twice daily, was shown to be superior to warfarin for the prevention of stroke and systemic embolism in the RE-LY trial.¹ However, at a cost of \$295.71 per month, dabigatran is significantly more expensive than warfarin (\$3.04 per month).² Previous research demonstrated dabigatran's cost-effectiveness when compared to standard warfarin therapy from a societal perspective. These studies estimated dabigatran's incremental cost-effectiveness ratio (ICER) at \$12,000, \$45,372 and \$25,000 per quality-adjusted life year (QALY) over warfarin from a societal viewpoint.³⁻⁵ Freeman et al used a higher market price estimate for dabigatran than its eventual market price, which may account for the larger ICER estimation (\$45,372). Because of the complexity of payment for the components of each therapy (e.g. drug monitoring versus medication cost) in the US healthcare system, individual payers (Medicare and patients) may have financially divergent preferred therapies not captured in a societal perspective analysis. A better understanding of the cost-effectiveness of therapies from societal, Medicare and patient perspectives can have significant impacts in patient decision-making and healthcare policy between the newer direct oral anticoagulants and warfarin therapy. Our study aimed to assess the unique perspectives of society, Medicare and the patient in the cost-effectiveness of dabigatran 150 mg twice daily versus dose-adjusted warfarin and the influence of varying risk for ischemic stroke and Medicare Part D coverage.

Methods

Our analysis used a state-transition model of anticoagulation therapy for ischemic stroke prophylaxis in individuals ≥ 65 years of age with NVAF at increased risk of stroke (CHADS₂ score ≥ 1 or equivalent), creatinine clearance ≥ 50 ml/min and no contraindications to anticoagulation therapy. Results were expressed in QALYs, 2013 US dollars and ICERs. Model analyses used TreeAge Pro Suite 2014 (TreeAge Software, Williamstown, Massachusetts) and Microsoft Excel 2007 (Microsoft, Redmond, Washington). This analysis was based on the results from previous studies and did not use human subjects, so IRB approval was not necessary.

Using a Markov model, we compared 2 strategies for the prevention of ischemic stroke in patients with NVAF: dose-adjusted warfarin with goal INR 2.0-3.0 and dabigatran 150 mg

twice daily. The base case scenario was a hypothetical cohort of patients 65 years of age with NVAF at increased risk of stroke (CHADS₂ score 1 or equivalent), renal clearance 50 ml/min, Medicare Parts A, B and D coverage and no contraindication to anticoagulation therapy. Quality adjusted life expectancy, net costs and adverse event risks were calculated for 35 years (to age 100) or until death. Patient movement between health states occurred in 1-month cycles. During each cycle, patients could be impacted by adverse events, with changes in utility and costs for each health state transition. The following health states were included in our model: healthy with NVAF; ischemic stroke (IS): transient ischemic attack (TIA), minor, major, or fatal; intracranial hemorrhage (ICH): minor, major, or fatal; extracranial hemorrhage (ECH): minor, major, or fatal; myocardial infarction (MI): non-fatal or fatal; and death (see Appendix Figures 1 and 2 for primary and secondary model branches). Adverse events incurred direct costs and reduced quality of life. Warfarin and dabigatran anticoagulation therapy also incurred indirect costs, including the economic cost of patient time for warfarin anticoagulation management.⁶ The estimated cost of clinic visits alone was used for patients receiving dabigatran.

Patients were age 65 to represent the age of Medicare coverage initiation. Adverse event risks were obtained from the RE-LY trial and related trials of warfarin therapy for ischemic stroke prevention as shown in Appendix Table 1.¹ Mortality, IS, ICH, and MI rates were adjusted for age beginning at age 65.⁷⁻⁹ Adjustments for IS and MI rates based on CHADS₂ scores (categorized 1, 2, or 3) were performed for comparative analyses.^{10,11} Relative risks of death for each health state, including NVAF, post-MI, post-ICH, post-ECH and post-IS were included.^{12,13} We assumed patients with ICH and major ECH discontinued anticoagulation therapy and initiated lifelong aspirin therapy. Minor ECH patients discontinued anticoagulation therapy for 2 days before reinitiating anticoagulation. Patients in a minor-adverse event state who accrued a second minor-adverse event were assumed to move into the major-adverse event state associated with the initial adverse event.

Age-related baseline mortality rates for patients with NVAF were obtained through the United States Life Tables 2004, as shown in Appendix Table 1.¹⁴ We assumed increased relative risk of mortality with history of MI, IS, and ICH (1.3, 1.4, 1.97, respectively).⁷⁻⁹ Baseline utilities were adjusted for age, anticoagulation treatment and NVAF diagnosis. Utilities associated with other health states based on previously-published research (see Appendix Table 1). Due to the inconvenience of warfarin therapy, being healthy on dabigatran was given a higher utility than being healthy on warfarin, consistent with prior analyses of warfarin and dabigatran.³⁻⁵ Future costs and benefits were discounted at a rate of 3% annually.

Our model adopted societal, Medicare and patient perspectives of inpatient, outpatient and prescription medical care. Indirect costs of the economic value of patient time were included using the human capital method and the national average wage, \$19.29 per hour as obtained by the June 2006 National Compensation Survey.¹⁵ For warfarin treatment, these costs were determined by variables including the frequency and length of time related to pharmacy trips, time for diet and food preparation, time for pharmacy trips, time for blood draws, and time waiting for and attending outpatient clinic visits, including companion time and varying numbers of visits per year. We converted annual costs to monthly costs. For dabigatran, the

economic value of patient time was determined by variables such as the frequency and length of time related to pharmacy trips as well as waiting for and attending outpatient clinic visits, including companion time and varying number of visits per year. Indirect costs associated with adverse events were not calculated. Outpatient clinic visits occurred once every 3 months, with direct costs equal to the cost of 1 anticoagulation clinic visit.¹⁶

Medicare coverage in our model included estimates of Parts A, B and D reimbursement rates. Medicare Part A covered 100% of in-hospital costs beyond the first \$1,184. Medicare Part B covered 80% of all outpatient clinic visits. Medicare Part D covered 75% of prescription drug costs with the assumption that patients did not fall within the Medicare Part D coverage gap. The cost of dabigatran was determined from wholesale price.² The cost of warfarin treatment was estimated as the cost of 20 annual INR draws and medication costs as well as 90-day anticoagulation management (Current Procedural Terminology [CPT] Code 99363).¹⁷ One-time event costs for ICH, MI, gastrointestinal (GI) hemorrhage, and IS were obtained from the Agency for Healthcare Research and Quality's Healthcare Cost and Utilization Project National Inpatient Sample Database and from previously established research (see Appendix Table 1). The cost of a major ECH event was estimated as the cost of diagnosis group-related cost (DGR: 378) for major GI bleed, given that major GI hemorrhages are the most frequent major ECH. The cost of a minor hemorrhage was estimated as the cost of an outpatient clinic visit (Current Procedural Terminology code: 99212). Costs were expressed in 2013 dollars.

In sensitivity analyses, cost ranges from previously published research were included (see Appendix Table 1). All costs are inflation-adjusted to 2013 dollars. Division of costs by payer is provided in Appendix Table 2. Analyses were performed for the various perspectives across varying CHADS₂ scores and Medicare Part D coverage. Model parameters were varied over ranges established in the RELY substudy and other published studies (see Appendix Table 1). A probabilistic sensitivity analysis varied all parameters simultaneously using a Monte Carlo simulation of 10,000 iterations. Distributions for parameter inputs are described in Appendix Table 3. Threshold analysis was conducted to determine the value for the cost of dabigatran that would lead to a change in the base case results, under a willingness-to-pay (WTP) threshold of \$100,000.^{4,18} Stratified analyses were conducted to assess how the results may be sensitive to changes in the underlying patient population. These stratifications included different adverse event risks and treatment costs based on CHADS₂ score and Medicare Part D coverage, respectively.

Results

Costs, QALYs and ICERs associated with the base case scenarios for the 3 payers are shown in Table 1. Figure 1a shows how the ICER changes depending on Medicare Part D coverage and payer type (society, Medicare, patient). Under the base-case scenario (CHADS₂ score 1 and Medicare Part D), dabigatran was a cost-effective alternative to warfarin from all perspectives (societal, Medicare and patient). From Medicare's perspective, the ICER comparing dabigatran with warfarin was \$35,311 per QALY gained for patients with Part D coverage and cost-saving for patients without Part D coverage. From the patient perspective,

the ICER comparing dabigatran with warfarin was cost-saving for patients with Part D coverage and \$63,884 per QALY for those without Part D coverage.

All 3 stroke risk groups (CHADS₂ = 1, CHADS₂ = 2 and CHADS₂ = 3) accrued more QALYs with dabigatran. From Medicare's perspective, the ICER comparing dabigatran with warfarin was markedly lower in patients with higher CHADS₂ scores, as displayed in Figure 1b. From the patient perspective, dabigatran improved health outcomes and saved money when compared to warfarin, regardless of CHADS₂ score.

One-way sensitivity analyses were conducted for multiple variables related to costs, utilities and age from a societal perspective. The financial value of patient time for both warfarin therapy and dabigatran therapy included minimums of 0 given the uncertainty of how patients value time. Varying the cost of dabigatran and the economic value of patient time for warfarin therapy had the greatest impact on the cost-effectiveness of dabigatran from a societal perspective, as shown in Appendix Figure 3. Varying the cost of dabigatran had the greatest impact on the cost-effectiveness of dabigatran for patients with Medicare Part D coverage from Medicare's perspective. Varying the cost of intracranial hemorrhage events had the greatest impact on the cost-effectiveness of dabigatran versus warfarin for patients with Medicare Part D coverage from Medicare's perspective. At a cost below \$7.43 per capsule (\$445.70 per month), dabigatran 150 mg twice daily has a total healthcare cost less than warfarin with greater QALY's (negative ICER) for patients with Medicare Part D coverage from the patient perspective. Varying the economic values of patient time for warfarin and dabigatran therapies had the greatest impacts on the cost-effectiveness of dabigatran versus warfarin from the patient perspective for patients with Medicare Part D coverage. Varying the monthly cost of dabigatran had the greatest impact on the cost-effectiveness of dabigatran versus warfarin for patients without Medicare Part D coverage from the patient perspective (ICER of \$22,900 to \$77,200 per QALY). Varying the economic value of patient time for warfarin and dabigatran therapies also had significant impacts on the cost-effectiveness of dabigatran versus warfarin.

The highest price at which dabigatran 150 mg twice daily remains cost-effective compared to warfarin from a societal perspective ranges from \$515 (CHADS₂ = 1) to \$789 (CHADS₂ = 3) per month (\$8.59 to \$13.16 per capsule) at a WTP threshold of \$100,000 (Figure 2).

The probabilistic sensitivity analysis varied all parameters simultaneously. With a WTP threshold of \$100,000 per QALY, dabigatran 150 mg twice daily was cost-effective in 100% of simulations in the base case scenario from a societal perspective (Figure 3a) and from the patient perspective. Dabigatran 150 mg twice daily was cost-effective for 95.3% of simulations in the base case scenario from Medicare's perspective. With a WTP threshold of \$100,000 per QALY, dabigatran 150 mg twice daily was cost-effective in 99.7% of simulations for patients without Medicare Part D coverage from the patient perspective (Figure 3b) and 100% of simulations from Medicare's perspective.

Discussion

This study demonstrates that for patients 65 years of age with NVAf at an increased risk of ischemic stroke, dabigatran 150 mg twice daily is a cost-effective alternative to dose-adjusted warfarin for ischemic stroke prophylaxis from societal and Medicare perspectives in all scenarios at a WTP of \$100,000. Of unique interest, at a WTP threshold of \$100,000, dabigatran remains a cost-effective alternative to warfarin therapy for society and payers irrespective of prescription coverage, but the cost per QALY varies greatly depending on prescription coverage. The availability of prescription drug coverage has significant impact in the cost trade-off between dabigatran and warfarin from the patient perspectives. For patients with Medicare Part D prescription drug coverage, dabigatran was cost-saving, but for patients without Medicare Part D coverage, dabigatran therapy was costly (\$63,884 per QALY), showing the considerable effect prescription coverage may have on cost-effectiveness for patients. In particular, adults without Medicare Part D coverage represent a vulnerable population that may have reduced financial resources and therefore may be more sensitive to the higher relative cost of dabigatran.

This high cost per QALY may affect short-term prescription coverage choices and long-term medical therapy for this population. Prescription drug coverage has noteworthy effects on the cost-effectiveness from the patient perspective. This situation is unusual: a recently-developed medication became cost-saving from the patient perspective when compared to a generic drug. This held under a variety of assumptions.

Ischemic stroke risk, as estimated by the CHADS₂ score, had a noticeable effect on the ICER of dabigatran. With increasing risk of IS, dabigatran becomes even more cost-effective from societal and Medicare viewpoints. Models by Davidson, Rognoni, Sorenson and Pink reported similar results from societal perspectives in Swedish, Italian, Canadian and United Kingdom healthcare systems, respectively.¹⁹⁻²¹ As the incidence of IS increases with CHADS₂ scores, the societal and Medicare costs associated with these events rise. Dabigatran remained cost-effective at WTP thresholds of \$100,000 and \$50,000 across the spectrum of CHADS₂ scores.

We determine dabigatran to have an ICER of \$21,980 per QALY from a societal perspective for the base case. Two recent studies support our finding that, from a societal viewpoint, dabigatran is a cost-effective alternative to warfarin at a WTP threshold of \$100,000. In a study published by Kamel et al, the ICER for dabigatran 150 mg twice daily compared to warfarin was \$25,000 per QALY from a societal perspective.⁴ Similarly, Freeman et al estimated an ICER of \$12,386 per QALY when American market cost of dabigatran was included³. However, neither study utilized indirect costs of patient time or costs of anticoagulation clinic visits, and these studies included patients initiating warfarin at age 70. A separate model developed by Shah and Gage yielded an ICER of \$86,000 per QALY, far greater than the values seen by Kamel and Freeman,⁵ perhaps because the former models stratified results by the effectiveness of INR control, modeled dyspepsia as a side effect of dabigatran, and included bleeding risks and stroke risk scores. We did not stratify results by the effectiveness of INR control because we compared the probability of adverse events for the entire cohort of patients receiving warfarin therapy in the RE-LY trial.

Alternatively, previous research by Pink et al under a United Kingdom health care perspective demonstrated an ICER of €26,700 (\$35,700) per QALY, with dabigatran only being cost-effective in patients at high risk of IS.²⁰ Our results may differ for several reasons. Pink's model did not include the economic value of patient time on either therapy option, which may account for the increased ICER. Furthermore, in Pink's model, the treatment cost for an ICH was assumed to be higher due to the lack of an appropriate reversal agent.

While our research used broadly similar model structure and estimates as previous research, our inclusion of recently-published costs and the economic value of patient time had significant effects on dabigatran's cost-effectiveness. This may account for the differences in our findings from previous studies.³⁻⁵ Including the economic value of patient time, the frequency of INR lab draws and anticoagulation clinic visits increased the cost of warfarin therapy relative to dabigatran. Similarly, given over-estimation of the cost of dabigatran in previous studies, overall cost of dabigatran therapy was improved in our study.

This model had a few limitations. Notably, the efficacy of medical therapy in the RE-LY clinical trial may vary from therapy in real-world settings, which generally have reduced adherence, reduced monitoring, and increased probability of comorbidities. In particular, bleeding rates may be higher in older, unselected adults treated with dabigatran compared to the bleeding rates observed in the RE-LY trial.²² Furthermore, important information related to the proportion of patients, average cost, and average length of stay for skilled nursing facilities and nursing home care for each adverse event was unavailable. Other models, similarly, did not include this information in their analysis.^{3-5, 19-21} Another limitation is the division of Medicare Part D coverage to either complete or incomplete coverage without including scenarios with Medicare supplemental (Medigap) plans, which could create significant variations in costs of medications, medical services and medical supplies. Similarly, we did not assess the effects of patients who enter into the Medicare Part D prescription coverage gap. However, the Affordable Care Act has established plans to eliminate this coverage gap by 2020. Lastly, our analysis assumes a baseline age of 65, which may not replicate clinical scenarios when NVAF develops in younger patients.

Our study had multiple notable strengths. This analysis is the first to examine multiple payer perspectives across a range of CHADS₂ scores while also examining the impact of Medicare Part D coverage. Furthermore, this study's model allows for future manipulation of multiple variables, notably the frequency of clinic visits, INR blood draws, and anticoagulation follow-up. This model also allows for future stratification of the study population by socioeconomic status, gender and a multitude of patient-related factors. Our model includes current, updated medical therapy costs and a mixture of previously published research models. Finally, this model can be utilized to compare warfarin and dabigatran therapy with other direct oral anticoagulants.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award 2-TL1-TR000435 and the National Heart, Lung and Blood Institute under award 2-T32-HL007853-16. The funding sources did not have a role in the design, development, conduction, data management, analysis, interpretation, review and publication for this study.

Dr. Froehlich has served as a consultant for Boehringer-Ingelheim, Merk, Bristol-Meyers and Jansen along with research grant funding from Blue Cross Blue Shield of Michigan and BMS/Pfizer (Ann Arbor, Michigan). Dr. Barnes has served as a consultant for Portola along with research grant funding from Blue Cross Blue Shield of Michigan and BMS/Pfizer (Ann Arbor, MI).

References

1. Connolly SJ, Ezekowitz MD, Yusuf S, Eikelboom J, Oldgren J, Parekh A, Pogue J, Reilly PA, Themeles E, Varrone J, Wang S, Alings M, Xavier D, Zhu J, Diaz R, Lewis BS, Darius H, Diener H, Joyner CD, Wallentin L. Dabigatran versus Warfarin in Patients with Atrial Fibrillation. *N Engl J Med*. 2009; 361:1139–1151. [PubMed: 19717844]
2. Reuters, Thomson. Red book. Thomson Reuters; Montvale, NJ: 2009. p. 1032
3. Freeman JV, Zhu RP, Owens DK, Garber AM, Hutton DW, Go AS, Wang PJ, Turakhia MP. Cost-effectiveness of dabigatran compared with warfarin for stroke prevention in atrial fibrillation. *Ann Intern Med*. 2011; 154:1–11. [PubMed: 21041570]
4. Kamel H, Johnston SC, Easton JD, Kim AS. Cost-effectiveness of dabigatran compared with warfarin for stroke prevention in patients with atrial fibrillation and prior stroke or transient ischemic attack. *Stroke*. 2012; 43:881–883. [PubMed: 22308255]
5. Shah SV, Gage BF. Cost-effectiveness of dabigatran for stroke prophylaxis in atrial fibrillation. *Circulation*. 2011; 123:2562–2570. [PubMed: 21606397]
6. Jonas DE, Bryant Shilliday B, Laundon WR, Pignone M. Patient time requirements for anticoagulation therapy with warfarin. *Med Decis Making*. 2010; 30:206–216. [PubMed: 19773584]
7. Ariesen MJ, Claus SP, Rinkel GJ, Algra A. Risk factors for intracerebral hemorrhage in the general population: a systematic review. *Stroke*. 2003; 34:2060–2065. [PubMed: 12843354]
8. Agency for Healthcare Research and Quality. 2003 Nationwide Inpatient Sample. Healthcare Cost and Utilization Project; Rockville, MD: 2003.
9. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials. *Arch Intern Med*. 1994; 154:1449–1457. [PubMed: 8018000]
10. Capodanno D, Capranzano P, Giacchi G, Calvi V, Tamburino C. Increasing CHADS2 scores may attenuate the benefit of novel oral anticoagulants versus warfarin in reducing intracranial bleeding. *Internat J Cardiol*. 2012; 161:176–177.
11. Oldgren J, Alings M, Darius H, Diener HC, Eikelboom MD, Kamensky G, Reilly PA, Yang S, Yusuf S, Wallentin L, Connolly SJ. Risks for stroke, bleeding, and death in patients with atrial fibrillation receiving dabigatran or warfarin in relation to the CHADS2 score: a subgroup analysis of the RELY trial. *Ann Intern Med*. 2011; 155:660–667. [PubMed: 22084332]
12. Van Wijk I, Kappelle LJ, van Gijn J, Koudstaal PJ, Franke CL, Vermeulen M, Gorter JW, Algra A. Long-term survival and vascular event risk after transient ischaemic attack or minor ischaemic stroke: a cohort study. *Lancet*. 2005; 365:2098–2104. [PubMed: 15964446]
13. Zhang Z, Fang J, Gillespie C, Wang G, Hong Y, Yoon PW. Age-specific gender differences in in-hospital mortality by type of acute myocardial infarction. *Am J Cardiol*. 2012; 109:1097–1103. [PubMed: 22245410]
14. Arias E. United States life tables, 2004. *Natl Vital Stat Rep*. 2007; 56:1–40. [PubMed: 18274319]
15. Jonas DE, Bryant Shilliday B, Laundon WR, Pignone M. Patient time requirements for anticoagulation therapy with warfarin. *Med Decis Making*. 2010; 30:206–216. [PubMed: 19773584]

16. Lafata JE, Martin SA, Kaatz S, Ward RE. The cost-effectiveness of different management strategies for patients on chronic warfarin therapy. *J Gen Intern Med.* 2000; 15:31–37. [PubMed: 10632831]
17. Davidson T, Husberg M, Janzon M, Oldgren J, Levin L-Å. Cost-effectiveness of dabigatran compared with warfarin for patients with atrial fibrillation in Sweden. *Eur Heart J.* 2012; 24:177–183. [PubMed: 22733833]
18. Neumann PJ, Cohen JT, Weinstein MC. Updating cost-effectiveness – the curious resilience of the \$50,000-per-QALY threshold. *N Engl J Med.* 2014; 371:796–797. [PubMed: 25162885]
19. Sorensen SV, Kansal AR, Connolly S, et al. Cost-effectiveness of dabigatran etexilate for the prevention of stroke and systemic embolism in atrial fibrillation: A Canadian payer perspective. *Thromb Haemost.* 2011; 105:908–919.
20. Pink J, Lane S, Pirmohamed M, Hughes DA. Dabigatran etexilate versus warfarin in management of non-valvular atrial fibrillation in UK context: quantitative benefit-harm and economic analyses. *BMJ.* 2011; 343:d6333. [PubMed: 22042753]
21. Hernandez I, Baik SH, Piñera A, Zhang Y. Risk of bleeding with dabigatran in atrial fibrillation. *JAMA Intern Med.* 2014; 175:18–24. [PubMed: 25365537]
22. Mark DB, Knight JD, Cowper PA, Davidson-Ray L, Anstrom KJ. Long-term economic outcomes associated with intensive versus moderate lipid-lowering therapy in coronary artery disease: results from the Treating to New Targets (TNT) Trial. *Am Heart J.* 2008; 156:698–705. [PubMed: 18926150]

Figure 1a. Incremental Cost-Effectiveness Ratio for Dabigatran Versus Warfarin, With and Without Medicare Part D, by Payer

Incremental Cost-Effectiveness Ratio (ICER) of dabigatran versus dose-adjusted warfarin for patients with and without Medicare Part D insurance, assessed by paper type (society, Medicare and patient). Cost-saving describes a scenario with higher quality-adjusted life years gained and lower overall costs when receiving treatment A versus receiving treatment B.

Figure 1b. Incremental Cost-Effectiveness Ratio For Dabigatran Versus Warfarin, Across Varying CHADS₂ Scores and Payers

Incremental Cost-Effectiveness Ratio (ICER) of dabigatran versus dose-adjusted warfarin for patients with Medicare Part D insurance across varying CHADS₂ scores. Cost-saving describes a scenario with higher quality-adjusted life years gained and lower overall costs when receiving treatment A versus receiving treatment B. Assume Part D Coverage

Figure 2. Maximal Societal Value of Dabigatran, by Willingness-to-Pay Threshold

Dollar value represents that maximum monthly cost of dabigatran at which dabigatran 150 mg twice daily remains cost-effective compared to warfarin from a societal perspective, using both \$50,000 and \$100,000 willingness-to-pay thresholds. Base CHADS₂ score reflects the mean CHADS₂ score of patients in the RE-LY trial.

Figure 3a.

Cost-Effectiveness Acceptability Curve – Patient with Medicare Part D, Societal Perspective

Figure 3b.
Cost-Effectiveness Acceptability Curve – Patient Without Medicare Part D, Patient Perspective

Table 1

Projected Costs and QALYs for Patients with Nonvalvular Atrial Fibrillation, Patient with Medicare Part D Coverage, by Varying Perspective

Payer	Therapy	Cost, \$	QALYs	Marginal Cost per QALY, \$
Society	Warfarin	95,528	8.998	
	Dabigatran	104,186	9.392	21,980
Medicare	Warfarin	61,586	8.998	
	Dabigatran	75,496	9.392	35,311
Patient	Warfarin	33,942	8.998	
	Dabigatran	28,691	9.392	*Cost-saving

QALY = quality-adjusted life year

* Cost-saving = higher quality-adjusted life years gained and lower overall costs when receiving treatment

A versus receiving treatment B