Published in final edited form as:

Popul Environ. 2015 September ; 37(1): 22–43. doi:10.1007/s11111-015-0230-y.

Local Variability in the Impacts of Residential Particulate Matter and Pest Exposure on Children's Wheezing Severity: A Geographically Weighted Regression Analysis of Environmental Health Justice

Sara E. Grineski [Associate Professor of Sociology],

Department of Sociology and Anthropology, University of Texas at El Paso, 500 W. University Ave. El Paso TX 79968, USA, segrineski@utep.edu, 915-747-8471 (tele), 915-747-5505 (fax)

Timothy W. Collins [Associate Professor of Geography], and

Department of Sociology and Anthropology, University of Texas at El Paso, 500 W. University Ave. El Paso TX 79968, USA

Hector A. Olvera [Research Assistant Professor]

Center for Environmental Resource Management & School of Nursing, University of Texas at El Paso, 500 W. University Ave. El Paso TX 79968, USA

Abstract

Two assumptions have underpinned environmental justice over the past several decades: 1) uneven environmental exposures yield correspondingly unequal health impacts and 2) these effects are stable across space. To test these assumptions, relationships for residential pest and PM_{2.5} exposures with children's wheezing severity are examined using global (ordinary least squares) and local (geographically weighted regression [GWR]) models using cross-sectional observational survey data from El Paso (Texas) children. In the global model, having pests and higher levels of PM_{2.5} were weakly associated with greater wheezing severity. The local model reveals two types of asthmogenic socio-environments where environmental exposures more powerfully predict greater wheezing severity. The first is a lower-income context where children are disproportionately exposed to pests and PM_{2.5} and the second is a higher-income socio-environment where children are exposed to lower levels of PM_{2.5}, yet PM_{2.5} is counterintuitively associated with more severe wheezing. Findings demonstrate that GWR is a powerful tool for understanding relationships between environmental conditions, social characteristics and health inequalities.

Keywords

Environmental health justice; health disparities; asthma; geographically weighted regression; El Paso; Texas

1. Introduction

Myriad studies have shown that minority and lower income children suffer disproportionately from a host of health problems (e.g., Mehta, Lee, & Ylitalo, 2013). Minority and lower income neighborhoods also tend to be burdened by the unequal distribution of hazard exposures (Downey, 2006). Studies addressing these two hypothetically linked phenomena have remained largely distinct and a key question remains unanswered: *What is the role of exposure to environmental hazards in health disparities?* Studies investigating this linkage using a spatial approach have relatively recently begun to emerge under the rubric of “Environmental Health Justice” (EHJ) (Chakraborty & Maantay, 2011; Corburn, 2005; Grineski, Collins, Chakraborty, & McDonald, 2013). This is in response to decades of environmental justice research that has extended from the poorly supported assumption that uneven environmental exposures yield correspondingly unequal health impacts and that these effects are stable across space.

EHJ researchers have reported that environmental degradation plays a role in predicting geographic inequalities in health outcomes (Grineski, 2007; Grineski et al., 2013; Jephcote & Chen, 2012, 2013; Pearce, Richardson, Mitchell, & Shortt, 2010, 2011; Richardson, Pearce, Tunstall, Mitchell, & Shortt, 2013), but not in the monotonic way that many researchers may have assumed it would. For example, in the UK, Pearce et al. (2010) found that the relationship between environmental deprivation and mortality was strongest in the most affluent areas and weakest in the poorest areas. In California, Hispanics experienced the greatest exposure to ozone and PM_{2.5} as compared to blacks and whites, but they did not have the highest excess attributable risk for hospitalizations due to pollution exposure (Hackbarth, Romley, & Goldman, 2011). Grineski et al. (2013) found a significant association between air toxics and children’s respiratory infections in El Paso neighborhoods after adjusting for relevant controls, but they did not find the same for asthma.

In addition to counterintuitive findings, EHJ work has been characterized by a reliance on secondary data sources. On the health side (see Wheeler & Ben-Shlomo, 2005 for an exception), studies have relied on hospitalization data (e.g., Grineski et al., 2013; Jephcote & Chen, 2012) and mortality records (e.g., Pearce et al., 2010), which is attributable to the public availability and low cost of these sources of health data. While generating important insights, this work has focused attention on serious health outcomes and has tended to require the analysis of aggregated data for areal units (e.g., neighborhoods). In terms of air quality, modeled criteria air pollution surfaces are publically available in the UK (Jephcote & Chen, 2012; Pearce et al., 2010). These types of surfaces are not available from US governmental agencies, and therefore they are rarely used by EHJ researchers working in the US (see Grineski, 2007 for an exception). This is a critical limitation given the deleterious health effects of these pollutants (Samet & Krewski, 2007).

Recent studies raise the possibility that pollution-health linkages might vary widely across space (Pearce et al., 2010; Richardson et al., 2013). This makes geographically weighted regression (GWR) particularly suited to investigating EHJ, even though it is an “underused EJ technique” (Jephcote and Chen, 2012, p. 141). Unlike spatial autoregressive models, which account for spatial autocorrelation in generating parameter estimates (Chakraborty,

2011), GWR models how relationships between variables vary across space (Fotheringham, Brundson, & Charlton, 2002). To our knowledge, only three previous EJ studies focused on environmental hazards have used GWR (Gilbert & Chakraborty, 2011; Jephcote & Chen, 2012; Mennis & Jordan, 2005). One study analyzed data from New Jersey and concluded that global models used by many are likely insufficient for modeling environmental injustice (Mennis & Jordan, 2005). Most recently, Jephcote and Chen (2012) found that PM_{10} and ethnic minority status were stronger predictors of children's asthma hospitalization rates in the inner city of Leicester (UK) than they were in outlying areas. These findings underscore the point that environmental exposures may not impact people's health the same way in all locations and that the ways in which exposures impact health may be surprising.

While the majority of EJ studies focus on outdoor environments (e.g., factory releases and criteria air pollution), indoor environmental exposures can also be considered environmental injustices (Grineski & Hernandez, 2010). Poor home environments have less often been considered in terms of environmental injustice in the past (see Kraft & Scheberle, 1995; Landrigan, Rauh, & Galvez, 2010 for exceptions), possibly because there is a tendency to assume that in-home conditions are products of independent household decisions, rather than power-laden products of social processes (Grineski & Hernandez, 2010). Certainly, indoor environmental exposures (e.g., roaches, rodents, mold) are more prevalent in substandard housing inhabited by the poor (Matte & Jacobs, 2000) and have been linked to respiratory symptoms at the individual (Lanphear, Aligne, Auinger, Weitzman, & Byrd, 2001) and neighborhood levels (Grineski, 2007).

As is the case with air pollution, there is preliminary evidence that in-home exposures may not impact all children the same way. Low-income white children in New York (state) had higher rates of indoor risk factors than did middle class white children, and these risk factors were correlated with elevated overnight epinephrine, norepinephrine, and cortisol, but *only* in the low-income sample (Evans & Marcynyszyn, 2004). Pooling five survey datasets, researchers found that the odds of exposure to household pests was significantly associated with asthma only for children born in the US, and not for children born outside the US (Woodin, Tin, Moy, Palella, & Brugge, 2011). Those studies examined variation based on social characteristics – specifically, income and nativity. It remains unclear how the strength of the association between indoor exposures and respiratory health might vary across urban space.

This study makes several advances upon previous studies. First, we utilize individual-level data, which is rare in EJ studies, thus avoiding the problem of the ecological fallacy. Second, as opposed to relying on secondary hospitalization or mortality records, we use a wheezing symptoms severity measure, thus capturing a more common and broadly relevant health problem. This enables us to address Jephcote and Chen's (2012, p. 142) recent call "for future EJ research to develop upon [previous] GWR studies, through applying measurements of actual health events and exploring a wider range of cardiorespiratory conditions influenced by short-term exposures." Asthma hospitalizations, which are more often used in these types of studies, are relatively rare events; for example the asthma hospitalization rate is 27 per 10,000 children in the US (Akinbami, 2007). Third, we utilize a $PM_{2.5}$ (particulate matter less than 2.5 micrometers in diameter) surface, generated through

primary data collection. This allows us to move beyond US EPA-provided data, and to analyze an important traffic-associated criteria air pollutant for which data are not currently publicly available in the US. The associations between $PM_{2.5}$ and respiratory problems have been well-documented and inhalation of this pollutant has been linked to inflammatory responses and pulmonary oxidative stress (Hansen et al., 2012). Fourth, we consider both indoor (pest exposure) and outdoor ($PM_{2.5}$) environmental conditions through an EHJ framework, which is rarely done.

As per prior EHJ research (Gilbert & Chakraborty, 2011), our analysis approach relies on both aspatial and spatial modeling. We answer the following two research questions: 1) What are the global relationships for residential pest exposure and $PM_{2.5}$ with children's wheezing severity adjusting for relevant controls? 2) What is the degree of local spatial variation in the contribution of both residential pest exposure and $PM_{2.5}$ to children's wheezing severity adjusting for the relevant controls?

2. Data & Methods

2.1 Study Context

The study took place in El Paso County, Texas, which has an estimated population of 830,000 residents. According to the US Bureau of the Census, in 2011, 81% of its residents were Hispanic (compared with 17% for the US and 38% for TX), while smaller percentages were non-Hispanic white (14%) and non-Hispanic black (4%). El Paso County had a lower median household income (2011 US \$36,333) than the State of Texas (2011 US \$49,391) and the US (2011 US \$50,502) with a poverty rate of 24%, which was higher than the national rate (16%). In previous studies in this city, researchers have found relatively modest associations between air pollutants (including $PM_{2.5}$) and respiratory health effects (Grineski, et al., 2011; Sarnat et al., 2011; Svendsen et al., 2012; Zora et al., 2013).

2.2 Survey Data Collection

Social and health data were collected through a cross-sectional, observational mail survey that was approved by our university's Institutional Review Board. The closed-ended questionnaire was sent to all primary caretakers (parents and guardians) of 4th and 5th graders attending school in the El Paso Independent School District (EPISD). With more than 64,000 students across 94 campuses, the EPISD is the 10th largest district in Texas and the 61st largest district in the US (EPISD, 2013). Children in the 4th and 5th grade from all 58 elementary schools are represented in the dataset.

Surveys were conducted to obtain the highest achievable response rates by personalizing communication, following-up with non-respondents, and offering incentives (Dillman, Smyth, & Christian, 2009). All survey materials were provided to households in English and Spanish in three waves during May of 2012. Ultimately, 6,295 primary caretakers received surveys at their home address and 1,904 surveys were returned for a 30% response rate. Respondents were primarily mothers (82%), with the next largest shares being fathers (10%) and grandparents (4%). Descriptive statistics for the percentages of surveyed children who are male (49.9% vs. 51.4% in EPISD), Hispanic (82.2% vs. 82.6% in EPISD) and

economically disadvantaged (60.4% vs. 71.1% in EPISD) indicate that the is sample is generally representative of the EPISD student population (EPISD, 2013).

2.3 Selection Criteria

Of the 1904 children surveyed, 1736 were selected for inclusion in this study; 162 were excluded due to missing data for the analysis variables and 6 were excluded as spatial outliers. We focus on children because of their sensitivity to air pollution. Childhood is a critical time in the development and maturation of the cardiorespiratory system, which is highly susceptible to the absorption of toxins (Jephcote & Chen, 2013). A child's lung surface area is significantly larger relative to body mass than an adult's; children can breathe up to 50% more air per kilogram of body weight. Children also tend to spend more time outdoors participating in activities that increase their breathing rates. When coupled with exposure to air pollutants, these factors create conditions conducive to damaging or stunting the development of children's cardiorespiratory systems, creating health problems which can prevail throughout adulthood (Schwartz, 2004).

2.4 Dependent Variable

The dependent variable is a composite measure of six wheezing measures based on data collected using International Study of Asthma and Allergies in Childhood (ISAAC) (ISAAC Steering Committee, 2012) and National Asthma Survey questions (O'Connor et al., 2008). From the ISAAC, we used the following questions: 1) In the last 12 months, has the wheezing ever been severe enough to limit your child's speech to only one or two words at a time between breaths? (1=yes, 0=no); 2) In the last 12 months, has the child had wheezing or whistling in the chest when he/she did not have a cold or the flu? (1=yes, 0=no); 3) In the last 12 months, has your child's sleep been disturbed due to wheezing? (1=yes, 0=no); 4) In the last 12 months, has your child had a dry cough at night apart from cough associated with a cold or chest infection? (1=yes, 0=no); and 5) Has the child ever been told by a doctor or health professional that he or she has asthma? (1=yes, 0=no). From the National Asthma Survey, we used this question: 6) Symptoms of asthma include coughing, wheezing, shortness of breath, chest tightness or phlegm production when someone does not have a cold or respiratory infection. How long has it been since your child had any symptoms of asthma? (1=less than 1 year ago, 0 = more than 1 year ago). We created a composite measure of current "wheezing severity" using principal components analysis on these six variables (see Table 1 which includes factor loadings), after standardizing all items. The Eigen value for the one component was 2.96 and it explained 49.4% of the variance. Due to skewness and kurtosis, we used a natural log transformation on this variable (after first adding 1 to make all values positive). Descriptive statistics are presented in Table 2.

2.5 Independent Variables

The primary variables of interest are residential pest (indoor) and PM_{2.5} (outdoor) exposure. Pest is a dichotomous variable which is coded 1 if the caretaker reported being troubled by rats (1%), ants (18%), mice (2%), spiders (6%), cockroaches (14%), termites (1%) and/or another pest (3%) inside the home in the past 12 months, and 0 if she did not (55%). Biologic agents, including allergens from cockroaches and rodents are among the most prominent environmental factors implicated in asthma morbidity and poor housing

conditions are associated with exposure to these asthma inducing biologic agents. Roaches and rodents are also associated with excess moisture in the home, which can also trigger wheezing and asthma (Derose, Bahney, Lurie, & Escarce, 2009). Indoor exposures play at least two roles in asthma: (1) as a risk factor for genetically susceptible individuals and (2) as a source of ongoing airway inflammation and hypersensitivity to other irritants (Rauh et al. 2008). In using this variable, we assume that reporting “being troubled” by one or more of the pests maps to a problem with pests in the home, and not just the occasional spider or ant and that it is associated with poor housing conditions more generally. Also, the “pest” variable was more strongly correlated with wheezing severity than was roaches alone.

Residential PM_{2.5} values for outdoor environments at each child’s home site were extracted from a 2012 exposure surface created via land use regression (LUR) modeling for this project (as per Jerrett et al., 2007; Olvera et al., 2012). PM_{2.5} measurements were collected via a 26 site monitoring network designed using a location-allocation approach (as per Kanaroglou et al., 2005). The location-allocation method used an existing exposure surface from 2006–2009 to determine an optimal monitoring network (Olvera et al., 2012). We collected one 14-day averaged PM_{2.5} sample at each site per season (n=4) during 2012. Monitoring began in May 2012 to coincide with the survey. PM_{2.5} samples were collected on Teflon filters with multi-stage impactors and concentrations were determined via gravimetric analysis (Olvera et al., 2012). The four seasonal PM_{2.5} concentrations were averaged to produce annual estimates, which are appropriate for use in LUR models (Gerard Hoek et al., 2002). A linear regression model was built with PM_{2.5} at each of the 26 sites as the dependent variable and surrounding land use, traffic, and physical characteristics as predictors (i.e., traffic counts, vehicle miles traveled, land use, property values, population density, distance to the international border, and elevation, summarized for circular areas around the monitoring locations).

The relative fit of the model was determined by a R Square of .458, which is good for these types of models (G. Hoek et al., 2008; Javier, Wise, & Mendoza, 2007; Olvera et al., 2012). The absolute fit represented by the root mean square error was obtained via a leave-one-out bootstrap analysis for 1000 samples (Isakov, Johnson, Touma, & Özkaynak, 2012; Javier et al., 2007; Rose, Cowie, Gillett, & Marks, 2010) and was found to be 0.93 (95% CI: 0.56–1.09). The samples were generated via a “leave some out” cross validation technique. Specifically, we randomly sampled 100 different observation sets from our 26 site observations by leaving out up to 3 sites each time. Technically we could have sampled 2600 different samples from our 26 sites, but for practical purposes we stopped at 1000. The samples were used to generate the model and the rest to test it. This validation method has been widely used in LUR studies and has been shown to be adequate and robust for such purposes (Isakov et al., 2012; Johnson, Isakov, Touma, Mukerjee, & Özkaynak, 2010; Parenteau & Sawada, 2010; Rose, Cowie, Gillett, & Marks, 2010). Considering a mean PM_{2.5} concentration of 7.2 µg/m³ across the region, the accuracy of the estimates is ±13% (95% CI: ±7.2%–±15%). Additional details of the monitoring procedures and the LUR modeling technique used can be found elsewhere (Jerrett et al., 2007; Olvera et al., 2012).

This approach generated PM_{2.5} values for 2193 points on a 500 m grid. We then used inverse distance weighting (IDW) with a distance decay function of 2 to create a continuous

PM_{2.5} surface (see Figure 1) and each child was given the value of PM_{2.5} corresponding to his or her home location. We used 2 as the coefficient because it produced a robust PM_{2.5} surface for the United States in a previous study (Al-Hamdan et al., 2006). Also, the IDW technique was used to interpolate PM_{2.5} values estimated via the LUR model over a grid of 500 m resolution. Hence, the value of 2 ensured that the contributions of more distant observations to the weighted interpolated value were very small. In the analysis, a natural log transformation was used to correct for skewness and kurtosis in PM_{2.5}; a standardized version of this variable is used in the models. The Pest and PM_{2.5} variables are summarized in Table 2.

2.6 Control Variables

Five control variables were selected based on relevant literature and contextual relevance to El Paso; we also prioritized selecting variables for which the proportions of missing data were lower being that GWR software cannot handle multiply imputed datasets at this time. These control variables include sex (Wright, Stern, Kauffmann, & Martinez, 2006); whether the biological mother has asthma and/or allergies (Hryhorczuk et al., 2009); and whether the child has allergies or not (Holt, Theall, & Rabito, 2013; Kocevar et al., 2005). We used an indicator of the family having postponed or avoided seeking health care for the child because of concerns about cost. We do not adjust for household income or parental education due to missing data. This variable represents a SES-related access to care indicator. Given the Hispanic majority context of this study and lower rates of asthma among this population, we include an indicator of Hispanic ethnicity (Padilla, Hamilton, & Hummer, 2009); we do not account for membership in any other racial/ethnic groups due to small counts. Descriptive statistics for all variables are included in Table 2.

3. Analysis Methods

These data are analyzed in two steps, following other GWR EHJ studies (Gilbert & Chakraborty, 2011; Jephcote & Chen, 2012) and in accordance with the two research questions. The starting point for development of a GWR model is the ordinary least squares (OLS) multiple regression equation that expresses the relationship between the dependent variable and a combination of independent variables simultaneously in a single model (Gilbert & Chakraborty, 2011). We first used OLS, an aspatial multivariate regression technique, to identify the important overall predictors of wheezing severity in El Paso. This provides what researchers who conduct spatial regression analysis term “global” findings because OLS models assume that the correlations are constant over space. This means that every independent variable has one regression coefficient that maps to the average situation for all the observations in the study area (Tu, Tu, & Tedders, 2012).

Second, we used geographically weighted regression (GWR) to uncover “local” spatial relationships. The significance of local spatial relationships is predicated on the concept of spatial non-stationarity, which means that “the measurement of a relationship depends... on where the measurement is taken” (Fotheringham et al., 2002, p. 9). To model non-stationarity, we use GWR in a geographic information system (GIS) to calculate individual regression equations for each data point, using the surrounding points (Mennis & Jordan,

2005). Points, in our case, are geocoded addresses of children participating in this study. GWR uses a distance decay function, which assumes that observations closer to a given point will have stronger influences on the local parameter than points further away (Tu et al., 2012).

Either a fixed or adaptive kernel bandwidths can be used to generate the local parameter estimates. Fixed kernels rely on a constant bandwidth for all the observations, while adaptive kernels modify the size of the bandwidth based on spatial variations in the density of observations. With adaptive kernels, longer bandwidths are used in areas where data points are sparser, and shorter bandwidths are employed in areas with a greater density of points (Tu et al., 2012). See Figure 2 for visual display of the approximate locations of children's home sites (points) under study. Clearly the density of observations varies over the study area; thus, the adaptive kernel bandwidth was used. Following others (Chalkias et al., 2013; Tu et al., 2012), the optimal bandwidth was determined by minimizing the corrected Akaike Information Criterion (AICc). Additional details regarding GWR (including equations) can be found in the literature (Fotheringham et al., 2002; Mennis & Jordan, 2005).

We used the readily available ArcGIS 10 Spatial Statistics tools to run both the global and local models. For OLS models, ArcGIS provides the parameter estimates, standard errors, t -statistics and p -values as well as robust versions of these measures; a measure of multicollinearity, the Koenker statistic (significance indicates robust p -values should be used and that relationships between some or all of the independent variables and the dependent variable are non-stationary); model fit statistics (e.g., R^2) and the ability to test model standardized residuals for spatial autocorrelation. In our case, these OLS diagnostics revealed the absence of a multicollinearity problem, non-normally distributed residuals, and a significant Koenker statistic. For that reason, robust p -values are presented in Table 3. The residuals did not exhibit significant autocorrelation based on a global univariate Moran's I test ($I=-0.008$, $p=0.866$) indicating that the OLS model was appropriate for these data. However, because of our theoretical interest in exploring non-stationarity (i.e., how relationships vary over space) as per research question 2, we ran a GWR model as the next step.

For the GWR results, ArcGIS produces a local parameter estimate, a local R^2 value and a local standardized residual for each point in the dataset; model fit statistics (e.g., R^2) are also provided for the model as a whole. In our model, a Moran's I test of the residuals revealed insignificant clustering ($I=-0.003$, $p=.959$) meaning that there were not spatial dependencies in the residuals. A notable aspect of running GWR in ArcGIS is the fact that no p -values are calculated for the individual parameter estimates. This contrasts with OLS where it is conventional to test whether parameter estimates are different from 0 using a t -test. Utilizing such tests in GWR raises the problem of multiplicity (Charlton & Fotheringham, 2009). It would be inappropriate to carry out 1736 individual significance tests for each of the 7 variables in the model since, at a 95% significance level, 5% ($n=608$) would hypothetically be significant at random. Fotheringham et al. (2002) have suggested a Bonferroni correction to the significance level, but Charlton and Fotheringham (2009) report

that Bonferroni is overly conservative and argue that the answer to the multiplicity problem is an avenue for continued research. This issue is beyond the scope of this study.

Setting aside the issue of statistical significance, we analyzed the GWR-generated parameter estimates for the Pest and PM_{2.5} variables in two ways. First, we used a standard deviation (SD) break (0.04196 for Pest and 0.0198 for PM_{2.5}) to map where the parameter estimates were relatively high (1 or more SD above the mean and positive) and low (1 or more SD below the mean and negative). This allowed us to visualize where these two predictors were more closely related to wheezing in the district. We then overlaid these data on a neighborhood map of mean household income from American Community Survey 2006–2011 block group data.

Second, to better understand the local conditions that give rise to our findings, following an approach taken by Chalkias et al. (2013), we characterized the attributes of children for whom PM_{2.5} or Pest was a relatively important positive predictor (1 or more SD above mean) and those for whom PM_{2.5}/Pest was a relatively important negative predictor (1 or more SD below mean) of wheezing severity. Chalkias et al. (2013) used this to determine the range of income, green space and population density that led to an education level coefficient (scaled so that high values matched lower levels of education) that was negative, slightly positive and positive in relation to obesity.

To better systematize this type of comparison, we employed two sets of independent samples t-tests (one for Pest and one for PM_{2.5}), grouping the cases based on whether the local parameter was 1 or more SD above mean or 1 or more SD below mean. This provides us with the ability to compare mean scores for the original variables for the two groups. We also conducted a post-hoc independent samples t-test within the “1 or more SD above the mean for PM_{2.5}” group because PM_{2.5} was a relatively important positive predictor in two distinct geographic areas of the district. Therefore, we compared these two PM_{2.5} risk groups to each other in terms of mean scores for the original variables.

4. Results

4.1 Global Model

In the OLS model (see Table 3A), the variables of interest – Pest and PM_{2.5} – exhibit modest (insignificant) positive relationships with greater wheezing severity. Being male, having postponed or not sought health care due to concerns about cost, having a mother with asthma and/or allergies, and having allergies were statistically significant risk factors.

4.2 Local Model

Parameter statistics for the GRW model are presented in Table 3B. We found that 55% of cases had a local R² that was higher than the global R² for the OLS model, showing that for over half of children, the GWR model showed improved performance over the OLS model. Overall, the model fit for the OLS and GWR models was quite similar (see Table 3, “Model Fit”). Considering the map of the local R² values (see Figure 2), we can see that the children for whom the GWR model fits best are in the “south side” and “northeast”. Model fit is worst among “west side” children, especially those closest to the mountains.

To visually isolate where Pest and residential PM_{2.5} are important predictors, standard deviation maps for the GWR parameter estimates are presented in Figures 3 and 4. In Figure 3, the black dots represent children for whom Pest was a relatively important positive predictor of wheezing; in other words, black dots indicate that Pest was a risk factor. White dots represent children for whom Pest was a relatively important negative predictor of wheezing; in this case, white dots illustrate that Pest was a protective factor. In Figure 4, the black dots represent children for whom PM_{2.5} was a relatively important positive predictor of wheezing; in this case, black dots indicate that PM_{2.5} was a protective factor. White dots represent children for whom PM_{2.5} was a relatively important negative predictor of wheezing; in other words, white dots illustrate that PM_{2.5} was a protective factor. Looking at these patterns overlaid on a neighborhood map of mean household income reveals that there is correspondence between children with a pest parameter that is 1 or more SD above the mean and the poorer areas of town: the “downtown”, “south side” and the “northeast”. The pest variable is actually “protective” (increased odds of pest, less wheezing) among some children living in the wealthier “west side”. PM_{2.5} is an important positive predictor among children on the “west side”, which is relatively affluent, and the “south side”, which is one of the poorer parts of the district. PM_{2.5} is “protective” in the farthest reaches of the “northeast,” where neighborhood incomes are moderate to high.

Table 4 shows results of the t-test analyses characterizing the attributes of children for whom their pest or PM_{2.5} parameter was 1 or more SD above the mean vs. 1 or more SD below the mean. Children for whom Pest was a relatively important positive predictor had significantly higher levels of PM_{2.5} and a significantly greater likelihood of being Hispanic (81% vs. 68%). They also had a greater likelihood of having pests (although this finding was not quite significant: 49% of children in the positive group vs. 38% of children in the negative group). Children for whom residential PM_{2.5} was a relatively important positive predictor had significantly higher levels of PM_{2.5} as well as greater odds of pest exposure (46% in the positive group as compared to 34% in the negative group). They were significantly more likely to be Hispanic (84% in the positive groups as compared to 46% in the negative group) and not to have allergies (51% in positive group vs 61% in the negative group) than children for whom residential PM_{2.5} was a negative predictor.

An examination of Figure 4 reveals that there are two areas in which PM_{2.5} was important and positive: a “western” risk group, where neighborhood incomes are relatively high, and an “eastern” risk group encompassing the “south side” and the adjacent southern portion of the “northeast,” where neighborhood incomes are low. Figure 3 does not reveal a similar pattern for pest exposure, as all children for whom Pest was a relatively important positive predictor are located in the eastern half of the district. Considering the two distinct zones wherein PM_{2.5} was a relatively powerful predictor of more severe wheezing, Table 5 presents results for a t-test comparing children in the two groups. Children in the “west side” group were significantly more likely to be non-Hispanic (78% Hispanic vs. 85% Hispanic), to have lower odds of having pests (36% vs. 54%), and to have lower levels of PM_{2.5} (below the mean vs. above the mean) as compared to children in the “eastern” group (see Table 5).

5. Discussion

In the global model, the Pest and PM_{2.5} variables were positively associated with wheezing severity, but not significantly. Being male, postponing health care due to concerns about cost, having allergies, and having a mother with asthma and/or allergies were significant risk factors for wheezing severity, which closely aligns with the literature (Carlson & Stroebel, 2001; Holt et al., 2013; Wright et al., 2006). In terms of situating the pest exposure findings within the extant literature, an analysis which pooled cross-sectional data from five studies revealed that exposure to any home pests was relatively weakly associated with asthma (Woodin et al., 2011). In comparing our PM_{2.5} findings to other El Paso studies, which used different methodological approaches, we found similar associations. A case crossover study found that a 10 unit increase in PM_{2.5} was associated with 1–2% (non-significant) daily increase in hospitalizations from asthma and bronchitis (Grineski et al., 2011). An insignificant but positive relationship between higher levels of PM_{2.5} and worse asthma control was also found among a sample of 36 asthmatic children at two elementary schools (Zora et al., 2013). At the same two El Paso schools two years earlier, researchers found that an interquartile increase in PM_{2.5} was associated with an insignificant <1% increase in airway inflammation for the 30 asthmatic children under study (Sarnat et al., 2011).

However, just because results in this study indicate that social and biological factors – such as postponing care due to concerns about cost or having a mother with asthma – were more important predictors of wheezing severity in the global model than indoor and outdoor environmental factors does not mean that environmental conditions are unrelated to health outcomes. Interpreting results from EHJ analyses must be done carefully. Insignificant and significant results can illuminate the social and environmental structure of health disparities in different contexts (Grineski et al., 2013). Insignificant regression findings do not necessarily mean that environmental exposures or social inequalities have no influence on health. In this case, insignificant findings for Pest and PM_{2.5} overall concealed heterogeneity in effects across the school district.

Local results suggest the working hypothesis that the association between air pollution and respiratory problems may vary based on socio-environmental context. This study reveals two types of asthmogenic socio-environments. The first is a lower-income context where children are disproportionately exposed to pests and PM_{2.5} and where both exposures are positively associated with wheezing, especially for Hispanic children. In this situation, pests and PM_{2.5} appear to synergistically amplify wheezing severity. This maps to a classic multiple jeopardies/environmental injustice model whereby mutually reinforcing social, environmental, and health disadvantages can be observed to co-locate in urban space. Something similar was also found in Leicester, where pollution and racial/ethnic minority status were stronger predictors of asthma hospitalization in the inner city than in the surrounding areas (Jepcohte and Chen 2012). While not focused on pests, there is evidence that air pollution can combine with other indoor exposures – such as in-home endotoxin levels in Cincinnati (Ryan et al., 2009) and smoking in Beijing (Xu & Wang, 1998) – to have synergistic effects on health.

The second is a higher-income socio-environment where children are less likely to have in-home pests or to be Hispanic; children in this situation are exposed to lower levels of $PM_{2.5}$, yet air pollution exposure is counterintuitively associated with more severe wheezing. We hypothesize that, in this context, the air pollution-wheezing linkage is attributable to residents' inability to control exposure to outdoor air pollution relative to other asthma triggers, including indoor pests. This runs counter to the multiple jeopardy model and, from an environmental injustice perspective, is unexpected. However, others have found positive relationships between environmental hazards and health effects in affluent/less polluted areas (Pearce et al., 2012; Tu, Tu and Tedders, 2012). In this type of socially advantaged context, air pollution may be a relatively important correlate of respiratory problems, even though levels of exposure are typically lower, because the relationship between pollution and health is not complicated by social deprivation and the challenges of poverty. Qualitative inquiry with parents of asthmatic children revealed that while wealthier parents demonstrated a much greater ability to control their children's home environments than poor parents, they found it quite difficult to protect their children from outdoor air pollution (Grineski, 2009). The emergence of two distinct types of asthmogenic socio-environments where outdoor air pollution exposure exerts a more powerful influence highlights the value of GWR for understanding fine-scale spatial variability in contextual determinants of health problems.

5.1 Limitations

While the GWR model can reveal spatial variations in the influence of variables on an outcome, a limitation of the technique is that it does not suggest the source of the variation (Wheeler & Páez, 2010). Interpretation must be done carefully, using contextual knowledge of the study area (Chalkias et al., 2013). The study relies only on a $PM_{2.5}$ exposure surface; we did not have access to other pollutant surfaces which limits our ability to generalize to other pollutants. Our inability to consider other environmental exposure surfaces (e.g., PM_{10}) may be reflected in the negative $PM_{2.5}$ parameters in the far "northeast" (see the white dots in Figure 4). In this newly developing, relatively affluent area of the city, air quality could hypothetically be worse further from roadways (the primary sources of $PM_{2.5}$) due disturbed desert crust. The predictive ability of LUR models to generate pollution surfaces is impacted by the number of monitoring sites employed in its identification (Javier et al., 2007). In this case, the small number of sites might have resulted in an overestimated predictive ability of the LUR model and thus affected its accuracy. Using only residential environmental indicators resulted in an incomplete exposure assessment, given children's space-time geographies. However, we did not have information about school-based pest exposure to include. The study uses a cross-sectional approach, which limits our ability to assess causal relationships. We also relied only on parental reports of children's symptoms using validated ISAAC items. No measures of lung function were collected as part of this study. Lastly, the survey sample is slightly less economically disadvantaged than the EPISD population, which could affect the results.

6. Conclusion

The answer to the first research question is that having pests and higher levels of $PM_{2.5}$ at a child's residence were modestly associated with greater wheezing severity. The answer to the second research question is more complex and includes two asthmogenic socio-environments. One is characterized by a situation of multiple jeopardy: pest and $PM_{2.5}$ exposures were important predictors of wheezing severity among children living in lower-income neighborhoods, among those who were more likely to be Hispanic, and among those with higher levels of $PM_{2.5}$ exposure at their home sites. The second ran counter to a multiple jeopardy model: $PM_{2.5}$ was an important positive predictor of wheezing severity in some wealthier neighborhoods, among non-Hispanic children, and among those for whom home site levels of outdoor $PM_{2.5}$ and odds of exposure to pests were lower. In more general terms, these results suggest that the assumption underpinning a good deal of research – that uneven environmental exposures produce correspondingly unequal health impacts and that these effects are stable across space – is not tenable.

Because of their capacity to identify specific areas at risk, GWR models lend themselves to locally relevant interventions. From a policy perspective, this research demonstrates particular areas in the city of El Paso where particulate matter and pests are important correlates of children's wheezing severity. This information could be used to aid local and state environmental and public health agencies to design targeted interventions in areas where they are most needed. In this case, GWR results suggest pest remediation as a tool to improve children's respiratory health in the "downtown," "south side," and "northeast" sides of El Paso. Pollution reduction efforts would benefit all children, given what is known about the deleterious health effects of air pollution and the positive coefficient in our global model. Increasing awareness of air pollution might be particularly important on the "west side", given that comparatively low levels of $PM_{2.5}$ play an important role in children's wheezing severity there. Given the apparently synergistic effects of exposure to pests and $PM_{2.5}$ in parts of the study area, addressing even just one of these factors would be advantageous to children's health. Results also suggest that GWR is a powerful tool that should be more widely used by researchers in their quest to understand complex relationships between environmental conditions, social characteristics, and health inequalities.

Acknowledgements

We acknowledge Bibi Mancera and Zuleika Ramirez at the Hispanic Health Disparities Center and the staff at the UTEP Campus Post Office for their assistance in carrying out the survey. The research participants are also gratefully recognized for taking the time to complete the survey. The work of student research assistants Anthony Jimenez, Marie Gaines, Stephanie Clark-Reyna, Alexander Balcazar, Paola Chavez-Payan, Young-an Kim, Omar Martinez, and Joe Valencia is gratefully recognized. This project was supported by Award Number P20 MD002287-05S1 from the National Institute on Minority Health and Health Disparities (NIMHD) and the Environmental Protection Agency (EPA) and Award Number CMMI-1129984 from the National Science Foundation (NSF). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIMHD, EPA, or NSF.

References

- Akinbami, Lara. Asthma Prevalence, Health Care Use and Mortality: United States, 2003–05. 2007. Retrieved July 22, 2008, from <http://www.cdc.gov/nchs/products/pubs/pubd/hestats/ashtma03-05/asthma03-05.htm>

- Al-Hamdan MZ, Crosson WL, Limaye AS, Rickman DL, Quattrochi DA, Estes MG Jr, Qualters JR, Sinclair AG, Tolsma DD, Adeniyi KA, Niskar AS. Methods for Characterizing Fine Particulate Matter Using Ground Observations and Remotely Sensed Data: Potential Use for Environmental Public Health Surveillance. *Journal of the Air & Waste Management Association*. 2006; 59(7):865–881. [PubMed: 19645271]
- Carlson, Joy E.; Stroebel, Carol. Childhood Asthma: A Growing Epidemic in the United States. *Population Today*. 2001; 29(4):10.
- Chakraborty, Jayajit. Revisiting Tobler's first law of geography: spatial regression models for assessing environmental justice and health risk disparities. In: Maantay, Juliana A.; McLafferty, Sara, editors. *Geospatial analysis of environmental health*. New York: Springer; 2011.
- Chakraborty, Jayajit; Maantay, Juliana. Proximity Analysis for Exposure Assessment in Environmental Health Justice Research. In: Maantay, JA.; McLafferty, S., editors. *Geospatial Analysis of Environmental Health*. New York: Springer; 2011. p. 111-138.
- Chalkias C, Papadopoulos AG, Kalogeropoulos K, Tambalis K, Psarra G, Sidossis L. Geographical heterogeneity of the relationship between childhood obesity and socio-environmental status: Empirical evidence from Athens, Greece. *Applied Geography*. 2013; 37:34–43.
- Charlton, Martin; Fotheringham, Stewart. Geographically Weighted Regression: A Tutorial on using GWR in ArcGIS 9.3. Maynooth, County Kildare, Ireland: National Centre for Geocomputation, National University of Ireland Maynooth; 2009. p. 1-27.
- Corburn, Jason. *Street Science: Community Knowledge and Environmental Health Justice*. Cambridge, MA: MIT Press; 2005.
- Deroose KP, Bahney BW, Lurie N, Escarce JJ. Immigrants and Health Care Access, Quality, and Cost. *Medical Care Research and Review*. 2009; 66(4):355–408. [PubMed: 19179539]
- Dillman, Don A.; Smyth, Jolene D.; Christian, Leah Melani. *Internet, Mail, and Mixed-mode Surveys: The Tailored Design Method*. 3rd ed.. Hoboken, New Jersey: John Wiley & Sons; 2009.
- Downey L. Environmental inequality in metropolitan America in 2000. *Sociological Spectrum*. 2006; 26(1):21–41. [PubMed: 21909177]
- EPISD. District information. 2013. Retrieved March 20, 2013, from http://www.episd.org/_district/
- Evans GW, Marcynyszyn LA. Environmental justice, cumulative environmental risk, and health among low- and middle-income children in upstate New York. *American Journal of Public Health*. 2004; 94(11):1942–1944. [PubMed: 15514234]
- Fotheringham, Stewart; Brundson, Chris; Charlton, Martin. *Geographically Weighted Regression: the analysis of spatially varying relationships*. West Sussex, England: John Wiley & Sons, LTD; 2002.
- Gilbert, Angela; Chakraborty, Jayajit. Using geographically weighted regression for environmental justice analysis: Cumulative cancer risks from air toxics in Florida. *Social Science Research*. 2011; 40(1):273–286.
- Grineski, Sara E. Incorporating Health Outcomes into Environmental Justice Research: The case of children's asthma and air pollution in Phoenix, Arizona. *Environmental Hazards*. 2007; 7(4):360–371.
- Grineski, Sara E. Human-Environment Interactions & Environmental Justice: How do diverse parents of asthmatic children minimize hazards? *Society and Natural Resources*. 2009; 22(8):727–743.
- Grineski, Sara E.; Collins, Timothy; Chakraborty, Jayajit; McDonald, Yolanda. Environmental Health Injustice: Exposure to Air Toxics and Children's Respiratory Hospital Admissions. *The Professional Geographer*. 2013; 65(1):31–46.
- Grineski, Sara E.; Hernandez, Alma A. Landlords, fear, and children's respiratory health: an untold story of environmental injustice in the central city. *Local Environment*. 2010; 15(3):199–216.
- Grineski, Sara E.; Staniswalis, Joan G.; Bulathsinhala, Priyangi; Peng, Yanlei; Gill, Thomas E. Hospital admissions for asthma and acute bronchitis in El Paso, Texas: Do age, sex, and insurance status modify the effects of dust and low wind events? *Environmental Research*. 2011; 111(8): 1148–1155. [PubMed: 21782162]
- Hackbarth AD, Romley JA, Goldman DP. Racial and ethnic disparities in hospital care resulting from air pollution in excess of federal standards. *Social Science & Medicine*. 2011; 73(8):1163–1168. [PubMed: 21893376]

- Hansen A, Bi P, Nitschke M, Pisaniello D, Ryan P, Sullivan T, Barnett AG. Particulate air pollution and cardiorespiratory hospital admissions in a temperate Australian city: A case-crossover analysis. *Science of the Total Environment*. 2012; 416:48–52. [PubMed: 22221868]
- Hoek, Gerard; Beelen, R.; Hoogh, K.; Vienneau, D.; Gulliver, J., et al. A review of land-use regression models to assess spatial variation of outdoor air pollution. *Atmospheric Environment*. 2008; 42(33):7561–7578.
- Hoek, Gerard; Meliefste, Kees; Cyrys, Josef; Lewné, Marie; Bellander, Tom; Brauer, Mike; Fischer, Paul; Gehring, Ulrike; Heinrich, Joachim; van Vliet, Patricia; Brunekreef, Bert. Spatial variability of fine particle concentrations in three European areas. *Atmospheric Environment*. 2002; 36(25): 4077–4088.
- Holt EW, Theall KP, Rabito FA. Individual, Housing, and Neighborhood Correlates of Asthma among Young Urban Children. *Journal of Urban Health-Bulletin of the New York Academy of Medicine*. 2013; 90(1):116–129. [PubMed: 22689297]
- Hryhorczuk D, Dardynskaia I, Lukyanova E, Matwyshyn-Fuoco M, Friedman L, Shkiryak-Nizhnyk Z, Zvinchuck A, Chislovska N, Antipkin Y. Risk factors for wheezing in Ukrainian children: Ukraine European Longitudinal Study of Pregnancy and Childhood Group. *Paediatric and Perinatal Epidemiology*. 2009; 23(4):346–351. [PubMed: 19523081]
- ISAAC Steering Committee. International Study of Asthma and Allergies in Children. 2012. Retrieved October, 2012, from <http://isaac.auckland.ac.nz/>
- Isakov, V.; Johnson, M.; Touma, J.; Özkaynak, H. Development and evaluation of land-use regression models using modeled air quality concentrations. In: Steyn, DGG.; Trini Castelli, ST., editors. *Air Pollution Modeling and its Application*. Vol. Vol. XXI. Netherlands: Springer; 2012. p. 717-722.
- Javier JR, Wise PH, Mendoza FS. The relationship of immigrant status with access, utilization, and health status for children with asthma. *Ambulatory Pediatrics*. 2007; 7(6):421–430. [PubMed: 17996835]
- Jephcote C, Chen HB. Environmental injustices of children's exposure to air pollution from road-transport within the model British multicultural city of Leicester: 2000–09. *Science of the Total Environment*. 2012; 414:140–151. [PubMed: 22154180]
- Jephcote C, Chen HB. Geospatial analysis of naturally occurring boundaries in road-transport emissions and children's respiratory health across a demographically diverse cityscape. *Social Science & Medicine*. 2013; 82:87–99. [PubMed: 23453321]
- Jerrett, Michael; Arain, M.; Kanaroglou, P.; Beckerman, B.; Crouse, D., et al. Modeling the intraurban variability of ambient traffic pollution in Toronto, Canada. *Journal of Toxicology and Environmental Health*. 2007; Part A 70(3):200–212. [PubMed: 17365582]
- Johnson M, Isakov V, Touma JS, Mukerjee S, Özkaynak H. Evaluation of land-use regression models used to predict air quality concentrations in an urban area. *Atmospheric Environment*. 2010; 44:3660–3668.
- Kanaroglou P, Jerrett M, Morrison J, Beckerman B, Arain M, Gilbert N, Brook J. Establishing an air pollution monitoring network for intra-urban population exposure assessment: a location–allocation approach. *Atmospheric Environment*. 2005; 39:2399–2409.
- Kraft ME, Scheberle D. Environmental Justice and the Allocation of Risk - the Case of Lead and Public-Health. *Policy Studies Journal*. 1995; 23(1):113–122.
- Landrigan PJ, Rauh VA, Galvez MP. Environmental Justice and the Health of Children. *Mount Sinai Journal of Medicine*. 2010; 77(2):178–187. [PubMed: 20309928]
- Lanphear, Bruce P.; Aligne, C. Andrew; Auinger, Peggy; Weitzman, Michael; Byrd, Robert S. Residential Exposures Associated with Asthma in US Children. *Pediatrics*. 2001; 107(3):505–511. [PubMed: 11230590]
- Matte TD, Jacobs DE. Housing and Health. *Journal of Urban Health*. 2000; 77(1):7–25. [PubMed: 10741839]
- Mehta, Neil K.; Lee, Hedwig; Ylitalo, Kelly R. Child health in the United States: Recent trends in racial/ethnic disparities. *Social Science & Medicine*. 2013; 95:6–15. [PubMed: 23034508]
- Mennis JL, Jordan L. The distribution of environmental equity: Exploring spatial nonstationarity in multivariate models of air toxic releases. *Annals of the Association of American Geographers*. 2005; 95(2):249–268.

- O'Connor KS, Osborn LL, Olson SJ, Blumberg Frankel MR, Srinath KP, Giambo P. Design and operation of the National Asthma Survey. *Vital and Health Statistics*. 2008; 46:1–122. [PubMed: 18958992]
- Olvera HA, Garcia M, Li WW, Yang H, Amaya MA, Myers O, Pingitore NE. Principal component analysis optimization of a PM_{2.5} land use regression model with small monitoring network. *Science of the Total Environment*. 2012; 425:27–34. [PubMed: 22464030]
- Padilla YC, Hamilton ER, Hummer RA. Beyond the Epidemiological Paradox: The Health of Mexican-American Children at Age Five. *Social Science Quarterly*. 2009; 90(5):1072–1088. [PubMed: 20072721]
- Parenteau M-P, Sawada M. The role of spatial representation in the development of a LUR model for Ottawa, Canada. *Air Quality, Atmosphere & Health*. 2010:1–13.
- Pearce JR, Richardson EA, Mitchell RJ, Shortt NK. Environmental justice and health: the implications of the socio-spatial distribution of multiple environmental deprivation for health inequalities in the United Kingdom. *Transactions of the Institute of British Geographers*. 2010; 35(4):522–539.
- Pearce JR, Richardson EA, Mitchell RJ, Shortt NK. Environmental justice and health: A study of multiple environmental deprivation and geographical inequalities in health in New Zealand. *Social Science & Medicine*. 2011; 73(3):410–420. [PubMed: 21726927]
- Richardson, Elizabeth A.; Pearce, Jamie; Tunstall, Helena; Mitchell, Richard; Shortt, Niamh K. Particulate air pollution and health inequalities: a Europe-wide ecological analysis. *International Journal of Health Geographics*. 2013; 12(34)
- Rose N, Cowie C, Gillett R, Marks GB. Validation of a Spatiotemporal Land Use Regression Model Incorporating Fixed Site Monitors. *Environmental Science & Technology*. 2010; 45:294–299. [PubMed: 21133418]
- Rose, Nectarios; Cowie, Christine; Gillett, Robert; Marks, Guy B. Validation of a Spatiotemporal Land Use Regression Model Incorporating Fixed Site Monitors. *Environmental Science & Technology*. 2010; 45(1):294–299. [PubMed: 21133418]
- Ryan PH, Bernstein DI, Lockey J, Reponen T, Levin L, Grinshpun S, LeMasters G. Exposure to Traffic-related Particles and Endotoxin during Infancy Is Associated with Wheezing at Age 3 Years. *American Journal of Respiratory and Critical Care Medicine*. 2009; 180(11):1068–1075. [PubMed: 19745206]
- Samet J, Krewski D. Health effects associated with exposure to ambient air pollution. *Journal of Toxicology and Environmental Health-Part a-Current Issues*. 2007; 70(3–4):227–242.
- Sarnat, Stefanie Ebel; Raysoni, Amit U.; Li, Wen-Whai; Holguin, Fernando; Johnson, Brent A.; Luevano, Silvia Flores; Garcia, Jose Humberto; Sarnat, Jeremy A. Air Pollution and Acute Respiratory Response in a Panel of Asthmatic Children along the U.S.–Mexico Border. *Environmental Health Perspectives*. 2011; 120(3)
- Sazonov Kocevar V, Thomas J III, Jonsson L, Valovirta E, Kristensen F, Yin DD, Bisgaard H. Association between allergic rhinitis and hospital resource use among asthmatic children in Norway. *Allergy*. 2005; 60(338–342)
- Schwartz, Joel. Air pollution and children's health. *Pediatrics*. 2012; 113(4):1037–1043. [PubMed: 15060197]
- Svendsen ER, Gonzales M, Mukerjee S, Smith L, Ross M, Walsh D, Neas LM. GIS-Modeled Indicators of Traffic-Related Air Pollutants and Adverse Pulmonary Health Among Children in El Paso, Texas. *American Journal of Epidemiology*. 2012; 176:S131–S141. [PubMed: 23035137]
- Tu J, Tu W, Tedders SH. Spatial variations in the associations of birth weight with socioeconomic, environmental, and behavioral factors in Georgia, USA. *Applied Geography*. 2012; 34:331–344.
- Wheeler BW, Ben-Shlomo Y. Environmental equity, air quality, socioeconomic status, and respiratory health: a linkage analysis of routine data from the Health Survey for England. *Journal of Epidemiology and Community Health*. 2005; 59(11):948–954. [PubMed: 16234422]
- Wheeler, DC.; Páez, A. Geographically weighted regression. In: Fischer, MM.; Getis, A., editors. *Handbook of applied spatial analysis: Software tools, methods and applications*. New York: Springer; 2010. p. 461–486.

- Woodin M, Tin AH, Moy S, Palella M, Brugge D. Lessons for Primary Prevention of Asthma: Foreign-Born Children Have Less Association of SES and Pests with Asthma Diagnosis. *Journal of Immigrant and Minority Health*. 2011; 13(3):462–469. [PubMed: 20953840]
- Wright AL, Stern DA, Kauffmann F, Martinez FD. Factors influencing gender differences in the diagnosis and treatment of asthma in childhood: the Tucson Children's Respiratory Study. *Pediatric Pulmonology*. 2006; 41(4):318–325.
- Xu XP, Wang LH. Synergistic effects of air pollution and personal smoking on adult pulmonary function. *Archives of Environmental Health*. 1998; 53(1):44–53. [PubMed: 9570308]
- Zora JE, Sarnat SE, Raysoni AU, Johnson BA, Li WW, Greenwald R, Sarnat JA. Associations between urban air pollution and pediatric asthma control in El Paso, Texas. *Science of the Total Environment*. 2013; 448:56–65. [PubMed: 23312496]

Figure 1.
PM_{2.5} Surface and Major Transportation Routes in El Paso County, Texas

Figure 2.

Study Area Map including GWR Model Performance Statistics (i.e., Local R^2)

Note: “West Side”, “Northeast”, “South Side”, and “Downtown” are used locally to refer to the labeled regions of the city.

Figure 3.
Standard Deviation Map of the Local Pest Parameters when Predicting Wheezing Severity

Figure 4.
Standard Deviation Map of the Local PM_{2.5} Parameters when Predicting Wheezing Severity

Table 1

Component loadings and individual variable means for the six wheezing variables included in the ‘Current wheezing severity’ measure

Variable	PCA Component Loadings	Individual Variable Mean
Doctor-diagnosed asthma	.682	.16
Asthma symptoms (including wheeze)	.833	.19
Night cough	.514	.24
Wheezing in sleep	.785	.07
Wheezing limited speech	.499	.02
Wheezing with no cold	.821	.08

N=1736

Table 2

Descriptive statistics of variables used in analysis

Variable (continuous)	Min.	Max.	Mean.	St. Dev.
Current wheezing severity (ln) ^I	-.73	1.71	-.29	.67
Residential PM _{2.5} (ln)	1.75	2.94	2.10	.25
Variable (dichotomous)	Response	Frequency	Percent	
Home has pest(s)	Yes (1)	781	45%	
	No (0)	955	55%	
Child is male	Yes (1)	868	50%	
	No (0)	868	50%	
Mother has asthma and/or allergies	Yes (1)	573	33%	
	No (0)	1163	67%	
Child has allergies	Yes (1)	885	51%	
	No (0)	851	49%	
Postponed or did not seek health care due to concerns about cost	Yes (1)	417	24%	
	No (0)	1319	76%	
Child is Hispanic	Yes (1)	1424	82%	
	No (0)	312	18%	

N=1736;

^I 1 was added to the wheezing measure before it was natural logged.

Table 3

OLS and GWR results: Predicting children's current wheezing severity

Variable	A. OLS MODEL				B. GWR MODEL			
	Adjusted R Square	β	Robust SE	Robust p-value	VIF	Mean β	Min β	Max β
Intercept		-0.647	0.043	0.001	-	0.605	-0.692	-0.512
Male (v. female)		0.091	0.029	0.002	1.011	0.084	0.033	0.135
Mom has asthma a/o allergies		0.130	0.037	0.001	1.147	0.125	0.009	0.207
Child has allergies		0.439	0.031	0.001	1.140	0.449	0.400	0.495
Postpone health care/cost concerns		0.120	0.037	0.002	1.020	0.110	0.047	0.167
Child is Hispanic (v. not Hispanic)		-0.004	0.041	0.922	1.070	-0.002	-0.070	0.060
Home has Pest(s)		0.040	0.030	0.177	1.014	0.030	-0.049	0.096
Residential PM (ln, Z)		0.012	0.015	0.411	1.073	0.017	-0.053	0.042

N=1736

T-test results characterizing children for whom their local GWR parameters for Pest and PM_{2.5} were 1 or more standard deviations above vs. below the mean

Table 4

	GWR Pest Parameter Group		Mean for Original Variable		GWR PM _{2.5} Parameter Group		Mean for Original Variable		Mean for Original Variable	
	N		N		N		N		N	
	p		p		p		p		p	
Male	1 SD Above	943	.500	1 SD Above	1141	.520	1 SD Above	1141	.520	
	1 SD Below	50	.480	1 SD Below	181	.490	1 SD Below	181	.490	
Mother has asthma and/or allergies	1 SD Above	943	.314	1 SD Above	1141	.334	1 SD Above	1141	.334	
	1 SD Below	50	.260	1 SD Below	181	.403	1 SD Below	181	.403	
Child has allergies	1 SD Above	943	.520	1 SD Above	1141	.510	1 SD Above	1141	.510	*
	1 SD Below	50	.540	1 SD Below	181	.610	1 SD Below	181	.610	
Problems with cost when seeking healthcare	1 SD Above	943	.249	1 SD Above	1141	.231	1 SD Above	1141	.231	
	1 SD Below	50	.140	1 SD Below	181	.232	1 SD Below	181	.232	
Child is Hispanic	1 SD Above	943	.840	1 SD Above	1141	.813	1 SD Above	1141	.813	*
	1 SD Below	50	.460	1 SD Below	181	.680	1 SD Below	181	.680	
Pest	1 SD Above	943	.480	1 SD Above	1141	.461	1 SD Above	1141	.461	*
	1 SD Below	50	.380	1 SD Below	181	.343	1 SD Below	181	.343	
Residential PM _{2.5} (ln, Z)	1 SD Above	943	.245	1 SD Above	1141	.099	1 SD Above	1141	.099	*
	1 SD Below	50	-1.239	1 SD Below	181	-.949	1 SD Below	181	-.949	

N=1736;

*
p<0.05

Table 5

T-test results characterizing the two spatial groupings of positive local GWR parameters for PM_{2.5}

	GWR PM_{2.5} Parameter Group (1+ SD Above Mean)	N	Mean for Original Variable	p
Male	“West”	527	.500	
	“Eastern”	639	.540	
Mother has asthma and/or allergies	“West”	527	.347	
	“Eastern”	639	.327	
Child has allergies	“West”	527	.520	
	“Eastern”	639	.500	
Problems with cost when seeking healthcare	“West”	527	.228	
	“Eastern”	639	.239	
Child is Hispanic	“West”	527	.776	*
	“Eastern”	639	.845	
Pest	“West”	527	.360	*
	“Eastern”	639	.540	
Residential PM _{2.5} (ln, Z)	“West”	527	-.454	*
	“Eastern”	639	.541	

N=1736;

*
p<0.05