Published in final edited form as:

Clin Nutr. 2016 June ; 35(3): 679–684. doi:10.1016/j.clnu.2015.04.016.

Egg consumption and risk of type 2 diabetes among African Americans: The Jackson Heart Study

Luc Djoussé^{1,2,3}, Andrew B Petrone¹, DeMarc A Hickson^{4,5}, Sameera A. Talegawkar⁶, Patricia M Dubbert⁷, Herman Taylor⁸, and Katherine L Tucker⁹

¹Brigham and Women's Hospital, Boston MA

²Boston VA Healthcare System, Boston, MA

³Harvard Medical School, Boston, MA

⁴University of Mississippi Medical Center, Jackson, MS

⁵My Brother's Keeper, Inc, Ridgeland, MS

⁶Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

⁷South Central VA MIRECC and Little Rock GRECC, Little Rock, AR

⁸Morehouse School of Medicine, Atlanta, GA

⁹University of Massachusetts Lowell, Lowell, MA

Abstract

Background and Aims—Type 2 diabetes (DM) disproportionally affects African Americans. Data on the association between egg consumption and risk of DM are sparse. We sought to examine whether egg consumption is associated with the prevalence and incidence of DM among African Americans.

Methods—We analyzed baseline data from 4,568 participants of the Jackson Heart Study. Egg consumption was obtained using a food frequency questionnaire designed for this population. We used generalized estimating equations to calculate adjusted prevalence ratios of DM and Cox regression to estimate hazard ratios of DM with corresponding 95% confidence intervals (CI).

Results—The average age was 55 ± 13 years and 64% of subjects were women. The median frequency of egg consumption was 2/week for men and 1/week for women. The prevalence of DM

Corresponding author: Luc Djoussé, MD, MPH, ScD, Division of Aging, Brigham and Women's Hospital, 1620 Tremont St. 3rd floor, Boston MA 02120, Ldjousse@rics.bwh.harvard.edu, Tel. (617) 525-7591, Fax. (617) 525-7739.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Author contribution:

Conception and design of the study (LD)

Acquisition of data (PMD, HT, and KLT)

Data analysis (LD, ABP)

Interpretation of data (LD, ABP, DAH, SAT, PMD, HT, and KLT)

Drafting the article (LD) or revising it critically for important intellectual content (LD, ABP, DAH, SAT, PMD, HT, KLT)

was 22% overall (21% of men and 23% of women). Multivariable adjusted prevalence ratio [PR(95% CI)] for DM were: 1.00 (ref), 1.14 (0.90–1.44), 1.33 (1.04–1.70), 1.33 (1.06–1.68), 1.26 (0.99–1.61), and 1.52 (1.17–1.97) for egg consumption of <1/month, 1–3/month, 1/week, 2/week, 3–4/week, and 5+/week, respectively, *p* for linear trend 0.0024. Corresponding multivariable adjusted hazard ratios were 1.00 (ref), 0.88 (0.65–1.19), 0.94 (0.68–1.30), 0.91 (0.66–1.25), 1.11 (0.81–1.52), and 1.17 (0.81–1.70), respectively, during a mean follow up of 7.3 years (*p* for linear trend 0.22).

Conclusions—While egg consumption was positively associated with prevalent DM, prospective analysis did not show an association of egg intake with incidence of DM among African Americans.

Keywords

Epidemiology; nutrition; eggs; type 2 diabetes; risk factors

Introduction

As a major risk factor for cardiovascular disease (CVD), type 2 diabetes (DM) is a major health issue worldwide^{1–3}. It is anticipated that 440 million individuals will be diagnosed with DM by 2030². Global healthcare expenditures due to DM are projected around 490 billion dollars by 2030³. The burden of DM disproportionately affects African-Americans. In 2009, the risk of death per 100,000 due to DM was 44.2 for black men (vs. 23.3 for white men) and 35.9 for black women (vs. 15.7 for white women)⁴.

Previous research has identified modifiable lifestyle factors such as diet and physical activity as important determinants of DM⁵. However, the association between different macronutrients and risk of DM is not consistent. For example, among saturated fatty acids, while short-chain saturated fatty acids have been associated with a higher risk of DM, odds chain and long-chain saturated fatty acids have been related with a lower risk of DM^{6–8}. In addition, the EPIC cohort reported that animal but not vegetable proteins were associated with a higher risk of DM⁹ whereas a meta-analysis of three studies showed no association of animal proteins with DM¹⁰.

Among individual foods, eggs could play a unique role among dietary factors affecting the risk of DM, partly because they are rich in proteins, cholesterol, and other nutrients. Current data on the association of eggs with DM have been inconsistent, ranging from no association in Japanese¹¹ and non-Hispanic white adults^{12,13} to a higher risk of DM with frequent intake of eggs in Chinese¹⁴ and other populations^{15–18}. In two previous meta-analyses, the highest category of egg consumption was associated with a 42% to 68% higher risk of DM^{19,20}; however most of those data were derived from non-Hispanic whites, and it is unclear whether egg consumption is associated with a higher risk of DM in African Americans. Thus, the current paper sought to examine the association of egg consumption with prevalent and incident DM among African American participants of the Jackson Heart Study (JHS).

Methods

Study population

The present analysis used JHS data collected during the baseline examination (2000–2004). Detailed description of the JHS has been previously published^{21–25}. Of the 5,301 JHS participants who completed a baseline clinic examination, we excluded 248 who had missing data on egg consumption; one person with type 1 DM; 16 with restricted informed consent; 422 with energy intake outside of normal ranges (600 to 4000 kilocalories per day); and 46 with missing information on DM status. Thus, current analyses are based on 4,568 individuals with complete data for cross-sectional analyses and 3,564 subjects for prospective analyses after exclusion of prevalent DM (n=1004). Each participant gave written informed consent and the study protocol was approved by the institutional review boards of all institutions involved.

Assessment of egg consumption

Egg consumption was assessed using a 158-item food frequency questionnaire (FFQ)²⁶ that has been validated in this cohort²⁷. Participants reported their frequency of egg consumption, including fried and scrambled eggs, as never; less than once per month; 1/month; 2–3/month; 1/week; 2/week; 3–4/week; 5–6/week; 1/day; or 2/day. Adjacent categories were combined into <1/month; 1–3/month; 1/week; 2/week; 3–4/week; and 5+/-week to allow for stable estimate in current analyses.

Ascertainment of DM

Prevalent DM was defined as fasting glucose ≥ 126 mg/dL, hemoglobin A1C $\geq 6.5\%$, or current use of insulin or oral hypoglycemic agents at baseline. Similar criteria were used to define incident DM during follow up among participants that were free of DM at baseline. Since we did not have information on the exact date of diagnosis of DM, we assigned the JHS visit date following the initial diagnosis of DM as date of DM occurrence.

Other variables

Assays and procedures for measuring fasting whole blood, plasma, and serum-derived measurements in the JHS have been previously described²¹. At JHS visit 1, we collected information on demographic and anthropometric variables, cigarette smoking, physical activity, education, alcohol consumption, history of hypertension, CVD, LDL- and HDL-cholesterol, triglycerides, and diet. Intakes of fruit and vegetables, red meat, fish, trans fat, total energy, dietary fiber, and dietary magnesium were used as covariates. Physical activity was measured as a sum of four index scores (active living, work/occupational, home life, and sports)²⁸.

Statistical Analysis

Baseline characteristics within each category of egg consumption are presented as mean $\pm$ standard deviation (SD) for continuous variables with Gaussian distribution (otherwise median with inter-quartile range), and proportions for categorical variables. We used generalized estimating equations to calculate prevalence ratio (PR) for cross-sectional

analyses and Cox proportional hazard model for prospective analyses along with corresponding 95% confidence limits. Person-time of follow-up was computed from baseline until the first occurrence of incident diabetes, death, or last contact. We built sequential models based on a priori knowledge. In model 1, we adjusted for age (<45, 45–55, 55–65, >65) and sex. Model 2 was additionally adjusted for BMI (<25, 25–30, 30–35, >35), smoking status (never, former, current), alcohol intake in the past 12 months (yes/no), physical activity score (tertiles), and education (less than high school/high school or some college/associates degree or higher). The final model adjusted for variables in model 2 plus energy intake (tertiles), red meat (tertiles), dietary fiber (tertiles), dietary magnesium (tertiles), fruit and vegetable intake (tertiles), dietary trans fat (tertiles), waist circumference (tertiles), history of hypertension (yes/no), and history of CVD (yes/no). We obtained a p-value for linear trend by including an ordinal variable that was assigned to frequency of egg consumption in the regression model. Cox proportional hazard model assumption were tested using product term of egg frequency and log-transformed person-time and were met (all $p > 0.05$).

In secondary analyses, we stratified the main analysis by sex, BMI, and age and also examined whether egg consumption was associated with baseline measures of insulin resistance and A1C by computing least squares means for logarithmic-transformed A1C, homeostatic model assessment-insulin resistance (HOMA-IR), and homeostatic model assessment-beta cell function (HOMA-B). Log-transformation was necessary due to non-Gaussian distribution of HOMA and A1C measures. All analyses were completed using Statistical Analysis Systems, version 9.3 (SAS Institute, Cary, North Carolina) with alpha level of 0.05.

Results

Baseline characteristics of the 4,568 African Americans (Table 1) show a mean age of 55 ± 13 years (range 21–95), with 64% women. The prevalence of DM was 22% (21% of men and 23% of women). Compared to egg consumption of < once per month, higher egg consumption was associated with male sex, current smoking, lower educational attainment, alcohol consumption, higher waist circumference, lower HDL, and consumption of red meat, fish, bacon, saturated and trans fats (Table 1). Egg consumption was correlated with saturated fat (Spearman $r=0.37$), energy intake ($r=0.32$), trans fatty acids ($r=0.25$), processed meats ($r=0.21$), dietary magnesium ($r=0.18$), and fiber ($r=0.13$).

Cross-sectional analyses (eggs with prevalent DM and biomarkers)

In a model adjusting for age and sex, prevalence ratios (95% CI) for DM were 1.00 (ref), 1.06 (0.85–1.32), 1.28 (1.02–1.60), 1.29 (1.05–1.60), 1.25 (1.00–1.56), and 1.54 (1.22–1.94) from the lowest to highest categories of egg consumption (p for linear trend 0.0001, Table 2). These findings persisted and were not substantially altered after additional adjustment for smoking status, alcohol consumption, BMI, physical activity score, education, energy intake, intake of red meat and fruit and vegetables, trans fat, dietary magnesium, and fiber, and history of hypertension and CVD (Table 2).

In secondary analysis, we observed similar egg-DM associations in men (p-trend 0.05) and women (p-trend 0.02), Table 3. Neither sex (p-interaction 0.53), nor age (p-interaction 0.88) or BMI (p-interaction 0.20) modified the association of egg consumption with prevalent DM. Lastly, egg consumption was not associated with mean A1C, HOMA-IR, or HOMA-B in multivariable adjusted models with a subsample of 3,178 participants without DM (Table 4).

Prospective analyses (eggs and incidence of DM)

During a mean follow up of 7.3 years, 531 new cases of DM occurred among 3,564 subjects free of DM at baseline. In both crude and multivariable adjusted analyses, egg consumption was not associated with the risk of developing DM (Table 5). Gender, age, and BMI did not modify the relation of egg consumption with incidence of DM (all p for interaction >0.58).

Discussion

In this large cohort, we found a positive and linear relation between frequency of egg consumption and baseline prevalence of DM in African Americans. However, no association was observed between frequency of egg consumption and incidence rate of DM in this cohort. In a secondary cross-sectional analysis, we found no evidence for an association of egg consumption with A1C concentration or with measures of insulin resistance (HOMA-IR and HOMA-B) in participants without DM at baseline.

To our knowledge, this is the first study to examine the relation of egg consumption with prevalent and incident DM in a large cohort of African Americans. Contrary to our cross-sectional results, we did not find evidence for an association between frequency of egg consumption and the incidence of DM during a mean follow up of seven years in this cohort. A lack of an association between egg consumption and incidence of DM is consistent with results from the Cardiovascular Health Study, where egg consumption was not associated with incidence of DM among older US adults who were predominantly Caucasians¹³. Similarly, neither a large Japanese cohort of 63,466 adult men and women¹¹ nor a Mediterranean cohort of 15,956 adult men and women¹² found an association of egg consumption with DM with comparable follow up times. Nevertheless, our prospective findings are contrary to the health professional data showing a positive relation between egg consumption and incident DM¹⁵. A case-control study in a Lithuanian out-patient clinic reported a three-fold increased odds of DM comparing 5 eggs/week to <1 egg/week¹⁷. Furthermore, a prospective study of about 3,000 women reported positive and graded relation of egg consumption with gestational DM (P for trend=0.008)¹⁶. A cross-sectional analysis of Chinese adults showed a two-fold higher odds of DM comparing 1+ eggs/d to <2 eggs/week (OR=2.28 (1.14–4.54))¹⁴, finding that is consistent with our cross-sectional results. In a meta-analysis using prospective data from the Physicians' Health Study, the Women's Health Study, the Cardiovascular Health Study (CHS), and the Adventist Health Studies, the highest category of egg consumption was associated with 42% higher risk of DM (95% CI: 9–86%) when compared to the lowest category¹⁹. Another meta-analysis, which included studies in China and Lithuania, also found a positive relation of egg consumption with risk of DM [pooled RR: 1.68 (95% CI: 1.41–2.00)]²⁰. Of note is that

above meta-analyses did not include large recent studies with negative findings (i.e., the Japan Public Health Center-based Prospective Study with more than 63,000 people).

Several factors could account for the inconsistency of these findings with our results. First, differences in overall dietary habits (i.e., adherence to Mediterranean diet known to lower DM risk²⁹) between cohorts reporting positive versus no association could partially explain the inconsistency, where the effect of a single food item such as eggs, may be insignificant when compared to the effects of an overall healthy dietary pattern. Second, differences in other lifestyle factors that are inherently associated with ethnic groups or geographic locations might account for some of the disparities. In the current study, we did not assign a high weight to cross-sectional results in the presence of prospective data since we cannot ascertain temporality in cross-sectional analyses. It is possible that subjects diagnosed with DM altered their diet voluntarily or per clinicians' advice to obtain their proteins from eggs rather than red or processed meats. Such a scenario could partially explain the positive relation observed between egg consumption and prevalent DM in our cohort.

Eggs are a good source of protein and other nutrients. However, a medium egg also contains up to 200 mg of dietary cholesterol³⁰. Qiu et al.¹⁶ reported a positive association between dietary cholesterol intake and risk of gestational DM. However, our group reported no evidence of association between dietary cholesterol and incident DM in older adults. In addition, a Japanese study¹¹ reported an inverse relation of dietary cholesterol with DM risk [OR=0.68 (95 % CI 0.49- 0.94), comparing the fourth to the first quartile of dietary cholesterol]. A human study found that 12 weeks of a high-protein diet with eggs improved fasting blood glucose (−0.5 mmol/l)³¹. Taken together, current data do not lend support to the hypothesis that dietary cholesterol is a major culprit for the development of DM. Could choline metabolites help explain positive results of egg-DM association?

Eggs are an important source of choline³², which can be metabolized by gut bacteria to generate trimethylamine, that can further be transformed to trimethylamine-N-oxide (TMAO) in the liver³³. Tang et al.³⁴ reported a positive association of egg consumption and TMAO concentration. In a randomized controlled trial of six volunteers, higher egg yolk intake led to a graded increase in plasma and urine TMAO³². TMAO may increase LDL oxidation and promote inflammation³⁵, a key component of the pathogenesis of DM³⁶. A targeted metabolomic approach suggested that diacyl-phosphatidylcholines C32:1, C36:1, C38:3, and C40:5 were independently associated with higher risk of DM in the EPIC Postdam cohort³⁷. However, prospective data relating plasma TMAO concentration with incident DM are lacking. In particular, we did not have information on TMAO in our cohort to further explore this hypothesis. We should note that other sources of TMAO or TMAO precursors like seafood³⁸ could confound any egg-disease relation. In a pilot study testing 46 various foods, fish intake but not eggs or other food items led to higher urinary TMAO excretion³⁸. This study further weakens the hypothesis of eggs-TMAO-DM risk. A lack of association between egg consumption and A1c, fasting glucose or insulin in our data is consistent with data from an egg feeding trial showing no effect of egg on fasting glucose in overweight men¹⁸.

Limitations of our study include self-reported information on egg consumption and a lack of information on the size of eggs consumed, type of egg preparation, quantity of yolks consumed per egg; and on mixed dishes containing eggs. As observational study, we cannot exclude unmeasured confounding in our data. The use of an all-African American cohort limits the generalizability of our findings. However, this concern is offset by previously reported null findings in other ethnic groups. On the other hand, strengths of our study include a large sample size, availability of data on several important covariates, and use of standardized procedures to collect data in this cohort.

In conclusion, our data do not lend support for a higher incidence rate of DM with frequency of egg consumption despite a positive relation of egg consumption with prevalent DM in the JHS cohort.

Acknowledgements

The authors thank the participants and staff in the Jackson Heart Study for their commitment to the study.

Funding: The Jackson Heart Study is supported by contracts HHSN268201300046C, HHSN268201300047C, HHSN268201300048C, HHSN268201300049C, HHSN268201300050C from the National Heart, Lung, and Blood Institute and the National Institute on Minority Health and Health Disparities. This project was also supported by grant R21 NR013231 from the National Health Institutes. Dr. Talegawkar was supported by grant 13BGIA17080036 from the American Heart Association.

References

1. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. *Diabetes Care*. 1998; 21:1414–1431. [PubMed: 9727886]
2. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract*. 2010; 87:4–14. [PubMed: 19896746]
3. Zhang P, Zhang X, Brown J, Vistisen D, Sicree R, Shaw J, Nichols G. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res Clin Pract*. 2010; 87:293–301. [PubMed: 20171754]
4. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Huffman MD, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Magid D, Marcus GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Nichol G, Paynter NP, Schreiner PJ, Sorlie PD, Stein J, Turan TN, Virani SS, Wong ND, Woo D, Turner MB. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation*. 2013; 127:e6–e245. [PubMed: 23239837]
5. Djoussé L, Driver JA, Gaziano JM, Buring JE, Lee IM. Association between modifiable lifestyle factors and residual lifetime risk of diabetes. *Nutr Metab Cardiovasc Dis*. 2013; 23:17–22. [PubMed: 21982361]
6. Forouhi NG, Koulman A, Sharp SJ, Imamura F, Kroger J, Schulze MB, Crowe FL, Huerta JM, Guevara M, Beulens JW, van Woudenberg GJ, Wang L, Summerhill K, Griffin JL, Feskens EJ, Amiano P, Boeing H, Clavel-Chapelon F, Dartois L, Fagherazzi G, Franks PW, Gonzalez C, Jakobsen MU, Kaaks R, Key TJ, Khaw KT, Kuhn T, Mattiello A, Nilsson PM, Overvad K, Pala V, Palli D, Quiros JR, Rolandsson O, Roswall N, Sacerdote C, Sanchez MJ, Slimani N, Spijkerman AM, Tjønneland A, Tormo MJ, Tumino R, van der AD, van der Schouw YT, Langenberg C, Riboli E, Wareham NJ. Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study. *Lancet Diabetes Endocrinol*. 2014; 2:810–818. [PubMed: 25107467]
7. Ma W, Wu JH, Wang Q, Lemaitre RN, Mukamal KJ, Djoussé L, King IB, Song X, Biggs ML, Delaney JA, Kizer JR, Siscovick DS, Mozaffarian D. Prospective association of fatty acids in the de

- novo lipogenesis pathway with risk of type 2 diabetes: the Cardiovascular Health Study. *Am J Clin Nutr.* 2015; 101:153–163. [PubMed: 25527759]
8. Lemaitre RN, Fretts AM, Sitlani CM, Biggs ML, Mukamal K, King IB, Song X, Djoussé L, Siscovick DS, McKnight B, Sotoodehnia N, Kizer JR, Mozaffarian D. Plasma phospholipid very-long-chain SFAs and incident diabetes in older adults: the Cardiovascular Health Study. *Am J Clin Nutr.* 2015
 9. Sluijs I, Beulens JW, van der AD, Spijkerman AM, Grobbee DE, van der Schouw YT. Dietary intake of total, animal, and vegetable protein and risk of type 2 diabetes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-NL study. *Diabetes Care.* 2010; 33:43–48. [PubMed: 19825820]
 10. Alhazmi A, Stojanovski E, McEvoy M, Garg ML. Macronutrient intakes and development of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *J Am Coll Nutr.* 2012; 31:243–258. [PubMed: 23378452]
 11. Kurotani K, Nanri A, Goto A, Mizoue T, Noda M, Oba S, Sawada N, Tsugane S. for the Japan Public Health Center-based Prospective Study Group. Cholesterol and egg intakes and the risk of type 2 diabetes: The Japan Public Health Center-based Prospective Study. *Br J Nutr.* 2014:1–8.
 12. Zazpe I, Beunza JJ, Bes-Rastrollo M, Basterra-Gortari FJ, Mari-Sanchis A, Martinez-Gonzalez MA, SUN P I. Egg consumption and risk of type 2 diabetes in a Mediterranean cohort; the sun project. *Nutr Hosp.* 2013; 28:105–111. [PubMed: 23808437]
 13. Djoussé L, Kamineni A, Nelson TL, Carnethon M, Mozaffarian D, Siscovick D, Mukamal KJ. Egg consumption and risk of type 2 diabetes in older adults. *Am J Clin Nutr.* 2010; 92:422–427. [PubMed: 20534749]
 14. Shi Z, Yuan B, Zhang C, Zhou M, Holmboe-Ottesen G. Egg consumption and the risk of diabetes in adults, Jiangsu, China. *Nutrition.* 2011; 27:194–198. [PubMed: 20471806]
 15. Djoussé L, Gaziano JM, Buring JE, Lee IM. Egg Consumption and Risk of Type 2 Diabetes in Men and Women. *Diabetes Care.* 2009; 32:295–300. [PubMed: 19017774]
 16. Qiu C, Frederick IO, Zhang C, Sorensen TK, Enquobahrie DA, Williams MA. Risk of gestational diabetes mellitus in relation to maternal egg and cholesterol intake. *Am J Epidemiol.* 2011; 173:649–658. [PubMed: 21324948]
 17. Radzeviciene L, Ostrauskas R. Egg consumption and the risk of type 2 diabetes mellitus: a case-control study. *Public Health Nutr.* 2012; 15:1437–1441. [PubMed: 22390963]
 18. Djoussé L. Relation of eggs with incident cardiovascular disease and diabetes: Friends or foes? *Atherosclerosis.* 2013; 229:507–508. [PubMed: 23755730]
 19. Shin JY, Xun P, Nakamura Y, He K. Egg consumption in relation to risk of cardiovascular disease and diabetes: a systematic review and meta-analysis. *Am J Clin Nutr.* 2013; 98:146–159. [PubMed: 23676423]
 20. Li Y, Zhou C, Zhou X, Li L. Egg consumption and risk of cardiovascular diseases and diabetes: A meta-analysis. *Atherosclerosis.* 2013; 229:524–530. [PubMed: 23643053]
 21. Carpenter MA, Crow R, Steffes M, Rock W, Heilbraun J, Evans G, Skelton T, Jensen R, Sarpong D. Laboratory, reading center, and coordinating center data management methods in the Jackson Heart Study. *Am J Med Sci.* 2004; 328:131–144. [PubMed: 15367870]
 22. Fuqua SR, Wyatt SB, Andrew ME, Sarpong DF, Henderson FR, Cunningham MF, Taylor HA Jr. Recruiting African-American research participation in the Jackson Heart Study: methods, response rates, and sample description. *Ethn Dis.* 2005; 15:S6–S29. [PubMed: 16317982]
 23. Payne TJ, Wyatt SB, Mosley TH, Dubbert PM, Guiterrez-Mohammed ML, Calvin RL, Taylor HA Jr, Williams DR. Sociocultural methods in the Jackson Heart Study: conceptual and descriptive overview. *Ethn Dis.* 2005; 15:S6–S48. [PubMed: 16317984]
 24. Taylor HA Jr, Wilson JG, Jones DW, Sarpong DF, Srinivasan A, Garrison RJ, Nelson C, Wyatt SB. Toward resolution of cardiovascular health disparities in African Americans: design and methods of the Jackson Heart Study. *Ethn Dis.* 2005; 15:S6–S17.
 25. Hickson DA, Waller LA, Gebreab SY, Wyatt SB, Kelly J, Antoine-Lavigne D, Sarpong DF. Geographic representation of the Jackson heart study cohort to the African-American population in Jackson, Mississippi. *Am J Epidemiol.* 2011; 173:110–117. [PubMed: 21076050]

26. Carithers T, Dubbert PM, Crook E, Davy B, Wyatt SB, Bogle ML, Taylor HA Jr, Tucker KL. Dietary assessment in African Americans: methods used in the Jackson Heart Study. *Ethn Dis.* 2005; 15:S6–S55.
27. Carithers TC, Talegawkar SA, Rowser ML, Henry OR, Dubbert PM, Bogle ML, Taylor HA Jr, Tucker KL. Validity and calibration of food frequency questionnaires used with African-American adults in the Jackson Heart Study. *J Am Diet Assoc.* 2009; 109:1184–1193. [PubMed: 19559135]
28. Dubbert PM, Carithers T, Ainsworth BE, Taylor HA Jr, Wilson G, Wyatt SB. Physical activity assessment methods in the Jackson Heart Study. *Ethn Dis.* 2005; 15:S6–S61.
29. Schwingshackl L, Missbach B, König J, Hoffmann G. Adherence to a Mediterranean diet and risk of diabetes: a systematic review and meta-analysis. *Public Health Nutr.* 2014;1–8. [PubMed: 24650538]
30. Song WO, Kerver JM. Nutritional contribution of eggs to American diets. *J Am Coll Nutr.* 2000; 19:556S–562S. [PubMed: 11023007]
31. Pearce KL, Clifton PM, Noakes M. Egg consumption as part of an energy-restricted high-protein diet improves blood lipid and blood glucose profiles in individuals with type 2 diabetes. *Br J Nutr.* 2011; 105:584–592. [PubMed: 21134328]
32. Miller CA, Corbin KD, da Costa KA, Zhang S, Zhao X, Galanko JA, Blevins T, Bennett BJ, O'Connor A, Zeisel SH. Effect of egg ingestion on trimethylamine-N-oxide production in humans: a randomized, controlled, dose-response study. *Am J Clin Nutr.* 2014; 100:778–786. [PubMed: 24944063]
33. Zeisel SH, Mar MH, Howe JC, Holden JM. Concentrations of choline-containing compounds and betaine in common foods. *J Nutr.* 2003; 133:1302–1307. [PubMed: 12730414]
34. Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, Wu Y, Hazen SL. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med.* 2013; 368:1575–1584. [PubMed: 23614584]
35. Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, Feldstein AE, Britt EB, Fu X, Chung YM, Wu Y, Schauer P, Smith JD, Allayee H, Tang WH, DiDonato JA, Lusis AJ, Hazen SL. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature.* 2011; 472:57–63. [PubMed: 21475195]
36. Julia C, Czernichow S, Charnaux N, Ahluwalia N, Andreeva V, Touvier M, Galan P, Fezeu L. Relationships between adipokines, biomarkers of endothelial function and inflammation and risk of type 2 diabetes. *Diabetes Res Clin Pract.* 2014; 105:231–238. [PubMed: 24931702]
37. Floegel A, Stefan N, Yu Z, Mühlenbruch K, Drogan D, Joost HG, Fritsche A, Haring HU, Hrabe dA, Peters A, Roden M, Prehn C, Wang-Sattler R, Illig T, Schulze MB, Adamski J, Boeing H, Pischon T. Identification of serum metabolites associated with risk of type 2 diabetes using a targeted metabolomic approach. *Diabetes.* 2013; 62:639–648. [PubMed: 23043162]
38. Zhang AQ, Mitchell SC, Smith RL. Dietary precursors of trimethylamine in man: a pilot study. *Food Chem Toxicol.* 1999; 37:515–520. [PubMed: 10456680]

Table 1

Characteristics of the 4,568 Jackson Heart Study participants by frequency of egg consumption *

	Egg Consumption Categories						p for linear trend
	<1/month (n=742)	1-3/month (n=958)	1/week (n=698)	2/week (n=881)	3-4/week (n=781)	5+/week (n=508)	
Age (y)	57 ± 12	54 ± 13	55 ± 12	55 ± 13	55 ± 12	57 ± 13	0.43
BMI (kg/m ²)	31 ± 7	31 ± 7	32 ± 8	32 ± 7	32 ± 7	32 ± 7	0.01
Male (%)	23	32	37	37	41	50	<0.0001
Smoking (%)							
Never	71	69	71	73	65	61	0.0013
Past	20	20	18	16	21	24	0.23
Current	7.6	12	11	11	14	15	0.002
Education (%)							
< High School	19	16	16	17	17	26	0.02
High School/GED/Some College	42	37	41	42	44	45	0.02
College/Assoc. deg or higher	39	47	43	42	39	29	<0.0001
Drank alcohol in past year (%)	33	49	46	45	48	46	0.0001
Hypertension (%)	67	58	61	63	66	69	0.02
CVD (%)	11.2	7.8	9.6	10	12	13	0.02
LDL (mg/dL)	128 ± 38	128 ± 37	127 ± 37	128 ± 36	125 ± 35	127 ± 35	0.19
HDL (mg/dL)	53 ± 15	53 ± 15	51 ± 14	52 ± 14	51 ± 15	51 ± 14	0.0002
Triglycerides (mg/dL)	107 ± 63	103 ± 69	104 ± 62	109 ± 71	103 ± 78	111 ± 80	0.25
Waist Circumference (cm)	99 ± 15	99 ± 16	100 ± 17	101 ± 17	102 ± 16	103 ± 15	<0.0001
Physical Activity Score	8.0 ± 2.5	8.5 ± 2.5	8.6 ± 2.5	8.3 ± 2.6	8.4 ± 2.6	8.0 ± 2.8	0.77
Fruit and Vegetables (servings/day)	2.8[1.8-4.3]	2.7[1.7-3.9]	2.8[1.8-4.1]	3.4[2.3-4.7]	3.2[2.2-4.5]	3.5[2.4-5.0]	<0.0001
Red Meat (g/day)	7.8 [3.0-20]	8.1[3.5-16]	12.4[5.0-24]	14.5 [7.2-27]	15.3[7.7-29]	21[10-38]	<0.0001
Fish (g/day)	8.6[3.0-20.7]	9.7[4.3-21.9]	13.8[5.3-27.4]	12.0[5.6-27.1]	13.8[5.3-27.1]	10.8[4.3-24.7]	<0.0001
Saturated Fat (g/day)	18[12-26]	19[14-27]	23[16-31]	26[18-35]	29 [21-38]	34[26-44]	<0.0001
Trans Fat (g/day)	3.2[2.1-4.8]	3.4[2.3-4.8]	3.7[2.7-5.7]	4.4[2.9-6.2]	4.6[3.2-6.6]	5.2[3.5-7.3]	<0.0001
Dietary Cholesterol (mg/day)	162[109-243]	190[145-255]	257[192-327]	334[260-433]	439[345-548]	657[478-804]	<0.0001
Dietary Magnesium (mg/day)	251[191-326]	244[191-319]	270[215-354]	286[225-359]	298[229-371]	315[238-388]	<0.0001
Dietary fiber (g/day)	13 ± 6	12 ± 6	14 ± 6	14 ± 6	14 ± 6	15 ± 6	<0.0001

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Egg Consumption Categories					
	<1/month (n=742)	1-3/month (n=958)	1/week (n=698)	2/week (n=881)	3-4/week (n=781)
Calories (kcal/d)	1644 ± 709	1655 ± 703	1869 ± 746	2041 ± 773	2154 ± 780
				2348 ± 786	
					p for linear trend <0.0001

* Mean±SD for continuous variables with Gaussian distribution or median [interquartile range] if not normally distributed

Table 2

Prevalence ratios (95% CI) of diabetes at baseline according to egg consumption in the Jackson Heart Study (n=4,568)

Egg Consumption	Cases/n [*]	Prevalence ratios (95% CI)			
		Crude	Model 1 [†]	Model 2 [‡]	Model 3 [§]
<1/month	145/742	1.00	1.00	1.00	1.00
1–3/month	180/958	0.96 (0.77–1.20)	1.06 (0.85–1.32)	1.11 (0.89–1.40)	1.14 (0.90–1.44)
1/week	156/698	1.14 (0.91–1.43)	1.28 (1.02–1.60)	1.30 (1.02–1.64)	1.33 (1.04–1.70)
2/week	206/881	1.20 (0.97–1.49)	1.29 (1.05–1.60)	1.31 (1.05–1.63)	1.33 (1.06–1.68)
3–4/week	172/781	1.13 (0.90–1.41)	1.25 (1.00–1.56)	1.23 (0.98–1.55)	1.26 (0.99–1.61)
5+/week	145/508	1.46 (1.16–1.84)	1.54 (1.22–1.94)	1.43 (1.11–1.82)	1.52 (1.17–1.97)
p for trend		0.0005	0.0001	0.0042	0.0024

^{*} Number of prevalent diabetes / total number of subjects in each category of egg consumption

[†] Model 1 adjusted for age and sex

[‡] Model 2 adjusted for age, sex, smoking, alcohol, BMI, physical activity score, and education

[§] Model 3 adjusted for variables in model 2 plus additional adjustment for energy intake, red meat (including bacon), fiber, dietary magnesium, fruit/vegetables, trans fat, waist circumference, history of hypertension, and history of CVD

Table 3

Prevalence ratios (95% confidence intervals) for prevalent type 2 diabetes by egg consumption stratified by gender*

Egg Consumption	Multivariable Adjusted Model	
	Male	Female
<1/month	1.00 (ref)	1.00 (ref)
1–3/month	1.78 (1.05–3.00)	0.99 (0.75–1.29)
1/week	1.68 (0.98–2.87)	1.26 (0.95–1.67)
2/week	1.71 (1.01–2.88)	1.26 (0.97–1.64)
3–4/week	1.65 (0.97–2.78)	1.19 (0.90–1.58)
5+/week	2.10 (1.24–3.57)	1.38 (1.00–1.90)
p for trend	0.048	0.019

* Adjusted for age, smoking, alcohol, BMI, physical activity score, education, energy intake, intake of red meat and fruits and vegetables, dietary trans fat, magnesium, and fiber, and history of hypertension and CVD; p for sex*egg interaction = 0.53.

Table 4

Least squares means (standard errors) for the logarithmic-transformed A1C, HOMA-IR, and HOMA-B in 3178 non-diabetic subjects*

Egg Consumption	A1C	HOMA-IR	HOMA-B
<1/month	1.71 (0.004)	1.10 (0.02)	5.25 (0.02)
1–3/month	1.70 (0.003)	1.11 (0.02)	5.28 (0.02)
1/week	1.70 (0.004)	1.12 (0.02)	5.28 (0.02)
2/week	1.70 (0.003)	1.10 (0.02)	5.26 (0.02)
3–4/week	1.70 (0.004)	1.08 (0.02)	5.27 (0.02)
5+/week	1.70 (0.01)	1.13 (0.03)	5.27 (0.03)
P linear trend	0.25	0.77	0.72

* Derived from generalized linear model adjusted for age, gender, smoking, BMI, alcohol, physical activity score, and education. A1C, HOMA-IR, and HOMA-B were all logarithmically transformed.

Hazard ratios (95% CI) of diabetes according to egg consumption in the Jackson Heart Study (n=3,564)

Table 5

Egg Consumption	Cases/n [*]	Hazard ratios (95% CI)			
		Crude	Model 1 [†]	Model 2 [‡]	Model 3 [§]
<1/month	92/597	1.00	1.00	1.00	1.00
1–3/month	101/778	0.82 (0.62–1.09)	0.85 (0.64–1.13)	0.83 (0.62–1.11)	0.88 (0.65–1.19)
1/week	79/542	0.92 (0.68–1.24)	0.95 (0.71–1.29)	0.90 (0.66–1.23)	0.94 (0.68–1.30)
2/week	101/675	0.95 (0.71–1.26)	0.97 (0.73–1.29)	0.88 (0.65–1.18)	0.91 (0.66–1.25)
3–4/week	102/609	1.08 (0.82–1.43)	1.12 (0.84–1.49)	1.07 (0.79–1.43)	1.11 (0.81–1.52)
5+/week	56/363	1.05 (0.75–1.47)	1.09 (0.78–1.53)	1.05 (0.74–1.48)	1.17 (0.81–1.70)
p for trend		0.20	0.15	0.34	0.22

^{*} Number of incident diabetes / total number of subjects at risk in each category of egg consumption

[†]Model 1 adjusted for age and sex

[‡]Model 2 adjusted for age, sex, smoking, alcohol, BMI, physical activity score, and education

[§]Model 3 adjusted for variables in model 2 plus additional adjustment for energy intake, red meat (including bacon), fiber, dietary magnesium, fruit/vegetables, trans fat, waist circumference, history of hypertension, and history of CVD