Received:
12 January 2015

Revised:
2 April 2015

Accepted:
13 April 2015

doi: 10.1259/bjr.20150036

Cite this article as:

Adebahr S, Collette S, Shash E, Lambrecht M, Le Pechoux C, Faivre-Finn C, et al. LungTech, an EORTC Phase II trial of stereotactic body radiotherapy for centrally located lung tumours: a clinical perspective. *Br J Radiol* 2015;88:20150036.

ADVANCES IN RADIOTHERAPY SPECIAL FEATURE: REVIEW ARTICLE

LungTech, an EORTC Phase II trial of stereotactic body radiotherapy for centrally located lung tumours: a clinical perspective

^{1,2}S ADEBAHR, ³S COLLETTE, ³E SHASH, ⁴M LAMBRECHT, ⁵C LE PECHOUX, ⁶C FAIVRE-FINN, ⁷D DE RUYSSCHER, ⁸H PEULEN, ⁸J BELDERBOS, ⁹R DZIADZIUSZKO, ¹⁰C FINK, ¹¹M GUCKENBERGER, ⁴C HURKMANS and ^{1,2}U NESTLE

¹Department of Radiation Oncology, University Medical Center Freiburg, Freiburg, Germany

²German Cancer Consortium (DKTK), Heidelberg, Partner Site Freiburg, Germany

³EORTC Headquarters, Brussels, Belgium

⁴Department of Radiation Oncology, Catharina Hospital, Eindhoven, Netherlands

⁵Department of Radiotherapy, Institut Gustave Roussy, Villejuif, France

⁶Institute of Cancer Sciences, Radiotherapy Related Research, The Christie NHS Foundation Trust and University of Manchester, Manchester, UK

⁷KU Leuven–University of Leuven, University Hospitals Leuven, Department of Radiation Oncology, Leuven, Belgium

⁸Department of Radiation Oncology, The Netherlands Cancer Institute Antoni van Leeuwenhoek Hospital, Amsterdam, Netherlands

⁹Department of Oncology and Radiotherapy, Medical University of Gdańsk, Gdańsk, Poland

¹⁰Department of Radiology, Allgemeines Krankenhaus Celle, Celle, Germany

¹¹Department of Radiation Oncology, University of Zurich, Zurich, Switzerland

Address correspondence to: Professor Ursula Nestle

E-mail: ursula.nestle@uniklinik-freiburg.de

ABSTRACT

Evidence supports stereotactic body radiotherapy (SBRT) as a curative treatment option for inoperable early stage non-small-cell lung cancer (NSCLC) resulting in high rates of tumour control and low risk of toxicity. However, promising results are mainly derived from SBRT of peripheral pulmonary lesions, whereas SBRT for the central tumours can lead to severe radiation sequelae owing to the spatial proximity to the serial organs at risk. Robust data on the tolerance of mediastinal structures to high-dose hypofractionated radiation are limited; furthermore, there are many open questions regarding the efficiency, safety and response assessment of SBRT in inoperable, centrally located early stage NSCLC, which are addressed in a prospective multicentre study [sponsored by the European Organization for Research and Treatment of Cancer (EORTC 22113-08113—LungTech)]. In this review, we summarize the current status regarding SBRT for centrally located early stage NSCLC that leads to the rationale of the LungTech trial. Outline and some essential features of the study with focus on a summary of current experiences in dose/fraction-toxicity coherences after SBRT to the mediastinal structures that lead to LungTech normal tissue constraints are provided.

Stereotactic body radiotherapy (SBRT) is a technique in which high doses of radiotherapy are very precisely delivered with steep dose gradients and a short overall treatment time (OTT). This achieves a very high biological dose.¹ Evidence supports SBRT as a curative treatment option for inoperable early stage non-small-cell-lung cancer (NSCLC).^{2–5} High-precision, hypofractionated dose delivery enables a significant reduction in both target volume size and exposure of normal tissue (NT) to high doses, resulting in a decreased toxicity risk and high rates of local control.^{6–8} However, promising results are mainly derived from the SBRT of small peripheral pulmonary lesions with low risk of treatment-related toxicity, as these tumours are surrounded by parallel organs at risk

(OARs), specifically lung tissue. In contrast, the SBRT for central tumours can lead to severe, potentially life-threatening radiation sequelae owing to the spatial proximity to serial OARs. As robust, prospective and multicentre data on the tolerance of mediastinal structures to high-dose hypofractionated radiation is limited, there is generally caution in implementing SBRT for central lung tumours. Current published data are rather inconsistent regarding the definition of “central tumour”, SBRT techniques, dose prescription and reporting, calculation algorithms, image guidance and evaluation of outcome. A recent systematic review has summarized the current limited evidence on this topic, mainly from single-centre experiences,⁹ but there are still many open questions

regarding patient selection, efficiency, safety and response assessment, which hamper the use of SBRT for central lung tumours in routine practice.¹⁰ A prospective study sponsored by the European Organization for Research and Treatment of Cancer (EORTC 22113-08113-LungTech) was launched in late 2014 and is expected to answer the question of the efficacy and toxicity of SBRT in inoperable and centrally located early stage NSCLC in a multicentre setting.

NSCLC: TREATMENT OPTIONS FOR EARLY STAGE NSCLC

The incidence and mortality of lung cancer in the European Union are 52/100.000 and 47/100.000 per year, respectively.¹¹ The NSCLC accounts for approximately 80%¹¹ of all cases with a poor 5-year survival rate of 16%, mainly owing to patients being diagnosed at advanced stages. If diagnosed at an earlier stage, 5-year survival for NSCLC can be as high as up to 80%.^{12,13} The current standard of care for early stage tumours is anatomic surgical resection in medically fit patients, consisting of lobectomy or pneumonectomy accompanied by a systematic mediastinal lymph node sampling or lymphadenectomy.^{13–15} Associated 5-year survival rates are commonly accepted to be 60–80% for Stage I and 40–50% for Stage II NSCLC.¹³ The efficacy of routine sublobar resections for tumours <2 cm is currently being investigated worldwide. This technique spares lung tissue and preserves pulmonary function when compared with lobectomy. However, the results of a randomized controlled trial indicate that this is associated with inferior oncological outcome.¹⁶ In contrast to peripheral tumours, those located centrally often show compression and/or invasion in vessels, major bronchi or other critical mediastinal structures and, therefore, require more extensive surgical procedures associated with higher mortality and morbidity.^{17–19} Approximately 20% of all patients with NSCLC Stage I are medically inoperable because of poor general condition or coexisting morbidities such as chronic obstructive pulmonary disease and/or heart disease,²⁰ and <50% of all patients with early stage NSCLC older than 75 years undergo surgery.²¹ It is expected that as the global population ages and lung cancer screening of high-risk populations is implemented,²² the proportion of inoperable patients with lung cancer with comorbidities will increase.^{23–25} In Stage I patients with NSCLC who refuse surgery and do not receive other treatments such as radiotherapy, 5-year overall survival (OS) and cancer-specific survival (CSS) are low, 6% and 16%, respectively.²⁶ The standard of care treatment option for medically inoperable patients with early stage NSCLC has been conventionally fractionated radiotherapy (CFRT), superior to best supportive care with CSS rates of about 30% after 5 years^{27,28} and 5-year OS of 29–37% for T1 tumours.²⁹ Evidence supports a clear dose effect for local control and survival in NSCLC,^{30,31} however, traditional CFRT-planning and delivery techniques are associated with low accuracy and broad safety margins resulting in increased toxicity rates,³² therefore, limiting the delivery of higher doses in inoperable patients with early stage NSCLC. Although advanced techniques and positron emission tomography (PET) staging allow modern three dimensional (3D)-CFRT the application of high biological doses, resulting in local control rates >85% in Stage I patients with NSCLC,^{33,34} the implementation of such 3D-CFRT regimes leads to longer OTT of approximately 3–6 weeks,^{33,34} whereas SBRT usually can be performed in OTT of days rather than weeks.

SBRT AS AN EFFICIENT TREATMENT MODALITY FOR PERIPHERALLY LOCATED NSCLC

In the past two decades, SBRT has been accepted as a curative treatment alternative for inoperable patients with small (<5 cm) peripheral early stage NSCLC.^{2–5} Several prospective studies reported excellent local tumour control rates of up to approximately 90%,^{5,6,8,35,36} in the same range as those obtained with surgery.^{37,38} Low rates of toxicity have been observed, including elderly, medically, inoperable patients with severe comorbidities.^{6,8,35,36,39,40} Survival after SBRT is modest (approximately 50% at 3 years) but has to be interpreted in the context of patients with multiple comorbidities. As expected, survival after SBRT is higher in fitter, operable patients who refused surgery.⁴¹ Furthermore, population-based studies indicated improved OS in medically inoperable patients following the introduction of SBRT.⁴² Based on these data, SBRT for patients with early stage NSCLC has been recommended as the standard of care for medically inoperable cases by the European Society For Medical Oncology, Lugano, Switzerland,⁴³ the National Comprehensive Cancer Network, Fort Washington, PA¹⁴ and by the Deutsche Gesellschaft für Radioonkologie e.V., Berlin, Germany.²⁰

Although histological or cytological confirmation of NSCLC is recommended before performing SBRT, biopsy is sometimes not feasible or associated with an unacceptably high risk because of the severe comorbidities of the patient. Thus, an inevitable percentage of patients are treated based on clinical suspicion and imaging criteria of malignancy only. However, in Western Europe, the likelihood of having NSCLC has been demonstrated to be well over 90% using a calculation based on clinical parameters, CT characteristics of malignancy and significant uptake of fluorine-18 fludeoxyglucose (¹⁸F-FDG) on PET scan.⁴³ Moreover, when malignancy is highly likely based on these imaging criteria, SBRT without pathological confirmation was shown to lead to the same outcome as in pathology-proven NSCLC.⁴⁴

With regard to dose/fractionation, there is vast heterogeneity in the SBRT regimes used routinely. Internationally, 3 × 18 Gy is one of the most frequently used regimes, whereas in Germany for small peripheral lesions, lower doses of 3 × 13.5–15.0 Gy are recommended²⁰ and have been demonstrated to result in local control rates of >90%.⁴⁵ Local tumour control has repeatedly been reported to show a dose-dependent increase with a minimum biologically effective dose (BED; α/β ratio, 10 Gy) of 100 Gy to the planning target volume (PTV) surrounding isodose resulting in local tumour control rates >90%,^{46,47} that even translates into improved OS.^{46,48} Interestingly, in a recent meta-analysis, Zhang *et al*⁴⁹ demonstrated a plateau effect with regard to OS with doses of 83.2–146.0 Gy BED and a detrimental effect on OS with doses >146 Gy.

SBRT FOR CENTRALLY LOCATED NSCLC

The major difference between peripherally and centrally located NSCLC is the spatial proximity to centrally located serial OARs, such as, main airways, large blood vessels, the heart, the oesophagus, the phrenic nerves or the brachial plexus, where hypofractionated high doses might lead to severe, potentially life-threatening consequences. Several articles have reported high rates of toxicity after SBRT of centrally located NSCLC. A prospective study by Timmerman *et al*⁵⁰ from the Indiana

University reported a 2-year freedom from severe toxicity of only 54% for central tumours compared with 83% for peripheral tumours, in 70 patients treated with 60–66 Gy total in three fractions. Furthermore, SBRT may have contributed to the events leading to the death of six patients (four patients from bacterial pneumoniae; one patient from pericardial effusion; and one patient owing to massive haemoptysis), of which four of the six patients had central tumours.⁵⁰ These deaths occurred after a median of 10.4 months following SBRT (range, 1–20 months). The 2-year incidence of toxicity grade ≥ 3 was 17% and 46% for peripherally and centrally tumours, respectively. The observation of severe toxicity after SBRT of centrally located NSCLC was subsequently confirmed by two studies: grade 5 toxicity was observed in 1 out of 17 patients after treatment with 60 Gy in 4 fractions³⁶ and in 1 out of 9 patients after irradiation with 48 Gy in 4 fractions⁵¹ delivered on consecutive days. Moreover, a case report was published by the University of Pennsylvania, Philadelphia, PA, reporting on a fatal central airway necrosis more than 8 months after 50 Gy in five fractions was delivered to a central lung tumour.⁵² Conversely, investigators from the Free University in Amsterdam demonstrated that the use of “risk-adapted SBRT” using a more fractionated regime (60 Gy in eight fractions) did not result in excess toxicity for centrally located early stage lung tumours ($n = 63$), and clinical outcomes were comparable with those seen for peripheral tumours.⁵³ However, none of the retrospective data included detailed dose–volume histogram/toxicity analyses. A PRISMA structured literature review published in 2013 identified 20 publications reporting outcomes for 563 central lung tumours treated with SBRT, including 315 patients with early stage NSCLC.⁹ The majority of these studies were retrospective and conducted at single institutions. Only four studies were prospective and reported on 68 patients with central tumours. Local control was 85% when the prescribed BED₁₀ was >100 Gy. Tumour location (central vs peripheral) did not impact on OS, and treatment-related mortality was 2.7% overall. Grade 3/4 toxicities following SBRT for central tumours were more common than for peripheral tumours, and occurred in $<9\%$ of patients. It should however be noted that there was heterogeneity in the common toxicity criteria used to define toxicities and treatment–response criteria. Furthermore, the follow-up of these studies was relatively short (median, 18 months), and long-term toxicity data are needed given the high dose and hypofractionated nature of the treatments delivered. The BED₁₀ delivered ranged between 60 and 180 Gy. It is important to note that published studies vary widely in their reporting of dose/fractionation as well as dose specifications and calculations. Only 11 of the 20 reviewed studies used tissue heterogeneity correction, and the prescription isodoses ranged between 50% and 100%.

In summary, there is a clear need for a prospective, multicentre study using standardized RT techniques, treatment–response criteria and toxicity assessment. As the maximum tolerated doses and optimal fractionation schemes for SBRT near mediastinal structures are currently unknown, collective knowledge of the complex relationship between the radiation dose and the volume of irradiated tissue for each mediastinal structure is a pre-requisite to the safe future routine use of SBRT for central tumours. The lack of high-quality evidence to support SBRT in centrally located tumours is reflected in

international guidelines that only recommend SBRT for peripheral and medically inoperable early stage NSCLC.^{43,54} We therefore advocate that treatment of central lesions should preferably be performed within the context of a well-conducted prospective trial.

EORTC 22113-08113 LungTech: CONCEPT OF THE TRIAL

The LungTech trial aims to evaluate the efficiency and toxicity of SBRT for patients with histologically or cytologically proven early stage, centrally located, inoperable NSCLC in a multicentre setting. The primary end point will be freedom from local progression at 3 years as assessed by serial CT scans and confirmed by ¹⁸F-FDG-PET/CT. If still unclear, biopsies or 3 monthly repeated imaging will be performed. Using one-sided 5% Type I error and 80% power, 150 patients (included so far from 23 planned centres in Belgium, France, Germany, Poland, Switzerland and UK) should be sufficient to reject the hypothesis of freedom from the local progression rate at 3 years of $\leq 80\%$ under the assumption that SBRT is expected to achieve a freedom from local progression rate of 90%. The trial will also investigate acute and late toxicities as well as patterns of local and distant recurrence, including mediastinal failure, assessed by serial CT scans and confirmed by ¹⁸F-FDG-PET/CT and subsequent repeat imaging or biopsy, if necessary. Moreover, this study offers a unique opportunity to evaluate the role of ¹⁸F-FDG-PET/CT to monitor disease progression and toxicity, as translational research end points, including staging comparison between 3D- and four-dimensional (4D) ¹⁸F-FDG-PET/CT assessment, possible impact of 4D ¹⁸F-FDG-PET/CT on target volume contouring and external validation of Huang et al⁵⁵ response criteria.

Recently, the Radiation Therapy Oncology Group (RTOG) has performed a dose-escalation Phase I/II study (RTOG 0813) in centrally located NSCLC.⁵⁶ Derived from the North American standard, the study is based on a five-fraction regime, the results are awaited. Although the RTOG study includes only T1/T2 tumours, the LungTech study will also include a limited population of T3 tumours and therefore supplement the available data pool. In Europe, owing to the risk of toxicity with three- to five-fraction regimes as described above, it was decided to investigate a less hypofractionated “risk-adapted” approach (60 Gy in eight fractions) as reported by the Free University in Amsterdam.⁵³ To ensure the feasibility and safety of the planned treatment, only those centres that have used SBRT for a period of at least 12 months and have treated a minimum of 20 patients will be eligible. The study will be conducted only within EORTC centres and will benefit from the EORTC and Radiation Oncology Group Quality Assurance infrastructure. A pooled data analysis of both studies maybe an interesting future option.

Some essential features of the protocol are outlined below.

Definition of central tumour

In current literature, the definition of what constitutes a “central” tumour varies widely and further hampers the comparison between trials. Only 11 out of the 20 studies reviewed by Senthil et al⁹ used the same definition as RTOG 0813;⁵⁶ Modh et al⁵⁷

even included two different definitions of “central tumours” within the same publication of their study.

Haasbeek *et al*⁵³ defined central tumours as located within a 1-cm zone from the mediastinal envelope. In RTOG 0236,⁵⁸ the following definition is used: a tumour within 2 cm of the proximal bronchial tree (carina, right and left main bronchi, and bronchial tree to the second bifurcation), whereas RTOG 0813 sticks to the definition of Timmerman *et al*:⁵⁰ a tumour located within 2 cm in all directions around the proximal bronchial tree and immediately adjacent to mediastinal or pericardial pleura with a PTV expected to touch or include the pleura. This definition has also been slightly modified in the LungTech trial: “Centrally located tumour defined as tumour within 2 cm or touching the zone of the proximal bronchial tree or tumour that is immediately adjacent to the mediastinal or pericardial pleura, with a PTV expected to touch or include the pleura.”

Some tumours might be “too central” to be safely treated with SBRT when applying BED >100 Gy. Even using the “risk-adapted” LungTech regime, there will be central tumours, for example, invading the proximal bronchial tree and/or hilar structures, which will not be treatable within the trial owing to a very high risk of severe toxicity (Figure 1). In current literature, a selection bias with regard to the inclusion of certain anatomical subgroups within the SBRT “danger zone” cannot be excluded. Indeed, the examples and description of patient cohorts, for example, in the Haasbeek’s study⁵³ show that “very central” tumours have not been included in the larger published series. However, a clear definition of what constitutes “too central tumours” is lacking. In the LungTech protocol, T3 tumours >7 cm, all T4 tumours and tumours abutting the oesophagus or presenting with separate tumour nodule(s) in the same lobe are excluded to increase the safety of this treatment. Furthermore, all potential LungTech cases will undergo a central expert review of eligibility relating to tumour anatomical location and treatment planning before enrolment in the study treatment, thus confirming central tumour localization and excluding tumours that are “too central” according to the eligibility criteria and dose constraints (e.g. proximity to the central bronchi, Figure 1).

Prescribed dose, normal tissue constraints

In order to define dose/fractionation and NT constraints for the LungTech trial, current literature and experiences in other trials were reviewed (Tables 1 and 2).^{9,36,50–53,56,57,59–75}

In terms of tumour dose, the LungTech protocol will investigate a medium hypofractionated approach, applying 8.0×7.5 Gy to a total dose of 60 Gy; this is equal to a BED of 105 Gy ($\alpha/\beta = 10$). Referring to report 83 of the International Commission on Radiation Units and Measurements (ICRU), 95% of the PTV has to receive at least the nominal fraction dose, and 99% of the PTV receives a minimum of 90% of the nominal dose. The maximum dose within the PTV should not be <110%, nor should it exceed 130% of the prescribed dose. Gross tumour volume (GTV) delineation is based on 4D-CT in treatment position, mandatory 3D-PET/CT scan and

supplementary clinical information, for example, results of bronchoscopy. To account for tumour motion during the breathing cycle, the protocol allows the individual internal target volume (ITV) and the average mean position approach. The PTV margin based on the ITV concept is primarily meant to take into account patient set-up uncertainties, thus requiring an isotropic ITV expansion of 3–5 mm. For PTV generation, based on the average mean position of the tumour, the margin should take into account both the set-up error and breathing-induced motion and should not be <3 mm. All patients treated in this trial will receive image-guided SBRT. Further details on the technical aspects of the radiotherapy planning including dose specification and radiation therapy quality assurance (RTQA) procedures will be reported elsewhere.

Defining NT constraints for thoracic OARs, in the context of SBRT, is a major challenge, as the maximum-tolerated doses and optimum fractionation for mediastinal structures are currently unknown. Another challenge is the assessment of treatment-related toxicity in patients with multiple comorbidities, which

Figure 1. (a) An 81-year-old patient with centrally located non-small-cell lung cancer (NSCLC), T1N0M0, meeting the eligibility criteria for inclusion in the LungTech trial. Delineation of tumour: gross tumour volume (GTV) (purple), internal target volume (ITV) (red), planning target volume (PTV) (pink) and organs at risk. (b) The proximity to the right central bronchus (cyan) does not allow planning according to the LungTech protocol without exceeding the dose constraint for the proximal bronchus tree (prox BT) 44 Gy [equivalent dose in 2 Gy fractions (EqD2), 74.8 Gy]—maximum dose for prox BT here 66.76 Gy (EqD2, 151.3 Gy)—leading to secondary exclusion from the trial.

Table 1. Literature review of severe radiation induced toxicities for central lung tumors for definition of normal tissue (NT) constraints for the LungTech trial

Study	Number of reported patients (treated tumours)	Number of centres	p/r	Results ^a
Bronchial tree/trachea (α/β , 3 Gy), potential side effects: fatal haemoptysis, fistula, stenosis, necrosis, atelectasis, pneumonia and abscess				
Timmerman et al ⁵⁰	70	1	p	Central lesions: 11× more likely to experience grade 3–5 lung toxicity than those with peripheral tumours, 1× local recurrence next to carina with subsequent fatal haemoptysis (19.5 months after SBRT) 3× 20–22 Gy = 60–66 Gy (EqD2: 276–330 Gy) ^b
Fakiris et al ⁵⁹	22	1	p	4-year results of above study: lung toxicity rates of 10.4% (peripheral) and 27.3% (central lesions); 1× fatal haemoptysis (same as above), three patients died of pneumonia
Li et al ⁶⁰	43	1	p	One case of fatal haemoptysis, 70 Gy, ten fractions, hilar $D_{\max}$ = 83 Gy (EqD2, 187.6 Gy)
Modh et al ⁵⁷	91	1	r	Two cases of fatal haemoptysis, 47 Gy (EqD2, 116 Gy) and 48 Gy (EqD2, 121 Gy), five fractions. Tumours were found involving the hilum and encasing the left superior segmental bronchus, respectively
Corradetti et al ⁵²	1	1	r	Central airway necrosis, 5 × 10 Gy = 50 Gy (EqD2, 130 Gy) ^b
Nishimura et al ⁶¹	133	1	r	Fatal haemoptysis in two cases with $D_{\max}$ >50 Gy (EqD2, 130 Gy) to the pulmonary artery /bronchus (five fraction regime)
Song et al ⁵¹	9	1	r	8× partial or complete bronchial strictures, 1× complete bronchial stricture with fatal consequences (bleeding, aspiration and pneumonia), 4 × 12 Gy = 48 Gy (EqD2, 144 Gy) ^b
Milano et al ⁶²	53	1	r	1× fatal haemoptysis (bronchus received a cumulative dose of 98 Gy; EqD2 not applicable)
Oshiro et al ⁶³	21	1	r	1× fatal haemoptysis (re-treatment: 1× 25 Gy; EqD2, 140 Gy) ^b
Bral et al ³⁶	17	1	p	1× bronchial stenosis and successively fatal haemoptysis after stent insert, 4 × 15 Gy = 60 Gy (EqD2, 216 Gy) ^b
Unger et al ⁶⁴	17	1	r	1× bronchial fistula, mainstem bronchus received a maximum point dose of 49 Gy (EqD2 not applicable)
Canon et al ⁶⁵	75 ^c	1	p	CFRT: EqD2 of 75×3 Gy predicting a 5% complication rate, 3× fatal haemoptysis, 85 and 75 Gy, 25 fractions (EqD2, 118 and 90, respectively), tumours encasing or abutting a mainstem or proximal lobar bronchus and partially local invasion of adjacent normal structures
Heart (α/β , 3 Gy), potential side effects: congestive heart failure, pericarditis, pericardial effusion and arrhythmia				
Bonomo et al ⁶⁸	16	1	r	Paracardiac and cardiac lesions treated with SBRT (36 Gy, three fractions, $D_{100\%}$: 51.4 Gy; EqD2, 108–204 Gy): no cardiological symptoms or electrocardiographic abnormalities, even months after SBRT

(Continued)

Table 1. (Continued)

Study	Number of reported patients (treated tumours)	Number of centres	p/r	Results ^a
Great vessels (aorta, vena cava superior and inferior brachiocephalic veins) (α/β , 3 Gy), potential side effects: haemoptysis and fatal bleeding				
Timmerman <i>et al</i> ⁵⁰	70	1	p	Single cases of haemoptysis and fatal bleeding with varying SBRT regimens (see also Brochial tree/trachea)
Senthi <i>et al</i> ⁹	(563)	20 ^d	r/p (4)	Single cases of haemoptysis and fatal bleeding with varying SBRT regimens (Song <i>et al</i> ⁵¹ , Milano <i>et al</i> ⁶² , Oshiro <i>et al</i> ⁶³ , Braley <i>et al</i> ³⁶)
Canon <i>et al</i> ⁶⁵	75 ^c	1	p	(see also Brochial tree/trachea)
Oesophagus (α/β , 3 Gy), potential side effects: fistula, stenosis, perforation, oesophagitis, ulcer, haemorrhage				
Onimaru <i>et al</i> ⁶⁹	45(57) ^c	1	p	One death owing to radiation-induced ulcer in the oesophagus 5 months after SBRT, 48 Gy, eight fractions (EqD2, 86.4 Gy) ^b , maximum dose of 50.5 Gy at the oesophagus (EqD2, 93.7 Gy)
Stephans <i>et al</i> ⁷⁰	52	1	r	Two cases of oesophageal fistula, when the oesophageal point dose >51 Gy and 1-cm ³ doses >48 Gy, EqD2 not applicable
Modh <i>et al</i> ⁵⁷	91	1	r	One fistula with an oesophageal $D_{\max}$ of 46 Gy in five fractions (EqD2, 112 Gy). Oesophageal toxicity $\geq$ grade 2: 12.8% (median $D_{\max}$ of 29.5 Gy for those patients with oesophageal toxicity), EqD2 could not be derived from those data
Spinal cord (α/β , 1-3 Gy), potential side effects: radiation-induced myelitis				
Kirkpatrick <i>et al</i> ⁷¹	335 ^g 1946 ^g 1400 ^f	5 ^d 12 ^d 9 ^d	r r r	CFRT of 2 Gy per day including the full cord cross-section: total dose of 50 Gy, 60 Gy and approximately 69 Gy (= EqD2) are associated with a 0.2, 6 and 50% rate of myelopathy. For partial cord irradiation as part of spine radiosurgery, maximum cord dose of 13 Gy in one fraction (EqD2, 48.8 Gy) or 20 Gy in three fractions (EqD2, 110 Gy) appear associated with a <1% risk of injury
Brachial plexus (α/β , 3 Gy), potential side effects: brachial plexopathy				
Forquer <i>et al</i> ⁷²	(37 ^e)	2	r	37 apical lesions treated to a median total dose of 57 Gy, median maximum brachial plexus dose of patients developing brachial plexopathy: 30 Gy (18–82 Gy) (EqD2 not applicable). 7/37 apical lesions developed grade 2–4 plexopathy. Brachial plexus maximum dose should be kept <26 Gy in three (EqD2, 59.9 Gy) or four (EqD2, 49.4 Gy) fractions
Lungs (α/β , 3 Gy), potential side effects: radiation-induced pneumonitis, fibrosis and decrease in lung function, atelectasis, pneumonia (see also Bronchial Tree /Trachea) and abscess				
Borst <i>et al</i> ⁷³	128 ^c	1	r	No difference between SBRT and CFRT for the relationship between the lung dose and the incidence of radiation-induced pneumonitis
Stanic <i>et al</i> / RTOG0236 ⁷⁴	55	43	p	No clinically significant changes in pulmonary function following SBRT for early stage peripheral NSCLC
Unger <i>et al</i> ⁶⁴	17	1	r	1 $\times$ grade 3 radiation pneumonitis (EqD2 not applicable)

(Continued)

Table 1. (Continued)

Study	Number of reported patients (treated tumours)	Number of centres	p/r	Results ^a
Chest wall (α/β , 3 Gy), potential side effects: chest wall pain and rib fractures				
Taremi et al ⁷⁵	46 (49 ^c)	1	r	Description of risk factors for radiation induced bone injury after SBRT: increasing age, female gender and high RT dose to 0.5 cm ³ of nearby ribs

CFRT, conventional radiotherapy; $D_{\max}$, maximum dose; EqD2, equivalent dose in 2 Gy fractions; NSCLC, non-small-cell lung cancer; p, prospective; r, retrospective; RT, radiotherapy; RTOG, Radiation Therapy Oncology Group; SBRT, stereotactic body radiotherapy.

The number within parenthesis for column "p/r" are the number of treated tumours.

^aMaximum point dose or dose/fractionation and EqD2 in Gray provided if possible.

^bDoses applied to the tumour.

^cIncluding peripheral and central tumours.

^dStudies, ^eapical of 276, ^ftreated with stereotactic fractionated radiotherapy (SBRT), ^gtreated with CFRT.

may be a confounding factor. Whilst toxicity for SBRT delivered to peripheral tumours is well documented, this is not the case for central tumours, and long-term data are necessary given that late effects can become apparent more than 1 year post treatment.⁶⁵ For this reason, patients will be prospectively assessed for at least 3 years after treatment in the LungTech study.

Although the linear quadratic model (LQM) has been criticized for not being applicable to SBRT,⁷⁶ data from Guckenberger et al⁷⁷ suggested accurate modelling of local tumour control in fractionated SBRT for Stage I NSCLC with the traditional linear-quadratic formalism. Brown et al⁷⁸ reported that there is compelling *in vitro* and *in vivo* NT evidence that the LQM

Table 2. Normal tissue (NT) constraints for the LungTech trial derived from literature review of severe toxicities (Table 1) and used NT constraints

Study	Recommendations/NT constraints
Bronchial tree/trachea (α/β , 3 Gy), potential side effects: fatal haemoptysis, fistula, stenosis, necrosis, atelectasis, pneumonia and abscess	
Timmerman et al ⁶⁶	Maximum point dose, 20.2 Gy (one fraction regime; EqD2, 93.7 Gy) 40 Gy (five fraction regime; EqD2, 88 Gy)
RTOG 0813 ⁵⁶	>105% PTV, EqD2 not applicable
Haasbeek et al ⁵³	Maximum dose restriction (0.5 cm ³), $8 \times 5.5 = 44$ Gy (EqD2, 74.8 Gy)
Nuytens et al ⁶⁷	Maximum dose restriction (0.5 cm ³), $6 \times 8 = 48$ Gy (EqD2, 105.6 Gy)
EORTC 22113-08113	Dose constraint, 8×5.5 Gy = 44 Gy (EqD2, 74.8 Gy)
Heart (α/β , 3 Gy), potential side effects: congestive heart failure, pericarditis, pericardial effusion and arrhythmia	
Timmerman et al ⁶⁶	Maximum point dose, 22 Gy (one fraction regime; EqD2, 110 Gy) 38 Gy (five fraction regime; EqD2, 80.6 Gy)
RTOG 0813 ⁵⁶	Maximum point dose: 63 Gy (five fraction regime; EqD2, 196 Gy) 60 Gy (ten fraction regime; EqD2, 108 Gy)
EORTC 22113-08113	No restrictions, but recording of DVH data for toxicity
Great vessels (aorta, vena cava superior and inferior, brachiocephalic veins) (α/β , 3 Gy), potential side effects: haemoptysis and fatal bleeding	
Timmerman et al ⁶⁶	Maximum point dose, 37 Gy (one fraction regime; EqD2, 296 Gy) 53 Gy (five fraction regime; EqD2, 144.2 Gy)
RTOG 0813 ⁵⁶	Maximum point dose, 63 Gy (five fraction regime; EqD2, 196.6 Gy) 75 Gy (ten fraction regime; EqD2, 157.5 Gy)
EORTC 22113-08113	No restrictions, but recording of DVH data for toxicity
Oesophagus (α/β , 3 Gy), potential side effects: fistula, stenosis, perforation, oesophagitis, ulcer, haemorrhage	
Timmerman et al ⁶⁶	Maximum point dose, 15.4 Gy (one fraction regime; EqD2, 56.7 Gy) 35 Gy (five fraction regime; EqD2, 70 Gy)

(Continued)

Table 2. (Continued)

Study	Recommendations/NT constraints
RTOG 0813 ⁵⁶	Maximum point dose, 63 Gy (five fraction regime; EqD2, 196 Gy) 50 Gy (ten fraction regime; EqD2, 80 Gy)
Haasbeek <i>et al</i> ⁵³	Maximum dose restriction (0.5 cm^3), $8 \times 5 = 40 \text{ Gy}$ (EqD2, 66 Gy)
Nuytens <i>et al</i> ⁶⁷	Maximum dose restriction (0.5 cm^3), $6 \times 6 = 36 \text{ Gy}$ (EqD2, 64.8 Gy)
EORTC 22113-08113	Dose constraint, $8 \times 5 \text{ Gy} = 40 \text{ Gy}$ (EqD2, 64 Gy)
Spinal cord (α/β , 1-3 Gy), potential side effects: radiation induced myelitis	
Timmerman <i>et al</i> ⁶⁶	Maximum point dose, 14 Gy (one fraction regime, EqD2: 56 Gy) 30 Gy (five fraction regime; EqD2, 60 Gy)
RTOG 0813 ⁵⁶ (56)	Maximum point dose, 25 Gy (four fractions regime; EqD2, 51.66 Gy) 30/40 Gy (five/ten fraction regime; EqD2, 60 Gy)
Haasbeek <i>et al</i> ⁵³	Maximum dose restriction (0.5 cm^3), $8 \times 3.5 = 28 \text{ Gy}$ (EqD2, 36 Gy)
Nuytens <i>et al</i> ⁶⁷	Maximum dose restriction (0.5 cm^3), $6 \times 4.5 = 27 \text{ Gy}$ (EqD2, 43.9 Gy)
EORTC 22113-08113	Dose constraint, $8 \times 4 \text{ Gy} = 32 \text{ Gy}$ (EqD2, 48 Gy)
Brachial plexus (α/β , 3 Gy), potential side effects: brachial plexopathy	
Timmerman <i>et al</i> ⁶⁶	Maximum point dose, 17.5 Gy (one fraction regime; EqD2, 71.8 Gy) 30.5 Gy (five fraction regime; EqD2, 55.5 Gy)
RTOG 0813 ⁵⁶	Maximum point dose, 32 Gy (five fraction regime; EqD2, 60.2 Gy) 55 Gy (five/ten fraction regime; EqD2, 93.5 Gy)
Haasbeek <i>et al</i> ⁵³	Maximum dose restriction (0.5 cm^3), $8 \times 4.5 = 36 \text{ Gy}$ (EqD2, 54 Gy)
Nuytens <i>et al</i> ⁶⁷	Maximum dose restriction (0.5 cm^3), $6 \times 8 = 48 \text{ Gy}$ (EqD2, 105.6 Gy)
EORTC 22113-08113	Dose constraint, $8 \times 4.75 \text{ Gy} = 38 \text{ Gy}$ (EqD2, 58.9 Gy)
Lungs (α/β , 3 Gy), potential side effects: radiation-induced pneumonitis, fibrosis and decrease in lung function, atelectasis, pneumonia (see also Brochial tree/trachea) and abscess	
RTOG 0813 ⁵⁶	$V12.5 > 1500 \text{ cm}^3$ and $V13.5 \text{ Gy} < 1000 \text{ cm}^3$; EqD2 not applicable
EORTC 22113-08113	No restrictions, but recording of DVH data for toxicity
Chest wall (α/β , 3 Gy), potential side effects: chest wall pain and rib fractures	
Timmerman <i>et al</i> ⁶⁶	Maximum point dose, 30 Gy (one fraction regime; EqD2, 198 Gy) 43 Gy (five fraction regime; EqD2, 99.8 Gy)
RTOG 0813 ⁵⁶	Maximum point dose 32 Gy (five fraction regime; EqD2, 60.2 Gy) 82 Gy (ten fraction regime; EqD2, 183 Gy)
EORTC 22113-08113	No restrictions, but recording of DVH data for toxicity

DVH, dose-volume histogram; EORTC, European Organization for Research and Treatment of Cancer; EqD2, equivalent dose in 2 Gy fractions; PTV, planning target volume.

provides a reasonable estimate of dose-response relationships including single high doses. Therefore, the NT constraints defined in LungTech have been derived from the available literature on SBRT coupled with LQM estimates related to CFRT experience (Tables 1 and 2).

While translating stated dose/fraction from the varying data to comparable equivalent dose in 2 Gy fractions (EqD2), wide ranges of allegedly comparable equivalent doses become apparent (Tables 1 and 2). This is most likely because different α/β ratios have been used and/or other factors than LQM estimates might have contributed to dose calculation or recommendation. This obvious heterogeneity underlines the need for systemically collected data for defining maximum-tolerated

doses and optimum fractionation for mediastinal structures, and thus robust NT constraints for thoracic OARs. The dose constraints and possible deviations chosen for the LungTech trial are presented in the EORTC 22113-8113RTQA guidelines (Table 3).

Primary end point assessment

Clinical symptoms of acute radiation-induced lung injury develop within approximately 3–6 months after treatment. A proportion of patients will subsequently develop radiation fibrosis (from 6 months after SBRT). The assessment of tumour response after SBRT by CT-based criteria [response evaluation criteria in solid tumour (RECIST)] is therefore challenging, as such changes are typically seen within the high

Table 3. EORTC 221133 LungTech trial: dose constraints for organs at risk (OARs)

OAR	α/β (Gy)	$D_{\max}$ (Gy)	EqD2 (Gy)	Acceptable variation (Gy)	Acceptable variation EqD2 (Gy)	Unacceptable variation (Gy)	Unacceptable variation EqD2 (Gy)
Trachea/main bronchus	3	$8 \times 5.5 = 44$	74.8	$<8 \times 5.81 = 46.68$	<81.9	$\geq 8 \times 5.81 = 46.68$	>81.9
Heart ^a	3						
Great vessels ^a	3						
Oesophagus	3	$8 \times 5 = 40$	64	$<8 \times 5.44 = 43.52$	<73.6	$\geq 8 \times 5.44 = 43.52$	≥ 73.6
Spinal cord ^b	2	$8 \times 4 = 32$	48			$>8 \times 4 = 32$	>48
Brachial plexus ^b	3	$8 \times 4.75 = 38$	58.9	$<8 \times 5.17 = 41.36$	<67.7	$\geq 8 \times 5.17 = 41.36$	≥ 67.7
Body-PTV ^b	3	$8 \times 7.5 = 60$	126	$<8 \times 7.785 = 62.28$	<134.2	$\geq 8 \times 7.785 = 62.28$	≥ 134.2
Lung-CTV ^a	3						
Chest wall ^a	3						

CTV, clinical target volume; DVH, dose-volume histogram; EORTC, European Organization for Research and Treatment of Cancer; EqD2, equivalent dose in 2 Gy fractions; PTV, planning target volume.

Source: EORTC 22113-0813-LungTech radiation therapy quality assurance guidelines.

^aNo restrictions are provided but recording of DVH data for toxicity evaluation is required.

^bFor $<0.5 \text{ cm}^3$.

dose area. However, some changes, independently of the tumour size change, have been identified as more reliable indicators of local recurrence (e.g. opacity with a convex border, disappearance of air bronchograms⁵⁵). In order to confirm a suspicion of local recurrence, it has been shown that ^{18}F -FDG-PET/CT may play an important role in increasing the sensitivity of the diagnosis. In a systematic review of the literature, Huang et al⁵⁵ proposed a set of criteria for the evaluation of local recurrence with CT and PET findings in patients with primary lung tumours or lung metastases treated with SBRT. A value of maximum standard value uptake ($\text{SUV}_{\max}$) > 5 was suggested^{79,80} for the differential diagnosis between local recurrence and post-treatment changes. However, the quantitative measurement criteria set by Huang et al,⁵⁵ are derived from various single-centre studies with heterogeneous PET imaging protocols. Therefore, Huang et al⁵⁵ recommend conducting all PET scans for a specific patient on the same machine and with a standardized scanner. This is obviously not possible in the context of a multicentre trial.

In the LungTech trial, ^{18}F -FDG-PET/CT will be requested in case of equivocal findings or progressive soft-tissue abnormalities. The integration of quantitative measurement criteria would require the standardization of all the ^{18}F -FDG-PET/CT scanners. Given that such standardization, e.g. the “European Association of Nuclear Medicine” (EANM) accreditation, is not used in all the participating centres, a modified Huang et al⁵⁵ criterion is being applied with local progression defined as “focal ^{18}F -FDG accumulation significantly above the mediastinal blood pool” (Figure 2). However, quantitative assessment ($\text{SUV}_{\max}$) of the treated tumour and regional lymph nodes will also be measured and collected in order to prospectively evaluate the robustness of the criteria set by Huang

et al⁵⁵ in a subset of centres that have EANM accreditation. In difficult cases, reimaging or biopsy will be requested to confirm progression if clinically indicated.

Toxicity assessment

An important secondary end point of the LungTech trial is the prospective evaluation of toxicity. As previously described, there is a paucity of prospective data regarding the relationship of dose fractionation, irradiated volume and toxicity of SBRT in centrally located tumours. One of the objectives of LungTech is to elaborate on such coherences and thus be able to provide robust recommendations on central NT constraints in patients treated with SBRT. In order to make recommendations that will be applicable to other SBRT dose/fractionation, pooling the data of LungTech and RTOG 0813⁵⁶ (five-fraction regime) is furthermore an interesting future option. Adverse events are assessed according to Common Terminology Criteria for Adverse Events (CTCAE) v. 4 at baseline, at the end of treatment, 6 weeks after SBRT, 3 monthly for the first 3 years, 6 monthly for up to 5 years and yearly thereafter. Additionally, CT imaging is performed and pulmonary function tests recommended. *Serious Adverse Event* and *Suspected Unexpected Serious Adverse Reactions* reporting are conducted according to the Good Clinical Practice.⁸¹

Radiation therapy quality assurance and imaging quality control

Differences in delivery of care between institutions may induce variations affecting trial outcome. These variations may be multifactorial such as failure to adhere to protocol guidelines or differences in the equipment quality and/or its use. To prevent these biases, a high-technology quality assurance (QA) procedure has been developed. The RTQA

Figure 2. Local progression evaluation based on modified Huang *et al*⁵⁵ criteria. ¹⁸F-FDG, fluorine-18 fludeoxyglucose; PET, positron emission tomography. yr, year. ^a Repeat imaging should be performed no later than 3 months after the first abnormal CT. ^b Biopsy should be considered to confirm outcome if patient is fit enough and in particular if radical treatment remains an option should isolated local progression be confirmed.

procedure aims to ensure for each treatment plan an acceptable level of conformity to the protocol guidelines.⁸² The process includes the requirement to submit a benchmark case

for review prior to entry of the first patient in the trial (Figure 3). During the recruitment phase, all treatment plans are sent for a prospective central review by a team of experts,

Figure 3. A 54-year-old patient with centrally located non-small-cell lung cancer, T1N0M0, meeting the eligibility criteria for inclusion in the LungTech trial (benchmark case). (a) Delineation of tumour: gross tumour volume (GTV) (purple), internal target volume (red), planning target volume (pink), brachiocephalic vein (orange), superior trachea (pink), chest wall (green), vertebral body (blue), oesophagus (orange) and lungs-GTV (cyan). (b) Stereotactic body radiotherapy planning with dose prescription and adherence of organ at risk dose constraints according to LungTech protocol, isodoses: 64.2 Gy (red), 57.0 Gy (yellow), 48.0 Gy (green), 30.0 Gy (blue) and 15.0 Gy (cyan).

Figure 4. Design of the LungTech trial [European Organization for Research and Treatment of Cancer (EORTC) 22113-08113]. 3D, three dimensional; 4D-CT, four-dimensional CT; ^{18}F -FDG, Fluorine-18 Fludeoxyglucose; CBCT, cone beam CT; fr, fractions; NSCLC, non-small-cell lung cancer; ORTA, online randomized trials access; PET, position emission tomography; RT, radiotherapy; RTQA, radiation therapy quality assurance; SBRT, stereotactic body radiotherapy; yrs, years.

who will decide whether the patient is allowed to be treated based on the treatment plan provided (Figure 4).⁸³ Another aspect of RTQA is the verification of the correct use of RT techniques within the trial. End-to-end tests will be

performed during site visits using a specific breathing phantom (carrying films and an ionization chamber). Owing to the moving nature of the target, the RTQA procedure also includes an evaluation of the PET/CT, CT and 4D-CT

Figure 5. European organization for research and treatment of cancer (EORTC) infrastructure to support new generation clinical trials. ORTA (online randomized trials access), web-based application designed to facilitate the registration and randomization of patients in EORTC clinical trials; RTQA, radiation therapy quality assurance; VISTA, web-based application designed for the completion of most case report forms; PRISMA (protocol repository and integrated project management application), the EORTC clinical trial management system provides a global overview of all projects and also very detailed information about specific protocols.

techniques to assess the impact of potential motion artefacts on target volume accuracy. Furthermore, the LungTech QA evaluation will also include the assessment of consistency of Hounsfield units and SUV values, which is of particular relevance given the use of PET/CT data in this study.

Integration of technology advancement techniques in prospective trials

With technology advancement, the conduct of modern clinical research requires a major upgrade of clinical research organizations and the infrastructure for performing such sophisticated trials. Indeed, implementing this high-technology research brings new challenges to international multicentre clinical trials in terms of QA and standardization. For instance, imaging technologies such as PET scans have existed for decades and are widely available, their full use in clinical research, however, remains challenging. The quality and comparability of images collected within international multicentre clinical trials are not always optimal. Reliable clinical research involving imaging or radiotherapy can only be achieved within quality-assured, multicentre trials supported by robust methodology and operational infrastructures, allowing the processing, storage and analysis of imaging or treatment plan data by experienced researchers to be fully integrated with clinical and biological data (Figure 5).

Maintaining quality-assured clinical trials infrastructure is a cornerstone of independent clinical research, which will guarantee consistent, long-term and reliable research to patients. To have the necessary academic clinical research infrastructure available, skilled, experienced staff must be retained.

PERSPECTIVES/CONCLUSIONS

In summary, the LungTech trial is expected to provide high-quality, prospective, multicentre data on the efficacy and toxicity of moderately hypofractionated SBRT for central early stage NSCLC. The data generated by this trial will inform future recommendations on technique, prescription, risk estimation and response assessment for the routine use of this promising new radiotherapy technology. This is of particular importance given the likelihood of an increase in the proportion of patients with lung cancer diagnosed at an earlier stage through screening programs.

ACKNOWLEDGMENTS

The authors thank Sally Falk for language support.

FUNDING

This publication was supported by the EORTC Academic Fund, the Deutsche Krebshilfe e.V., the EORTC Charitable Trust and Cancer Research UK.

REFERENCES

- De Ruysscher D, Nakagawa K, Asamura H. Surgical and nonsurgical approaches to small-size non-small cell lung cancer. *Eur Respir J* 2014; **44**: 483–94. doi: [10.1183/09031936.00020214](https://doi.org/10.1183/09031936.00020214)
- Lagerwaard FJ, Versteegen NE, Haasbeek CJ, Slotman BJ, Paul MA, Smit EF, et al. Outcomes of stereotactic ablative radiotherapy in patients with potentially operable stage I non-small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2012; **83**: 348–53. doi: [10.1016/j.ijrobp.2011.06.2003](https://doi.org/10.1016/j.ijrobp.2011.06.2003)
- Guckenberger M. What is the current status of stereotactic body radiotherapy for stage I non-small cell lung cancer? *J Thorac Dis* 2011; **3**: 147–9. doi: [10.3978/j.issn.2072-1439.2011.06.04](https://doi.org/10.3978/j.issn.2072-1439.2011.06.04)
- Duncker-Rohr V, Nestle U, Momm F, Prokic V, Heinemann F, Mix M, et al. Stereotactic ablative radiotherapy for small lung tumors with a moderate dose. Favorable results and low toxicity. *Strahlenther Onkol* 2013; **189**: 33–40. doi: [10.1007/s00066-012-0224-y](https://doi.org/10.1007/s00066-012-0224-y)
- Timmerman R, Paulus R, Galvin J, Michalski J, Straube W, Bradley J, et al. Stereotactic body radiation therapy for inoperable early stage lung cancer. *JAMA* 2010; **303**: 1070–6. doi: [10.1001/jama.2010.261](https://doi.org/10.1001/jama.2010.261)
- Baumann P, Nyman J, Hoyer M, Wennberg B, Gagliardi G, Lax I, et al. Outcome in a prospective phase II trial of medically inoperable stage I non-small-cell lung cancer patients treated with stereotactic body radiotherapy. *J Clin Oncol* 2009; **27**: 3290–6. doi: [10.1200/JCO.2008.21.5681](https://doi.org/10.1200/JCO.2008.21.5681)
- Onishi H, Shirato H, Nagata Y, Hiraoka M, Fujino M, Gomi K, et al. Stereotactic body radiotherapy (SBRT) for operable stage I non-small-cell lung cancer: can SBRT be comparable to surgery? *Int J Radiat Oncol Biol Phys* 2011; **81**: 1352–8. doi: [10.1016/j.ijrobp.2009.07.1751](https://doi.org/10.1016/j.ijrobp.2009.07.1751)
- Ricardi U, Filippi AR, Guarneri A, Giglioli FR, Ciammella P, Franco P, et al. Stereotactic body radiation therapy for early stage non-small cell lung cancer: results of a prospective trial. *Lung Cancer* 2010; **68**: 72–7. doi: [10.1016/j.lungcan.2009.05.007](https://doi.org/10.1016/j.lungcan.2009.05.007)
- Senthi S, Haasbeek CJ, Slotman BJ, Senan S. Outcomes of stereotactic ablative radiotherapy for central lung tumours: a systematic review. *Radiother Oncol* 2013; **106**: 276–82. doi: [10.1016/j.radonc.2013.01.004](https://doi.org/10.1016/j.radonc.2013.01.004)
- Schanne DH, Nestle U, Allgauer M, Andrasschke N, Appold S, Dieckmann U, et al. Stereotactic body radiotherapy for centrally located stage I NSCLC: a multicenter analysis. *Strahlenther Onkol* 2015; **191**: 125–32. doi: [10.1007/s00066-014-0739-5](https://doi.org/10.1007/s00066-014-0739-5)
- D'Addario G, Felip E; ESMO Guidelines Working Group. Non-small-cell lung cancer: ESMO clinical recommendations for diagnosis, treatment and follow-up. *Ann Oncol* 2009; **20**(Suppl. 4): 68–70. doi: [10.1093/annonc/mdp132](https://doi.org/10.1093/annonc/mdp132)
- Gibbs IC, Loo BW Jr. CyberKnife stereotactic ablative radiotherapy for lung tumors. *Technol Cancer Res Treat* 2010; **9**: 589–96. doi: [10.1177/153303461000900607](https://doi.org/10.1177/153303461000900607)
- Scott WJ, Howington J, Feigenberg S, Movsas B, Pisters K; American College of Chest Physicians. Treatment of non-small cell lung cancer stage I and stage II: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest* 2007; **132**(Suppl. 3): 234S–42S.
- The National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: non-small cell lung cancer. Version 3, 2012. [Updated 15 August 2012.] Available from: <http://www.nccn.com>.
- Goeckenjan G, Sitter H, Thomas M, Branscheid D, Flentje M, Griesinger F, et al. Prevention, diagnosis, therapy, and follow-up of lung cancer interdisciplinary guideline of the German Respiratory Society and the German Cancer Society—abridged version. [In German.] *Pneumologie* 2011; **65**: e51–75. doi: [10.1055/s-0030-1256562](https://doi.org/10.1055/s-0030-1256562)
- Ginsberg RJ, Rubinstein LV. Randomized trial of lobectomy versus limited resection for T1 N0 non-small cell lung cancer. *Lung Cancer*

- Study Group. *Ann Thorac Surg* 1995; **60**: 615–22; discussion 22–3. doi: [10.1016/0003-4975\(95\)00537-U](https://doi.org/10.1016/0003-4975(95)00537-U)
17. Lausberg HF, Graeter TP, Tscholl D, Wendler O, Schäfers HJ. Bronchovascular versus bronchial sleeve resection for central lung tumors. *Ann Thorac Surg* 2005; **79**: 1147–52; discussion 1147–52. doi: [10.1016/j.athoracsur.2004.09.009](https://doi.org/10.1016/j.athoracsur.2004.09.009)
 18. Chen C, Bao F, Zheng H, Zhou YM, Bao MW, Xie HK, et al. Local extension at the hilum region is associated with worse long-term survival in stage I non-small cell lung cancers. *Ann Thorac Surg* 2012; **93**: 389–96. doi: [10.1016/j.athoracsur.2011.09.079](https://doi.org/10.1016/j.athoracsur.2011.09.079)
 19. Spigel L, Ferguson MK. Sleeve lobectomy versus pneumonectomy for lung cancer patients with good pulmonary function. In: Ferguson MK, ed. *Difficult decisions in thoracic surgery: an evidence-based approach*. 3rd edn. London, UK: Springer; 2007. pp. 103–9.
 20. Guckenberger M, Andratschke N, Alheit H, Holy R, Moustakis C, Nestle U, et al; Deutschen Gesellschaft für Radioonkologie (DEGRO). Definition of stereotactic body radiotherapy: principles and practice for the treatment of stage I non-small cell lung cancer. *Strahlenther Onkol* 2014; **190**: 26–33. doi: [10.1007/s00066-013-0450-y](https://doi.org/10.1007/s00066-013-0450-y)
 21. Haasbeek CJ, Palma D, Visser O, Lagerwaard FJ, Slotman B, Senan S. Early-stage lung cancer in elderly patients: a population-based study of changes in treatment patterns and survival in the Netherlands. *Ann Oncol* 2012; **23**: 2743–7. doi: [10.1093/annonc/mds081](https://doi.org/10.1093/annonc/mds081)
 22. National Lung Screening Trial Research Team; Church TR, Black WC, Aberle DR, Berg CD, Clingan KL, et al. Results of initial low-dose computed tomographic screening for lung cancer. *N Engl J Med* 2013; **368**: 1980–91. doi: [10.1056/NEJMoa1209120](https://doi.org/10.1056/NEJMoa1209120)
 23. Janssen-Heijnen ML, Smulders S, Lemmens VE, Smeenk FW, van Geffen HJ, Coebergh JW. Effect of comorbidity on the treatment and prognosis of elderly patients with non-small cell lung cancer. *Thorax* 2004; **59**: 602–7. doi: [10.1136/thx.2003.018044](https://doi.org/10.1136/thx.2003.018044)
 24. Asmis TR, Ding K, Seymour L, Shepherd FA, Leighl NB, Winton TL, et al; National Cancer Institute of Canada Clinical Trials Group. Age and comorbidity as independent prognostic factors in the treatment of non-small-cell lung cancer: a review of National Cancer Institute of Canada Clinical Trials Group trials. *J Clin Oncol* 2008; **26**: 54–9. doi: [10.1200/JCO.2007.12.8322](https://doi.org/10.1200/JCO.2007.12.8322)
 25. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011; **61**: 69–90. doi: [10.3322/caac.20107](https://doi.org/10.3322/caac.20107)
 26. Raz DJ, Zell JA, Ou SH, Gandara DR, Anton-Culver H, Jablons DM. Natural history of stage I non-small cell lung cancer: implications for early detection. *Chest* 2007; **132**: 193–9. doi: [10.1378/chest.06-3096](https://doi.org/10.1378/chest.06-3096)
 27. Sibley GS, Jamieson TA, Marks LB, Anscher MS, Prosnitz LR. Radiotherapy alone for medically inoperable stage I non-small-cell lung cancer: the Duke experience. *Int J Radiat Oncol Biol Phys* 1998; **40**: 149–54.
 28. Jeremic B, Classen J, Bamberg M. Radiotherapy alone in technically operable, medically inoperable, early stage (I/II) non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2002; **54**: 119–30. doi: [10.1016/s0360-3016\(02\)02917-6](https://doi.org/10.1016/s0360-3016(02)02917-6)
 29. Rowell NP, Williams CJ. Radical radiotherapy for stage I/II non-small cell lung cancer in patients not sufficiently fit for or declining surgery (medically inoperable). *Cochrane Database Syst Rev* 2001; CD002935. doi: [10.1002/14651858.CD002935](https://doi.org/10.1002/14651858.CD002935)
 30. Willner J, Baier K, Caragiani E, Tschammler A, Flentje M. Dose, volume, and tumor control prediction in primary radiotherapy of non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2002; **52**: 382–9. doi: [10.1016/s0360-3016\(01\)01823-5](https://doi.org/10.1016/s0360-3016(01)01823-5)
 31. Martel MK, Ten Haken RK, Hazuka MB, Kessler ML, Strawderman M, Turrisi AT, et al. Estimation of tumor control probability model parameters from 3-D dose distributions of non-small cell lung cancer patients. *Lung Cancer* 1999; **24**: 31–7. doi: [10.1016/s0169-5002\(99\)00019-7](https://doi.org/10.1016/s0169-5002(99)00019-7)
 32. Seppenwoolde Y, Lebesque JV, de Jaeger K, Belderbos JS, Boersma LJ, Schilstra C, et al. Comparing different NTCP models that predict the incidence of radiation pneumonitis. Normal tissue complication probability. *Int J Radiat Oncol Biol Phys* 2003; **55**: 724–35. doi: [10.1016/S0360-3016\(02\)03986-X](https://doi.org/10.1016/S0360-3016(02)03986-X)
 33. Bogart JA, Hodgson L, Seagren SL, Blackstock AW, Wang X, Lenox R, et al. Phase I study of accelerated conformal radiotherapy for stage I non-small-cell lung cancer in patients with pulmonary dysfunction: CALGB 39904. *J Clin Oncol* 2010; **28**: 202–6. doi: [10.1200/jco.2009.25.0753](https://doi.org/10.1200/jco.2009.25.0753)
 34. van Baardwijk A, Wanders S, Boersma L, Borger J, Ollers M, Dingemans AM, et al. Mature results of an individualized radiation dose prescription study based on normal tissue constraints in stages I to III non-small-cell lung cancer. *J Clin Oncol* 2010; **28**: 1380–6. doi: [10.1200/jco.2009.24.7221](https://doi.org/10.1200/jco.2009.24.7221)
 35. Nagata Y, Takayama K, Matsuo Y, Norihisa Y, Mizowaki T, Sakamoto T, et al. Clinical outcomes of a phase I/II study of 48 Gy of stereotactic body radiotherapy in 4 fractions for primary lung cancer using a stereotactic body frame. *Int J Radiat Oncol Biol Phys* 2005; **63**: 1427–31. doi: [10.1016/j.ijrobp.2005.05.034](https://doi.org/10.1016/j.ijrobp.2005.05.034)
 36. Bral S, Gevaert T, Linthout N, Versmissen H, Collen C, Engels B, et al. Prospective, risk-adapted strategy of stereotactic body radiotherapy for early stage non-small-cell lung cancer: results of a Phase II trial. *Int J Radiat Oncol Biol Phys* 2011; **80**: 1343–9. doi: [10.1016/j.ijrobp.2010.04.056](https://doi.org/10.1016/j.ijrobp.2010.04.056)
 37. Crabtree TD, Denlinger CE, Meyers BF, El Naqa I, Zoole J, Krupnick AS, et al. Stereotactic body radiation therapy versus surgical resection for stage I non-small cell lung cancer. *J Thorac Cardiovasc Surg* 2010; **140**: 377–86. doi: [10.1016/j.jtcvs.2009.12.054](https://doi.org/10.1016/j.jtcvs.2009.12.054)
 38. Grills IS, Mangona VS, Welsh R, Chmielewski G, McInerney E, Martin S, et al. Outcomes after stereotactic lung radiotherapy or wedge resection for stage I non-small-cell lung cancer. *J Clin Oncol* 2010; **28**: 928–35. doi: [10.1200/JCO.2009.25.0928](https://doi.org/10.1200/JCO.2009.25.0928)
 39. Haasbeek CJ, Lagerwaard FJ, Antonisse ME, Slotman BJ, Senan S. Stage I nonsmall cell lung cancer in patients aged > or =75 years: outcomes after stereotactic radiotherapy. *Cancer* 2010; **116**: 406–14. doi: [10.1002/cncr.24759](https://doi.org/10.1002/cncr.24759)
 40. Louie AV, Rodrigues G, Hannouf M, Lagerwaard F, Palma D, Zaric GS, et al. Withholding stereotactic radiotherapy in elderly patients with stage I non-small cell lung cancer and co-existing COPD is not justified: outcomes of a Markov model analysis. *Radiother Oncol* 2011; **99**: 161–5. doi: [10.1016/j.radonc.2011.04.005](https://doi.org/10.1016/j.radonc.2011.04.005)
 41. Shelkey J, Fakiris A, DeCamp MM, Medford-Davis LN, Flickinger JC, Recht A, et al. Matched-pair comparison of outcome of patients with clinical stage I non-small cell lung cancer treated with resection or stereotactic radiosurgery. *J Clin Oncol* 2012; **30**.
 42. Palma D, Visser O, Lagerwaard FJ, Belderbos J, Slotman BJ, Senan S. Impact of introducing stereotactic lung radiotherapy for elderly patients with stage I non-small-cell lung cancer: a population-based time-trend analysis. *J Clin Oncol* 2010; **28**: 5153–9. doi: [10.1200/JCO.2010.30.0731](https://doi.org/10.1200/JCO.2010.30.0731)
 43. Vansteenkiste J, De Ruysscher D, Eberhardt WE, Lim E, Senan S, Felip E, et al; ESMO Guidelines Working Group. Early and locally advanced non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013; **24**(Suppl. 6): vi89–98. doi: [10.1093/annonc/mdt241](https://doi.org/10.1093/annonc/mdt241)
 44. Versteegen NE, Lagerwaard FJ, Haasbeek CJ, Slotman BJ, Senan S. Outcomes of stereotactic ablative radiotherapy following a clinical diagnosis of stage I NSCLC: comparison

- with a contemporaneous cohort with pathologically proven disease. *Radiother Oncol* 2011; **101**: 250–4. doi: [10.1016/j.radonc.2011.09.017](https://doi.org/10.1016/j.radonc.2011.09.017)
45. Guckenberger M, Allgauer M, Appold S, Dieckmann K, Ernst I, Ganswindt U, et al. Safety and efficacy of stereotactic body radiotherapy for stage I non-small-cell lung cancer in routine clinical practice: a patterns-of-care and outcome analysis. *J Thorac Oncol* 2013; **8**: 1050–8. doi: [10.1097/JTO.0b013e318293dc45](https://doi.org/10.1097/JTO.0b013e318293dc45)
 46. Onishi H, Araki T, Shirato H, Nagata Y, Hiraoka M, Gomi K, et al. Stereotactic hypofractionated high-dose irradiation for stage I nonsmall cell lung carcinoma: clinical outcomes in 245 subjects in a Japanese multiinstitutional study. *Cancer* 2004; **101**: 1623–31. doi: [10.1002/cncr.20539](https://doi.org/10.1002/cncr.20539)
 47. Guckenberger M, Wulf J, Mueller G, Krieger T, Baier K, Gabor M, et al. Dose–response relationship for image-guided stereotactic body radiotherapy of pulmonary tumors: relevance of 4D dose calculation. *Int J Radiat Oncol Biol Phys* 2009; **74**: 47–54. doi: [10.1016/j.ijrobp.2008.06.1939](https://doi.org/10.1016/j.ijrobp.2008.06.1939)
 48. Onimaru R, Fujino M, Yamazaki K, Onodera Y, Taguchi H, Katoh N, et al. Steep dose–response relationship for stage I non-small-cell lung cancer using hypofractionated high-dose irradiation by real-time tumor-tracking radiotherapy. *Int J Radiat Oncol Biol Phys* 2008; **70**: 374–81. doi: [10.1016/j.ijrobp.2007.06.043](https://doi.org/10.1016/j.ijrobp.2007.06.043)
 49. Zhang J, Yang F, Li B, Li H, Liu J, Huang W, et al. Which is the optimal biologically effective dose of stereotactic body radiotherapy for Stage I non-small-cell lung cancer? A meta-analysis. *Int J Radiat Oncol Biol Phys* 2011; **81**: e305–16. doi: [10.1016/j.ijrobp.2011.04.034](https://doi.org/10.1016/j.ijrobp.2011.04.034)
 50. Timmerman R, McGarry R, Yiannoutsos C, Papiez L, Tudor K, DeLuca J, et al. Excessive toxicity when treating central tumors in a phase II study of stereotactic body radiation therapy for medically inoperable early stage lung cancer. *J Clin Oncol* 2006; **24**: 4833–9. doi: [10.1200/JCO.2006.07.5937](https://doi.org/10.1200/JCO.2006.07.5937)
 51. Song SY, Choi W, Shin SS, Lee SW, Ahn SD, Kim JH, et al. Fractionated stereotactic body radiation therapy for medically inoperable stage I lung cancer adjacent to central large bronchus. *Lung Cancer* 2009; **66**: 89–93. doi: [10.1016/j.lungcan.2008.12.016](https://doi.org/10.1016/j.lungcan.2008.12.016)
 52. Corradetti MN, Haas AR, Rengan R. Central-airway necrosis after stereotactic body-radiation therapy. *N Engl J Med* 2012; **366**: 2327–9. doi: [10.1056/NEJMc1203770](https://doi.org/10.1056/NEJMc1203770)
 53. Haasbeek CJ, Lagerwaard FJ, Slotman BJ, Senan S. Outcomes of stereotactic ablative radiotherapy for centrally located early stage lung cancer. *J Thorac Oncol* 2011; **6**: 2036–43. doi: [10.1097/jto.0b013e31822e71d8](https://doi.org/10.1097/jto.0b013e31822e71d8)
 54. Howington JA, Blum MG, Chang AC, Balekian AA, Murthy SC. Treatment of stage I and II non-small cell lung cancer: diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013; **143**(Suppl. 5): e278S–313S. doi: [10.1378/chest.12-2359](https://doi.org/10.1378/chest.12-2359)
 55. Huang K, Dahele M, Senan S, Guckenberger M, Rodrigues GB, Ward A, et al. Radiographic changes after lung stereotactic ablative radiotherapy (SABR)—can we distinguish recurrence from fibrosis? A systematic review of the literature. *Radiother Oncol* 2012; **102**: 335–42. doi: [10.1016/j.radonc.2011.12.018](https://doi.org/10.1016/j.radonc.2011.12.018)
 56. Radiation Therapy Oncology Group, RTOG0813. NCT00750269 [Updated 9 September 2009.] Available from: <http://rpc.mdanderson.org/rpc/credentialing/files/0813-Master-2-9-11.pdf>.
 57. Modh A, Rimmer A, Williams E, Foster A, Shah M, Shi W, et al. Local control and toxicity in a large cohort of central lung tumors treated with stereotactic body radiation therapy. *Int J Radiat Oncol Biol Phys* 2014; **90**: 1168–76. doi: [10.1016/j.ijrobp.2014.08.008](https://doi.org/10.1016/j.ijrobp.2014.08.008)
 58. Radiation Therapy Oncology Group, RTOG0236. A Phase II trial of stereotactic body radiation therapy (SBRT) in the treatment of patients with medically inoperable stage I/II non-small cell lung cancer. NCT000874. [Updated 9 September 2009.] Available from: <http://www.rtog.org/Clinical-Trials/ProtocolTable/StudyDetails.aspx?study0236>
 59. Fakiris AJ, McGarry RC, Yiannoutsos CT, Papiez L, Williams M, Henderson MA, et al. Stereotactic body radiation therapy for early stage non-small-cell lung carcinoma: four-year results of a prospective phase II study. *Int J Radiat Oncol Biol Phys* 2009; **75**: 677–82. doi: [10.1016/j.ijrobp.2008.11.042](https://doi.org/10.1016/j.ijrobp.2008.11.042)
 60. Li Q, Swanick CW, Allen PK, Gomez DR, Welsh JW, Liao Z, et al. Stereotactic ablative radiotherapy (SABR) using 70 Gy in 10 fractions for non-small cell lung cancer: exploration of clinical indications. *Radiother Oncol* 2014; **112**: 256–61. doi: [10.1016/j.radonc.2014.07.010](https://doi.org/10.1016/j.radonc.2014.07.010)
 61. Nishimura S, Takeda A, Sanuki N, Ishikura S, Oku Y, Aoki Y, et al. Toxicities of organs at risk in the mediastinal and hilar regions following stereotactic body radiotherapy for centrally located lung tumors. *J Thorac Oncol* 2014; **9**: 1370–6. doi: [10.1097/jto.0000000000000260](https://doi.org/10.1097/jto.0000000000000260)
 62. Milano MT, Chen Y, Katz AW, Philip A, Schell MC, Okunieff P. Central thoracic lesions treated with hypofractionated stereotactic body radiotherapy. *Radiother Oncol* 2009; **91**: 301–6. doi: [10.1016/j.radonc.2009.03.005](https://doi.org/10.1016/j.radonc.2009.03.005)
 63. Oshiro Y, Aruga T, Tsuboi K, Marino K, Hara R, Sanayama Y, et al. Stereotactic body radiotherapy for lung tumors at the pulmonary hilum. *Strahlenther Onkol* 2010; **186**: 274–9. doi: [10.1007/s00066-010-2072-y](https://doi.org/10.1007/s00066-010-2072-y)
 64. Unger K, Ju A, Oermann E, Suy S, Yu X, Vahdat S, et al. CyberKnife for hilar lung tumors: report of clinical response and toxicity. *J Hematol Oncol* 2010; **3**: 39. doi: [10.1186/1756-8722-3-39](https://doi.org/10.1186/1756-8722-3-39)
 65. Cannon DM, Mehta MP, Adkison JB, Khuntia D, Traynor AM, Tomé WA, et al. Dose-limiting toxicity after hypofractionated dose-escalated radiotherapy in non-small-cell lung cancer. *J Clin Oncol* 2013; **31**: 4343–8. doi: [10.1200/jco.2013.51.5353](https://doi.org/10.1200/jco.2013.51.5353)
 66. Timmerman R, Heinzerling J, Abdulrahman R, Choy H, Meyer JL. Stereotactic body radiation therapy for thoracic cancers: recommendations for patient selection, setup and therapy. *Front Radiat Ther Oncol* 2011; **43**: 395–411. doi: [10.1159/000322503](https://doi.org/10.1159/000322503)
 67. Nuytens JJ, van der Voort van Zyp NC, Praag J, Aluwini S, van Klaveren RJ, Verhoef C, et al. Outcome of four-dimensional stereotactic radiotherapy for centrally located lung tumors. *Radiother Oncol* 2012; **102**: 383–7. doi: [10.1016/j.radonc.2011.12.023](https://doi.org/10.1016/j.radonc.2011.12.023)
 68. Bonomo P, Livi L, Rampini A, Meattini I, Agresti B, Simontacchi G, et al. Stereotactic body radiotherapy for cardiac and paracardiac metastases: University of Florence experience. *La Radiologia medica* 2013; **118**: 1055–65. doi: [10.1007/s11547-013-0932-0](https://doi.org/10.1007/s11547-013-0932-0)
 69. Onimaru R, Shirato H, Shimizu S, Kitamura K, Xu B, Fukumoto S, et al. Tolerance of organs at risk in small-volume, hypofractionated, image-guided radiotherapy for primary and metastatic lung cancers. *Int J Radiat Oncol Biol Phys* 2003; **56**: 126–35. doi: [10.1016/s0360-3016\(03\)00095-6](https://doi.org/10.1016/s0360-3016(03)00095-6)
 70. Stephans KL, Djemil T, Diaconu C, Reddy CA, Xia P, Woody NM, et al. Esophageal dose tolerance to hypofractionated stereotactic body radiation therapy: risk factors for late toxicity. *Int J Radiat Oncol Biol Phys* 2014; **90**: 197–202. doi: [10.1016/j.ijrobp.2014.05.011](https://doi.org/10.1016/j.ijrobp.2014.05.011)
 71. Kirkpatrick JB, van der Kogel AJ, Schultheiss TE. Radiation dose-volume effects in the spinal cord. *Int J Radiat Oncol Biol Phys* 2010; **76**(3 Suppl.): S42–9. doi: [10.1016/j.ijrobp.2009.04.095](https://doi.org/10.1016/j.ijrobp.2009.04.095)
 72. Forquer JA, Fakiris AJ, Timmerman RD, Lo SS, Perkins SM, McGarry RC, et al. Brachial

- plexopathy from stereotactic body radiotherapy in early stage NSCLC: dose-limiting toxicity in apical tumor sites. *Radiother Oncol* 2009; **93**: 408–13. doi: [10.1016/j.radonc.2009.04.018](https://doi.org/10.1016/j.radonc.2009.04.018)
73. Borst GR, Ishikawa M, Nijkamp J, Hauptmann M, Shirato H, Onimaru R, et al. Radiation pneumonitis in patients treated for malignant pulmonary lesions with hypofractionated radiation therapy. *Radiother Oncol* 2009; **91**: 307–13. doi: [10.1016/j.radonc.2009.02.003](https://doi.org/10.1016/j.radonc.2009.02.003)
 74. Stanic S, Paulus R, Timmerman RD, Michalski JM, Barriger RB, Bezjak A, et al. No clinically significant changes in pulmonary function following stereotactic body radiation therapy for early-stage peripheral non-small cell lung cancer: an analysis of RTOG 0236. *Int J Radiat Oncol Biol Phys* 2014; **88**: 1092–9. doi: [10.1016/j.ijrobp.2013.12.050](https://doi.org/10.1016/j.ijrobp.2013.12.050)
 75. Taremi M, Hope A, Lindsay P, Dahele M, Fung S, Purdie TG, et al. Predictors of radiotherapy induced bone injury (RIBI) after stereotactic lung radiotherapy. *Radiat Oncol* 2012; **7**: 159. doi: [10.1186/1748-717X-7-159](https://doi.org/10.1186/1748-717X-7-159)
 76. Strigari L, Benassi M, Sarnelli A, Polico R, D'Andrea M. A modified hypoxia-based TCP model to investigate the clinical outcome of stereotactic hypofractionated regimes for early stage non-small-cell lung cancer (NSCLC). *Med Phys* 2012; **39**: 4502–14. doi: [10.1118/1.4730292](https://doi.org/10.1118/1.4730292)
 77. Guckenberger M, Klement RJ, Allgauer M, Appold S, Dieckmann K, Ernst I, et al. Applicability of the linear-quadratic formalism for modeling local tumor control probability in high dose per fraction stereotactic body radiotherapy for early stage non-small cell lung cancer. *Radiother Oncol* 2013; **109**: 13–20. doi: [10.1016/j.radonc.2013.09.005](https://doi.org/10.1016/j.radonc.2013.09.005)
 78. Brown JM, Carlson DJ, Brenner DJ. The tumor radiobiology of SRS and SBRT: are more than the 5 Rs involved? *Int J Radiat Oncol Biol Phys* 2014; **88**: 254–62. doi: [10.1016/j.ijrobp.2013.07.022](https://doi.org/10.1016/j.ijrobp.2013.07.022)
 79. Takeda A, Kunieda E, Takeda T, Tanaka M, Sanuki N, Fujii H, et al. Possible misinterpretation of demarcated solid patterns of radiation fibrosis on CT scans as tumor recurrence in patients receiving hypofractionated stereotactic radiotherapy for lung cancer. *Int J Radiat Oncol Biol Phys* 2008; **70**: 1057–65. doi: [10.1016/j.ijrobp.2007.07.2383](https://doi.org/10.1016/j.ijrobp.2007.07.2383)
 80. Vahdat S, Oermann EK, Collins SP, Yu X, Abedalthagafi M, Debrito P, et al. CyberKnife radiosurgery for inoperable stage IA non-small cell lung cancer: 18F-fluorodeoxyglucose positron emission tomography/computed tomography serial tumor response assessment. *J Hematol Oncol* 2010; **3**: 6. doi: [10.1186/1756-8722-3-6](https://doi.org/10.1186/1756-8722-3-6)
 81. Available from: www.ich.org/fileadmin/Public_Web_Site/ICH_Products/Guidelines/Safety/Safety_Guidelines.zip
 82. Melidis C, Bosch WR, Izewska J, Fidarova E, Zubizarreta E, Ulin K, et al. Global harmonization of quality assurance naming conventions in radiation therapy clinical trials. *Int J Radiat Oncol Biol Phys* 2014; **90**: 1242–9. doi: [10.1016/j.ijrobp.2014.08.348](https://doi.org/10.1016/j.ijrobp.2014.08.348)
 83. Fairchild A, Aird E, Fenton PA, Gregoire V, Gulyban A, Lacombe D, et al. EORTC Radiation Oncology Group quality assurance platform: establishment of a digital central review facility. *Radiother Oncol* 2012; **103**: 279–86. doi: [10.1016/j.radonc.2012.04.015](https://doi.org/10.1016/j.radonc.2012.04.015)