Published in final edited form as:

Coron Artery Dis. 2015 December ; 26(8): 678–685. doi:10.1097/MCA.0000000000000303.

CAC Score as Possible Criteria for Administration of Angiotensin Converting Inhibitors and/or Angiotensin Receptor Blockers: The Multi Ethnic Study of Atherosclerosis

Sirous Darabian¹, Yanting Luo¹, Arman Homat², Khashayar Khosraviani³, Nathan Wong⁴, Irfan Zeb⁵, Khurram Nasir⁶, and Matthew J. Budoff¹

¹Los Angeles Biomedical Research Institute

²Advanced Cardio-diagnostic clinic, ON, Canada

³Albert Einstein College of Medicine/Jacobi Medical Center, Bronx, NY

⁴University of California, Irvine College of Medicine

⁵Bronx-Lebanon Hospital, Bronx, NY

⁶Baptist Health South Florida, Miami, FL

Keywords

Angiotensin Converting Enzyme Inhibitor; Coronary artery calcium score

Introduction

The Renin-Angiotensin-Aldosterone System (RAAS) plays an important role in blood pressure (BP), cardiovascular disease and renal function. Angiotensin converting enzyme inhibitors (ACEIs) and Angiotensin receptor blockers (ARBs) are part of medications that were developed to attenuate the effect of The Renin-Angiotensin-Aldosterone System (RAAS). Currently, these medications are widely utilized for treatment of congestive heart failure (CHF), myocardial infarction, hypertension, and prevention of diabetes's macro and micro-vascular complications. Also, they have been shown to slow progression of chronic kidney disease (CKD) to end stage renal disease. In 2009, ACE inhibitors were the fourth most utilized drug class in the United States, and they were prescribed 162.8 million times by US physicians.¹

Both ACEIs and ARBs are medications that have proven effects to lower mortality and morbidity rates². Among patients with congestive heart failure, the SOLVD trial showed that enalapril significantly improved survival and reduced further hospitalization rates.² Moreover, in the Survival And Ventricular Enlargement (SAVE) trial, captopril was able to decrease risk of recurrent myocardial infarction.³ In the Heart Outcomes Prevention Evaluation (HOPE) study, among high risk patients, ramipril lowered the risk of

atherosclerotic events including all stroke, myocardial infarction and cardiovascular death. Benefits were independent of ramipril's effect on blood pressure and cardiac systolic function.⁴ However, in the Captopril Prevention Project (CAPPP) study of 10,985 patients, there was no difference in cardiovascular morbidity and mortality rates. In this study, Captopril failed to show any advantages over conventional therapy in preventing cardiovascular morbidity and mortality.⁵ The discrepancies regarding the effect of these medications on the rate of cardiovascular events across different trials suggest that we need to find better criteria to predict who may benefit from these medications. The evaluation of Framingham Risk Score (FRS) has been a poor predictor of whom will benefit from ACE inhibition. In this study, we sought to examine if the Coronary Artery Calcium (CAC) score can determine which individuals would benefit from ACEI or ARB effects (in the form of fewer cardiovascular events). There are no guidelines for instituting ACEI or ARB therapies based on CAC score or Framingham risk. The goal of this study is to identify a subgroup of primary prevention patients who may benefit from ACE inhibition for reducing CV risk, not reducing complications of DM (albuminuria) or hypertension (left ventricular hypertrophy).

Methodology

Population characteristics

The Multi Ethnic Study of Atherosclerosis (MESA) is a prospective study intended to evaluate development and progression of subclinical cardiovascular disease in clinically asymptomatic individuals among different ethnic groups. In this study, participants were enrolled from four different ethnicities (white, African American, Hispanic, and Asian, predominantly of Chinese descent) at 6 Field Centers (Forsyth County, NC; Northern Manhattan and the Bronx, NY; Baltimore City and Baltimore County, MD; St. Paul, MN; Chicago and the village of Maywood, IL; and Los Angeles County, CA.). All participants underwent extensive evaluation at baseline, including clinical history, physical examination, and many laboratory tests. The MESA protocol, including information about the details of populations and recruitment method, details of inclusion and exclusion criteria, investigators' contact information, and other detailed information, is available on the World Wide Web at www.mesa-nhlbi.org. Out of 6,814 the MESA population, we enrolled all 2,906 participants who never used ACEIs or ARBs (n=2,457, 84.5 %) or were taking any of these medications (n=449, 15.5%) during the baseline and all follow up years. Participants that were taking these medications intermittently were excluded. They were followed for an average of 8.0 ± 1.7 years (range 0.02 to 10.9 years). Our selected population, with 49.0% male and the average age of 60.1 ± 9.7 years (ranged 45–84) had no apparent clinical cardiovascular disease (CVD) at the baseline. (See table-1) All participants underwent a non-contrast enhanced cardiac computerized tomography (Cardiac CT) and evaluated for CAC score. According to the participants CAC score and self-reported utilization of ACEIs and ARBs, they were categorized into six different groups; Zero CAC with ACEI/ARBs (n=163), Zero CAC without ACEI/ ARBs (n=1,440), intermediate CAC scores (1 to 399) with ACEI/ARBs (n=203), intermediate CAC scores (1 to 399) without ACEI/ARBs (n=865), higher CAC scores (≥ 400) with ACEI/ARBs (n=83), and high CAC scores without ACEI/ARBs (n=152). Moreover, to examine superiority of the CAC score over the Framingham risk score (FRS), we also divided participants into the six groups of three FRS

Groups (low, intermediate and high) and with or without ACE/ARB use. Low FRS with ACEI/ARBs constituted 92 participants, Low FRS without ACEI/ARBs (n=1,531), intermediate FRS with ACEI/ARBs (n=126), Intermediate FRS without ACEI/ARBs (n=575), high FRS with ACEI/ARBs (n=231), and high FRS without ACEI/ARBs (n=351). Out of all participants, 55.2% (n=1,603) had zero CAC score, and 8.1% (n=235) had CAC score of more than 400 (See Table-2 and 3)

Data Collection

All participants signed informed consent and were evaluated at baseline examination between July 2000 and July 2002. Also they were evaluated at four follow up examinations as follows: a second examination between July 2002 and January 2004, a third examination between January 2004 and July 2005, a forth examination between July 2005 and July 2007 and a fifth examination between April 2010 and April 2012. The components of each examination are described in table-3. Participant's demographic data, medical condition, family history of CVD, and current use of prescription and non-prescription medication were collected by questionnaires. During the examination, height, weight, and waist and hip circumferences are measured. Resting blood pressure was measured three times in the seated position using a Dinamap model Pro 100 automated oscillometric sphygmomanometer (Critikon, Tampa, Florida).

Coronary artery calcification score

Cardiac CT was performed by utilizing either a cardiac-gated electron-beam CT scanner (the Chicago, Los Angeles, and New York field centers) (Imatron C-150; Imatron, San Francisco, California) or a prospectively electrocardiogram-triggered scan acquisition at 50% of the R-R interval with a multi-detector CT, acquiring a block of four 2.5-mm slices for each cardiac cycle in a sequential or axial scan mode (the Baltimore, Forsyth County, and St. Paul field centers) (Lightspeed, General Electric Medical Systems, Waukesha, Wisconsin; or Volume Zoom, Siemens, Erlanger, Germany). All participants were scanned over phantoms with known physical density. CAC scores were evaluated centrally at the Los Angeles Biomedical Research Center at Harbor-UCLA in Torrance, California.

Cardiovascular events

Events were defined as all CVD events including myocardial infarction, resuscitated cardiac arrest, definite and probable angina (only if followed by revascularization), stroke, stroke death, coronary heart disease death, other atherosclerotic death as well as all other cardiovascular deaths. Silent myocardial infarctions identified using criteria applied to the follow-up electrocardiogram. Resuscitated cardiac arrest and other CVD endpoints were identified from medical records. In order to classify CVD events occurring during follow-up time, information have been collected from death certificates, hospital medical records, autopsy reports, interviews with participants, and for those who died out of the hospital, interviews with or questionnaires were administered to physicians, relatives, or friends.

Statistical analysis methods

To address the objective of the study, a number of statistical analyses were conducted. Subjects were classified into groups of using and not using ACEIs and/or ARBs. To assess risk factors, the chi-square test (for categorical variables) and 1-way analysis of variance (for continuous variables) were performed. Continuous variables were presented as mean $\pm$ SD, categorical variables were presented as frequency/percentage. Mann-Whitney U test (Wilcoxon signed-rank test) was performed to compare the Framingham risk score distribution between using and not using ACEIs and/or ARBs groups. Framingham risk score values were presented by median (interquartile range (IQR)). The chi-square test of proportion was also performed to examine the relationships between the distribution of CAC score (low 0, intermediate 0–400, high >400), FRS (low 0–10, intermediate 10–20, high >20), and all cardiovascular events. Subjects were then cross-classified into six groups according to their CAC score levels (low, intermediate, high)/FRS levels (low, intermediate, high) and ACEIs/ARBs use. We assessed the cardiovascular events rate per 1,000 person-year of both ACEIs/ARBs user and non-user groups, stratified by CAC score or FRS levels. In addition to the univariate Cox regression analysis, unadjusted and adjusted multivariate Cox proportional hazard model were used to calculate hazard ratios (HR) in the relationship among ACEI/ARBs usage stratified by CAC score/FRS levels, and all cardiovascular events. We considered the user group as the references for these analyses. Further adjustments were performed in two steps. Covariates were entered into the model based on their potential roles as confounders. The first step was performed with demographics, smoking, diabetes, hypertension, hyperlipidemia, and family history of premature CVD; in the second step, aspirin and statin usage were added to the model. Kaplan-Meier plots were presented for all cardiovascular events over ACEIs and/or ARBs and CAC score/FRS groups. All statistical analysis were performed using SAS software, version 9.3 (SAS Institute Inc., Cary, NC), and statistical significance levels of all the analysis were set at $p=0.05$ (2-sided).

Results

In total, 2,906 participants (52.6% female) with the mean age of 60.1 (ranged 45–84) were followed up for an average of 8.0 ± 1.7 years (ranged 0.02–10.9 years). Of these participants, 449 (49.0% female, mean age of 64.4 years) had taken ACEIs and/or ARBs through the all follow-up years and non-users had never utilized these medications (mean age of 59.3, and 53.2% female). In average, 96.7% of participants in the ACEI/ARB group had hypertension, whereas, only 20.4% of non-ACE/ARB participants had hypertension. Similarly, diabetes was significantly more common among ACE/ARB group (32.1%) while only 3.4% of non-user participants reported having diabetes. (See Table-1)

In total, 248 (8.5%) CVD events were recorded. Event rates were greater among the groups with greater levels of CAC and/or FRS (see Figure-1, table-4 and table -5). Moreover, the ACEIs and/or ARBs user population had significantly greater rate of CVD events than the non-user group (17.1% and 7.0%, in order) (See Table 2).

Among the patients with normal CAC scores, unadjusted Cox regression model analysis showed that compared to the user group, the non-users had significantly lower risk of events

(HR=0.37, 95% CI: 0.18–0.78, $P<0.05$). However, when they were adjusted by major cardiovascular disease risk factors (adjusted model 1) or major cardiovascular disease risk factors plus statin and aspirin use (adjusted model 2), no significant difference persisted.

In contrast, among the patients with intermediate CAC score, both of adjusted models showed a significantly greater event rate among those who did not used ACEi/ARBs. In this group, the non-ACEi/ARBs users showed a significantly higher hazard ratio for all CVD events (HR = 1.6, 95%CI: 1.0–2.6, $P<0.05$ in model one and HR=3.3, 95% CI:1.1- 10.4, $P<0.05$ in model two). (See figures 2, 3 and 4)

Among the patients with a high CAC score, in spite of significant differences in unadjusted models, adjusted models failed to show any significant difference over the use of ACEi/ARBs. (See table-6)

In contrast to the CAC score, no significant difference was observed between groups when they were classified over FRS levels. (Table-7)

Discussion

RAAS inhibitors reduce pro-thrombotic activity, modify anti-inflammatory mediators, and reduce plaque rupture. In addition, ACEIs have been shown to improve coronary artery endothelial function. There are many studies demonstrating a significant improvement in both survival and quality of life with ACEIs and/or ARBs among patients with heart failure.⁶ However, appropriate selection of patients for treatment with ACEIs and/or ARBs is challenging. Currently many patients without heart failure are using these medications. Despite data that indicate ACEIs may decrease event rates in patients with hypertension, it is not clear which patients will benefit. Furthermore, some trials were not able to show any overall benefits.⁷ The PEACE trial, which enrolled 8290 participants with normal or near normal left ventricular ejection fraction, reported that cardiovascular event rates did not differ when they were adjusted for major cardiovascular risk factors.

In our study, patients who over the all follow up years received ACEIs and/or ARBs had greater average of both FRS and CAC scores. Therefore, generally, patients who took ACEIs and/or ARBs had greater risk and event rates, most likely due to the underlying indications for ACE and ARB use (See figures 1, 2 and 3). Similarly, when we divided patients based on their FRS, the non ACE/ARB group had lower events than ACEi or ARBs users, but this effect was attenuated by the presence of diabetes and other non FRS risk factors.

When we classified our population based on their CAC scores, even after adjustment with risk factors or risk factors plus statin and aspirin use, patients with intermediate CAC scores had fewer events if they were on ACEIs or ARBs. In contrast, in the groups with zero CAC score or CAC scores higher than 400, our results showed there was no significant difference between the participants taking ACE/ARB and those not. It has been demonstrated that patients with zero CAC have very low CV event rates. Therefore, it is difficult to show benefit from further risk reduction. Regarding CAC >400, it is possible that these participants, while at high cardiovascular risk, have more established atherosclerosis that

may be less amenable to different therapies.^{8,9,10} Some clinical and biomechanical studies revealed that plaque composition has an important role in acute clinical events. It is possible that more calcified lesions are less vulnerable to rupture and spotty calcification and mixed plaques are more likely to be associated with future CV events.^{11,12,13} Interestingly, in contrast to the CAC score, our study shows that Framingham risk score did not help in stratification of whom would benefit from ACE inhibitors or ARBs.. No FRS levels were associated with lower cardiovascular event rates between ACEI/ARB users and the non-users. (See table-7)

Limitations

We excluded participants with temporary use of ACEIs and/or ARBs, as the follow up reflected both time on and time off these medications. We considered all participants who were consistently taking ACEI/ARBs from the baseline visit and continued all through the years of follow up. Of course, a randomized trial in patients with intermediate CAC score would be most definitive, but this is unlikely to be performed.

Conclusion

Participants with intermediate CAC scores (defined as scores 1–399), who were on ACEIs and/or ARBs, experienced less future cardiovascular events than those not taking these therapies. Also, Framingham risk score could not define patients who may or may not benefit from ACE/ARBs in this observational study. These CAC results can be obtained on a dedicated heart scan or thoracic imaging for lung cancer screening.¹⁴ While a randomized prospective trial would be confirmatory, this study adds critical information that the cardioprotective effects of agents may be dependent on baseline CV risk, and the prognostic information received with CAC scores may help determine optimal medical regimens.

Acknowledgement

This research was supported by R01-HL071739 contracts N01-HC-95159 through N01-HC-95169 from the National Heart, Lung, and Blood Institute and by grants UL1-RR-024156 and UL1-RR-025005 from NCRR. The authors thank the other investigators, the staff, and the participants of the MESA study for their valuable contributions. A full list of participating MESA investigators and institutions can be found at <http://www.mesa-nhlbi.org>.

REFERENCES

1. Bian MS, Boyang; Kelton PhD, Christina ML.; Guo PhD, Jeff J.; Wigle PharmD, Patricia R. ACE Inhibitor and ARB Utilization and Expenditures in the Medicaid Fee-For-Service Program from 1991 to 2008. *J Manag Care Pharm*. 2010; 16(9):671–679. [PubMed: 21067253]
2. The SOLVD investigators. Effect of Enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med*. 1991; 325:293–302. [PubMed: 2057034]
3. Pfeffer MA, Braunwald E, Moye LA, et al. on behalf of the SAVE investigators. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction: results of the Survival And Ventricular Enlargement (SAVE) trial. *N Engl J Med*. 1992; 327:669–677. [PubMed: 1386652]
4. The heart outcomes prevention evaluation study investigators. Effects of an Angiotensin Converting Enzyme Inhibitor, Ramipril, on cardiovascular events in high risk patients. *N Engl J Med*. 2000; 342:145–153. [PubMed: 10639539]

5. Hansson, Lennart; Lindholm, Lars H.; Niskanen, Leo; Lanke, Jan; Hedner, Thomas; Niklason, Anders; Luomanmäki, Kimmo; Dahlöf, Björn; de Faire, Ulf; Mörlin, Claes; Karlberg, Bengt E.; Wester, PO.; Björck, Jan-Erik. for the Captopril Prevention Project (CAPPP) study group. Effect of angiotensin-converting-enzyme inhibition compared with conventional therapy on cardiovascular morbidity and mortality in hypertension: the Captopril Prevention Project (CAPPP) randomized trial. *Lancet*. 1999; 353:611–616. [PubMed: 10030325]
6. Donnelly R, Manning G. Angiotensin-converting enzyme inhibitors and coronary heart disease prevention. *JRAAS*. 2007; 8:13–22. [PubMed: 17487822]
7. van Vark LC, Bertrand M, Akkerhuis KM, Brugts JJ, Fox K, Mourad JJ, Boersma E. Angiotensin-converting enzyme inhibitors reduce mortality in hypertension: a meta-analysis of randomized clinical trials of renin-angiotensin-aldosterone system inhibitors involving 158,998 patients. *Eur Heart J*. 2012 Aug; 33(16):2088–2097. [PubMed: 22511654]
8. Budoff MJ, McClelland RL, Nasir K, Greenland P, Kronmal RA, Kondos GT, Shea S, Lima JA, Blumenthal RS. Cardiovascular events with absent or minimal coronary calcification: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am Heart J*. 2009 Oct; 158(4):554–561. [PubMed: 19781414]
9. Blaha MJ, Blumenthal RS, Budoff MJ, Nasir K. Understanding the utility of zero coronary calcium as a prognostic test: a bayesian approach. *Circ Cardiovasc Qual Outcomes*. 2011; 4(2):253–256. [PubMed: 21406674]
10. Shareghi S, Ahmadi N, Young E, Gopal A, Liu ST, Budoff MJ. Prognostic significance of zero coronary calcium scores on cardiac computed tomography. *J Cardiovasc Comput Tomogr*. 2007; 1:155–159. [PubMed: 19083900]
11. Motoyama S, Kondo T, Sarai M, et al. Multislice computed tomographic characteristics of coronary lesions in acute coronary syndromes. *J Am Coll Cardiol*. 2007; 50:319–326. [PubMed: 17659199]
12. Pundziute G, Schuijf JD, Jukema JW, et al. Evaluation of plaque characteristics in acute coronary syndromes: non-invasive assessment with multi-slice computed tomography and invasive evaluation with intravascular ultrasound radiofrequency data analysis. *Eur Heart J*. 2008; 29:2373–2381. [PubMed: 18682447]
13. Kitagawa T, Yamamoto H, Horiguchi J, et al. Characterization of Noncalcified Coronary Plaques and Identification of Culprit Lesions in Patients With Acute Coronary Syndrome by 64-Slice Computed Tomography. *J Am Coll Cardiol.: CARDIOVASCULAR IMAGING*. 2009; 2(2)
14. Budoff MJ, Nasir K, Kinney GL, Hokanson JE, Barr RG, Steiner R, Nath H, Lopez-Garcia C, Black-Shinn J, Casaburi R. Coronary artery and thoracic calcium on noncontrast thoracic CT scans: Comparison of ungated and gated examinations in patients from the COPD Gene cohort. *J Cardiovasc Comput Tomogr*. 2011; 5(2):113–118. [PubMed: 21167806]

Figure 1.
Event Rates over the CAC Scores and Usage of ACEi/ARBs

Figure 2.
Distribution of all Cardiovascular Events Among Framingham Risk Scores and Presence of Coronary Artery Calcification

Figure 3.
Kaplan-Meier Plot for all Cardiovascular Events Over ACEi/ARBs and Coronary Artery Calcium Score

Figure 4.
Kaplan-Meier Plot for all Cardiovascular Events Over ACEi/ARBs Use and Framingham Risk Score

Table 1

Components of the Multi-Ethnic Study of Atherosclerosis by Examinations

	Exam 1 Jul00– Aug02 (24 months)	Exam 2 Sep02– Feb04 (18 Months)	Exam 3 Mar04– Sep05 (18 Months)	Exam 4 Sep05– May07 (21 Months)	Exam 5 Apr10– Dec11 (21 Months)
	N=6814	N=6232	N=5939	N=5704	N=4651
Questionnaires					
Personal Hx	X	X	X	X	X
Medical Hx	X	X	X	X	X
Medications	X	X	X	X	X
Family Hx		X			
Procedures/Assessments					
Anthropometry	X	X	X	X	X
Urine collection	X	X	X		X
Blood Pressure	X	X	X	X	X
Electrocardiogram (ECG)	X ₆₇₆₅				X ₄₆₂₅
CT Coronary (Chest)	X ₆₈₁₄	X ₂₉₅₅	X ₂₈₀₅	A ₁₄₀₅	A ₃₂₀₀

X = MESA Classic Cohort (all available assumed unless number provided)
A = Ancillary Study procedure

Counts provided above are reflective data available in datasets. Some variation in the number available for a specific analysis is expected.

Table-2

Distribution of Major Cardiovascular Risk Factors

Factors	Total (n=2,906, 100%)	ACEIs and/or ARBs (n=449, 15.5%)	NO ACEIs and/or ARBs (n=2,457, 84.5%)	p Value
Age, year	60.1±9.7	64.4±9.0	59.3±9.7	<.0001
Female, n (%)	1527(52.6)	220(49.0)	1307(53.2)	0.1015
Smoking, n (%)				0.5186
Non Smoker	1496 (51.2)	231(51.5)	1265(51.6)	
Former Smoker	1053(36.3)	170(37.9)	883(36.0)	
Current Smoker	353(12.2)	48(10.7)	305(12.4)	
Hypertension, n (%)	935(32.2)	434(96.7)	501(20.4)	<.0001
Diabetes Mellitus, n (%)	227(7.8)	144(32.1)	83(3.4)	<.0001
Hyperlipidemia, n (%)	1680(58.0)	240(53.8)	1440(58.7)	0.0542
Family History of Premature	777(26.7)	151(33.6)	626(25.5)	0.0003

Table-3

Distribution of Framingham Risk Score and Coronary Artery Calcification Score

Factors	Total (n=2,906, 100%)	ACEI/ARB (n=449, 15.5%)	NO ACEI/ARBs (n=2,457, 84.5%)	p Value
Framingham Risk Score [£]	9.4 (4.5, 18.4)	21.5 (11.7, 30.0)	7.9 (4.5, 13.7)	<.0001
FRS categories, n (%) [£]				<.0001
Low FRS	1623(55.8)	92(20.5)	1531(62.3)	
Moderate FRS	701(24.1)	126(28.1)	575(23.4)	
High FRS	582(20.0)	231(51.4)	351(14.3)	
CACs categories, n (%) [£]				<.0001
CACs = 0	1603(55.2)	163(36.3)	1440(58.6)	
0 < CACs 400	1068(36.8)	203(45.2)	865(35.2)	
CACs > 400	235(8.1)	83(18.5)	152(6.2)	
CVD, all, n (%) [£]	248(8.5)	77(17.1)	171(7.0)	<.0001

[£] p<0.0001

[£] Presented by median (interquartile range)

FRS = Framingham Risk Score; CACs = Coronary Artery Calcification score; CVD = Cardiovascular Disease

Table-4

Distribution of Coronary Artery Calcification by Framingham Risk Score

	Low FRS (<10%)	Moderate FRS (10%–20%)	High FRS (>20%)	p Value
CACs = 0	1179(72.6)	289(41.2)	135(23.2)	<.0001
0 < CACs 400	416(25.6)	340(48.5)	312(53.6)	
CACs > 400	28(1.7)	72(10.3)	135(23.2)	
Total	1623 (100.0)	701 (100.0)	582 (100.0)	

FRS = Framingham Risk Score; CACs = Coronary Artery Calcification score

Table-5

Prevalence of All Cardiovascular Events across Different Levels of Framingham Risk Score and CAC scores

	Low FRS (<10%)	Moderate FRS (10%-20%)	High FRS (>20%)	p Value
CVD, all	38(2.3)	67(9.6)	143(24.6)	<.0001
Total	1623 (100.0)	701 (100.0)	582 (100.0)	
	CACs = 0	0 < CACs < 400	CACs > 400	p Value
CVD, all	39(2.4)	129(12.1)	80(34.0)	<.0001
Total	1603 (100.0)	1068 (100.0)	235 (100.0)	

FRS = Framingham Risk Score; CACs = Coronary Artery Calcification score; CVD = Cardiovascular Disease

Table-6

Event Rate per 1000 person years and Hazard Ratios (HR) for ACEIs/ARBs and Coronary Artery Calcification Score Groups in Adjusted and Unadjusted Cox Proportional Hazard Regression Analysis for All Cardiovascular Events

	Event Rate Per 1000 Person Years	HR (95% CI)		
		Unadjusted Model (N=2,906)	Adjusted Model 1 (N=2,886)	Adjusted Model 2 (N=1,367)
ACEI/ARBs and CACs = 0	6.9[9/163]	Reference	Reference	Reference
ACEIs and/or ARBs and 0 < CACs < 400	19.8[30/203]	2.84(1.35,5.99) **	2.12(1.00,4.49) *	0.67(0.17,2.74)
ACEI/ARBs, CACs>400	77.8[38/83]	10.90(5.27,22.55) ***	5.52(2.62,11.66) ***	3.51(0.97,12.76)
No ACEI/ARBs and CACs = 0	2.5[30/1440]	0.37(0.18,0.78) **	1.03(0.47,2.27)	0.49(0.12,2.07)
No ACEI/ARBs and 0 < CACs < 400	14.5[99/865]	2.11(1.07,4.18) *	3.48(1.70,7.13) ***	2.23(0.69,7.19)
No ACEI/ARBs and CACs >= 400	39.6[42/152]	5.64(2.74,11.59) ***	6.41(3.01,13.68) ***	6.97(2.03,23.93) **
ACEI/ARBs and CACs = 0	6.9[9/163]	0.35(0.17,0.74) **	0.47(0.22,0.99) *	1.48(0.37,6.02)
ACEIs/ARBs and 0 < CACs < 400	19.8[30/203]	Reference	Reference	Reference
ACEI/ARBs and CACs >= 400	77.8[38/83]	3.83(2.37,6.19) ***	2.608(1.60,4.26) ***	5.21(1.46,18.65) *
No ACEI/ARBs and CACs = 0	2.5[30/1440]	0.13(0.08,0.22) ***	0.488(0.27,0.87) *	0.72(0.17,3.09)
No ACEIs/ARBs and 0<CACs<400	14.5[99/865]	0.74(0.49,1.12)	1.64(1.05,2.59) *	3.30 (1.05,10.38) *
No ACEI/ARBs and CACs >= 400	39.6[42/152]	1.98(1.24,3.17) **	3.03(1.83,5.02) ***	10.34(3.16,33.76) ***
ACEI/ARBs and CACs = 0	6.9[9/163]	0.09(0.04,0.19) ***	0.18(0.09,0.38) ***	0.28(0.08,1.03)
ACEI/ARB and 0 < CACs < 400	19.8[30/203]	0.26(0.16,0.42) ***	0.38(0.24,0.63) ***	0.19(0.05,0.69) *
ACEI/ARBs and CACs >= 400	77.8[38/83]	Reference	Reference	Reference
No ACEI/ARB and CACs = 0	2.5[30/1440]	0.03(0.02,0.05) ***	0.19(0.10,0.34) ***	0.14(0.03,0.56) **
No ACEI/ARBs and 0<CAC< 400	14.5[99/865]	0.19(0.13,0.28) ***	0.63(0.40,0.99) *	0.63(0.22,1.83)
No ACEI/ARBs and CACs >= 400	39.6[42/152]	0.52(0.33,0.80) **	1.16 (0.71,1.91)	1.98(0.65,6.02)

*
p<0.05,

**
p<0.01,

p<0.001 compared to reference group;

CACs = Coronary Artery Calcification score; CVD = Cardiovascular Disease

Adjusted model 1 adjusted for age, gender, ethnicity, cigarette smoking, hypertension, diabetes, hyperlipidemia, and family history of premature CVD.

Adjusted model 2; adjusted for age, gender, ethnicity, cigarette smoking, hypertension, diabetes, hyperlipidemia, and family history of premature CVD, and utilizing statin and aspirin

Table-7

Event Rate per 1000 person years and Hazard Ratios (HR) for ACEIs/ARBs and Framingham Risk Score Groups in Adjusted and Unadjusted Cox Proportional Hazard Regression Analysis for All Cardiovascular Events

	Event Rate Per 1000 Person Years	HR (95% CI)		
		Unadjusted Model (N=2,906)	Adjusted Model 1 (N=2,886)	Adjusted Model 2 (N=1,367)
ACEI/ARBs and Low FRS	5.3[4/92]	Reference	Reference	Reference
ACEI/ARBs and Moderate FRS	15.6[15/126]	2.90(0.96,8.73)	1.77(0.58,5.41)	3.08(0.36,26.53)
ACEI/ARBs and High FRS	36.2[58/231]	6.68(2.42,18.39)***	2.10(0.71,6.25)	2.06(0.22,19.58)
No ACEI/ARBs and Low FRS	2.7[34/1531]	0.51(0.18,1.42)	1.00 (0.34,2.90)	1.39(0.16,12.02)
No ACEI/ARBs and Mod FRS	11.3[52/575]	2.13(0.77,5.88)	1.87(0.65,5.37)	4.91(0.61,39.52)
No ACEI/ARBs and High FRS	33.9[85/351]	6.26(2.30,17.07)***	2.98(1.02,8.75)*	5.04(0.58,43.83)
ACEI/ARBs and Low FRS	5.3[4/92]	0.34(0.12,1.04)	0.56(0.19,1.72)	0.33(0.04,2.80)
ACEI/ARBs and Moderate FRS	15.6[15/126]	Reference	Reference	Reference
ACEI/ARBs and High FRS	36.2[58/231]	2.3(1.31,4.07)**	1.18(0.64,2.19)	0.67(0.21,2.18)
No ACEI/ARBs and Low FRS	2.7[34/1531]	0.17(0.10,0.32)***	0.56(0.28,1.12)	0.45(0.12,1.65)
No ACEI/ARBs and Mod FRS	11.3[52/575]	0.73(0.41,1.30)	1.06(0.57,1.95)	1.60(0.56,4.53)
No ACEI/ARBs and High FRS	33.9[85/351]	2.16(1.25,3.74)**	1.68(0.92,3.07)	1.64(0.57,4.75)
ACEI/ARBs and Low FRS	5.3[4/92]	0.15(0.05,0.41)***	0.48(0.16,1.42)	0.49(0.05,4.61)
ACEI/ARBs and Moderate FRS	15.6[15/126]	0.43(0.25,0.77)**	0.84(0.46,1.56)	1.49(0.46,4.87)
ACEI/ARBs and High FRS	36.2[58/231]	Reference	Reference	Reference
No ACEI/ARBs and Low FRS	2.7[34/1531]	0.08(0.05,0.12)***	0.47(0.24,0.92)*	0.67(0.15,3.09)
No ACEI/ARBs and Mod FRS	11.3[52/575]	0.32(0.22,0.46)***	0.89(0.54,1.47)	2.39(0.77,7.37)
No ACEI/ARBs and High FRS	33.9[85/351]	0.94(0.67,1.31)	1.42(0.97,2.10)	2.45(0.95,6.30)

* p<0.05,

** p<0.01,

*** p<0.001 compared to reference group;

FRS = Framingham Risk Score

Adjusted model 1 adjusted for age, gender, ethnicity, cigarette smoking, hypertension, diabetes, hyperlipidemia, and family history of premature CVD.

Adjusted model 2; adjusted for age, gender, ethnicity, cigarette smoking, hypertension, diabetes, hyperlipidemia, and family history of premature CVD, and utilizing statin and aspirin