Published in final edited form as:

Dev Psychopathol. 2012 August ; 24(3): 755–769. doi:10.1017/S0954579412000351.

Externalizing problems, attention regulation, and household chaos: A longitudinal behavioral genetic study

Zhe Wang,

Virginia Polytechnic Institute and State University

Kirby Deater-Deckard,

Virginia Polytechnic Institute and State University

Stephen A. Petrill, and

The Ohio State University

Lee A. Thompson

Case Western Reserve University

Abstract

Previous research has documented a robust link between difficulties in self-regulation and development of externalizing problems (i.e., aggression and delinquency). In the current study, we examined the longitudinal additive and interactive genetic and environmental covariation underlying this well-established link using a twin design. The sample included 131 pairs of monozygotic twins and 173 pairs of same-sex dizygotic twins who participated in three waves of annual assessment. Mothers and fathers provided reports of externalizing problems. Teacher report and observer rating were used to assess twin's attention regulation. The etiology underlying the link between externalizing problems and attention regulation shifted from a common genetic mechanism to a common environmental mechanism in the transition across middle childhood. Household chaos moderated the genetic variance of and covariance between externalizing problems and attention regulation. The genetic influence on individual differences in both externalizing problems and attention regulation was stronger in more chaotic household. However, higher levels of household chaos attenuated the genetic link between externalizing problems and attention regulation.

Keywords

externalizing problems; attention regulation; household chaos; longitudinal; behavioral genetics

Address correspondence to: Kirby Deater-Deckard, Department of Psychology, Virginia Tech, 109 Williams Hall (0436), Blacksburg, VA, 24061. kirbydd@vt.edu.

Zhe Wang and Kirby Deater-Deckard are in the Department of Psychology, Virginia Polytechnic Institute and State University, Blacksburg, VA. Stephen A. Petrill is in the Department of Human Development and Family Science, The Ohio State University, Columbus, OH. Lee A. Thompson is in the Department of Psychological Sciences, Case Western Reserve University, Cleveland, OH.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Child Health And Human Development or the National Institutes of Health.

There is an established literature pointing to the importance of children's systems of self-regulation in deterring the development of externalizing behavior problems such as aggression and delinquency. Children with more difficulties in self-regulation show more externalizing problems concurrently and longitudinally, suggesting a synergy between the two constructs. Yet, the underlying etiology of individual differences in this developmental process is unclear. In the current study, we examined the concurrent and longitudinal associations between externalizing problems and behavioral indicators of attention regulation to estimate additive and interactive genetic and nongenetic influences using a twin behavioral genetic design.

Externalizing Problems and Attention Regulation

There is mounting evidence that good attention regulation plays a central role in a healthy developmental trajectory that is relatively free from behavioral and emotional problems. Difficulties with regulating attention are associated with more aggressive and delinquent behaviors, both concurrently and longitudinally (Kochanska & Knaack, 2003; Murray & Kochanska, 2002; Olson, Sameroff, Kerr, Lopez, & Wellman, 2005; Zhou, et al., 2007). Children with better attention regulation are more able and flexible in choosing the "to-be-attended-to" information in the external world and in their internal thought processes. Refocusing attention can be an effective strategy in modulating anger, thereby reducing its contribution to antisocial behavior (Kim & Deater-Deckard, 2011; Morris, Silk, Steinberg, Terranova, & Kithakye, 2010; Rothbart, Ellis, Rueda, & Posner, 2003). Attention regulation also can be used to inhibit impulsive inappropriate actions (Posner & Rothbart, 2006).

Behavioral genetic studies have been conducted to clarify the genetic and nongenetic etiology underlying the link between externalizing problems and attention regulation. Univariate behavioral genetic studies have suggested that individual differences in externalizing problems are attributable to moderate genetic, modest shared environmental (i.e., nongenetic factors that contribute to family member similarity) and moderate nonshared environmental (i.e., nongenetic factors that do not contribute to family member similarity) sources of variance (Miles & Carey, 1997; Rhee & Waldman, 2002). The longitudinal stability of individual differences in externalizing problems is mainly accounted for by genetic and shared environmental factors (Bartels et al., 2007). Individual differences in sustained attentive behavior are attributable to moderate genetic variance and moderate nonshared environmental variance (Deater-Deckard & Wang, 2012; Fan, Wu, Fossella, & Posner, 2001). The longitudinal stability of attention is accounted for by stable genetic influences, while changes in attention over time appear to be mediated by nonshared environmental influences such as differential parental sensitivity in caregiving behavior (Deater-Deckard, Petrill, Thompson, & DeThorne, 2006).

But what is the etiology of the longitudinal connections between attention and externalizing problems? This question has not been addressed in previous research, and answering it provides the main rationale for current investigation. To answer this question, multivariate quantitative genetic models are needed to reveal the underlying structure of variance and covariance through which externalizing problems and attention regulation are linked. Using the first wave from the current longitudinal study, we used a multivariate model to conduct a

cross-sectional analysis and found a common genetic factor underlying the covariation between poorer attention and more conduct problems above and beyond any underlying genetic influences involving negative affect (Deater-Deckard, Petrill, & Thompson, 2007). Other behavioral genetic studies have taken a similar approach, and the results are mixed. Some have suggested a common genetic factor (Dick, Viken, Kaprio, Pulkkinen, & Rose, 2005; Nadder, Rutter, Silberg, Maes, & Eaves, 2002), but others have pointed to a common underlying shared environmental factor (Burt, Krueger, McGue, & Iacono, 2001; Burt, McGue, Krueger, & Iacono, 2005) or nonshared environmental factor (Deater-Deckard et al., 2007).

There are several features of the behavioral genetic studies in the literature that could explain the mixed results. First, the samples vary in terms of children's ages. This matters, because there is evidence of growth in heritable variance and dissipation of shared environmental variance in attention over the transition to and through middle childhood (Deater-Deckard & Wang, 2012). By comparison, over the same developmental period, externalizing problems show stable nonshared environmental variance but there are potential fluctuations in levels of heritable and shared environmental variance over time (Bartels et al., 2007). Thus, the mechanism that links externalizing problems and attention regulation may also vary across time. To our knowledge, only one prior study has examined longitudinally the genetic and nongenetic etiology of the link between attention and externalizing—and in that case, the emphasis was on attention deficit disorder (ADHD) and oppositional defiant disorder (ODD) symptoms (Nadder et al., 2002). That study found a genetic factor that captured all the covariation between ADHD and ODD, across two time points. However, the sample spanned a wide age range (8 to 16 years), resulting in a potential washing out of any discernible developmental shifts that might be evident over smaller age spans and shorter periods of development in middle childhood and adolescence. In the current study, we used a cross-sectional and longitudinal behavioral genetic design that spans a narrower age range (5 to 8 years old in the first wave) and includes three annual assessments in middle childhood, prior to puberty.

A second factor that might contribute to the mixed results in the literature lies in the intrinsic limitation of the quasi-experimental twin design, which is relatively underpowered for detecting shared environment effects compared to the higher statistical power for detecting genetic effects (Plomin, DeFries, McClearn, & McGuffin, 2008). Depending on sample size and whether non-twin siblings were also included in the analysis, prior studies have varied in their ability to detect significant shared environmental variance in, and covariance between, externalizing and attention. Although we used a twin design in the current study, we attempted to partially ameliorate this power limitation by examining correlated constructs (i.e., attention and externalizing) in longitudinal multivariate models—an approach that increases power for detecting shared environment effects (Schmitz, Cherny, & Fulker, 1998).

Third, genetic and nongenetic variance and covariance estimates are influenced by the methods that are used to assess attention regulation and externalizing problems, so differences in studies in methodology can influence results. Typically, larger shared environment effects and smaller genetic effects are obtained when observations are used, but

smaller shared environment effects and larger genetic effects are obtained when questionnaires are used (Ghodsian-Carpey & Baker, 1987; Leve, Winebarger, Fagot, Reid, & Goldsmith, 1998). Furthermore, evidence of a common shared environmental factor in the covariation between two constructs (such as attention and externalizing) is more likely to be found if the same method and informant is used to assess both constructs, very likely reflecting a shared method effect. To our knowledge, nearly all of the prior behavioral genetic studies of attention and externalizing have this monomethod limitation. In the current study we used a multi-informant multi-method approach so that the covariation between the two constructs would include minimal amounts of overlapping variance arising from method and informant. Mothers and fathers rated their children's externalizing problems, and observers and teachers rated children's sustained attentive behaviors.

The cognitive neuroscience literature has emphasized measurement of attention regulation using task-based performance, typically in a laboratory setting. However, this is not necessary when the goal is to examine the behavioral manifestations of attention regulation. Observational and questionnaire-based assessments of sustained attentive behavior such as those used in the current study are more feasible in large longitudinal designs compared to laboratory-based task assessments, and are reliable (though distal) indicators of underlying cognitive mechanisms of attention regulation (Deater-Deckard & Wang, 2012; Rothbart, 2007; Willcutt, Doyle, Nigg, Faraone, & Pennington, 2005). These broader behavioral measures of attention are suitable when the goal is to examine links with other aspects of behavior (such as externalizing problems), as is the case in the current study. However, the caveat is that these kinds of measures do not precisely specify the sub-components of attention regulation that can be assessed with task-based laboratory performance measures.

Gene by Environment Interaction: Household Chaos as Context

An assumption that is required in the interpretation of genetic and nongenetic factors in the variance in, and covariance between, attention and externalizing problems is that these effects are additive. However, the soundness of this assumption has been increasingly called into question based on more recent empirical findings and methodological advances. For example, Leve et al. (2010) used a parent-offspring adoption design and reported a statistical interaction between adoptive parent anxious/depressive symptoms and adopted infants' genetic risk for externalizing problems, in the prediction of those children's subsequent heightened attention to frustrating events (Leve et al., 2010). Traditional behavioral genetic analyses cannot detect this kind of interaction, with the gene-by-environment interaction variance being included in the heritable and nonshared environmental variance estimates. Additive genetic and nongenetic covariation between externalizing problems and attention regulation has been detected in prior studies (Deater-Deckard et al., 2007; Dick et al., 2005; Nadder et al., 2002), but the presence of interactive effects has been untested. Therefore, in the current study we examined this possibility by incorporating household chaos as a candidate environmental factor into the traditional behavioral genetic model.

Household chaos reflects the level of disorganization in the family environment, including noise, crowding, and unpredictability in daily routines. It was chosen in the current study as the environmental attribute of interest because it is one of the most important proximal

environment factors through which other environmental factors exert their influence on a variety of socioemotional and psychopathology outcomes (Evans, Gonnella, Marcynyszyn, Gentile, & Salpekar, 2005). Higher levels of household chaos have been shown to predict children's poorer cognitive performance, more limited attention focusing, and more behavioral and emotional problems concurrently and longitudinally, even after controlling for other confounding environmental factors such as parental education and IQ, parental warmth and negativity, parental stress, and other neighborhood attributes (Deater-Deckard et al., 2009; Dumas et al., 2005; Pike, Iervolino, Eley, Price, & Plomin, 2006). Chaos also functions as a moderator in the link between parenting or classroom environments and children's conduct problems, whereby it exacerbates the effect of poor parenting and classroom environments on children's behavioral problems (Asbury, Dunn, Pike, & Plomin, 2003; Coldwell, Pike, & Dunn, 2006; Oliver, Pike, & Plomin, 2008). In the current study, we tested whether household chaos interacted with genetic variance and covariance in explaining the link between attention and externalizing behavior problems over middle childhood. To summarize, advances in theory and empirical research on the development of psychopathology require the examination of multiple levels of analysis—in the current study, from the home environment (i.e., chaos) to behavioral assessments of siblings to behavioral genetic estimates of heritable and nonheritable factors—to elucidate the developmental processes that cause growth and minimization of antisocial behavior problems in childhood and adolescence (Cicchetti & Valentino, 2007). We addressed the following questions:

1. At the “phenotypic” level, are externalizing problems and attention regulation correlated when covariance from shared method and informant effects is minimized? Furthermore, does the correlation differ as a function of household chaos?
2. At the behavioral genetic level, what is the genetic and environmental etiology underlying the phenotypic correlation? Is it attributable to additive genetic and environmental effects, or an interactive gene x environment interplay involving household chaos?
3. Longitudinally, does the underlying genetic and environmental etiology of the phenotypic correlation change across middle childhood?

Method

Participants

The data were from the first three annual waves of a longitudinal twin study: the Western Reserve Reading Project (WRRP). Families were recruited from the Cleveland, Columbus, and Cincinnati metropolitan areas. There were 131 pairs of monozygotic twins (MZ twins; 39% male) and 173 pairs of dizygotic twins (DZ twins; 44% male). The average age was 6.07 (SD = .68, range = 4.33 to 7.92) in the first wave, 7.15 (SD = .67, range = 6.00 to 8.83) in the second wave, and 8.30 (SD = .75, range = 6.50 to 10.00) in the third wave. Ninety-one percent of the sample was Caucasian, 5% African-American, and 2% Asian. Twenty-five percent of the parents had some post-graduate education or degree, 31% bachelor's degree,

23–27% some college, and 12–17% high school or less. Ninety percent of the parents were married and lived together, 2% were unmarried but cohabited, and 6% were single.

Procedure

In each annual assessment wave, informed consent was obtained and parents were informed that they would receive \$100 for their participation. Parents and twins completed a series of behavioral and cognitive assessments during a three-hour home visit. At the end of the home visit, twins' behaviors during the cognitive assessments were rated by trained testers. Teachers completed questionnaires via mail.

Measures

Externalizing problems—We used the externalizing problems syndrome score (including delinquent behavior and aggression) from the Child Behavior Checklist (CBCL; Achenbach, 1991) to measure child externalizing problems. The items were rated by mothers and fathers on a 3-point Likert-type scale (0 = not true, 1 = somewhat or sometimes true, 2 = very true or often true). Correlations between mother and father reports ranged from .44 to .58, depending on twin and longitudinal wave. Mother and father reports were averaged, when applicable, to obtain a more reliable parent-rated externalizing problem score. When only mother or father report was available, a single parent report was used. Among the 608 children, 554 children had mother reports on externalizing problems in the first wave, 501 in the second wave, and 435 in the third wave. Three hundred and seven children had father reports on externalizing problems in the first wave, 274 in the second wave, and 189 in the third wave. Three hundred and three children had both parents' reports in the first wave, 263 in the second wave, and 185 in the third wave. Higher scores represented more externalizing problems.

Attention regulation—Three items were selected from the Teacher Report Form (TRF; Achenbach, 1991). The TRF is a widely-used teacher's ratings of children's behavioral and emotional problems. The items are rated on a 3-point Likert-type scale (0 = not true, 1 = somewhat or sometimes true, 2 = very true or often true). Three items were chosen to assess children's difficulties in attention regulation: 4-fails to finish things; 8-cannot concentrate or pay attention for long; 78-inattentive, easily distracted. Each child was rated by three different teachers at three different waves. In the first wave, 46.5% of the twin pairs were rated independently by two different teachers, 64.8% in the second wave, and 63.4% in the third wave. The percentage of twin pairs who were rated independently by two different teachers at each wave was the same for MZ and DZ twins.

Two items were selected from Bayley's Behavior Record (BBR; Bayley, 1969). Trained testers rated child's behavior after observing him/her during a three-hour cognitive assessment using the BBR. Twin siblings were rated independently by two different testers at each wave. A different tester rated each child at each wave. The BBR includes 26 items rated on a 5-point Likert-type scale. Two items relevant to attention regulation were included: attention to task (1 = constantly off task, does not attend, 3 = off task half the time, 5 = constantly attends) and persistence in attempting to complete task (1 = consistently lacks persistence, 3 = lacks persistence half the time, 5 = consistently persistent).

Attention regulation was tested as a composite score that included the three teacher-rated items and two tester-rated items. In each of the three waves, principal component analyses showed acceptable internal consistency (variance explained: from 46% to 55%; loadings: from .40 to .86). Therefore, these items were standardized, averaged, and standardized again to obtain an attention regulation composite z-score, separately for each wave. The composite scores were scaled such that higher scores represented more difficulties in regulating attention.

Household chaos—A shortened version of the Chaos, Hubbub and Order Scale (CHAOS, Matheny, Wachs, Ludwig, & Philips, 1995) was used to measure household chaos at all three waves. Items in the short version are rated on a 5 point Likert-type scale (1 = definitely untrue, 3 = neither untrue nor true, 5 = definitely true) instead of the original binary (yes/no) scale. The short version has been used in several previous studies (Coldwell et al., 2006, $\alpha = .56$; Deater-Deckard et al., 2009, mother-father rater agreement and test-retest reliabilities in the .6 to .8 range; Pike et al., 2006, $\alpha = .63$). The short version includes six items from the original scale: ‘I have a regular morning routine’ (reverse scored), ‘You can’t hear yourself think in our home’, ‘It’s a real zoo in our home’, ‘We are usually able to stay on top of things’ (reverse scored), ‘There is usually a television turned on somewhere in our home’, and ‘The atmosphere in our house is calm’ (reverse scored). Mother and father reports were averaged, when applicable, to obtain a more reliable chaotic home environment score, with higher scores indicating more household chaos. When only mother or father report was available, a single parent report was used. Among the 304 families, 276 had mother reports on household chaos in the first wave, 251 in the second wave, and 218 in the third wave. One hundred and fifty-one families had father reports on household chaos in the first wave, 137 in the second wave, and 96 in the third wave. One hundred and forty-eight families had both mother and father reports on household chaos in the first wave, 132 in the second wave, and 94 in the third wave.

Missing Data

Because the current study involved three annual assessments, we examined whether there were mean differences in the initial assessment of the main study variables (i.e., externalizing problems, attention regulation, and household chaos) between children who did and did not have missing values on these variables assessed in the third wave. We used the main study variables assessed in the 1st wave as outcome variables and whether or not there were missing values on each of the three variables in the 3rd wave (0 = yes; 1 = no for each variable) as predictors in a multivariate analysis of variance, separately for each twin. For both twins, there were no mean differences on externalizing problems, attention regulation, and household chaos assessed in the 1st wave between children who did and did not have missing values on all three variables assessed two years later ($F = .47$ to 2.09 , $p = .10$ to $.70$). Therefore, in the following analyses, full information maximum likelihood (FIML) was used for all the model fitting procedures.

Results

Descriptive statistics are shown in Table 1 on the diagonal. The average level of externalizing problems decreased over the three annual assessments from 7.18 to 6.29. Because attention scores were composite Z-scores, they had means of 0 and standard deviations of 1 at each wave, and any potential mean-level changes could not be determined. We used a repeated-measure multivariate analysis of variance to test whether this decrease was significant, and whether this pattern differed as a function of sex or zygosity. Neither the interaction between time and sex, $F(2, 398) = .91, p < .40$, nor the interaction between time and zygosity, $F(2, 398) = 2.49, p < .10$, was significant, suggesting that longitudinal mean level changes in externalizing problems did not differ by gender or zygosity. Regardless of sex and zygosity, externalizing problems significantly decreased over time, $F(2, 398) = 8.18, p < .001$. The same analysis was conducted for household chaos, and the results suggested that the average level of household chaos did not significantly decrease over time, $F(2, 200) = 2.35, p > .05$.

Question 1: Phenotypic Analyses

Our first question was: At the “phenotypic” (i.e., directly observed) level, are externalizing problems and attention regulation correlated when covariance from shared method and informant effects is minimized? Furthermore, does the correlation differ as a function of household chaos?

Bivariate Pearson “phenotypic” (i.e., directly observed) correlations between study variables were estimated and are shown in Table 1. These correlations were tested for twin 1 and twin 2 together, with significance tests adjusted via the equation $Z = r_{xy} * \sqrt{2N / (1 + r_{xx'} * r_{yy'} + r_{xy'}^2)}$, to control for bias in standard errors arising from the non-independence of the twin data within families (Griffin & Gonzalez, 1995). Attention regulation, externalizing problems and household chaos all showed moderate to substantial stability. Stability correlations for consecutive assessments were .35 to .47 for attention regulation, .68 to .76 for externalizing problems, and .72 to .80 for household chaos. Concurrent correlations between difficulties in attention and externalizing problems were modest to moderate and significant, ranging from .14 to .23. Difficulties in attention regulation and externalizing problems both were concurrently correlated with household chaos in all three waves ($r = .14$ to .17, $p < .05$ for attention, and $r = .28$ to .32, $p < .001$ for externalizing).

Next, we tested whether household chaos level moderated the phenotypic links between attention regulation and externalizing problems at each wave. We first computed the interaction terms between attention regulation and chaos for each wave by multiplying the standardized (for centering) attention and chaos scores. We then applied the actor-partner interdependence model (APIM; Kenny, 1996; see Figure 1) to test for the additive as well as the interactive effect of attention regulation and chaos on externalizing problems. The APIM is preferred over standard regression for two reasons. First, its structural equation modeling approach allows us to examine the additive and the interactive effects of attention regulation and chaos on externalizing problems together for both twins while controlling for biased standard errors arising from twin non-independence (Olsen & Kenny, 2006). Second, the APIM can be estimated in Amos 18.0 which has the added advantage of permitting use of

full information maximum likelihood (FIML) for estimating effects with missing data—an approach that has been shown to be more appropriate than traditional methods (e.g., mean substitution) for handling missing data (Enders & Bandalos, 2001).

As shown in Figure 1, b1 and b2 paths represent main effects of attention regulation and chaos on externalizing problems respectively, and b3 paths represent interactive effects between attention regulation and chaos on externalizing problems, above and beyond main effects. To test for significance of the main and interactive effects, constrained models in which main effects or interactive effects were fixed at 0 were compared against the full model using χ^2 difference tests. A significant χ^2 difference suggested that fixing a certain path at 0 in the constrained model resulted in a decrease in model fit, which in turn indicated that the fixed path made significant contribution in explaining the variance in externalizing problems and therefore should be freely estimated. To test for main effects of attention regulation, b1 paths and b3 paths were both fixed at 0. To test for main effects of chaos, b2 and b3 paths were fixed at 0. To test for interactive effects between attention and chaos, b3 paths were fixed at 0. Table 2 shows the model fit indices and the standardized path estimates from the full model. For all three waves, dropping main effects of attention regulation and chaos both led to significant decreases in model fit ($\chi^2 = 11.11$ to 68.46, $df = 2$, $p = .004$ to .000), indicating statistical significance of main effects of attention regulation and chaos on externalizing problems. Poorer attention regulation and higher chaos independently predicted more externalizing problems at all three waves.

Dropping the interactive effect between attention regulation and chaos led to a marginally significant decrease in model fit for the 1st wave ($\chi^2 = 3.78$, $df = 1$, $p = .052$), a significant decrease in model fit for the 2nd wave ($\chi^2 = 8.44$, $df = 1$, $p = .004$), and no significant decrease in model fit for the 3rd wave ($\chi^2 = 1.56$, $df = 1$, $p = .211$). This suggested that for the 2nd wave and possibly the 1st wave, attention regulation and chaos also interactively predicted externalizing problems: for the 1st wave, externalizing = $.14 * \text{attention} + .25 * \text{chaos} - .06 * \text{attention} * \text{chaos} = (.14 - .06 * \text{chaos}) * \text{attention} + .25 * \text{chaos}$; for the 2nd wave, externalizing = $.07 * \text{attention} + .29 * \text{chaos} - .09 * \text{attention} * \text{chaos} = (.07 - .09 * \text{chaos}) * \text{attention} + .29 * \text{chaos}$. Therefore, for both waves, the higher the level of chaos was, the smaller the effect of attention regulation on externalizing problems was.

Question 2: Cross-sectional Behavioral Genetic Analyses

Our second question was: At the behavioral genetic level, what is the genetic and environmental etiology underlying the phenotypic correlation? Is it attributable to additive genetic and environmental effects, or an interactive gene x environment interplay involving household chaos? To examine the etiology of the link between externalizing problems and attention regulation within each of the three annual waves, we applied a bivariate Cholesky decomposition model. On the left side of Figure 2 (shown for only one twin in the pair for simplicity), a traditional Cholesky decomposition model is presented. In this basic model, the observed phenotypic variance and covariance were decomposed into latent “overlapping” or common additive genetic (A), common shared environmental (C), and common nonshared environmental (E) variance, as well as latent residual or “unique” genetic (a), residual shared environmental (c), and residual nonshared environmental (e)

variance. On average, monozygotic (MZ) twins share all of their alleles and dizygotic (DZ) twins share 50% of their segregating alleles identical by descent. Therefore, the correlation for genetic similarity between twins was set as 1 and .5 for MZ and DZ twins, respectively. The correlations for shared environmental (i.e., environmental factors that lead to twin similarity) and nonshared environmental (i.e., environmental factors that do not lead to twin similarity) variance between twins were set as 1 and 0, respectively for both MZ and DZ twins.

A modified Cholesky decomposition model is presented on the right side of Figure 2 (Dick & York, 2010; Purcell, 2002), again shown only for one twin in the pair. Chaos was tested as a continuous moderator of the overlapping (A) and unique (a) genetic components. In this modified Cholesky model, the genetic variance was no longer a constant, but a linear function of the moderator (i.e., chaos), expressed as $A_{\text{total}} = A' + \beta M$. In this equation, A' indicated the magnitude of additive genetic influences. The value of M was the measured chaos for each family, and thus varied from family to family. β represented the magnitude of the moderating effect to be estimated from the data. The paths A_{att} , A_{cov} , and a_{ext} represented total genetic variance for attention regulation, genetic covariance between attention regulation and externalizing problems, and residual genetic variance for externalizing problems, respectively. Accordingly, β_{att} , β_{cov} , and β_{ext} represented the extent to which chaos changed the degree of total genetic influences on attention regulation, genetic covariance between attention regulation and externalizing problems, and unique genetic variance for externalizing problems, respectively.

We tested the modified bivariate Cholesky model for each wave separately. All models were fit to raw continuous data using maximum likelihood model-fitting procedures in Mx (Neale, 1997). The significance of the moderating effect was tested by dropping the corresponding moderation parameter and examining the change in $-2 \log$ likelihood (-2ll) between the full model and the constrained model. The difference in -2ll follows a chi square distribution. Therefore, a significant difference in -2ll suggested that dropping the moderation parameter led to a significant decrease in model fit, which in turn suggested that the moderation effect on that particular path was statistically significant.

Externalizing problems and chaos scores were standardized (for centering purposes) for the following analyses. Table 3 shows model fit and comparison results. For all three waves, dropping the moderation effect on total genetic variance in attention regulation and unique genetic variance in externalizing problems led to significant decreases in model fit ($-2 \log$ likelihood = 6.79 to 23.85, $df = 1$, $p = .009$ to $.000$), suggesting a significant moderating effect of chaos on the total genetic variance of attention regulation and unique genetic variance in externalizing problems. The moderation effect of chaos on the genetic covariance between attention regulation and externalizing problems was only significant at the 1st wave, as suggested by a significant decrease in model fit resulting from constraining this effect ($-2\text{ll} = 4.91$, $df = 1$, $p = .027$). This moderation effect on the genetic covariance between attention and externalizing was not significant in the 2nd and 3rd waves ($-2\text{ll} = 2.07$ to 2.28 , $df = 1$, $p = .130$ to $.150$).

In order to compare models between the three annual waves, we have presented results from the full model for each wave. The unstandardized variance components, and genetic and nongenetic correlations at three levels of the moderator (i.e., 1SD below the mean, the mean, and 1SD above the mean of chaos scores), are reported in Table 4. In each wave, genetic variance of both attention regulation and externalizing problems increased as the chaos level increased. The genetic correlation between attention control and externalizing problems became smaller, as chaos scores were higher, only in the first wave.

The genetic, shared environmental, and nonshared environmental variance components were all consistent across the three waves. However, the genetic correlation gradually decreased over time, from .45 to .01 at average levels of chaos, while the shared environmental correlation and the nonshared environmental correlation gradually increased over time—from .02 to .84 for shared environment correlation, and from .06 to .32 for nonshared environment correlation. Overall, it seems that the mechanisms through which attention regulation and externalizing problems were linked gradually shifted from genetic correlation to nongenetic correlations, independent of any potential moderating effects of household chaos.

Question 3: Longitudinal Behavioral Genetic Analyses

Although the cross-sectional analyses for question 2 allowed us to examine genetic and nongenetic statistical effects in each of three waves, those analyses did not address longitudinal continuity and change in genetic and nongenetic variance of and covariance between attention and externalizing problems—that is, the analyses did not address whether the genetic and nongenetic factors found in each wave also accounted for stability and change in individual differences over the three year period. Therefore, a longitudinal multivariate behavioral genetic analysis was used to address the third and final question of our study. We were especially interested in learning whether there are: a) overlapping genetic, shared environmental, and nonshared environmental factors that would statistically account for the longitudinal phenotypic covariance between attention and externalizing over three years; b) wave-specific genetic, shared environmental and nonshared environmental covariances between attention and externalizing; and c) unique variances remaining for both constructs at each time point after controlling for all the overlapping sources of covariance between the two constructs. However, the Cholesky decomposition model used above does not address those important sub-questions. Therefore, we instead used a multivariate common-factor independent pathway model. As shown in Figure 3 (shown for one twin in the pair), phenotypic variance of and covariance between attention regulation and externalizing problems were decomposed into latent genetic (A), shared environmental (C), and nonshared environmental (E) variance and covariance. Latent A1, C1, and E1 represented overlapping additive genetic, shared environmental, and nonshared environmental covariance, respectively, between attention regulation and externalizing problems across all three waves. A2, C2, and E2 represented any temporally emerging additive genetic, shared environmental, and nonshared environmental covariances, respectively, between attention regulation and externalizing problems that overlapped across the 2nd and 3rd waves, and that was independent of the 1st wave. A3, C3, and E3 represented the additive genetic, shared environmental, and nonshared environmental covariances,

respectively, between attention regulation and externalizing problems appearing only in the 3rd wave. Residual genetic, shared environmental, and nonshared environmental variances unique to each variable at each time point were represented by a, c, and e, respectively. The pathways between genetic, shared environmental, and nonshared environmental variance and covariance across twins were set in the exactly the same manner as in the modified Cholesky decomposition models used in previous analyses.

Nested models were systematically tested by fixing certain paths at 0, and difference in -2ll were used to select the best fitting models among all competing models to balance the loss in fit with the difference in degrees of freedom. Model trimming continued until no more paths could be fixed at 0 without worsening model fit. Compared to the full model ($-2\text{ll} = 7324.34$, $df = 3053$), the final trimmed model ($-2\text{ll} = 7332.71$, $df = 3080$) did not have a significant decrease in fit ($-2\text{ll} = 8.37$, $df = 27$, $p > .05$), and therefore was chosen to be interpreted (see Table 5).

First, we examined this model by focusing on univariate continuity and change. Results suggested that there was a general genetic factor and a general nonshared environmental factor that explained the longitudinal continuity in externalizing problems across all three waves, represented by significant paths from A1 (.88, .66, and .64) and E1 (.34, .34, and .30). Yet, there was also a new set of genetic influences emerging at the 2nd wave that accounted for the continuity in externalizing problems from just wave 2 to wave 3 (paths from A2: .45 and .52). Changes in externalizing problems were mainly attributable to unique genetic variance at the 2nd and 3rd waves (paths from a: .35 and .40), as well as unique nonshared environmental variance at all three waves (paths from e: .33, .35, and .26). Shared environmental variance was only detected at the 3rd wave (paths from C1: .09). Turning to attention regulation, continuity across three waves was attributable to shared environmental influences, represented by significant path estimates from C1 (.56, .66, and .54). There also was a general genetic factor that explained continuity across the first two waves (paths from A1: .18 and .19), and a general nonshared environmental factor that explained continuity across the second two waves (paths from E2: .38 and .27). Unique genetic variance was found for the 1st and 3rd waves (paths from a: .37 and .45), and unique nonshared environmental variance was found for all three waves (paths from e: .72, .62, and .57), accounting for changes in attention regulation.

Next, we focused on bivariate longitudinal relations between attention regulation and externalizing problems. The bivariate phenotypic correlation between attention regulation and externalizing problems was accounted for by a general genetic factor across the first two waves, represented by significant path estimates from A1 (.18, .88, .19, and .66). The bivariate phenotypic correlation between attention regulation and externalizing problems at the 3rd wave was attributable to shared and nonshared environmental covariance, represented by significant path estimates from C1 (.54 and .09) and E1 (.34 and .30).

Discussion

There is a growing literature documenting the strong connection between children's difficulties in self-regulation and their externalizing behavioral problems (Kochanska &

Knaack, 2003; Pitzer, Esser, Schmidt, & Laucht, 2009; Prior, Smart, Sanson, & Oberklaid, 2001). Attention regulation is an essential core component of self-regulatory capacity (Posner & Rothbart, 2006), and has been demonstrated to deter the development of behavioral problems through inhibiting impulsive responses, redirecting attention to less negative internal as well as external stimuli, and dampening other negative environmental influences or facilitating positive environmental influences (Belsky, Fearon, & Bell, 2007; Eisenberg et al., 2005; Kim & Deater-Deckard, 2011; Morris et al., 2010). The aim of the current study was to extend this literature by applying a multi-informant multi-method approach to investigate the underlying genetic and environmental etiology of the link between attention regulation and externalizing behavior problems as well as any transition in the etiology over three years in middle childhood.

Externalizing Problems and Attention Regulation

Our first question was: At the “phenotypic” level, are externalizing problems and attention regulation correlated when covariance from shared method and informant effects is minimized? Furthermore, does the correlation differ as a function of household chaos? Descriptive analyses indicated that both externalizing problems and attention regulation showed considerable stability over time. The correlation between the two constructs was modest and comparable across the three waves. Yet, the magnitude of the correlations (.14 to .23) appeared somewhat lower than those found in other studies (e.g., Eisenberg et al., 2005). We argued that this is likely due to the removal of any shared method effect (e.g., instrumental, situational, and informant overlap) between the two constructs in the current study: mothers and fathers provided reports on externalizing problems whereas teachers and observers provided reports on attention regulation. Other studies examining the correlation between effortful sustained attentive behavior and externalizing problems using a multi-informant multi-method approach have reported similar effect sizes (Eisenberg et al., 2004; Kim & Deater-Deckard, 2011).

Our second question was: At the behavioral genetic level, what is the genetic and environmental etiology underlying the phenotypic correlation? Is it attributable to additive genetic and environmental effects or an interactive gene x environment interplay involving household chaos? To address this question, we investigated the genetic and environmental variance of and covariance between externalizing problems and attention regulation in each of the three waves. Cross-sectional results suggested that the genetic, shared environmental, and nonshared environmental variance components in both attention regulation and externalizing problems were all relatively consistent across the three years. However, household chaos moderated the genetic variance in both attention regulation and externalizing problems in all three waves, whereby genetic influences on both externalizing behavior problems and difficulties in attention regulation were more substantial in more chaotic environments.

As early as the 1980s, gene environment interaction effects had been proposed and initially demonstrated in the etiology of various externalizing behavior problems (Cadoret, Cain, & Crowe, 1983). Cadoret et al. used an adoption design to show that a genetic “risk” for antisocial behavior statistically interacted with adversity in the adoptive home environment

(e.g., parental psychiatric problems, marital discord) to promote higher levels of antisocial behavior in adolescence. This finding was consistent with another study showing a similar effect for antisocial behavior and criminality (Mednick, Gabrielli, & Hutchings, 1984). More recent research suggests that the predictive effects on externalizing problems of several candidate genes including dopamine receptor genes DRD4 and DRD5 and transporter gene DAT1 are enhanced when they are coupled with prenatal risks such as maternal nicotine and alcohol use (Brookes et al., 2006; Langley et al., 2008; Thapar et al., 2005). Other research points to growth in genetic variance in attentive behavior over middle childhood, fueled in part by interaction between DRD4 and maternal sensitivity (Berry, Deater-Deckard, McCartney, Wang, & Petrill, 2013; Deater-Deckard & Wang, 2012). The findings from the current study implicate household chaos as another important candidate environmental factor to be considered as a moderator of genetic influences on antisocial behavior and attention regulation.

Turning to the bivariate link between attention regulation and externalizing problems, in the phenotypic analyses (Question 1) we found that both attention regulation and externalizing problems were correlated with chaos, whereby higher levels of chaos were associated with more difficulties in attention regulation and more externalizing problems. This finding is consistent with previous literature (Coldwell et al., 2006; Deater-Deckard et al., 2009; Pike et al., 2006). In addition, chaos also statistically moderated the association between attention regulation and externalizing problems in the 2nd wave, and was a marginally significant moderator in the 1st wave. Poor sustained attentive behavior was a stronger statistical predictor of externalizing problems in less chaotic families. In regard to the behavioral genetic analyses, levels of household chaos also moderated the genetic correlation between externalizing problems and attention regulation in the 1st wave, whereby the genetic correlation was lower at higher levels of chaos.

When taken together with the phenotypic results, the behavioral genetic findings suggest that higher levels of household chaos overwhelm the regulatory processes involving attention, reflected both at the observed phenotypic level of analysis and at the underlying latent behavioral genetic level of analysis. Executive regulation of attention is likely to depend at least in part on the broader family context. Chronic exposure to uncertainty and unpredictability in chaotic household heavily taxes executive regulation capacity (Evans, Hygge, & Bullinger, 1995) and hence may compromise its development, potentially via the neuroendocrine system (Erickson, Drevets, & Schulkin, 2003). In addition, constant distraction in a chaotic household interferes with effortful regulation processes of behavior (Blair et al., 2007). These regulatory processes may operate in a relatively automatic and more efficient fashion when no such distracters or stressors are present (Macrae & Bodenhausen, 2000). Therefore, the stress and inconsistency of daily life in chaotic households may disrupt the development of self-regulatory capacity and skills that are important to minimizing growth in behavior problems. This idea is consistent with broader theoretical and empirical work implicating chaos as a source of impairment in healthy, normative developmental processes (Evans & Wachs, 2009).

There are candidate genes and neurological pathways that are thought to account for the association between deficits in attention regulation and conduct problems (Holmes et al.,

2002; Raine, 2002). However, to our knowledge no theories or empirical research have addressed whether and how environmental adversities—including household chaos—constrain these genetic and neurophysiological links between attention and externalizing behavior problems. The current study suggests that answering these crucial questions is an important next step for future studies.

It is worth emphasizing that the moderating effect of chaos on the genetic correlation between attention and externalizing was only significant in the 1st wave. This could be due in part to the dissipation of the genetic correlation over the three waves (from .45 in wave 1 to essentially 0 by wave 3)—at the same time that we saw a similar magnitude increase in shared environment correlation across the three waves. We address this interesting developmental pattern in more detail below when interpreting the longitudinal data analyses.

Longitudinal Influences on Stability and Change

Our third and final question was: Longitudinally, does the underlying genetic and environmental etiology of the phenotypic correlation change across middle childhood? Specifically, do the genetic and nongenetic factors found in each wave also accounted for stability and change in individual differences over the three year period? We tested a longitudinal multivariate common-factor independent pathway model to answer this question, by addressing externalizing, then attention, then the covariation between them.

The stability of individual differences in externalizing problems was mainly accounted for by the same genetic influences over the three-year period. This is consistent with previous studies of younger and older children (Eley, Lichtenstein, & Moffitt, 2003; van Beijsterveldt, Bartels, Hudziak, & Boomsma, 2003). Not all of the variance in externalizing problems was explained by this single underlying “stability” genetic factor—however, the unique residual genetic factors at waves 2 and 3 were modest in effect size by comparison. The clear implication for longitudinal analysis of candidate genes is that any effects that are found to differentiate individuals at one point in childhood are very likely to continue doing so over childhood and into adolescence.

Although weaker by comparison to genetic influences over time, nonshared environmental influences also contributed to some of the longitudinal stability of individual differences in externalizing problems. This was surprising, given that nonshared environmental variance typically has not been shown to contribute to stability in behavior problems in prior studies (Eley et al., 2003; van Beijsterveldt et al., 2003). The idea remains feasible in theory nonetheless—nonshared environmental influences could be stable in their effects over time. Candidate nonshared environment influences have been identified, including sibling differential parental treatment and classroom environments (Asbury et al., 2003; Mullineaux, Deter-Deckard, Petrill, & Thompson, 2009; Oliver et al., 2008). However, the field has not yet demonstrated that these nonshared influences persist either at the phenotypic or behavioral genetic level of analysis—an important gap in knowledge that should be addressed in future research.

Consistent with the data on externalizing problems, for attentive behavior there was a set of common genetic factors that explained stability of individual differences over the first two

annual waves (a finding originally reported in Deater-Deckard et al., 2006, when only two waves of data were available). This result is consistent with a previous twin study (Rietveld, Hudziak, Bartels, van Beijsterveldt, & Boomsma, 2004). In addition, two other longitudinal studies have implicated the dopamine receptor DRD4 candidate gene in explaining some of the temporally stable genetic influence on attention over middle childhood, for attention deficit symptoms in boys (El-Faddage, Laucht, Maras, Vohringer, & Schmidt, 2004) as well as typical variation in sustained attentive behavior in boys and girls (Deater-Deckard & Wang, 2012).

However, there also was a major distinction in the results for attention compared to externalizing problems, in that a substantial portion of the longitudinal stability of attentive behavior was accounted for by stable shared environment influence. This suggests that the nongenetic factors that contribute to sibling similarity in attentive behavior—above and beyond any similarity arising from genetic factors—are stable in their influences over this three-year period spanning middle childhood. Given that the informants (teachers and observers) were different at each of the three waves, and all twin pairs were rated independently by two different observers at each wave, it is unlikely that this shared environment effect on the stability of attentive behavior could be attributed to a method effect. Therefore, the overlapping shared environment factor that captured the developmental stability of individual differences in attention regulation was more likely to be genuine than reflecting artifacts of measurement. That aside, the results contradict the previous longitudinal study of attention which found that only genetic influences contributed to longitudinal stability (Rietveld et al., 2004). The differing results between the two studies could be due in part to differences in methods and informants; Rietveld et al. used parents' ratings, and in the current study we examined observers' and teachers' ratings.

The longitudinal analyses on the link between externalizing and attention produced results that were very similar to those gleaned from the cross-sectional analyses reported for Question 2. The phenotypic correlation between attention regulation and externalizing problems was accounted for by a single underlying genetic correlation in the 1st and 2nd waves, but through shared and nonshared environmental correlations at the 3rd wave. The emerging evidence from the current cross-sectional and longitudinal analyses points to an important transition in middle childhood that may stem from increased involvement in antisocial activities involving peers and siblings that are close in age (Farrington, 1995; Quinton, Pickles, Maughan, & Rutter, 1993; Rowe, Rodgers, & Meseck-Bushey, 1992). As children get older, these social experiences serve to reinforce growth in many aspects of aggressive and nonaggressive conduct problems in childhood and adolescence, and can reflect socialization and selection mechanisms that emerge as sources of shared environmental variance and longitudinal stability in behavioral genetic models (Gilson, Hunt, & Rowe, 2001). The current study's results contribute to this literature, by demonstrating that the impact of these shared environmental processes may become more important over middle childhood, and may operate on self-regulatory process involving attention.

Caveats, Future Directions, and Conclusion

In the current study, we extended the literature on the process linking externalizing problems and difficulties in attention regulation by examining their longitudinal genetic and nongenetic effects, as well as the potential moderating influence of household chaos. However, there are some limitations and new directions that need to be addressed in future research. First, we are confident that the sustained attentive behavior scores in the current study capture well the underlying attention regulation mechanisms, based on the temperament and ADHD literatures (Deater-Deckard & Wang, 2012; Rothbart, 2007; Willcutt et al., 2005), but we did not directly assess the specific components of attention (e.g., orienting, alerting, controlling) that function in cognitive self-regulation of behavior. We acknowledge that there remains some debate about the underlying structure and function of these various sub-components of attention and their behavioral manifestations (e.g., Watson, Kotov, & Gamez, 2006). Second, our reliance on parent reports of household chaos was not ideal. Although the brief CHAOS questionnaire has been shown to be a reliable and valid instrument (Coldwell et al., 2006; Deater-Deckard et al., 2009), it does not cover the full range of indicators of household chaos. In the future, structured interviews and observations should be used to complement parent report to assess household chaos more thoroughly and comprehensively. In addition, the same parent provided reports on externalizing problems and chaos across three waves, which may yield potential rater bias and artificially inflate the stability of each construct as well as the correlations between them. That being said, this design limitation probably has a minimal impact on estimation of statistical moderation and bivariate longitudinal effects, given that shared method variance was minimized in the links with attention which was measured using different informants. Third, a community twin sample was used in the current study. Essentially all of the children were below any clinical threshold for conduct problems. This potentially limits the generalizability of the current results to the clinical range of problem behaviors, or to families of non-twin children. Fourth, because the statistical effect explaining the link between externalizing problems and attention regulation shifted from genetic correlation to shared and nonshared environmental correlation over middle childhood, future studies should also examine whether environment x environment interaction exists. It also would be ideal to integrate moderation into longitudinal analyses. We were not able to do that, because the current study's sample size was underpowered for such an analysis. A related future direction would be to integrate candidate genes into these behavioral genetic models to examine similar questions. Fifth and finally, because the current study is the first one to examine the longitudinal transition of genetic and nongenetic association between externalizing problems and attention regulation, more studies are needed to test for replication of our results.

In conclusion, the developmental context for children's self-regulation and their behavioral problems is complex. This complex nature automatically calls for a collaborative effort from multiple domains, integrating multiple methods including behavior and molecular genetics, neurobiological and psychosocial approaches to fully understand the mechanisms through which one can optimize his development. Any study trying to explain the underlying etiology via a snapshot or a simple additive genetic and environmental story would not be conclusive. Despite the limitations, the current study addressed several shortcomings in

previous studies and investigated this intricate link by examining gene x environment interplay mechanism from a developmental perspective. Results suggested that these endeavors are necessary and still much needed.

Acknowledgments

We thank the study participants and research staff. This research was supported by NICHD grants HD38075.

References

- Achenbach, TM. Manual for the Teacher's Report Form and 1991 Profile. Burlington, VT: University of Vermont, Department of Psychiatry; 1991.
- Asbury K, Dunn JF, Pike A, Plomin R. Nonshared environmental influences on individual differences in early behavioral development: A monozygotic twin differences study. *Child Development*. 2003; 74(3):933–943. [PubMed: 12795399]
- Bartels M, van Beijsterveldt CEM, Derks EM, Stroet TM, Polderman TJC, Hudziak JJ, Boomsma DI. Young Netherlands Twin Register (Y-NTR): A longitudinal multiple informant study of problem behavior. *Twin Research and Human Genetics*. 2007; 10(1):3–11. [PubMed: 17539360]
- Bayley, N. Bayley Scales of Infant Development. New York: Psychological Corp; 1969.
- Belsky J, Fearon RMP, Bell B. Parenting, attention and externalizing problems: Testing mediation longitudinally, repeatedly and reciprocally. *Journal of Child Psychology and Psychiatry*. 2007; 48(12):1233–1242. [PubMed: 18093029]
- Berry D, Deater-Deckard K, McCartney K, Wang Z, Petrill SA. Gene–environment interaction between DRD4 7-repeat VNTR and early maternal sensitivity predicts inattention trajectories across middle childhood. *Development & Psychopathology*. 2013; 25(2):291–306. [PubMed: 23627945]
- Blair K, Smith B, Mitchell D, Morton J, Vythilingam R, Pessoa L, Blair R. Modulation of emotion by cognition and cognition by emotion. *Neuroimage*. 2007; 35:430–440. [PubMed: 17239620]
- Brookes K, Mill J, Guindalini C, Curran S, Xu X, Knight J, Asherson P. A common haplotype of the dopamine transporter gene associated with attention-deficit/hyperactivity disorder and interacting with maternal use of alcohol during pregnancy. *Archives of General Psychiatry*. 2006; 63:74–81. [PubMed: 16389200]
- Burt SA, Krueger RF, McGue M, Iacono WG. Sources of covariation among attention-deficit/hyperactivity disorder, oppositional defiant disorder, and conduct disorder: The importance of shared environment. *Journal of Abnormal Psychology*. 2001; 110:516–525. [PubMed: 11727941]
- Burt SA, McGue M, Krueger RF, Iacono WG. Sources of covariation among the child-externalizing disorders: Informant effects and the shared environment. *Psychological Medicine*. 2005; 35(8): 1133–1144. [PubMed: 16116939]
- Cadoret RJ, Cain CA, Crowe RR. Evidence for gene-environment interaction in the development of adolescent antisocial behavior. *Behavior Genetics*. 1983; 13(3):301–310. [PubMed: 6615382]
- Cicchetti, D.; Valentino, K. Toward the application of a multiple-levels-of-analysis perspective to research in development and psychopathology. In: Masten, AS., editor. *Multilevel dynamics in developmental psychopathology: Pathways to the future*. New York: Taylor & Francis/Erlbaum; 2007. p. 243–284.
- Coldwell J, Pike A, Dunn J. Household chaos-links with parenting and child behaviour. *Journal of Child Psychology and Psychiatry*. 2006; 47(11):1116–1122. [PubMed: 17076750]
- Deater-Deckard K, Mullineaux PY, Beekman C, Petrill SA, Schatschneider C, Thompson L. Conduct problems, IQ, and household chaos: A longitudinal multi-informant study. *Journal of Child Psychology and Psychiatry*. 2009; 50(10):1301–1308. [PubMed: 19527431]
- Deater-Deckard K, Petrill SA, Thompson LA. Anger/frustration, task persistence, and conduct problems in childhood: A behavioral genetic analysis. *Journal of Child Psychology and Psychiatry*. 2007; 48(1):80–87. [PubMed: 17244273]
- Deater-Deckard K, Petrill SA, Thompson LA, DeThorne LS. A longitudinal behavioral genetic analysis of task persistence. *Developmental Science*. 2006; 9:498–504. [PubMed: 16911452]

- Deater-Deckard, K.; Wang, Z. Development of temperament and attention: Behavioral genetic approaches. In: Posner, M., editor. *Cognitive neuroscience of attention*. 2. New York: Guildford; 2012. p. 331-342.
- Dick DM, Viken RJ, Kaprio J, Pulkkinen L, Rose RJ. Understanding the covariation among childhood externalizing symptoms: Genetic and environmental influences on conduct disorder, attention deficit hyperactivity disorder, and oppositional defiant disorder symptoms. *Journal of Abnormal Child Psychology*. 2005; 33(2):219–229. [PubMed: 15839499]
- Dick, D.; York, T. Gene-environment interaction & correlation. 2010. [PowerPoint slides]. Retrieved from <http://www.vipbg.vcu.edu/vipbg/HGEN619/GE%20presentation%2010-19-10.pdf>
- Dumas JE, Nissley J, Nordstrom A, Smith EP, Prinz RJ, Levine DW. Home chaos: Sociodemographic, parenting, interactional, and child correlates. *Journal of Clinical Child and Adolescent Psychology*. 2005; 34(1):93–104. [PubMed: 15677284]
- Eisenberg N, Spinrad TL, Fabes RA, Reiser M, Cumberland A, Shepard SA, Murphy B. The relations of effortful control and impulsivity to children's resiliency and adjustment. *Child Development*. 2004; 75(1):25–46. [PubMed: 15015673]
- Eisenberg N, Zhou Q, Spinrad TL, Valiente C, Fabes RA, Liew J. Relations among positive parenting, children's effortful control, and externalizing problems: A three-wave longitudinal study. *Child Development*. 2005; 76(5):1055–1071. [PubMed: 16150002]
- Eley TC, Lichtenstein P, Moffitt TE. A longitudinal behavioral genetic analysis of the etiology of aggressive and nonaggressive antisocial behavior. *Development and Psychopathology*. 2003; 15:383–402. [PubMed: 12931834]
- El-Faddagh M, Laucht M, Maras A, Vohringer L, Schmidt MH. Association of dopamine D4 receptor (DRD4) gene with attention-deficit /hyperactivity disorder (ADHD) in a high-risk community sample: A longitudinal study from birth to 11 years of age. *Journal of Neural Transmission*. 2004; 111:883–889. [PubMed: 15206004]
- Enders CK, Bandalos DL. The relative performance of full information maximum likelihood estimation for missing data in structural equation models. *Structural Equation Modeling: A Multidisciplinary Journal*. 2001; 8(3):430–457.
- Erickson K, Drevets W, Schulkin J. Glucocorticoid regulation of diverse cognitive functions in normal and pathological emotional states. *Neuroscience and Biobehavioral Review*. 2003; 27:233–246.
- Evans GW, Gonnella C, Marcynyszyn LA, Gentile L, Salpekar N. The role of chaos in poverty and children's socioemotional adjustment. *Psychological Science*. 2005; 16(7):560–565. [PubMed: 16008790]
- Evans GW, Hygge S, Bullinger M. Chronic noise and psychological stress. *Psychological Science*. 1995; 6(6):333–338.
- Evans, GW.; Wachs, TD. *Chaos and its influence on children's development: An ecological perspective*. Washington, DC: American Psychological Association; 2009.
- Fan J, Wu Y, Fossella JA, Posner MI. Assessing the heritability of attention networks. *BMC Neuroscience*. 2001; 2:14. [PubMed: 11580865]
- Farrington DP. The development of offending and antisocial behaviour from childhood: Key findings from the Cambridge Study in Delinquent Development. *Journal of Child Psychology and Psychiatry*. 1995; 360:929–964. [PubMed: 7593403]
- Ghodsian-Carpey J, Baker LA. Genetic and environmental influences on aggression in 4 to 7-year old twins. *Aggressive Behavior*. 1987; 13:173–186.
- Gilson MS, Hunt CB, Rowe DC. The friends of siblings: A test of social homogamy vs. peer selection and influence. *Marriage & Family Review*. 2001; 33:205–223.
- Griffin D, Gonzalez R. Correlational analysis of dyad-level data in the exchangeable case. *Psychological Bulletin*. 1995; 118(3):430–439.
- Holmes J, Payton A, Barrett J, Harrington R, McGuffin P, Owen M, Thapar A. Associations of DRD4 in children with ADHD and comorbid conduct problems. *American Journal of Medical Genetics*. 2002; 114B:150–153. [PubMed: 11857575]
- Kenny DA. Models of non-independence in dyadic research. *Journal of Social and Personal Relationships*. 1996; 13:279–294.

- Kim J, Deater-Deckard K. Dynamic changes in anger, externalizing and internalizing problems: Attention and regulation. *Journal of Child Psychology and Psychiatry*. 2011; 52(2):156–166. [PubMed: 20825522]
- Kochanska G, Knaack A. Effortful control as a personality characteristic of young children: Antecedents, correlates, and consequences. *Journal of Personality*. 2003; 71(6):1087–1112. [PubMed: 14633059]
- Langley K, Turic D, Rice F, Holmans P, van den Bree MBM, Craddock N, Thapar A. Testing for gene x environment interaction effects in attention deficit hyperactivity disorder and associated antisocial behavior. *American Journal of Medical Genetics*. 2008; 147B:49–53. [PubMed: 17579368]
- Leve LD, Kerr DC, Shaw D, Ge X, Neiderhiser JM, Scaramella LV, Reiss D. Infant pathways to externalizing behavior: Evidence of genotype X environment interaction. *Child Development*. 2010; 81(1):340–356. [PubMed: 20331671]
- Leve LD, Winebarger AA, Fagot BI, Reid JB, Goldsmith HH. Environmental and genetic variance in children's observed and reported maladaptive behavior. *Child Development*. 1998; 69:1286–1298. [PubMed: 9839416]
- Macrae CN, Bodenhausen GV. Social cognition: Thinking categorically about others. *Annual Review of Psychology*. 2000; 51:93–120.
- Matheny AP, Wachs TD, Ludwig JL, Phillips K. Bringing order out of chaos: Psychometric characteristics of the Confusion, Hubbub, and Order Scale. *Journal of Applied Developmental Psychology*. 1995; 16:429–444.
- Mednick SA, Gabrielli WF, Hutchings B. Genetic factors in criminal behavior: Evidence from an adoption cohort. *Science*. 1984; 224:891–893. [PubMed: 6719119]
- Miles DR, Carey G. Genetic and environmental architecture of human aggression. *Journal of Personality and Social Psychology*. 1997; 7:207–217. [PubMed: 9008382]
- Morris AS, Silk JS, Steinberg L, Terranova AM, Kithakye M. Concurrent and longitudinal links between children's externalizing behavior in school and observed anger regulation in the mother-child dyad. *Journal of Psychopathology and Behavioral Assessment*. 2010; 32:48–56.
- Mullineaux PY, Deater-Deckard K, Petrill SA, Thompson L. Parenting and child behaviour problems: A longitudinal analysis of non-shared environment. *Infant and Child Development*. 2009; 18(2): 133–148. [PubMed: 19436771]
- Murray KT, Kochanska G. Effortful control: Factor structure and relation to externalizing and internalizing behaviors. *Journal of Abnormal Child Psychology*. 2002; 30(5):503–514. [PubMed: 12403153]
- Nadder TS, Rutter M, Silberg JL, Maes HH, Eaves LJ. Genetic effects on the variation and covariation of attention deficit-hyperactivity disorder (ADHD) and oppositional defiant disorder/conduct disorder (ODD/CD) symptomatologies across informant and occasion of measurement. *Psychological Medicine*. 2002; 32:39–53. [PubMed: 11883729]
- Neale, M. Mx: Statistical modeling. 4. Richmond, VA: Department of Psychiatry; 1997.
- Oliver BR, Pike A, Plomin R. Nonshared environmental influences on teacher-reported behaviour problems: Monozygotic twin differences in perceptions of the classroom. *Journal of Child Psychology and Psychiatry*. 2008; 49(6):646–653. [PubMed: 18355217]
- Olsen JA, Kenny DA. Structural equation modeling with interchangeable dyads. *Psychological Methods*. 2006; 11(2):127–141. [PubMed: 16784334]
- Olson SL, Sameroff AJ, Kerr DCR, Lopez NL, Wellman HM. Developmental foundations of externalizing problems in young children: The role of effortful control. *Developmental and Psychopathology*. 2005; 17:25–45.
- Pike A, Iervolino AC, Eley TC, Price TS, Plomin R. Environmental risk and young children's cognitive and behavioral development. *International Journal of Behavioral Development*. 2006; 30(1):55–66.
- Pitzer M, Esser G, Schmidt MH, Laucht M. Temperamental predictors of externalizing problems among boys and girls: A longitudinal study in a high-risk sample from ages 3 months to 15 years. *European Archives of Psychiatry and Clinical Neuroscience*. 2009; 259:445–458. [PubMed: 19387718]

- Plomin, R.; DeFries, J.C.; McClearn, G.E.; McGuffin, P. Behavioral genetics. 5. New York: Worth Publishers; 2008.
- Posner, M.I.; Rothbart, M.K. Educating the human brain. Washington, DC: American Psychological Association; 2006.
- Prior M, Smart D, Sanson A, Oberklaid F. Longitudinal predictors of behavioural adjustment in pre-adolescent children. Australian and New Zealand Journal of Psychiatry. 2001; 35:297–307. [PubMed: 11437802]
- Purcell S. Variance components models for gene-environment interaction in twin analysis. Twin Research and Human Genetics. 2002; 5(6):554–571.
- Quinton D, Pickles A, Maughan B, Rutter M. Partners, peers, and pathways: Assortative pairing and continuities in conduct disorder. Development and Psychopathology. 1993; 5:763–783.
- Raine A. The role of prefrontal deficits, low autonomic arousal, and early health factors in the development of antisocial and aggressive behavior in children. Journal of Child Psychology and Psychiatry. 2002; 43(4):417–434. [PubMed: 12030589]
- Rhee SH, Waldman ID. Genetic and environmental influences on antisocial behavior: A meta-analysis of twin and adoption studies. Psychological Bulletin. 2002; 128:490–529. [PubMed: 12002699]
- Rietveld MJH, Hudziak JJ, Bartels M, van Beijsterveldt CEM, Boomsma DI. Heritability of attention problems in children: Longitudinal results from a study of twins, age 3 to 12. Journal of Child Psychology and Psychiatry. 2004; 45(3):577–588. [PubMed: 15055376]
- Rothbart M.K. Temperament, development, and personality. Current Directions in Psychological Science. 2007; 16(4):207–212.
- Rothbart M.K., Ellis L.K., Rueda M.R., Posner M.I. Developing mechanisms of temperamental effortful control. Journal of Personality. 2003; 71:1113–1143. [PubMed: 14633060]
- Rowe DC, Rodgers JL, Meseck-Bushey S. Sibling delinquency and the family environment: Shared and unshared influences. Child Development. 1992; 63:59–67. [PubMed: 1551330]
- Schmitz S, Cherny SS, Fulker DW. Increase in power through multivariate analyses. Behavior Genetics. 1998; 28(5):357–363. [PubMed: 9926617]
- Thapar A, Langley K, Fowler T, Rice F, Turic D, Whittinger N, O'Donovan M. Catechol O-Methyltransferase gene variant and birth weight predict early-onset antisocial behavior in children with attention-deficit/hyperactivity disorder. Archives of General Psychiatry. 2005; 62:1275–1278. [PubMed: 16275815]
- van Beijsterveldt CEM, Bartels M, Hudziak JJ, Boomsma DI. Causes of stability of aggression from early childhood to adolescence: A longitudinal genetic analysis in Dutch Twins. Behavior Genetics. 2003; 33(5):591–605. [PubMed: 14574135]
- Watson, D.; Kotov, R.; Gamez, W. Basic dimensions of temperament in relation to personality and psychopathology. In: Krueger, R.F.; Tackett, J.L., editors. Personality and Psychopathology. New York: Guilford Press; 2006. p. 7-38.
- Willcutt EG, Doyle AE, Nigg JT, Faraone SV, Pennington BF. Validity of the executive function theory of attention-deficit/hyperactivity disorder: A meta-analytic review. Biological Psychiatry. 2005; 57:1336–1346. [PubMed: 15950006]
- Zhou Q, Hofer C, Eisenberg N, Reiser M, Spinrad TL, Fabes RA. The developmental trajectories of attention focusing, attention and behavioral persistence, and externalizing problems during school-age years. Developmental Psychology. 2007; 43:369–385. [PubMed: 17352545]

Figure 1.

Actor-Partner Interdependence Model (APIM): Additive and Interactive Effects of Attention Regulation and Chaos on Externalizing Problems. APIM was used to test for additive and interactive effects of attention regulation and household chaos on externalizing problems phenotypically. C: correlation. B: regression coefficient. M: mean. V: variance. E: error variance.

Figure 2.

Bivariate ACE Cholesky Decomposition: with (right) and without (left) Chaos as a Moderator. Bivariate ACE Cholesky decomposition model was used to partition the variance of and covariance between attention regulation and externalizing problems and to test for potential moderation effect of household chaos on genetic variance of and covariance between attention regulation and externalizing problems at each wave. The left side shows the basic bivariate Cholesky decomposition, where latent variables A, C, and E represent overlapping additive genetic, shared environmental, and nonshared environmental covariance between attention regulation and externalizing problems, respectively. Latent variables a, c, and e represent residual additive genetic, shared environmental, and nonshared environmental variance of externalizing problems. The right side shows the bivariate Cholesky decomposition with a continuous moderator (i.e., household chaos), where the additive genetic variance and covariance is a function of the moderator $A_{total} = A' + \beta M$. The above figure was only shown for one twin in the pair.

Figure 3.

Longitudinal common-factor independent-pathway model. Longitudinal common-factor independent-pathway model was used to partition the variance of and covariance between attention regulation and externalizing problems across three waves. Latent variables A1, C1, and E1 represent overlapping additive genetic, shared environmental, and nonshared environmental covariance, respectively, between attention regulation and externalizing problems across all three waves. A2, C2, and E2 represent additive genetic, shared environmental, and nonshared environmental covariance, respectively, between attention regulation and externalizing problems shared across the 2nd and 3rd waves. A3, C3, and E3 represent the additive genetic, shared environmental, and nonshared environmental covariance, respectively, between attention regulation and externalizing problems only at the 3rd wave that is not overlapped with the 1st and 2nd waves. Residual genetic, shared environmental, and nonshared environmental variance unique to each variable at each time point is represented by a, c, and e, respectively. The above figure was only shown for one twin in the pair. Ext: externalizing problems. Att: attention regulation.

Table 1
Descriptive Statistics and Correlations between Attention Regulation, Externalizing Problems, and Chaos

	1 st att	2 nd att	3 rd att	1 st ext	2 nd ext	3 rd ext	1 st chaos	2 nd chaos	3 rd chaos
1 st att	.00(1.00)								
2 nd att	.47***	.00(1.00)							
3 rd att	.35***	.44***	.00(1.00)						
1 st ext	.18***	.16***	.14**	7.18 (5.54)					
2 nd ext	.12*	.14**	.16**	.71***	6.46 (5.23)				
3 rd ext	.20***	.20***	.23***	.68***	.76***	6.29 (5.51)			
1 st chaos	.14**	.11*	.05	.28***	.20***	.21***	2.38(.61)		
2 nd chaos	.13*	.17***	.15**	.32***	.32***	.29***	.76***	2.32(.58)	
3 rd chaos	.17**	.21***	.17**	.28***	.25***	.29***	.72***	.80***	2.32(.64)

Note. Numbers on the diagonal are shown as means (standard deviations) of study variables; numbers off diagonal are correlations between study variables. att = attention regulation; ext = externalizing problems. N ranged from 413 to 600.

* $p < .05$;
** $p < .01$;
*** $p < .001$.

Table 2
Actor-Partner Interdependence Model (APIM): Standardized Path Estimates from the Full Models and Model Fit Indices

	b1	b2	b3	Full model (χ^2, df)	No attention main effect (χ^2, df)	No chaos main effect (χ^2, df)	No interactive effect (χ^2, df)
wave 1	.14***	.25***	-.06 ⁺	32.19, 4	51.58, 6	83.21, 6	35.97, 5
wave 2	.07**	.29***	-.09**	54.10, 4	65.21, 6	122.56, 6	62.54, 5
wave 3	.16***	.25***	.04	35.10, 4	67.65, 6	72.06, 6	36.66, 5

Note. A significant path estimate suggested that fixing that path at 0 resulted in a significant decrease in model fit compared to the full model. b1: main effect of attention regulation on externalizing problems; b2: main effect of chaos on externalizing problems; b3: interactive effect between attention regulation and chaos on externalizing problems. N = 608.

⁺ $p = .052$;
* $p < .05$;
** $p < .01$;
*** $p < .001$.

Table 3

Bivariate Cholesky Decomposition: Fit Indices and Model Comparisons between Full Models and Constraint Models

	-2 log likelihood	df	-2 log likelihood	df
1 st wave				
full model	2865.42	1094	-	-
no $\beta_{\text{attention}}$	2885.42	1095	20.00***	1
no $\beta_{\text{externalizing}}$	2878.33	1095	12.91***	1
no β_{cov}	2870.42	1095	4.91*	1
2 nd wave				
full model	2664.19	1006	-	-
no $\beta_{\text{attention}}$	2670.98	1007	6.79**	1
no $\beta_{\text{externalizing}}$	2688.04	1007	23.85***	1
no β_{cov}	2666.26	1007	2.07	1
3 rd wave				
full model	2237.92	865	-	-
no $\beta_{\text{attention}}$	2257.34	866	19.42***	1
no $\beta_{\text{externalizing}}$	2249.18	866	11.26***	1
no β_{cov}	2240.20	866	2.28	1

Note. A significant -2 log likelihood suggested that fixing a particular path at 0 in that model resulted in a significant decrease in model fit compared to the full model. $\beta_{\text{attention}}$: the coefficient of the moderating effect of chaos on total genetic variance in attention regulation; $\beta_{\text{externalizing}}$: the coefficient of the moderating effect of chaos on unique genetic variance in externalizing problems; β_{cov} : the coefficient of the moderating effect of chaos on genetic covariance between attention regulation and externalizing problems.

* $p < .05$;

** $p < .01$;

*** $p < .001$.

Bivariate Cholesky Decomposition: Unstandardized Variance Estimates, Genetic and Nongenetic Correlations between Attention Regulation and Externalizing Problems at Three Different Levels of Chaos from the Full Model

Table 4

	A _{attention}	C _{attention}	E _{attention}	A _{externalizing}	C _{externalizing}	E _{externalizing}	R _A	R _C	R _E
1 st wave									
–1SD	.10	.15	.44	.47	.12	.22	.76	.02	.06
Mean	.31			.58			.45		
+1SD	.63			.87			.17		
2 nd wave									
–1SD	.28	.08	.44	.30	.12	.24	.47	.56	.00
Mean	.46			.57			.18		
+1SD	.67			1.01			.02		
3 rd wave									
–1SD	.05	.15	.43	.49	.11	.14	.00 ^a	.84	.32
Mean	.32			.77			.01		
+1SD	.81			1.18			.17		

Note. A: genetic variance. C: shared environmental variance. E: nonshared environmental variance. R_A: genetic correlation. R_C: shared environmental correlation. R_E: nonshared environmental correlation. –1SD and +1SD: moderator = 1 standard deviation below and above the mean of chaos, respectively. Mean: moderator = the mean of chaos.

^a we fixed this particular genetic correlation at 0 because computational result suggests a negative value which makes no theoretical sense.

Table 5

Longitudinal Common-Factor Independent-Pathway Model: Standardized Path Estimates from the Best-Fitting Model

	A1	A2	A3	C1	C2	C3	E1	E2	E3	a	c	e
1 st wave Attention	.18*			.56*			-			.37	-	.72*
1 st wave Externalizing	.88*			-			.34*			-	-	.33*
2 nd wave Attention	.19*	-		.66*	-		-	.38*		-	-	.62
2 nd wave Externalizing	.66*	.45*		-	-		.34*	-		.35	-	.35*
3 rd wave Attention	-	-	-	.54*	-	-	.34*	.27*	-	.45	-	.57
3 rd wave Externalizing	.64*	.52*	-	.09*	-	-	.30*	-	-	.40	-	.26*

Note. A1, C1, and E1 are common genetic, shared environmental, and nonshared environmental pathways between attention regulation and externalizing problems across three waves; A2, C2, and E2 are common genetic, shared environmental, and nonshared environmental pathways between attention regulation and externalizing problems across the 2nd and 3rd waves; A3, C3, and E3 are common genetic, shared environmental, and nonshared environmental pathways between attention regulation and externalizing problems at the 3rd wave. a, c, and e are residual genetic, shared environmental, and nonshared environmental pathways unique to attention regulation and externalizing problems at each wave. - = path fixed to 0.

* $p < .05$.