CORRESPONDENCE

Persistent Vegetative State and Minimally Conscious State: A systematic review and meta-analysis of diagnostic procedures

by PD Dr. med. Andreas Bender et al. in issue 14/2015

Additions

The diagnosis and differential diagnosis of (persistent) vegetative state, (P)VS, requires special expertise and experience (1). Recently, the term (P)VS has been substituted by the medical term *Unresponsive Wakefulness Syndrome* (UWS), as correctly mentioned by the authors (2). Our aim should be to reduce the unacceptably high rate of misdiagnosis in patients with (P)VS (37–43%) by improving quality management. One can only agree with the statement that above all the diagnosis of VS (UWS) is based on a qualified and standardized clinical neurological examination. Semantically, VS (UWS) and minimally conscious state (MCS) denote two functional transitory syndromes which can be clearly differentiated based on clinical findings. The reliability and validity of the German Coma Remission Scale (KRS) in identifying coma, PVS and MCS in early neurological rehabilitation has convinced specialists, health insurers and politicians (German Social Insurance Code (SGB) IX). With regard to the studies analyzed in the article, we had wished for a more adequate critical discussion on the evaluation and evidence of the way the respective neurological examination procedures were applied and what this meant for the rate of misdiagnosis. Have our recommendations of the European VS guidelines been followed in those studies (1)? Our recommendations have not been discussed in the review (2). Evidence-based, bed-side examination techniques were not mentioned. We recommended a 3-year further training program in a specialized department for VS patients. This qualification has been shown to provide medical-neurological expertise and to reduce the rate of misdiagnosis of VS by precisely allocating the typical symptoms observed with this transitory functional syndrome to the correct diagnosis. Supplementary tables on etiology, the clinical picture of full-blown VS (UWS) and clinical dynamics during the stages of regression are missing (3); these have been compiled with modifications in our guidelines (1) and continue to be valid. In UWS and MCS, not only neurotraumatologists are especially interested in the prognostic relevance of the cause, location and extent of the underlying brain damage and its functional changes over time, apart from patient age. Additional information about functional imaging is provided by Zakharova et al. (4).

DOI: 10.3238/arztebl.2015.0680a

REFERENCES

1. von Wild, K, Gerstenbrand F, Dolce G, et al.: Guidelines for quality management of apallic syndrome/vegetative state. *Eur J Trauma Emerg Surg* 2007; 3: 268–92.
2. Bender A, Jox RJ, Grill E, Straube A, Lulé D: Persistent vegetative state and minimally conscious state—a systematic review and meta-analysis of diagnostic procedures. *Dtsch Arztebl Int* 2015; 112: 235–42.
3. Gerstenbrand F: The symptomatology of the apallic syndrome. In: Dalle Ore G, Gerstenbrand F (eds.): *The apallic syndrome*. Heidelberg, New York 1977: 14–21.
4. Zakharova N, Kornienko V, Potapov A, Pronin I: *Neuroimaging of traumatic brain injury*. Cham, Heidelberg, New York, Dordrecht, London: Springer 2014: 69–123.

Prof. Prof. h.c. mult. Dr. med. Dr. h.c. Klaus R.H. von Wild
Neurochirurgie, Neurorehabilitation, Medizinische Fakultät der WW-Universität Münster,
kvw@neurosci.de

Prof. emer. Dr. Dr. hc. mult. Franz Gerstenbrand
Neurologie und Psychiatrie, Wien

Prof. Alexander Potapov, MD, PhD
Neurosurgery
The BURDENKO Neurosurgery Institute,
Moscow RUSSIA

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We like to thank our colleagues who have played a crucial role in shaping our understanding of PVS over the past decades for their important additions. Our focus was on the systematic meta-analysis of studies describing technical methods that help to differentiate between PVS and MCS (1). Unfortunately, we could not discuss all aspects and all important preparatory work related to this complex topic—not least because of the limited space available. We share the view that the German Coma Remission Scale (KRS) is an important instrument for evaluating the clinical course of patients with disturbances of consciousness. The use of an appropriate clinical examination method to differentiate between PVS and MCS was one key inclusion criterion for our analysis. The majority of the studies we identified were from non-German-speaking countries and used the Coma Recovery Scale-Revised (CRS-R) as an evidence-based, bed-side examination method with a good test quality profile and clearly operationalized differentiation between PVS and MCS (2). However, the use of CRS-R was by no means compulsory. It is crucial to use internationally established assessment tools to advance this area of research in Germany, too; in fact, this approach has already been adopted by German-speaking working groups (3). We have critically discussed (1) that the use of the CRS-R is associated with significant methodological risks and evaluated the quality of the studies especially based on the clinical examination method used (eTable in the article). To reduce the rate of clinical misdiagnosis, we firmly support the expert group's call for improved advanced training and continuing education for physicians in the assessment of disturbances of consciousness. Major neurological textbooks should have a special chapter exclusively dedicated to this topic. It should be ensured that seeing and managing this type of patients in a clinical setting is an integral part of specialist training.

DOI: 10.3238/arztebl.2015.0680b

REFERENCES

1. Bender A, Jox RJ, Grill E, Straube A, Lulé D: Persistent vegetative state and minimally conscious state—a systematic review and meta-analysis of diagnostic procedures. *Dtsch Arztebl Int* 2015; 112: 235–42.
2. La Porta F, Caselli S, Ianes AB, et al.: Can we scientifically and reliably measure the level of consciousness in vegetative and minimally conscious States? Rasch analysis of the coma recovery scale-revised. *Arch Phys Med Rehabil* 2013; 94: 527–35.
3. Kotchoubey B, Yu T, Mueller F, Vogel D, Vesper S, Lang S: True or false? Activations of language-related areas in patients with disorders of consciousness. *Curr Pharm Des* 2014; 20: 4239–47.

On behalf of the authors:

Prof. Dr. med. Andreas Bender
Therapiezentrum Burgau
a.bender@therapiezentrum-burgau.de

Conflict of interest statement

PD Dr. Bender received an honorarium from the Covidien company for lecturing at a stroke symposium.