CORRESPONDENCE

The Differential Diagnosis and Interdisciplinary Treatment of Anal Carcinoma

by Dr. Dimitrios Raptis, Prof. Dr. med. Ignaz Schneider,
Prof. Dr. med. Klaus E. Matzel, Prof. Dr. med. Dr. h.c. Werner Hohenberger,
PD Dr. med. Oliver Ott, Prof. Dr. med. Rainer Fietkau in issue 14/2015

Screening Patients at High Risk

As colorectal surgeons we read this important article with great interest (1). The authors correctly report that the incidence of anal cancer rises in HIV positive patients. The increased incidence in patients who are medically immunocompromised (for example, transplant patients) is not mentioned in the text in addition to other risk factors. The authors write: "There are no international, globally accepted, guidelines for screening in high-risk groups." In September 2013 the Association of the Scientific Medical Societies in Germany (AWMF) published a "German-Austrian" guideline (2), in which screening in this high-risk group (HIV positive patients) was described. In our opinion, this guideline is the only German-language guideline that deals with human papillomaviruses (HPV), anal dysplasias, and anal carcinomas. We are not aware of any general German guideline for anal cancers.

With regard to surgical treatment, the authors report that only T1 cancers of the anal margin should be excised. The guideline of the American Society of Colon and Rectal Surgeons (ASCRS), however, refers only to "small lesions," without providing any information on tumor size or safety margin. In the AWMF guideline mentioned earlier, local excision of T1 and T2 tumors is an option in HIV patients. For dermatologists, R0 excision of squamous cell carcinomas measuring 3 cm (=T2) outside the 5 cm anal margin is entirely justified (3). For this reason it is confusing if the authors of the present article recommend automatic radiochemotherapy for such T2 tumors within the anal margin (3).

The authors report that during follow-up care, incision biopsy is not permitted if relapse is suspected, "as the risk of incontinence or severely delayed wound healing is extremely high." Radiochemotherapy has follow-on effects over time, and confirmation of a recurrence by means of a biopsy should be avoided in the first three months. In suspected superficial local recurrence, we think that a small superficial incision biopsy is possible without damaging the anal sphincter.

DOI: 10.3238/arztebl.2015.0738a

REFERENCES

1. Raptis D, Schneider I, Matzel KE, Ott O, Fietkau R, Hohenberger W: The differential diagnosis and interdisciplinary treatment of anal carcinoma. *Dtsch Arztebl Int* 2015; 112: 243–9.

2. Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften: S2k-Leitlinie: Anale Dysplasien und Analkarzinome bei HIV-Infizierten: Prävention, Diagnostik und Therapie. www.awmf.org/leitlinien/detail/II/055-007.html (last accessed on 10 April 2015).

3. Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften: S2-Leitlinie: Kurzleitlinie – Plattenepithelkarzinom der Haut. www.awmf.org/leitlinien/detail/II/032-022.html (last accessed on 11 April 2015).

Dr. med. Johannes Jongen

Jonas Schumacher

Prof. Dr. med. Volker Kahlke

Proktologische Praxis Kiel, Abteilung Proktologische Chirurgie,
Park-Klinik Kiel
j.jongen@kielnet.net

Conflict of interest statement

The authors declare that no conflict of interest exists.

Unmentioned Side Effects

The review article (1) does not mention a crucial aspect of side effects associated with treatment, and in our own clinical experience, this topic is often not broached with women with anal carcinoma: sexual dysfunction after pelvic radiotherapy.

Obviously, sexual dysfunction is always a multi-causal and multidimensional problem, but radiochemotherapy of the pelvis is an unambiguous risk factor for the development of physiological and anatomical exchanges to the genital area, which may materially affect sexual functioning even years after the treatment was administered.

It has been well known for 60 years, from retrospective (2) as well as prospective (3) data collections, that radiotherapy of cervical cancers leads to sexual dysfunction. About 50% of women thus affected are no longer sexually active (in contrast to 26% in the control group of healthy women). The most common symptoms were dyspareunia, stenosis of the vaginal canal, a tendency to bleed, anorgasmia, and vaginal dryness. Clinically, the same sequelae have also been observed in anal cancer.

The LENT SOMA Score (Late Effects of Normal Tissues; SOMA = Subjective, Objective, Management and Analytic Categories) (4) is a suitable instrument in persons with sexual dysfunction for determining symptoms before and after treatment and to determine the therapeutic need. Unfortunately this is done too rarely in routine clinical practice. We can only assume that feelings of shame on the part of both patients and doctors is responsible for this omission.

Informing women about the regular use of vaginal dilators, as early as possible and also, if possible, during treatment, but at the latest after the inflammation has subsided results in a notable decrease in sexual dysfunction.

DOI: 10.3238/arztebl.2015.0738b

REFERENCES

1. Raptis D, Schneider I, Matzel KE, Ott O, Fietkau R, Hohenberger W: The differential diagnosis and interdisciplinary treatment of anal carcinoma. *Dtsch Arztebl Int* 2015; 112: 243–9.
2. Andersen BL, Van der Does J: Surviving gynecologic cancer and coping with sexual morbidity: an international problem. *Int J Gynecol Cancer* 1994; 4: 225–40.
3. Bergmark K, Avall-Lundquist E, Dickman PW, et al.: Vaginal changes and sexuality in women with a history of cervical cancer. *N Engl J Med* 1999; 340: 1383–9.
4. Seegenschmiedt MH (ed.): Nebenwirkungen in der Onkologie. Internationale Systematik und Dokumentation. Berlin, Heidelberg: Springer 1998.

PD Dr. med. Carsten Ziske

Praxisnetzwerk Hämatologie/Internistische Onkologie,
Onkologie St. Josef Hospital Troisdorf
ziske@onkologie-rheinsieg.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Prevention of Anal Carcinoma

The article by Raptis and colleagues provides an excellent overview on the current state of the diagnostic evaluation and therapy of anal carcinoma (1). We wish to comment on three particular aspects in the article.

Similar to cervical cancer, anal cancer is potentially preventable by means of screening examinations. The authors mention that so far, no internationally accepted guidelines have been issued on the screening of high risk groups. Although that is correct, a German-Austrian guideline has existed since 2013, which was published by the Association of the Scientific Medical Societies in Germany (AWMF), on the prevention, diagnostic evaluation, and therapy of anal dysplasias and carcinomas in HIV-infected individuals (2). This guideline recommends, in addition to clinical inspection of the anogenital region and digital rectal examination, anal cytology testing and, in case of a pathological findings or in patients at particularly high risk, they additionally recommend high-resolution anoscopy with biopsy, in analogy to colposcopy.

The authors mention that so far, no recommendations exist with regard to vaccination against anal cancers. A large controlled study showed the efficacy of the prophylactic quadrivalent HIV vaccine in preventing anal dysplasias in men who have sex with men (MSM). For this reason, as well as for the purpose of preventing genital warts, vaccinating MSM is explicitly recommended in the United Kingdom, since MSM do not benefit from the herd immunity conferred by vaccinating girls (3).

The authors say that according to US and European guidelines, primarily surgical treatment is indicated only in well differentiated cancers of the anal margin in stage T1. A recently published study showed, however, that T1 cancers localized in the anal canal can also be treated curatively by using excision and a safety resection margin <5 mm (4). It was thus possible to avoid radiotherapy, as is otherwise recommended in early invasive cancers of the anal canal. DOI: 10.3238/arztebl.2015.0739a

REFERENCES

1. Raptis D, Schneider I, Matzel KE, Ott O, Fietkau R, Hohenberger W: The differential diagnosis and interdisciplinary treatment of anal carcinoma. *Dtsch Arztebl Int* 2015; 112: 243–9.
2. Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften: S2k-Leitlinie: Anale Dysplasien und Analkarzinome bei HIV-Infizierten: Prävention, Diagnostik und Therapie. www.awmf.org/leitlinien/detail/II/055-007.html (last accessed on 23 2015).
3. Kirby T: UK committee recommends HPV vaccination for MSM. *Lancet Oncol* 2015; 16: e7.
4. Berry JM, Jay N, Cranston RD, et al.: Progression of anal high-grade squamous intraepithelial lesions to invasive anal cancer among HIV-infected men who have sex with men. *Int J Cancer* 2014; 134: 1147–55.

Prof. Dr. med. Alexander Kreuter

Klinik für Dermatologie, Venerologie und Allergologie,
HELIOS St. Elisabeth Klinik Oberhausen
alexander.kreuter@helios-kliniken.de

Dr. med. Stefan Esser

Klinik für Dermatologie und Venerologie,
Universität Duisburg-Essen

Prof. Dr. med. Ulrike Wieland

Institut für Virologie,
Nationales Referenzzentrum für Papillom- und Polyomaviren,
Uniklinik Köln

Conflict of interest statement

Prof Kreuter has received lecture honoraria for lectures and consultancy fees (advisory board) from MSD Sanofi Pasteur. Dr Esser has participated as a local investigator in Germany in the "Human Papillomavirus Quadrivalent Vaccine Study" of Merck and MSD Sanofi Pasteur. Prof Wieland has received lecture honoraria from ViiV Healthcare, Becton Dickinson, and A&O Labor Delitzsch.

In Reply:

Anal cancers are rare disease entities. Their treatment requires interdisciplinary multimodal therapy in experienced and specialized centers. Many studies have shown remarkable results in terms of individual aspects of diagnosis and treatment, but in the context of a review article addressing a general medical readership, it was not possible always to reflect these in detail.

For HIV patients, the 2013 German-language guidelines from the Association of the Scientific Medical Societies in Germany (AWMF) apply (1). However, as these guidelines are not widely accepted internationally, and as our article is available in an English version, it seemed more appropriate in the given context to explain the international guidelines. In the guidelines of the National Comprehensive Cancer Network (NCCN), screening in HIV positive patients and homosexual men is the subject of controversial discussion, since to date, no randomized study has evaluated the effectiveness of a screening program in terms of a reduced incidence and mortality in anal carcinoma in HIV positive patients (2). The European Society for Medical Oncology (ESMO) expresses similar sentiments in its guidelines (3).

We explicitly highlighted immunosuppression as a risk factor in our article in Box 1. The effectiveness of the quadrivalent HPV vaccine given prophylactically in order to prevent anal dysplasia in men who have sex with men is unequivocally confirmed. The results of the larger study from England were presented. In addition

to England, vaccination against HPV is recommended for both sexes in Canada, Austria, and Australia (4).

According to current guidelines, only the well differentiated T1 lesions of the anal margin should be treated by primary excision. In T2 carcinomas on the anal margin (measuring 2–5 cm at their largest diameter), neoadjuvant radiochemotherapy is the preferred option because of deleterious effects of surgery on sphincter function and continence. The American Society of Colon and Rectal Surgeons (ASCRS) in its guideline recommends excision unequivocally for T1 (≤ 2 cm), N0, G1 anal margin carcinomas with a safety resection margin of 1 cm. The European guidelines recommend a 5 mm safety margin for adequate resection. In larger cancers of the anal margin and in tumors of the anal canal in any stage, primary local excision is contraindicated (2, 3).

In suspected recurrence, incision biopsies should—in our experience—be avoided, if possible, since only more extensive excisions would find possible diffuse or scattered/dispersed tumor islands in the case of incomplete remission or relapse. Excisions in the anal canal should also be avoided at initial diagnosis as the risk of incontinence and vastly delayed wound healing in the radiation field are is very high.

In addition to disrupted wound healing, sexual dysfunction can occur after irradiation of the female pelvic region, owing to hormonal ovarian endocrine insufficiency and trophic changes to the vagina, some of which require treatment. These aspects of treatment are

covered in the regular information given to patients; the standardized information for patients unambiguously mentions ovarian endocrine insufficiency as well as changes to the vaginal epithelium as adverse effects.

DOI: 10.3238/arztebl.2015.0739b

REFERENCES

1. AWMF: Anale Dysplasien und Analkarzinome bei HIV-Infizierten: Prävention, Diagnostik und Therapie. www.awmf.org/leitlinien/detail/II/055-007.html (last accessed on 20 May 2015).
2. NCCN: NCCN Guidelines Anal Carcinoma Version 2.2015. www.nccn.org/professionals/physician_gls/f_guidelines.asp#anal (last accessed on 20 May 2015).
3. Glynn-Jones R, Nilsson PJ, et al.: Anal cancer: ESMO-ESSO-ESTRO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014; 25 Suppl 3:iii10-iii20.
4. Stanley M: HPV vaccination in boys and men. *Hum Vaccin Immunother* 2014; 10: 2109–11.
5. Raptis D, Schneider I, Matzel KE, Ott O, Fietkau R, Hohenberger W: The differential diagnosis and interdisciplinary treatment of anal carcinoma. *Dtsch Arztebl Int* 2015; 112: 243–9.

Dr. Dimitrios Raptis,
Prof. Dr. med. Ignaz Schneider,
Prof. Dr. med. Klaus E. Matzel,
Prof. Dr. med. Dr. h.c. Werner Hohenberger,
 Chirurgische Klinik, Universitätsklinikum Erlangen
dimitrios.raptis@uk-erlangen.de

PD Dr. med. Oliver Ott,
Prof. Dr. med. Rainer Fietkau
 Strahlenklinik, Universitätsklinikum Erlangen

Conflict of interest statement

The authors declare that no conflict of interest exists.