ERRATUM

Erratum for Uda et al., Establishment of an *In Vitro* D-Cycloserine-Synthesizing System by Using O-Ureido-L-Serine Synthase and D-Cycloserine Synthetase Found in the Biosynthetic Pathway

Narutoshi Uda, Yasuyuki Matoba, Takanori Kumagai, Kosuke Oda, Masafumi Noda, Masanori Sugiyama

Department of Molecular Microbiology and Biotechnology, Graduate School of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

Volume 57, no. 6, p. 2603–2612, 2013. Page 2609, right column, first full paragraph, line 6: “ $2.1 \times 10^5 \pm 0.3 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ ” should read “ $2.1 \times 10^2 \pm 0.3 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$.”

Page 2610: The values in Table 2 should appear as shown below.

TABLE 2 Kinetic parameters of DcsG

Substrate	Mean K_m (mM) ^a ± SD	Mean k_{cat} (s ^{−1}) ^a ± SD	Mean k_{cat}/K_m (M ^{−1} s ^{−1}) ± SD	Mean K_m^{app} for ATP (μM) ^b ± SD
D-OUS	10 ± 1	2.2 ± 0.2	210 ± 30	73 ± 8
β-Aminooxy-D-alanine	ND ^c	ND ^c	14 ± 1	37 ± 3
D-2,4-Diaminobutyrate	ND ^c	ND ^c	2.1 ± 0.1	45 ± 7
D-Ornithine	ND ^c	ND ^c	0.67 ± 0.04	45 ± 2
D-Homocysteine	19 ± 3	0.86 ± 0.06	45 ± 9	23 ± 6
D-Homoserine	ND ^c	ND ^c	6.1 ± 0.3	54 ± 10

^a K_m and k_{cat} were determined in the presence of 5 mM ATP.

^b K_m^{app} for ATP was determined in the presence of 50 mM D-amino acid.

^c Since K_m was too high, K_m and k_{cat} were not determined (ND).

Citation Uda N, Matoba Y, Kumagai T, Oda K, Noda M, Sugiyama M. 2015. Erratum for Uda et al., Establishment of an *in vitro* D-cycloserine-synthesizing system by using o-ureido-L-serine synthase and D-cycloserine synthetase found in the biosynthetic pathway. Antimicrob Agents Chemother 59:7926.
doi:10.1128/AAC.02364-15.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.