Published in final edited form as:

Bull Math Biol. 2015 October ; 77(10): 1833–1853. doi:10.1007/s11538-015-0106-4.

Dynamics of simple food webs

Tomas Gedeon and

Department of Mathematics, Montana State University, Bozeman, MT 59717

Patrick Murphy

Department of Mathematics, Montana State University, Bozeman, MT 59717

Abstract

We consider a simple food web with commensal relationship, where organisms utilize both external resources, and resources produced by other organisms. We show that in such a community with no competition there is at most one possible equilibrium for each fixed set of surviving species, and develop a natural condition that determines which species survive based on available resource. Our main result shows that among all possible communities described by equilibria, the one which is stable has the largest number of surviving species, largest combined biomass and hence maximizes utilization of available resources.

Keywords

Microbial Consortia; Chemostat; Commensal relationship

1 Introduction

From van Leeuwenhoek's earliest observations of the microverse to contemporary interest in human microbiomes, it has been apparent that microbes do not exist as monocultures. Naturally occurring ecosystems, optimized by eons of evolution, are almost exclusively organized in communities. In fact, a general positive correlation has been established between community diversity and productivity (Kassen et al, 2000; Venail et al, 2008). Recent advances in metagenomic techniques have given us the tools to estimate the diversity of naturally occurring microbial communities. In a wide range of samples from soil (Fierer and Jackson, 2006), to the ocean (Venter et al, 2004), to the human gut (Gill et al, 2006), it has been found that microbial communities are incredibly diverse, often consisting of thousands of interacting species. Due to these interactions, studying the behavior of individual microbes in isolation does not capture their behavior in a natural community.

Subsets of these communities form *consortia* that act together to enhance their capabilities. The interactions in these consortia lead to emergent behaviors, allowing the systems to perform advanced functions that the individual microbes are not capable of (Eiteman et al, 2008). Emergent properties of microbial interactions are known to be important in diverse areas including medical infections (e.g. diabetic ulcers), biofuels synthesis (e.g. biodiesel

production (Zuroff and Curtis, 2012; Peralta-Yahya et al, 2012)), environmental nutrient cycling (e.g. CO₂ sequestering), bioprocessing (Shong et al, 2012) and wastewater treatment (Seitz et al, 1990a,b; Schink, 1997).

Natural consortia are often organized either as *syntrophic* or *commensal* consortia around the sequential degradation of complex compounds like lignocellulosic material. In these systems, one species catabolizes the available substrate, oxidizing it to produce a byproduct that the second species can consume. If the byproduct inhibits the growth of the producer species then the interaction is syntrophic, if it has no effect on the producer, the interaction is commensal. The syntrophic chain community has been studied mathematically, and it can be shown that for n species in the chain, the coexistence state is stable (Reilly, 1974; Powell, 1985, 1986). This system can be modified to include other forms of inhibition, external toxins, multiple substrates, and other forms of mutualism. In all cases a stable coexistence steady state is found (Aota and Nakajima, 2001; Elkhader, 1991; Burchard, 1994; Katsuyama et al, 2009; Sari et al, 2012), indicating that this is a good candidate system for producing stable consortia.

The commensal chain can be seen as a way of dividing up the steps involved in degrading the available substrate, thereby allowing for the maximum utilization rate of the available energy, in agreement with the maximum power principle (MPP). Originally formulated by Lotka (1922) and further developed by Odum and Pinkerton (1955), the MPP states that biological systems capture and use energy to build and maintain structures and gradients, allowing additional capture and utilization of energy. Some arguments have been put forward to explain how such a system might naturally evolve and reach a steady state (de Mazancourt and Schwartz, 2010; Doebeli, 2002; Pfeiffer and Bonhoeffer, 2004; Bull and Harcombe, 2009; Estrela and Gudejl, 2010; Beardmore et al, 2011). An alternative *resource ratio* theory (de Mazancourt and Schwartz, 2010; Tilman, 1982) can describe cooperative populations by accounting for mutualistic resource exchange. Cooperating populations that exchange limiting resources can exist in a wider range of resource environments than is possible for either population individually. This highlights an evolutionary advantage of cooperation that has been observed in natural ecosystems (de Mazancourt and Schwartz, 2010).

In experiments where a wild-type *E. coli* was allowed to evolve for many generations, it was found to lead to a similar system, where one strain consumed the glucose substrate and another consumed the acetate the first strain produced. However, there was an additional secondary scavenger species that preferentially consumed glycerol (another byproduct of the primary glucose consumer) (Rosenzweig et al, 1994). This *hub* system of a primary producer and multiple scavengers has been found to evolve repeatably and to be robust to system perturbations (Helling et al, 1987; Treves et al, 1998; Rozen and Lenski, 2000), and thus is another good candidate for producing stable consortia.

Our goal in this paper is to develop a general theory for a commensal food web of arbitrary size, where the product of one species is consumed by another species. Since we want to concentrate on commensal relationships, we exclude competition from consideration. The food webs that we consider include both the food chains, where a resource is sequentially

degraded by a set of organism, and a hub-type food webs, where the principal resource is degraded to a set of secondary resources which then support a set of specialist species.

We now briefly summarize our results. We show under very general assumptions that a $n + m$ -dimensional consortium model which includes n species and m essential resources has an n -dimensional invariant manifold. It follows that the system can always be reduced to an n -dimensional system. We compute the reduced equations on this manifold. We then consider a narrower class of *simple food webs* with no competition for resources and commensal relationships between species. We show that there is at most one equilibrium with a given set of species surviving, and we provide a condition in terms of available resources that guarantees the survival of a particular species in such a community equilibrium.

Our main result concerns the stability of the equilibria. We show that there is unique stable equilibrium in the system which corresponds to the largest community that can be supported by the available resources. In other words, out of all existing equilibria, the one that is stable has the greatest number of species. Furthermore, this equilibrium maximizes biomass over all other equilibria. This is in agreement with the MPP principle which suggests that if coexistence occurs, the resulting communities should have higher power than either species could have alone, or other less effective communities (DeLong, 2008). Indeed, the stable equilibrium corresponds to the consortium that transforms more of the resources into biomass, and hence utilizes more of the available energy, than any other consortium in the system.

2 General System

We consider a chemostat model with n microbes and m substrates, which are both consumed and produced by the microorganisms.

$$\begin{aligned}\dot{x}_1 &= (f_1 - D)x_1 \\ \dot{x}_2 &= (f_2 - D)x_2 \\ &\vdots \\ \dot{x}_n &= (f_n - D)x_n\end{aligned}\tag{1}$$

$$\dot{S}_1 = (S_1^{\text{in}} - S_1)D - \sum_j \alpha_{1j} f_j x_j + \sum_j \beta_{1j} f_j x_j$$

$$\dot{S}_2 = (S_2^{\text{in}} - S_2)D - \sum_j \alpha_{2j} f_j x_j + \sum_j \beta_{2j} f_j x_j$$

$$\begin{aligned}&\vdots \\ \dot{S}_m &= (S_m^{\text{in}} - S_m)D - \sum_j \alpha_{mj} f_j x_j + \sum_j \beta_{mj} f_j x_j\end{aligned}$$

Here S_i^{in} is the influx rate of the substrate S_i into the chemostat, x_i is a concentration of the i^{th} microorganism, and D is a dilution rate, which is assumed to be the same for all substrates and species. The growth rate $f_i = f_i(S_1, \dots, S_m)$ of each microbe x_i may depend on every other substrate S_i , and the effect of a substrate may be positive when a substrate is consumed, or detrimental when the substrate is poisonous and decreases the growth rate of x_i . We denote consumption yield coefficients by $\alpha_{ij} > 0$ where i is the substrate and j is the consumer. On the other hand when a microbe j produces a substrate i , we denote the corresponding conversion, or yield coefficient, by β_{ij} . We assume that both types of yield coefficients are constant.

In vector form these equations can be written as

$$\dot{\mathbf{x}} = (\mathbf{F} - D\mathbf{I})\mathbf{x}$$

$$\dot{\mathbf{S}} = (\mathbf{S}^{\text{in}} - \mathbf{S})D - \mathbf{Y}\mathbf{F}\mathbf{x} \quad (2)$$

where $\mathbf{S}^{\text{in}} - \mathbf{S}$ is a vector with elements $S_i^{\text{in}} - S_i$, $\mathbf{Y}$ is the net consumption matrix with (i, j) th element $y_{ij} := \alpha_{ij} - \beta_{ij}$, and the matrix $\mathbf{F} = \mathbf{F}(\mathbf{S})$ is a $n \times n$ diagonal matrix with $f_i(\mathbf{S})$ being the (i, i) element on the diagonal.

Following Smith and Li (2003), our first observation is that this system admits a globally attracting affine n -dimensional manifold M . To see this, consider a new set of variables

$$w_i = S_i + \sum_j \alpha_{ij} x_j - \sum_j \beta_{ij} x_j, \quad i = 1, \dots, m$$

which summarize the influx and outflow of substrate i . We write this change of variables in the vector form

$$\mathbf{w} = \mathbf{Y}\mathbf{x} + \mathbf{S},$$

where $\mathbf{w}$, $\mathbf{S}$ and $\mathbf{x}$ are vectors of the corresponding variables

Then the $\mathbf{S}$ equations in (2) can be replaced by m equations in new variables $\mathbf{w}$

$$\dot{\mathbf{w}} = \mathbf{Y}\dot{\mathbf{x}} + \dot{\mathbf{S}} = \mathbf{Y}(\mathbf{F} - D\mathbf{I})\mathbf{x} + (\mathbf{S}^{\text{in}} - \mathbf{S})D - \mathbf{Y}\mathbf{F}\mathbf{x} = -D\mathbf{Y}\mathbf{x} + (\mathbf{S}^{\text{in}} - \mathbf{S})D = (\mathbf{S}^{\text{in}} - \mathbf{w})D \quad (3)$$

Therefore the system (2) is transformed into

$$\dot{\mathbf{w}} = (\mathbf{S}^{\text{in}} - \mathbf{w})D$$

$$\dot{\mathbf{x}} = (\mathbf{F}(\mathbf{w} - \mathbf{Y}\mathbf{x}) - D\mathbf{I})\mathbf{x} \quad (4)$$

Observe that this system has a globally attracting n -dimensional affine invariant manifold defined by

$$M := \{(\mathbf{x}, \mathbf{w}) \in \mathbb{R}^{n+} \times \mathbb{R}^{m+} | \mathbf{w} = \mathbf{S}^{\text{in}}\} = \{(\mathbf{x}, \mathbf{S}) \in \mathbb{R}^{n+} \times \mathbb{R}^{m+} | \mathbf{S} = \mathbf{S}^{\text{in}} - \mathbf{Y}\mathbf{x}\}$$

where $\mathbb{R}^{n+}$ denotes the non-negative orthant in $\mathbb{R}^n$. The dynamics of the original system on the manifold M have the form

$$\dot{\mathbf{x}} = (\mathbf{F}(\mathbf{S}^{\text{in}} - \mathbf{Y}\mathbf{x}) - D\mathbf{I})\mathbf{x} \quad (5)$$

where the dependence of $\mathbf{F}$ on $\mathbf{S}$ is replaced by a dependence on $\mathbf{S}^{\text{in}} - \mathbf{Y}\mathbf{x}$.

Note that the dynamics on the invariant manifold M depends on the yield matrix $\mathbf{Y}$.

3 Simple Food webs

In what follows we will put additional simplifying assumptions that will allow us to analyze the system, yet that are general enough to include interesting examples, some of which are analyzed in the following section. To describe the set of assumptions we will use the language of graph theory.

Let $G(V, E)$ be an oriented graph, where each vertex is labeled by a species x_i , and each edge is labeled by a resource S_j . Each edge connects a producer to a consumer of the resource labeling the edge. In other words, an edge starting at a node x_i corresponds to a resource S_j that is produced by x_i , and an edge that terminates in x_k corresponds to a resource that is consumed by x_k . The influx of external resources to chemostat is represented by edges with only a terminal node, and labeled by a particular resource.

Definition 1

A simple food web is a collection of n species and m resources, represented by an oriented graph $G(V, E)$ where

1. there are no non-trivial oriented cycles in G , i.e. no cycles except possibly when a species produces and consumes the same substrate;
2. each species x_i consumes a dedicated substrate S_j . This eliminates the competition in the system. After a change of indices, if necessary, we will assume that species x_i consumes substrate S_i . This means that all edges that terminate at a vertex x_i must have the same label, and that these labels are distinct for different vertices;
3. the growth functions f_i , which by (2) depend on a single substrate S_i , are monotonically increasing.
4. If species i produces S_j and consumes the same S_j , then the substrate-from-biomass yield β_{ji} is smaller than the biomass-from-substrate yield α_{ji} . The opposite case would allow a net production of biomass without any external resource.

We now discuss several consequences of our assumptions.

- Without loss of generality we can restrict our attention on simple food webs that are represented by a connected graph; if this is not the case we can restrict our attention to each connected component separately.
- Assumption (2) implies that $m \geq n$, i.e. the number of resources m is greater or equal to number of species n . If $m > n$, then the extra resources are not consumed, their dynamic behavior does not affect the rest of the system. Therefore we can restrict our attention to the core system of n species and n resources that are consumed by these species. If some species produce resources that are not consumed, their behavior can be determined after the behavior of the core system is identified.
- Our assumptions imply a stratification of G into food chain layers $X_0, X_1, \dots, X_k$. We denote the set of all vertices i with $S_i^{in} > 0$ by X_S . A depth $d(x_i)$ of the species x_i in the food chain is defined to be the length of the longest path in G from x_i to some vertex in X_S . Again, by assumption (1) depth is well defined for each species. The j -th layer of the simple food chain is the set X_k of those species x_i with $d(x_i) = k$.

We now prove that simple food webs have the following property.

Lemma 1

In the core system of any simple food web, the n species and n resources can be relabeled in such a way that the matrix $\mathbf{Y}$ is lower triangular and invertible.

Proof—We order all species according to their depth, starting with X_0 , species with depth 0. Since each substrate S_i is assigned to a unique species x_i , we order substrates in the same order the species are in. Therefore $\alpha_{ij} = 0$ for all $i < j$, and $\alpha_{ii} > 0$ for all i .

Now we examine yields β_{ij} . Observe that a species x_i with depth $d(x_i) = k$ cannot depend on resource S_j if $d(x_j) < k$. If this was the case, then there will be a path in G from X_S to x_i through x_j with length at least $k + 1$. This contradicts the fact that x_i has depth k . Therefore β_{ij} , the measure of production for substrate i by species j , is 0 for $i < j$.

Finally, by (4) the diagonal entries $y_{ii} = \alpha_{ii} - \beta_{ii}$ are greater than zero. Hence the yield matrix $\mathbf{Y}$ representing the core system is square and lower triangular with nonzero entries along the diagonal. Therefore $\mathbf{Y}$ is invertible.

4 Existence of Equilibria

Let $N := \{1, \dots, n\}$ and let $\mathcal{P}(N)$ be the collection of all subsets of N . Then the phase space $\mathbb{R}^{n+}$ can be decomposed into disjoint subsets C_U , parameterized by the sets $U \in \mathcal{P}(N)$, and defined by

$$C_U = \{x \in \mathbb{R}^{n+} \mid x_i > 0 \text{ for } i \in U, x_i = 0 \text{ for } i \notin U\}.$$

Definition 2

The necessary resource for a species x_i is the value S_i^{nec} implicitly defined by

$$f_i(S_i^{nec}) = D.$$

The proof of the next Theorem requires conditions that are significantly weaker than those in Definition 1. In particular, we only require monotonicity of each f_i , not that f_i is an increasing function.

Theorem 1

Assume that every $f_i(S_i)$ is monotonically increasing, or monotonically decreasing. Let $\mathbf{Y}_U$ be a principal minor of $\mathbf{Y}$ specified by the index set $U \in \mathcal{P}(N)$; that is $\mathbf{Y}_U$ is submatrix of $\mathbf{Y}$ that is constructed by deleting all rows and columns i where $i \notin U$.

Then, if $\det \mathbf{Y}_U \neq 0$, then the system (5) can have at most one equilibrium in C_U .

This equilibrium, if it exists, is determined by equations

$$x_i = 0 \text{ for } i \notin U; \quad x_i > 0 \text{ for } i \in U; \quad \mathbf{S}_U^{in} - \mathbf{S}_U^{nec} = \mathbf{Y}_U \mathbf{x}_U,$$

where $\mathbf{x}_U$, $\mathbf{S}_U^{in}$ and $\mathbf{S}_U^{nec}$ are vectors of $\mathbf{x}$, $\mathbf{S}^{in}$ and $\mathbf{S}^{nec}$ restricted to indices $i \in U$, respectively.

Proof—In a given set U , the equations that determine the equilibria are

$$x_i = 0 \text{ for } i \notin U; \quad f_i(S_i^{nec}) = D \text{ for } i \in U,$$

which follows immediately from (5) and the fact that at an equilibrium where $x_i = 0$, we must have $f_i(S_i) = D$. By monotonicity of f_i there is at most one solution S_i^{nec} of $f_i(S_i) = D$ for any $i \in U$. If for some $i \in U$ such S_i^{nec} does not exist, then C_U will not contain an equilibrium. With the vector $\mathbf{S}_U^{nec}$ assembled we compute the x_i components of the equilibrium by solving (see (5)) $\mathbf{S}^{nec} - \mathbf{S}^{in} = \mathbf{Y}\mathbf{x}$ restricted to the species x_i , $i \in U$. This yields equation

$$\mathbf{S}_U^{in} - \mathbf{S}_U^{nec} = \mathbf{Y}_U \mathbf{x}_U$$

which has a unique solution when $\mathbf{Y}_U$ is invertible. If this solution has a positive component $x_i > 0$ for all $i \in U$, then C_U contains an equilibrium.

We have the following Corollary for a simple food web.

Corollary 1

In a simple food web, each C_U contains at most one equilibrium.

Proof—Since $\mathbf{Y}$ is lower triangular with nonzero diagonal elements, each principal minor is invertible.

At this point a natural question is if there is a simple criterion to determine if a given C_U contains an equilibrium. We will need a concept of an *available resource* at $\mathbf{x} = (x_1, \dots, x_n)$.

Definition 3

Given a location in phase space $\mathbf{x} = (x_1, \dots, x_n)$, the available resource for species i is

$$S_i^{av}(\mathbf{x}) := S_i^{in} - [\mathbf{Y}\mathbf{x}]_i$$

where $[\mathbf{Y}\mathbf{x}]_i$ is the i^{th} entry of the vector $\mathbf{Y}\mathbf{x}$.

Definition 4

For each species x_i the set of predecessors P_i is the set of species j such that x_j produces the essential resource needed by x_i . In other words, $j \in P_i$ if, and only if, there is an edge from $j \rightarrow i$ in the graph G .

For each vector $\mathbf{e} = (e_1, \dots, e_n)$, let $\hat{\mathbf{e}}_j$ be a vector

$$\hat{e}_{ji} = \begin{cases} e_i & \text{if } i \in P_i \setminus \{j\} \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 2

A vector $\mathbf{e} = (e_1, \dots, e_n) \in \mathbb{R}^{n+}$ is a equilibrium if, and only if, for each i , either $e_i = 0$ or, if $e_i > 0$, then

$$e_i = \frac{S_i^{av}(\hat{\mathbf{e}}_i) - S_i^{nec}}{y_{ii}}. \quad (6)$$

Proof—($\Rightarrow$) We assume that $\mathbf{e}$ is an equilibrium in $\mathbb{R}^{n+}$. Then by (5) either $e_i = 0$, or, if $e_i > 0$, then $f_i(S_i^{nec}) = D$ and $S_i^{nec} = S_i^{in} - [\mathbf{Y}\mathbf{e}]_i$.

This implies

$$S_i^{nec} = S_i^{in} - [\mathbf{Y}\mathbf{e}]_i = S_i^{in} - \sum_{k \in P_i} y_{ik}e_k - y_{ii}e_i = S_i^{av}(\hat{\mathbf{e}}_i) - y_{ii}e_i,$$

from which (6) follows.

($\Leftarrow$) Observe that $e_i = 0$ always satisfies (5). Suppose that if $e_i > 0$ then (6) holds. Then we have

$$S_i^{in} - [\mathbf{Y}\mathbf{e}]_i = S_i^{in} - \sum_{k \in P_i} y_{ik}e_k - y_{ii}e_i = S_i^{av}(\hat{\mathbf{e}}_i) - y_{ii}e_i = S_i^{nec}$$

and therefore (5) is satisfied for component i . This shows that (5) holds for all components, hence $\mathbf{e}$ is an equilibrium.

Definition 5

A set of species corresponding to a set of vertices I is independent if $i \notin P_j$ for any two indices $i, j \in I$.

The next Corollary gives an inductive way to construct the set of all possible equilibria, after taking into account that $\mathbf{0} = (0, \dots, 0)$ is always an equilibrium.

Corollary 2

Let $\mathbf{e} = (e_1, \dots, e_n)$ be an equilibrium contained in some C_U . Let I be an independent set of species with $U \cap I = \emptyset$, satisfying $S_i^{av}(\mathbf{e}) > S_i^{nec}$ for all $i \in I$. Then there exists an equilibrium $\mathbf{E} = (E_1, \dots, E_n)$ in $C_{U \cup I}$ with $E_i > 0$ for all $i \in I \cup U$.

Proof—Pick an arbitrary $i_1 \in I$. We observe that since $I \cap U = \emptyset$ we have $e_{i_1} = 0$ and thus

$$S_{i_1}^{av}(\hat{\mathbf{e}}_{i_1}) = S_{i_1}^{av}(\mathbf{e}) = S_{i_1}^{in} - \sum_{k \in P_{i_1} \cap U} y_{ik}e_k.$$

We construct a vector $\mathbf{E}^1$ where $E_j^1 = e_j$ for all $j \neq i_1$ and replace $e_{i_1} = 0$ by

$$E_{i_1}^1 = \frac{S_{i_1}^{av}(\hat{\mathbf{e}}_{i_1}) - S_{i_1}^{nec}}{y_{i_1 i_1}} = \frac{S_{i_1}^{av}(\mathbf{e}) - S_{i_1}^{nec}}{y_{i_1 i_1}} > 0,$$

then by Theorem 2 the resulting vector $\mathbf{E}^1$ is an equilibrium with $\mathbf{E}^1 \in C_{U \cup \{i_1\}}$.

Select now an arbitrary $i_2 \in I \setminus \{i_1\}$. Since I is an independent set of species, $i_1 \notin P_{i_2}$ and therefore the i_1^{th} component of $\hat{\mathbf{e}}_{i_2}$ is zero. Therefore

$$S_{i_2}^{av}(\hat{\mathbf{e}}_{i_2}) = S_{i_2}^{av}(\mathbf{e}) = S_{i_2}^{av}(\mathbf{E}^1) = S_{i_2}^{in} - \sum_{k \in P_{i_2} \cap U} y_{ik}e_k.$$

As before, we construct a vector $\mathbf{E}^2$ such that $E_j^2 = E_j^1$ for all $j \neq i_2$ and replace $e_{i_2} = 0$ by

$$E_{i_2}^2 = \frac{S_{i_2}^{av}(\hat{\mathbf{e}}_{i_2}) - S_{i_2}^{nec}}{y_{i_2 i_2}} = \frac{S_{i_2}^{av}(\mathbf{E}^1) - S_{i_2}^{nec}}{y_{i_2 i_2}} > 0.$$

By Theorem 2 the resulting vector $\mathbf{E}^2$ is an equilibrium with $\mathbf{E}^2 \in C_{U \cup \{i_1, i_2\}}$. Repeating the argument until we exhaust the index set I finishes the proof.

Lemma 2

Consider any equilibrium $\mathbf{e} = (e_1, \dots, e_n)$. Then, for each i such that $e_i > 0$, there is an oriented path p in the graph G connecting a vertex in X_S with external resource input to the vertex x_i , such that $e_k > 0$ for all $k \in p$.

Stated more strongly, for each equilibrium $\mathbf{e}$ there is a set of species $x_{i_1}, \dots, x_{i_k}$ and (not necessarily disjoint) oriented paths $p_{i_1}, \dots, p_{i_k}$ such that p_{i_j} connects a vertex in X_S to vertex x_{i_j} , with the property that $e_i > 0$ if, and only if, $i \in \cup_j p_{i_j}$.

Proof—Let $e_i > 0$. Consider the equation

$$S_i^{nec} = S_i^{in} - [\mathbf{Y}\mathbf{e}]_i = S_i^{in} - \sum_{k \in P_i} y_{ik}e_k - y_{ii}e_i = S_i^{in} - \sum_{k \in P_i} \beta_{ik}e_k - y_{ii}e_i \quad (7)$$

Note that since $S_i^{nec} > 0$, in order for (7) to hold there must be either $S_i^{in} > 0$, and hence $x_j \in X_S$, or there must be at least one $k \in P_i$ with $e_k > 0$. Repeating this argument for every such k , we see that there must be at least one path in G from X_S to vertex x_i such that $e_k > 0$ for all k along that path.

To show the second statement of the Theorem, we start with some $e_i > 0$. Then we enumerate all the paths that connect X_S to x_i and for which each x_k along these paths satisfies $e_k > 0$. If this exhausts the non-zero entries of $\mathbf{e}$, we are done; if there is an $e_j > 0$ that is not accounted for by the paths already selected, we repeat the argument for e_j . Since the number of components of $\mathbf{e}$ is finite, this process will terminate in finitely many steps.

Theorem 2 and Lemma 2 motivate the following definition.

Definition 6

Fix the structure of simple food web, the yield matrix $\mathbf{Y}$, and the set of growth functions f_i . The set of *feasible equilibria*

$$E = E(S^{in}, D)$$

is defined to be the set of equilibria of the system (5) at a given level of inputs S^{in} and dilution rate D .

Corollary 2 gives an algorithm how to build the set E from the bottom up by starting with the zero equilibrium $(0, \dots, 0)$ and adding equilibria to E based on sufficiency of available resources. On the other hand Lemma 2 gives a recursive characterization of equilibria in E . Since the growth functions f_i are monotone, the sets of feasible equilibria are nested as a function of external resources, or the dilution rate D . In particular, if $D_1 < D_2$ then

$$E(\mathbf{S}^{in}, D_2) \subseteq E(\mathbf{S}^{in}, D_1) \text{ for any fixed } \mathbf{S}^{in}$$

Similar containment holds for external resources. If $\mathbf{S}^{in} \prec \bar{\mathbf{S}}^{in}$ stands for partial order of vectors in the positive orthant (that is $S_i^{in} < \bar{S}_i^{in}$ for at least one i and $S_j^{in} \leq \bar{S}_j^{in} \forall j \neq i$) then

$$E(S^{in}, D) \subseteq E(\bar{S}^{in}, D) \text{ for any fixed } D.$$

The structure of feasible sets of equilibria is not apriori clear in simple food webs that are not chains. As an example, consider simple food web in Figure 1. As we increase available external resources S_1 and S_2 a possible sequence of sets of feasible equilibria may be

$$E_1 = \{(0, 0, 0, 0, 0), (e_1^1, 0, 0, 0, 0)\},$$

$$E_2 = \{(0, 0, 0, 0, 0), (e_1^2, 0, 0, 0, 0), (e_{21}^2, e_{22}^2, 0, 0, 0)\},$$

$$E_3 = \{(0, 0, 0, 0, 0), (e_1^3, 0, 0, 0, 0), (e_{21}^3, e_{22}^3, 0, 0, 0), (e_{31}^3, 0, e_{33}^3, 0, 0)\},$$

where we assume that all $e_j^i > 0$. However, as we will show in Theorem 3, the set E_3 is not feasible. If equilibria $(e_{21}^3, e_{22}^3, 0, 0, 0)$ and $(e_{31}^3, 0, e_{33}^3, 0, 0)$ exist, there also must be an equilibrium of the form $(e_{41}^3, e_{42}^3, e_{43}^3, 0, 0)$.

To study the structure of feasible equilibria we introduce a set L of all subspaces that support an equilibrium.

Definition 7

Let

$$L = \{U \in \mathcal{P}(N) \mid \exists \mathbf{e} \in C_U \text{ such that } \mathbf{e} \text{ is an equilibrium of (5)}\}.$$

where L is partially ordered by inclusion.

We then have the following Theorem.

Theorem 3

If $U, W \in L$, then $U \cup W \in L$.

Proof—Let $\mathbf{e}$ be an equilibrium in C_U and let $\mathbf{v}$ be an equilibrium in C_W . Let $\mathbf{0} = (0, \dots, 0)$ be the zero equilibrium. We construct, by induction on the depth of coordinates in $U \cup W$, a set of equilibria $\mathbf{E}^0, \mathbf{E}^1, \dots, \mathbf{E}^n$, where n is the maximal depth of any vertex in $U \cup W$. Let A_j

$\coloneqq \{i \in U \cup W \mid x_i \in X_j\}$ be the stratification of vertices in $U \cup W$ according to their depth. We construct the equilibria $\mathbf{E}^j$ in such a way that they satisfy the following properties;

a. the i -th component of $\mathbf{E}^j$

$$\begin{cases} E_i^j > 0 & \text{for } i \in \cup_{k \leq j} A_k \\ 0 & \text{otherwise.} \end{cases}$$

b. Furthermore, we have the following inequalities

$$\begin{cases} E_i^j \geq e_i & \text{for } i \in A_j \cap U \\ E_i^j \geq v_i & \text{for } i \in A_j \cap W \end{cases}$$

Clearly, when $j = n$ condition (a) implies the statement of the Theorem, since we will then have an equilibrium whose set of positive components is exactly $U \cup W$.

To start the induction, consider first $A_0 \coloneqq \{i \in U \cup W \mid x_i \in X_0\}$. Since species corresponding to vertices in A_0 depend exclusively on the external resources, we have

$$S_i^{av}(\mathbf{0}) = S_i^{in} = S_i^{av}(\mathbf{e}) \text{ if } i \in A_0 \cap U \quad (8)$$

$$S_i^{av}(\mathbf{0}) = S_i^{in} = S_i^{av}(\mathbf{v}) \text{ if } i \in A_0 \cap W$$

By assumption, the equilibrium $\mathbf{e} \in C_U$ exists, and therefore by Theorem 2 we have $S_i^{av}(\mathbf{e}) > S_i^{nec}$ which implies

$$S_i^{av}(\mathbf{0}) > S_i^{nec} \text{ for all } i \in A_0 \cap U.$$

Similarly, since $\mathbf{v} \in C_W$ exists we have $S_i^{av}(\mathbf{v}) > S_i^{nec}$ which implies

$$S_i^{av}(\mathbf{0}) > S_i^{nec} \text{ for all } i \in A_0 \cap W.$$

Since species in $I \coloneqq A_0$ are clearly independent, by Corollary 2 with $I \coloneqq A_0$ there is an equilibrium $\mathbf{E}^0$ where

– if $i \in A_0$ then

$$E_i^0 \coloneqq \frac{S_i^{av}(\mathbf{0}) - S_i^{nec}}{y_{ii}} > 0;$$

– if $i \notin A_0$ then $E_i^0 \coloneqq 0$.

This proves (a) for $j = 0$. Furthermore, since by Theorem 2

$$e_i = \frac{S_i^{av}(\mathbf{e}) - S_i^{nec}}{y_{ii}}$$

for $i \in U$, (8) implies that

$$E_i^0 = e_i \text{ for all } i \in A_0 \cap U.$$

A similar argument for $i \in W$ implies statement (b) for $j = 0$.

We now proceed with the inductive step. Let

$$B_j = \bigcup_{0 \leq k \leq j} A_k.$$

Assume that (a) and (b) holds for index j .

Recall that since consumption yields α_{ii} only lie on the diagonal of matrix Y , we have $y_{ik} = -\beta_{ik}$ for $i \neq k$. We compute for an arbitrary index i

$$S_i^{av}(\mathbf{E}^j) = S_i^{in} - \sum_{k \in B_j \cap U} y_{ik} E_k^j - \sum_{k \in B_j \cap W} y_{ik} E_k^j = S_i^{in} + \sum_{k \in B_j \cap U} \beta_{ik} E_k^j + \sum_{k \in B_j \cap W} \beta_{ik} E_k^j \geq S_i^{in} + \sum_{k \in B_j \cap U} \beta_{ik} e_k + \sum_{k \in B_j \cap W} \beta_{ik} v_k, \quad (9)$$

where we used the inductive hypothesis (b) in the last line. Since at the equilibrium $\mathbf{e}$, the species in $A_{j+1} \cap U$ only depend on resources produced by species in $B_j \cap U$, we have that

$$S_i^{av}(\mathbf{e}) = S_i^{in} + \sum_{k \in B_j \cap U} \beta_{ik} e_k \text{ for all } i \in A_{j+1} \cap U.$$

Therefore (9) implies

$$S_i^{av}(\mathbf{E}^j) \geq S_i^{av}(\mathbf{e}) \text{ if } i \in A_{j+1} \cap U \quad (10)$$

A similar argument with equilibrium $\mathbf{v}$ yields

$$S_i^{av}(\mathbf{E}^j) \geq S_i^{av}(\mathbf{v}) \text{ if } i \in A_{j+1} \cap W \quad (11)$$

As before, since the equilibria $\mathbf{e} \in C_U$ and $\mathbf{v} \in C_W$ exist we have $S_i^{av}(\mathbf{e}) > S_i^{nec}$ and $S_i^{av}(\mathbf{v}) > S_i^{nec}$, which imply

$$S_i^{av}(\mathbf{E}^j) > S_i^{nec} \text{ for all } i \in A_{j+1} \cap U; S_i^{av}(\mathbf{E}^j) > S_i^{nec} \text{ for all } i \in A_{j+1} \cap W.$$

Therefore by Corollary 2 with $I := A_{j+1}$ there is an equilibrium $\mathbf{E}^{j+1}$ where

- if $i \in A_{j+1}$ then

$$E_i^{j+1} := \frac{S_i^{av}(\mathbf{E}^j) - S_i^{nec}}{y_{ii}} > 0;$$

- if $i \in B_j$ then $E_i^{j+1} := E_i^j$;
- if $i \notin B_{j+1}$ then $E_i^{j+1} := 0$.

This proves (a) for the inductive step.

We observe that (10) and (11), together with definition of E_i^{j+1} implies

$$E_i^1 \geq e_i \text{ for all } i \in A_1 \cap U \text{ and } E_i^1 \geq v_i \text{ for all } i \in A_1 \cap W.$$

This proves (b) for the inductive step and thus finishes the proof.

Corollary 3

Let $\mathbf{e}$ be an equilibrium in C_U , $\mathbf{v}$ be an equilibrium in C_W and let $\mathbf{q}$ be an equilibrium in $C_{U \cup W}$. Then the total biomass $\sum_{i=1}^n q_i$ at equilibrium $\mathbf{q}$ is larger than a total biomass at $\mathbf{e}$ and a total biomass at $\mathbf{v}$:

$$\sum_{i=1}^n q_i \geq \sum_{i=1}^n e_i, \text{ and } \sum_{i=1}^n q_i \geq \sum_{i=1}^n v_i.$$

Proof—This is a direct corollary of uniqueness of equilibria in each C_U (Corollary 1) and the inductive statement (b) in the proof of Theorem 3.

Theorem 3 and Corollary 3 illustrate two important aspects about simple food webs. If the microbes in a community do not harm each other directly or indirectly, and the growth rate functions are monotone, then increasing either the external resources or the number of microbes that produce resources internally will result in the existence of equilibria that represent a larger community in number of species Theorem 3, or overall biomass Corollary 3. Naturally this leads to the question of stability in the class of feasible equilibria.

5 Stability Analysis

We now offer a complete characterization of the stability of all feasible equilibria $E(\mathbf{S}^{\text{in}}, D)$ for the system.

Theorem 4

The unstable manifold of an equilibrium $\mathbf{e} \in C_U$ of the system (5) has dimension

$$k = \#\{i \notin U \mid S_i^{av}(\mathbf{e}) > S_i^{nec}\}.$$

Therefore an equilibrium $\mathbf{e} \in C_U$ is stable if $k = 0$, which corresponds to $S_i^{av}(\mathbf{e}) \leq S_i^{nec}$ for all $i \notin U$.

Proof—We denote the by J_{ij} the (i, j) entry of the Jacobian $\mathbf{J}$. Using the Chain Rule we evaluate Jacobian at a point x to get

$$J_{ii}(x) = f_i(\mathbf{S}) - D + \nabla f_i(\mathbf{S}) \cdot \frac{\partial \mathbf{S}}{\partial x_i} x_i. \quad (12)$$

and for $i \neq j$,

$$J_{ij} = \nabla f_i(\mathbf{S}) \cdot \frac{\partial \mathbf{S}}{\partial x_j} x_i.$$

Since we have $\mathbf{S} = \mathbf{S}^in - \mathbf{Y}\mathbf{e}$, then

$$\frac{\partial \mathbf{S}}{\partial x_j} = -\mathbf{y}_j$$

where $\mathbf{y}_j$ is the j^{th} column of the yield matrix $\mathbf{Y}$. The Jacobian matrix $\mathbf{J}$ can then be written as

$$\mathbf{J} = - \begin{bmatrix} \nabla f_1(\mathbf{S}) \\ \vdots \\ \nabla f_k(\mathbf{S}) \end{bmatrix} \mathbf{Y} \mathbf{X} + \mathbf{F}(\mathbf{S}) - D\mathbf{I}$$

where $\mathbf{X}$ is a diagonal matrix with ii^{th} entry x_i . With our assumptions on resource consumption, we can write the Jacobian as

$$\mathbf{J} = -\mathbf{F}_s \mathbf{Y} \mathbf{X} + \mathbf{F}(\mathbf{S}) - D\mathbf{I} \quad (13)$$

where $\mathbf{F}_s$ is a diagonal matrix with ii^{th} entry $\frac{\partial f_i}{\partial S_i}$. These entries are all positive since f_i is a monotone increasing function. The matrices $\mathbf{X}$, $D\mathbf{I}$, and $\mathbf{F}(\mathbf{S})$ have non-negative entries as well. Recall that by Definition 1, assumption (4), the diagonal entries in $\mathbf{Y}$ are positive.

Now we evaluate Jacobian at an equilibrium $\mathbf{e} \in C_U$. Since $\mathbf{J}$ is lower triangular, the eigenvalues are the diagonal entries of the Jacobian. We note that if $i \in U$, and hence $e_i > 0$, then we have $f_i(S_i) = D$. By inspection of (12) we have

$$J_{ii}(\mathbf{e}) = -y_{ii} \frac{df_i}{dS_i} e_i < 0.$$

It follows that $\mathbf{e}$ is always stable within the subspace $C_U \subset \mathbb{R}^{n+}$.

If $i \notin U$ then $e_i = 0$ and from (12)

$$J_{ii}(\mathbf{e}) = f_i(S_i^{av}) - D.$$

Therefore the positive eigenvalues correspond to those $i \notin V$ with

$$f_i(S_i^{av}(\mathbf{e})) > D.$$

Since $D = f_i(S_i^{nec})$ and f_i is monotonically increasing function, this is equivalent to

$$S_i^{av}(\mathbf{e}) > S_i^{nec}$$

completing the proof.

We are ready for the proof of the main theorem, which states that every simple food web has a unique stable equilibrium. Furthermore, this equilibrium represents the most diverse consortium that can survive in the chemostat. In addition to maximizing diversity, this equilibrium also maximizes biomass of the system.

Theorem 5

Any simple food web has a unique stable equilibrium $\mathbf{E}_s = (e_1, \dots, e_n)$ of (5). The stable equilibrium solves two independent optimization problems over the set of feasible equilibria $E(S^{in}, D)$:

1. E_s has the maximal number of non-zero components ($e_i > 0$) i.e. the maximal number of species that are present;
2. E_s has the maximal overall biomass $\sum_{i=1}^n e_i$.

Proof—By Theorem 3 if C_{U_1} and C_{U_2} contains equilibria, then also $C_{U \cup W}$ contains an equilibrium. In other words we have shown that the partially ordered set L , which is a subset of the lattice of all subsets $P(N)$ of index set $N = \{1, \dots, n\}$, is closed under the join operation in that lattice. Since L is finite, this implies that L has a unique maximal element Z . Let $\mathbf{e}$ be the unique equilibrium in C_Z .

We now show that $\mathbf{e}$ is stable. Assume by contradiction that $\mathbf{e}$ is not stable. This implies that there is $i \notin Z$ such that $S_i^{av}(\mathbf{e}) \leq S_i^{nec}$. By Corollary 2 this implies there is an equilibrium in $C_{Z \cup \{i\}}$. This contradicts maximality of the set Z and therefore $\mathbf{e} \in C_Z$ is stable.

Since Z is maximal in L , and L is closed under join operation, every other set $U \subsetneq Z$ in the lattice L is a subset of another set that belongs to L . Let V be an immediate successor of U in the lattice ordering, i.e. $U \subsetneq V$ and there is no set Q with $U \subsetneq Q \subsetneq V$. Then $V \setminus U = \{j\}$ for some j . Let $\mathbf{e} \in C_U$, $\mathbf{v} \in C_V$ be the equilibria in C_U and C_V respectively. Then $S_j^{av}(\mathbf{e}) = S_j^{av}(\hat{\mathbf{v}}) > S_j^{nec}$, which implies by Theorem 4 that $\mathbf{e}$ has at least a one-dimensional unstable manifold.

This shows that equilibria in C_U for $U \subsetneq Z$, are unstable and the dimension of the unstable manifold is equal to the difference in cardinality $|Z| - |U|$. As a consequence, the system has a unique stable equilibrium in C_Z . Maximization of nonzero entries follows directly from the fact that U is maximal in L , and the maximization of biomass follows from Corollary 3.

Remark

We can interpret Theorem 5 as a statement that the stable equilibrium of the system corresponds to the most diverse population that is sustainable on a given set of resources. The condition $S_i^{av}(\mathbf{e}) \leq S_i^{nec}$ means that the supply of the resource needed to support the growth of x_i is insufficient for its survival.

6 Examples

The theory we have developed for our restricted system (5) can be applied to several systems that have a specific interaction graph. We will look at two archetypical examples: consortium with hub-like graph of interactions and a consortium with a chain-like graph.

6.1 Hub consortium

Our first example is a consortium with hub-like structure, where one species produces all the substrates that other species feed on. The primary motivation is the evolved consortium studied by (Rosenzweig et al, 1994) and described in the introduction.

Only the initial substrate S_1 which feeds species x_1 is externally fed into the system.

Consider the system

$$\begin{aligned}
 \dot{x}_1 &= (f_1(S_1) - D)x_1 \\
 \dot{S}_1 &= (S_1^{\text{in}} - S_1)D - \alpha_{11}f_1(S_1)x_1 \\
 \dot{x}_2 &= (f_2(S_2) - D)x_2 \\
 \dot{S}_2 &= -S_2D - \alpha_{22}f_2(S_2)x_2 + \beta_{21}f_1(S_1)x_1 \\
 &\vdots \\
 \dot{x}_k &= (f_k(S_k) - D)x_k \\
 \dot{S}_k &= -S_kD - \alpha_{kk}f_k(S_k)x_k + \beta_{k1}f_1(S_1)x_1.
 \end{aligned} \tag{14}$$

As always $\mathbf{0} = (0, \dots, 0)$ is an equilibrium. By Corollary 2 there are two possibilities. If

$$S_1^{in} = S_1^{av}(\mathbf{0}) > S_1^{nec}$$

then there is an equilibrium $\mathbf{e} = (e_1, 0, \dots, 0)$; if $S_1^{in} \leq S_1^{nec}$ the only equilibrium is $\mathbf{0}$ (which is also then stable).

As an initial check, since $S_k^{in} = 0$ for $k > 1$ it follows from (5) that at an equilibrium $\mathbf{E}$ with species beyond the first present we need to have $[-Y\mathbf{e}]_i = S_i^{nec}$. This in turn implies $\beta_{k1}E_1 - \alpha_{kk}E_k = S_i^{nec} > 0$ and thus it follows for any equilibrium with $E_k > 0$ also must have $E_1 > 0$.

If $\mathbf{e}$ exists, then by Corollary 2, C_U with $U \neq \emptyset$ will contain an equilibrium $\mathbf{e}_U$ if, and only if

1. $1 \in U$; and
2. for every $i \in U, i \neq 1$ we have $S_i^{av}(\mathbf{e}) > S_i^{nec}$.

Thus if we set $Q := \{i > 1; | S_i^{av}(\mathbf{e}) > S_i^{nec}\}$ then any C_U with $U = \{1\} \cup B$, for any $B \subset Q$, contains a unique equilibrium $\mathbf{e}_U$.

By Theorem 5 the only stable equilibrium will be that which correspond to $U = \{1\} \cup Q$ which is the one where the most possible species survive.

6.2 Chain consortium

We now analyze systems with a chain-like interaction structure, where each species beyond the first is dependent on the resource produced by its predecessor in the chain. Again, we are assuming that only the substrate S_1 is fed externally into the system.

We consider the equations

$$\dot{x}_1 = (f_1(S_1) - D)x_1$$

$$\dot{S}_1 = (S_1^{in} - S_1)D - \alpha_{11}f_1(S_1)x_1$$

$$\dot{x}_2 = (f_2(S_2) - D)x_2$$

$$\begin{aligned} \dot{S}_2 &= -S_2D - \alpha_{22}f_2(S_2)x_2 + \beta_{21}f_1(S_1)x_1 \\ &\vdots \\ \dot{x}_k &= (f_k(S_k) - D)x_k \end{aligned} \tag{15}$$

$$\dot{S}_m = -S_mD - \alpha_{mm}f_m(S_m)x_m + \beta_{m,n-1}f_{n-1}(S_{n-1})x_{n-1}$$

It follows from (5) and the chain structure of the equations that if an equilibrium $\mathbf{E}$ with $E_i > 0$ exists, then we must have $E_{i-1} > 0$ and, by induction, $E_j > 0$ for all $j < i$. Therefore the indexing sets U for which C_U contains an equilibrium, are nested. In other words, there is a maximal k such that for all sets $U_s = \{1, \dots, s\}$ for $s \leq k$, C_{U_s} contains an equilibrium $\mathbf{E}_s$. In the case of the zero equilibrium, $k = 0$. By Theorem 5 the equilibrium $\mathbf{E}_k$ in C_{U_k} is stable.

To illustrate these ideas in more detail it is instructive to make explicit calculations. By Corollary 2 the equilibrium $\mathbf{E}_1$ in C_{U_1} exists if, and only if,

$$S_1^{av}(\mathbf{0}) = S_1^{in} > S_1^{nec}.$$

We compute the available resource at $\mathbf{E}_1$ for species x_2

$$S_2^{av}(\mathbf{E}_1) = \frac{\beta_{21}}{\alpha_{11}}(S_1^{in} - S_1^{nec}) \quad (16)$$

Applying Corollary 2 to $\mathbf{E}_1$, if $S_1^{av}(\mathbf{E}_1) \leq S_2^{nec}$, then there is no equilibrium $\mathbf{E}_2$ with both first and second components greater than zero. On the other hand, if $S_1^{av}(\mathbf{E}_1) > S_2^{nec}$, then $\mathbf{E}_2$ exists and we can calculate available resource at $\mathbf{E}_2$ for species x_3

$$S_3 = -\alpha_{33}x_3 + \beta_{32}x_2 = -\alpha_{33}x_3 + \frac{\beta_{32}}{\alpha_{22}}(\beta_{21}x_1 - S_2) = -\alpha_{33}x_3 + \frac{\beta_{32}}{\alpha_{22}}\left(\frac{\beta_{21}}{\alpha_{11}}(S_1^{in} - S_1^{nec}) - S_2\right)$$

which implies

$$S_3^{av}(\mathbf{E}_2) = \frac{\beta_{32}}{\alpha_{22}}\left(\frac{\beta_{21}}{\alpha_{11}}(S_1^{in} - S_1^{nec}) - S_2^{nec}\right) \quad (17)$$

By Corollary 2 $S_3^{av}(\mathbf{E}_2) > S_3^{nec}$, then $\mathbf{E}_3$ exists and we can continue by induction.

To make the induction easier, we will make the change of variables

$$s_1^{nec} := \frac{S_1^{nec}}{S_1^{in}}$$

$$s_i^{nec} = \frac{\alpha_{i-1,i-1}\alpha_{i-2,i-2} \cdots \alpha_{11}S_1^{in}}{\beta_{i,i-1}\beta_{i-1,i-2} \cdots \beta_{2,1}}S_i^{nec} \text{ for } i \geq 2.$$

With these new variables a short calculation shows that

$$S_i^{av}(\mathbf{E}_{i-1}) = \frac{\beta_{i,i-1}\beta_{i-1,i-2} \cdots \beta_{2,1}}{\alpha_{i-1,i-1}\alpha_{i-2,i-2} \cdots \alpha_{11}S_1^{in}}\left(1 - \sum_{j=1}^{i-1}s_j^{nec}\right). \quad (18)$$

This formula allows us to sequentially calculate how far down the chain the species survive. The chain will end at the first species i which satisfies

$$S_{i+1}^{av}(\mathbf{E}_i) \leq S_{i+1}^{nec}.$$

7 Discussion

In this paper we formulate and study simple food webs, where each microbial species depends on a dedicated resource that is either supported externally or by other species. Although the real consortia are much more complex, involve mutualistic as well as antagonistic relationships, and often have multiple alternative food sources, our analysis allows a rather complete understanding of which consortia can be supported in a simple food web.

Our motivation comes from trying to understand co-existence principles that govern natural, evolved (Rosenzweig et al (1994); Helling et al (1987); Treves et al (1998); Rozen and Lenski (2000)) and synthetic (Bernstein et al, 2012) microbial consortia. Synthetic consortia allows to test experimentally in simplified settings principles that apply in much more complex interactions in microbial communities, as well as to test predictions of mathematical theory.

We showed that there is at most one consortium of each type, that is with the same set of microbial species present (Corollary 1). Furthermore, which communities are sustainable depend on a simple condition that summarizes sufficiency of supplied resources (Theorem 2). Finally, we show that the only stable community is the one that has maximum number of species present for given supply of resources (Theorem 5). We also show that such a community maximizes the overall biomass over all sustainable communities, which supports the maximal power principle (Lotka, 1922; Odum and Pinkerton, 1955; de Mazancourt and Schwartz, 2010; Doebeli, 2002; Pfeiffer and Bonhoeffer, 2004; Bull and Harcombe, 2009; Estrela and Gudejl, 2010; Beardmore et al, 2011).

Our results apply to simple prototypes of food webs: chains and fan-like food webs. For both we derive conditions that characterize the stable equilibrium in each system.

Real consortia and microbial communities are clearly more complex than those studied here; however, we believe that the framework developed in this paper can be used to study communities with syntrophic, as well as indirect antagonistic interactions.

Acknowledgements

We would like to thank Jeff Heys and Ross Carlson for many stimulating discussions on microbial consortia. We would also like to thank anonymous referees whose comments significantly improved the presentation of the paper.

This work was partially supported by NSF grant DMS-1361240.

References

- Aota Y, Nakajima H. Mutualistic relationships between phytoplankton and bacteria caused by carbon excretion from phytoplankton. *Ecological Research*. 2001; 16:289–299.
- Beardmore R, Gudelj I, Lipson D, Hurst L. Metabolic trade-offs and the maintenance of the fittest and the flattest. *Nature*. 2011; 472:342–346. [PubMed: 21441905]
- Bernstein H, Paulson S, Carlson R. Synthetic *Escherichia coli* consortia engineered for syntrophy demonstrate enhanced biomass productivity. *Journal of Biotechnology*. 2012; 157:159–166. [PubMed: 22015987]
- Bull J, Harcombe W. Population dynamics constrain the cooperative evolution of cross-feeding. *PLoS One*. 2009; 4:e4115. [PubMed: 19127304]
- Burchard A. Substrate degradation by a mutualistic association of two species in the chemostat. *Journal of Mathematical Biology*. 1994; 32:465–489.
- DeLong JP. The maximum power principle predicts the outcomes of two-species competition experiments. *Oikos*. 2008; 117(9):1329–1336. URL <http://dx.doi.org/10.1111/j.0030-1299.2008.16832.x>.
- Doebeli M. A model for the evolutionary dynamics of cross-feeding polymorphisms in microorganisms. *Population Ecology*. 2002; 44:59–70.
- Eiteman M, Lee S, Altman E. A co-fermentation strategy to consume sugar mixtures effectively. *Journal of Biological Engineering*. 2008; 2:3. [PubMed: 18304345]
- Elkhader A. Global stability in a syntrophic chain model. *Mathematical Biosciences*. 1991; 104:203–245. [PubMed: 1804461]
- Estrela S, Gudejl I. Evolution of cooperative cross-feeding could be less challenging than originally thought. *PLoS One*. 2010; 5:e14, 121.
- Fierer N, Jackson R. The diversity and biogeography of soil bacterial communities. *PNAS*. 2006; 103:626–631. [PubMed: 16407148]
- Gill S, Pop M, DeBoy R, Eckburg P, Turnbaugh P, Samuel B, Gordon J, Relman D, Fraser-Liggett C, Nelson K. Metagenomic analysis of the human distal gut microbiome. *Science*. 2006; 312:1355–1359. [PubMed: 16741115]
- Helling R, Vargas C, Adams J. Evolution of *Escherichia coli* during growth in a constant environment. *Genetics*. 1987; 116:349–358. [PubMed: 3301527]
- Kassen R, Buckling A, Bell G, Rainey P. Diversity peaks at intermediate productivity in a laboratory microcosm. *Nature*. 2000; 406:508–512. [PubMed: 10952310]
- Katsuyama C, Nakaoka S, Takeuchi Y, Tago K, Hayatsu M, Kato K. Complementary cooperation between two syntrophic bacteria in pesticide degradation. *Journal of Theoretical Biology*. 2009; 256:644–654. [PubMed: 19038271]
- Lotka AJ. Contribution to the energetics of evolution. *Proc Natl Acad Sci USA*. 1922; 8:147, 151. [PubMed: 16576642]
- de Mazancourt C, Schwartz MW. A resource ratio theory of cooperation. *Ecology Letters*. 2010; 13:349–359. [PubMed: 20455920]
- Odum HT, Pinkerton RC. Times speed regulator: the optimal efficiency for maximum power output in physical and biological systems. *Am Sci*. 1955; 43:321, 343.
- Peralta-Yahya P, Zhang F, del Cardayre S, Keasling J. Microbial engineering for the production of advanced biofuels. *Nature*. 2012; 488:320–328. [PubMed: 22895337]
- Pfeiffer T, Bonhoeffer S. Evolution of cross-feeding in microbial populations. *The American Naturalist*. 2004; 163:E126–E135.
- Powell G. Stable coexistence of syntrophic associations in continuous culture. *Journal of Chemical Technology and Biotechnology*. 1985; 35B:46–50.
- Powell G. Stable coexistence of syntrophic chains in continuous culture. *Theoretical Population Biology*. 1986; 30:17–25.
- Reilly P. Stability of commensalistic systems. *Biotechnology and Bioengineering*. 1974; 16:1373–1392. [PubMed: 4429794]

- Rosenzweig F, Sharp R, Treves D, Adams J. Microbial evolution in a simple unstructured environment: Genetic differentiation in *Escherichia coli*. *Genetics*. 1994; 137:903–917. [PubMed: 7982572]
- Rozen D, Lenski R. Long-term experimental evolution in *Escherichia coli*. VIII. Dynamics of a balanced polymorphism. *The American Naturalist*. 2000; 155:24–35.
- Sari T, El Hajji M, Harmand J. The mathematical analysis of a syntrophic relationship between two microbial species in a chemostat. *Mathematical Biosciences and Engineering*. 2012; 9:627–645. [PubMed: 22881029]
- Schink B. Energetics of syntrophic cooperation in methanogenic degradation. *Microbiology and molecular biology reviews*. 1997; 61:262–280. [PubMed: 9184013]
- Seitz H, Schink B, Pfennig N, Conrad R. Energetics of syntrophic ethanol oxidation in defined chemostat cocultures- 1. Energy requirement for H₂ production and H₂ oxidation. *Archives of Microbiology*. 1990a; 155:82–88.
- Seitz H, Schink B, Pfennig N, Conrad R. Energetics of syntrophic ethanol oxidation in defined chemostat cocultures- 2. Energy sharing in biomass production. *Archives of Microbiology*. 1990b; 155:89–93.
- Shong J, Diaz M, Collins C. Towards synthetic microbial consortia for bioprocessing. *Current Opinion in Biotechnology*. 2012; 23:1–5. [PubMed: 22244690]
- Smith H, Li B. Competition for essential resources: A brief review. *Fields Institute Communications*. 2003; 3:213–227.
- Tilman, D. *Resource Competition and Community Structure*. Princeton, N.J., U.S.A.: Princeton University Press; 1982.
- Treves D, Manning S, Adams J. Repeated evolution of an acetate-crossfeeding polymorphism in long-term populations of *Escherichia coli*. *Molecular Biology and Evolution*. 1998; 15:789–797. [PubMed: 9656481]
- Venail P, MacLean R, Bouvier T, Brockhurst M, Hochberg M, Mouquet N. Diversity and productivity peak at intermediate dispersal rate in evolving metacommunities. *Nature*. 2008; 452:210–214. [PubMed: 18337821]
- Venter J, Remington K, Heidelberg J, Halpern A, Rusch D, Eisen J, Wu D, Paulsen I, Nelson K, Nelson W, Fouts D, Levy S, Knap A, Lomas M, Nealson K, White O, Peterson J, Hoffman J, Parsons R, Baden-Tillson H, Pfannkock C, Rogers Y, Smith HO. Environmental genome shotgun sequencing of the Sargasso Sea. *Science*. 2004; 304:66–74. [PubMed: 15001713]
- Zuroff T, Curtis W. Developing symbiotic consortia for lignocellulosic biofuel production. *Applied Microbiology and Biotechnology*. 2012; 93:1423–1435. [PubMed: 22278256]

Fig. 1.
An Example of a Simple Food web.

Fig. 2.
A Chain Consortium (left) and a Hub Consortium (right).