Published in final edited form as:

Psychiatry Res. 2015 December 15; 230(2): 553–560. doi:10.1016/j.psychres.2015.10.003.

Population heterogeneity of trait anger and differential associations of trait anger facets with borderline personality features, neuroticism, depression, Attention Deficit Hyperactivity Disorder (ADHD), and alcohol problems

Gitta H. Lubke^{a,b}, Klaasjan G. Ouwens^b, Marleen H.M. de Moor^c, Timothy J. Trull^d, and Dorret I. Boomsma^b

^aDepartment of Psychology, University of Notre Dame, USA ^bBiological Psychology, VU University Amsterdam, The Netherlands ^cClinical Child and Family Studies, VU University Amsterdam, The Netherlands ^dPsychological Sciences, University of Missouri

Abstract

Anger is an emotion consisting of feelings of variable intensity ranging from mild irritation to intense fury. High levels of trait anger are associated with a range of psychiatric, interpersonal, and health problems. The objectives of this study were to explore heterogeneity of anger as measured by the Spielberger Trait Anger Scale (STAS), and to assess the association of the different anger facets with a selection of psychiatric disorders covering externalizing and internalizing problems, personality disorders, and substance use. Factor mixture models differentiated between a high and low scoring class (28% vs. 72%), and between three factors (anger-temperament, anger-reaction, and immediacy of an anger response). Whereas all psychiatric scales correlated significantly with the STAS total score, regressing the three STAS factors on psychiatric behaviors model showed a more detailed pattern. Only borderline affect instability and depression were significantly associated with all three factors in both classes whereas other problem behaviors were associated only with 1 or 2 of the factors. Alcohol problems were associated with immediacy only in the high scoring class, indicating a non-linear relation in the total sample. Taking into account these more specific associations is likely to be beneficial when investigating differential treatment strategies.

Keywords

factor mixture modeling; heterogeneity; associations of anger factors with psychiatric outcomes

Corresponding Author: Gitta Lubke, 110 Haggard Hall, University of Notre Dame, Notre Dame, IN 46556, glubke@nd.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Anger is an emotion consisting of feelings of variable intensity ranging from mild irritation to intense fury or rage (Spielberger et al., 1983). Anger can be either suppressed or directed at self (anger-in) or expressed outwardly through aggressive or confrontational behavior (anger-out). High levels of trait anger are associated with a range of psychiatric, interpersonal, and health problems, and in some cases anger can lead to aggression and violence (McDermut et al., 2009; Owen, 2011). Although clinically relevant in its own right, high levels of anger are significantly associated with a range of psychiatric disorders, including mood disorders, anxiety disorders, personality, and substance use disorders (DiGiuseppe et al., 2012; Kassino and Tafrate, 2006; McDermut et al., 2009). Furthermore, anger is a central feature and diagnostic criterion for borderline personality disorder (American Psychiatric Association, 2013).

Anger has been described in terms of cognitive, emotional, and behavioral features, and several anger questionnaires attempt to capture these different aspects (Eckhardt et al., 2004; Owen, 2011; Smith et al., 2004). However, most investigations of the clinical correlates of anger have limited their focus to those correlates of aggregate measures of trait anger, failing to examine whether there are specific aspects of anger that show differential relations to common psychiatric problems. The present study focuses on aspects of anger measured by the Spielberger Trait Anger Scale (STAS, Spielberger et al., 1983). The STAS measures the disposition to have angry feelings. Factor analyses of the STAS have consistently shown that this scale is not unidimensional, but consists of two distinct facets, namely anger temperament and anger reaction. The STAS items measuring anger temperament are designed to refer to more frequent and intense feelings of anger that occur in a broader range of situations whereas anger reaction more narrowly refers to experiencing anger when negatively evaluated or mistreated by others (Spielberger et al., 1983). Analyses of the Dutch version point to the possibility of a third factor measuring the immediacy of an anger response, which is most likely due to the fact that the Dutch version is not a literal translation (van der Ploeg et al., 1982). Three items of the English STAS have been replaced, and two of these items are designed to tap into the immediacy of an anger response (van der Ploeg et al., 1982). The developers report a two-factor structure of the Dutch STAS, with a distinction between anger-temperament and anger-reaction, and suggest the two items concerning the immediacy of an anger response to be part of anger-temperament for subjects with high levels of Trait Anger but part of anger-reaction for subjects with low levels Trait Anger (van der Ploeg et al., 1982). These differences between subgroups were not directly investigated since statistical methods designed for this purpose were not available at the time, however, this issue is addressed in the current study.

One aim of the present study is to explore potentially different associations of the different STAS anger factors with a selection of psychiatric problems that are known to co-occur with anger, or feature anger as a symptom. Specifically, we investigated whether the anger factors show differential associations to a range of psychiatric problems from each of the following domains: externalizing and internalizing problems (ADHD and depression, respectively), substance use (alcohol problems), and maladaptive personality (borderline personality affect instability, identity problems, self-harm, negative relations, and

neuroticism). Although by no means comprehensive, this selection provides an excellent opportunity to investigate the potentially differential associations of anger temperament, anger reaction, and the immediacy of an anger response. Since neuroticism is highly associated with a total score of borderline personality features (Distel et al., 2009), we also included neuroticism to further explore similarities and differences in the association with anger factors compared to the different aspects of borderline personality features.

Our second goal was to investigate potential heterogeneity. Population heterogeneity refers to the presence of different clusters in the population that differ qualitatively (e.g., differential profiles on aspects of anger, differential comorbidities) and/or quantitatively (e.g., severity of symptomology) (Lubke and Muthén, 2005). Studies of population heterogeneity of anger are very limited (but see He et al., 2013). Given the importance of anger as a health risk factor (Chida and Steptoe, 2009), a detailed understanding of the presentation of anger is critical for clinical research. To investigate population heterogeneity of the different aspects of anger it is necessary to carry out item-level analyses rather than using total scores. In this study we modeled item-level data of the Dutch version of the STAS with factor mixture models, and assessed differential associations of the different anger facets in each of the latent classes. Factor mixture modeling is a recently developed statistical framework that permits estimating a specific factor structure within different clusters (aka latent classes) of subjects in the population (Lubke et al., 2013, Tueller and Lubke, 2010). Factor mixture modeling combines latent class analysis and factor analysis in a single modeling framework (Lubke and Muthén, 2005). Latent class analysis attributes overall covariation between questionnaire items entirely to mean differences between latent classes while constraining within class covariances of items to be zero. Factor analysis attributes overall covariation to continuous underlying factors that represent severity, and assumes a single homogeneous population. Factor mixture models are hybrids that permit mean differences between latent classes as well as severity differences in factors within each class. Most importantly, the clusters or classes reflect a grouping of subjects with similar response profiles that is data-driven. The factor structure and the response profiles of the items can differ across clusters. The classes can be compared with respect to qualitative differences (characteristic response profiles, differential associations with covariates), and quantitative differences (severity of symptomology).

In summary, in order to examine the differential associations of a range of psychiatric problems with different aspects of anger, we assessed the heterogeneity of anger, as measured by the STAS, using factor mixture modeling in a large population sample drawn from the Netherlands Twin Register (NTR) (Willemsen et al., 2013). We also addressed the question whether the Dutch version of the STAS is more adequately modeled with two or with three factors. Next, we investigated differences between classes regarding the association of the different STAS anger factors with borderline personality features, ADHD, depression, neuroticism, and alcohol problems.

2. Methods

2.1 Participants

The subjects were participants of the Netherlands Twin Register (NTR), a population-based sample, which is ascertained based on the presence of twins or higher-order multiples in the family (Willemsen et al., 2013). Data collection in 2 to 5-year intervals started 25 years ago. Most data included in the present study were collected between 2004 and 2005 (survey 7), with the exception of depression, which was collected in survey 8 (between 2009 and 2012, see below). The total number of participants with valid data for analysis was 3821 men, and 6451 women.

2.2 Measures and Descriptives

Trait anger was measured with the Dutch adaptation of the STAS (Spielberger et al., 1983; van der Ploeg et al., 1982). As described by van der Ploeg et al. (1982), the scale items are formulated to explicitly tap into a more general disposition rather than into feelings of anger at a given time. Three items of the Dutch version are not literal translations of the English version. Based on factor analyses and validity studies during the adaptation of the STAS, items “angry when slowed by others”, “annoyed when not given recognition”, and “feel like hitting someone” were replaced by “I feel quickly angry”, “I am hot-tempered”, and “I am quickly irritated”. The developers report a two-factor structure distinguishing between anger-temperament (items 1,2,3,5,6) and anger-reaction (items 7, 8 and 10), which mirrors the structure of the English version of the STAS. Items 4 and 9 (“I feel quickly angry”, “I am quickly irritated”)¹ measure the immediacy of an anger response. In our analyses we compared the suggested 2-factor structure as well as a 3-factor structure where items 4 and 9 load on a separate factor. By specifying 2 or, alternatively, 3 factors within each class, and leaving factor variances to be class-specific, we can obtain detailed information about this aspect of anger in the different latent classes.

Summary statistics of the ten items are presented in Tables 1 and 2. Responses were moderately skewed, and rates of the 4 response categories of each item were similar for males and females (see Table 1). The STAS has good discriminant and convergent validity as well as clinical utility, and has been administered across a wide range of subject populations and psychological domains (Chida and Steptoe, 2009; Eckhardt et al., 2004; van der Ploeg et al., 1982).

As suggested by the developers of the Dutch STAS our analyses are limited to subjects who had answered at least 7 of the 10 items although personality and health items in the NTR are relatively unbiased with respect to non-response (Distel et al., 2007; van der Ploeg et al., 1982). Age ranged from 16 to 90 years in males (median 42 years), and from 16.5 to 88 years (median 42 years) for females. Correlations of the anger items with age were small but in our large sample nevertheless significant. On average, age explained 0.3% of the variance of anger items in males, and 6% of the variance in females. This is consistent with previous

¹Note that the Dutch wording is much less similar than implied by the English translation.

results (Hakulinen et al., 2013; Okuda et al., 2015; Phillips et al., 2006). Age was therefore included as a covariate in all analyses.

Borderline Personality Disorder (BPD) features were measured with the Personality Assessment Inventory Borderline Features Scale (PAI-BOR, Morey, 1991). The 24 PAI-BOR items are grouped into 4 subscales measuring affect instability (AI), identity problems (IP), negative relations (NR), and self-harm (SH). The test-retest reliability of the PAI-BOR in the Dutch data is 0.78 (Distel et al., 2009). We used the four subscales because BPD features are known to be heterogeneous (Hallquist & Pilkonis, 2012; Lenzenweger et al., 2008; Lubke et al., 2014; Skodol et al., 2002). ADHD was measured with the ADHD index, which aggregates the 12 items of the Conner's Adult ADHD Rating Scale corresponding to DSM symptoms. The 6-week test-retest reliability is 0.67 (Boomsma et al., 2010). Neuroticism was measured with the neuroticism subscale of the NEO five-factor inventory (NEO-FFI), which consists of 12 items (Costa and MacCrae, 1992). Reliability of the NEO-FFI scales in a Dutch speaking population is between 0.68 and 0.86 (de Fryt & Mervielde, 1998). Alcohol Problems were assessed with the four CAGE items, which have a reported reliability of 0.85–0.95 (Mayfield et al., 1974; Dhalla and Kopec, 2007). We used a dichotomized version of the aggregate CAGE score, as a score of 2 or higher is indicative of alcoholism (Ewing, 1984; Geels et al., 2013). Finally, depression was measured with 10 items of the Adult Self Report that correspond to DSM depression symptoms (Achenbach, 1997). This measure was obtained during wave 8 of the NTR data collection, and has a reliability of 0.85 (de Moor et al. 2006).

Correlations between the 10 STAS items are shown in Table 2, and range between 0.22 and 0.62. Initial exploratory factor models fitted separately to data from males and females resulted in almost identical factor structures, and favored a 3-factor solution. The three factors also showed a very similar pattern of associations with the psychiatric problems for males and females. All models were therefore fitted to data of the two genders jointly.

The correlations among PAI-BOR scales, neuroticism, ADHD, depression, and alcohol problems were generally small to moderate, and, therefore, were not expected to cause multi-collinearity problems (see Table 3). The correlation table includes a sum-score of the 10 STAS items to show the overall association of anger with these behaviors. The mixture analyses described below aimed at providing a more detailed picture regarding aspects of anger as they relate to different psychopathology.

2.3 Analyses

2.3.1. Type 1 error control—The investigation of heterogeneity involved fitting multiple mixture models to the data, and the association assessment involved estimating multiple regression coefficients. Although in the area of mixture modeling it is not yet widely recognized, type 1 error is inflated if coefficients are estimated in a selected model (Hurvich and Tsai, 1990). The inflation of type 1 error due to model selection is comparable to multiple stage clinical trials, where well-performing drugs are selected for subsequent stages of a study. Not accounting for the selection implies capitalizing on chance, which in turn leads to type 1 error inflation. The model selection has therefore to be taken into account to keep type 1 error under control. We therefore adopted the following analytic strategy.

Benefitting from our extremely large sample, we randomly selected $N=2,499$ individuals, and fitted a series of alternative latent class and factor mixture models with 1–3 factors within class and an increasing number of classes. The best-fitting model was selected, and then fitted to the data in the remaining sample in order to estimate the regression coefficients of the factors on the psychiatric behaviors.

The Bayesian Information Criterion, BIC, has been shown to be one of the best indices to compare alternative mixture models (Schwarz, 1978; Nylund et al., 2007). We note that model selection based on fit indices such as the BIC is to some extent arbitrary since all fit indices depend on sample size, and hard cut-offs therefore do not exist (Joreskog, 1993). Our model selection is based on extensive prior simulation work of our team (Lubke and Neale, 2006, 2008, Tueller and Lubke, 2010) as power simulations using the current models and sample size are beyond computational feasibility. Model selection and estimation of regression coefficients was done after splitting the data into two separate sets, thus avoiding capitalizing on chance. We controlled for testing multiple covariates by using a more stringent significance level. A procedure to obtain an adequate significance level for correlated tests was implemented in the freely available program matSpD (Nyholt, 2004). Non-independence of observations due to family membership (e.g., our data included twins) was taken into account by using a sandwich estimator in all analyses (Freedman, 2006). This is implemented in Mplus with option “mixture complex”, and using a family indicator as grouping variable, Muthén and Muthén, 1998–2012). Age was used as class-predicting covariate to correct for age influence on class membership (Phillips et al., 2006; Hakulinen et al., 2013; Okuda et al., 2015).

2.3.2 Analysis Part 1: Assessment of population heterogeneity of anger using a random subsample of the data ($N=2,499$)—The analysis plan comprised of fitting latent class models and factor mixture models to the 10 Dutch STAS items observed in male and female participants. The 2-factor structure was based on previous studies using the Dutch version of the STAS, and the 3-factor structure, which was based on initial exploratory factor analyses, had items 4 and 9 (“quickly angry”, and “quickly irritated”) load on a separate additional factor. The number of random starts was set to 200, with 20 sets iterating to final convergence. If the likelihood was not replicated we increased the number of starts.

2.3.3 Analysis Part 2: Associations between anger factors and BPD features, ADHD, neuroticism, depression, and alcohol problems in the remaining sample with available covariate data—We planned to fit the best fitting model from step 1 of our analysis to the remaining sample to investigate the associations with covariates while keeping Type 1 error under control. In this model, the factors were regressed on the selected psychiatric behaviors within each class. Regression coefficients were estimated specific for each class. The sample with data for BPD features, neuroticism, ADHD, and alcohol problems was $N=9,763$. The sample with available data for depression was smaller, $N=5,384$. Note that we did not correct for potential item overlap (e.g., between anger, borderline features, and neuroticism scales) because the aim was to investigate whether

different aspects of anger are differentially associated with behaviors such as neuroticism *as commonly defined and measured*.

All models were fitted in Mplus 7.11 (Muthén, and Muthén, 1998–2013). We conducted appropriate Wald tests to assess whether the class differences in the associations of anger factors and covariates were significant.

3. Results

3.1 Analysis Part 1: Population Heterogeneity of Anger

The results of the models that were fitted to select the best fitting model using a subset of the data (N=2,499) are presented in Table 4. As expected, latent class analysis models (LCA models) did not fit the data as well as the factor mixture models, which allow for severity differences in anger.

All factor mixture models were fitted with class-specific factor variances and covariances. We also permitted class-specific loadings and thresholds to allow for deviations from measurement invariance, which is difficult to test in the context of mixture models (Millsap and Tein, 2004, Lubke and Neale, 2008). We found that factor mixture models with 3 factors provided a better fit than 2-factor models, thus separating the facets anger temperament, anger reaction, and immediacy of an anger response. The best fitting models were 3-factor single class and 3-factor 2-class models. Although the BIC was somewhat smaller for the single class model we expected the 2-class model to have superior fit in the larger confirmatory sample due to the much better resolution for the detection of classes in larger samples (see Lubke and Neale, 2006, 2008). We therefore selected the 3-factor 2-class model as the more adequate final model. This model was then fitted to the data not used for model selection in analysis Part 2 (see below).

The class proportions were 28% for the smaller high scoring class, and 72% for the lower scoring majority class. The response profiles on the ten items are presented in Figure 1, which shows that the profiles are reasonably parallel for the probability of scoring zero. However, the probability of scoring either zero or one differs across classes (see Figure 1, middle panel). This is the result of differences in item thresholds, which were estimated as class-specific. The higher scoring class provides more information concerning a differential pattern across the ten items for the higher response categories as the lower scoring class does not endorse categories higher than 1 on any of the items. There were no substantial differences in factor variances and covariances across the two classes. Factor correlations within class ranged between 0.35 and 0.68, with the lowest correlation being between temperament and reaction, and the highest between temperament and immediacy. The regression on age was significant (standardized regression coefficient $\beta=8.399$, p -value<0.001); younger participants were more likely members of the higher scoring class.

3.2 Analysis Part 2: Differences between classes regarding BPD features, ADHD, neuroticism, alcohol problems, and depression

A 2-class 3-factor model was fitted to the remaining data. In this model the three factors were regressed on the covariates in both classes. Although factor variances and covariances

showed only small differences between classes in the first part of the analysis, these parameters were kept as class-specific to avoid unnecessary bias in the estimation of regression coefficients. All reported regression coefficients were estimated simultaneously in this model. The resulting class proportions and within class parameters of this model were very similar to the model fit using $N=2,499$. (e.g., loadings correlated 0.95, see Table 5).

The results of the regression of the three anger factors on the psychiatric behaviors are shown in Table 6. The estimated effects were selected based on initial exploratory analyses in the smaller sample where the three factors were regressed on all covariates. The cutoff was a regression coefficient of 0.05. The selection was done to avoid the risk of including many nonsignificant covariate effects, which can lead to instability and unreliable results (Tueller and Lubke, 2010). The significance level was adjusted for multiple testing as implemented in the freely available program *matSpD* to $\alpha=0.0036$ (Nyholt, 2004). With the exception of alcohol problems and borderline negative relations (see below), the *differential pattern of associations* was the same in the high scoring minority class and the lower scoring majority class.

Only borderline affect instability and depression had consistent effects across the three anger factors. All other behaviors showed specific pattern of associations with the three different anger factors. Borderline self-harm was associated with anger temperament in both classes, whereas borderline negative relations was associated with anger reaction. The latter effect was observed only in the lower scoring majority class, but it should be noted that the low scoring class was about three times as large, thus affording higher statistical power. Borderline identity problems were not significantly associated with any anger facets. Neuroticism was only associated with the immediacy of an anger response, and the effect was somewhat more pronounced in the high scoring class. ADHD was related to anger reaction in both classes, and also to anger immediacy in the lower scoring majority class. Alcohol problems were related to the immediacy of an anger response. This effect was observed in the high scoring class, but was absent in the low scoring majority class. This result indicates that the overall effect is non-linear, that is, the effect varies with severity of the immediacy of an anger response.

We conducted Wald tests to investigate whether the class differences in associations of the three anger factors were significant. This was done using the Mplus option “Model Test”, which evaluates whether constraining the regression coefficients to be the same across classes is tenable. For the immediacy factor, regression coefficients were significantly different across classes (Wald statistic=21.68, p -value= 0.0014). For the other two factors the differences were not significant, which is consistent with the more similar pattern of associations shown in Table 6. We also tested whether class was moderating the associations, and found that the regressions of class on the covariates were all close to zero and non-significant. Therefore we can conclude that class is not moderating the associations.

4. Discussion

To our knowledge this is the first study that explores the associations between the different STAS anger factors and PAI-BOR subscales, neuroticism, depression, ADHD, and alcohol

problems in more detail in a population sample. These behaviors are known to be associated with anger, and were chosen as examples of personality problems, externalizing and internalizing problems, and substance use problems. Comparing the overall correlations of these behaviors with the STAS total score (see Table 3) with the more detailed results from our analysis (see Table 6) reveals a more distinct structure of associations that is hidden when considering anger as a total score.

4.1. Anger presents as a heterogeneous phenotype

The factor mixture analysis of the 10 STAS items revealed two latent classes. The classes differed quantitatively with respect to the severity of the scores on the 10 STAS items. The smaller, higher-scoring class comprised about 28% of the sample. The classes differed qualitatively with respect to the within class associations of the STAS anger factors with the psychiatric behaviors. In addition to replicating the two factors anger temperament and anger reaction that are known to the STAS items, we found that separating a third factor measured by the two items concerning the immediacy of an anger response (“I feel angry quickly”, “I am quickly irritated”) benefitted the model fit. It should be noted that the selection of a best fitting model is a matter of power. Especially when analyzing categorical outcomes a model with an additional class requires the estimation of a much larger number of parameters, which can lead to rejection when using the BIC for model comparisons (Lubke and Neale, 2008). Given that with respect to anger severity the classes differed mainly quantitatively, it will depend on the goals of future research whether it is useful to account for the apparent grouping into higher and lower scoring participants. In this context it is noteworthy that the response profiles on the 10 STAS items in the two classes showed that the higher scoring class differed especially with respect to one of the anger immediacy items (see Figure 1), and that the classes differed significantly with respect to the associations of the immediacy factor (see below).

4.2. Anger factors are differentially related to multiple psychiatric problems

The patterns of associations shown in Table 6 are consistent with both clinical observations and empirical findings for these psychiatric problems. First, the PAI_BOR affect instability scale and the depression scale reflecting DSM depression symptoms were the only scales with consistent effects across all three anger factors. Anger is central feature of Borderline Personality Disorder (BPD), and is a core emotion that manifests itself in the affective instability characterizing BPD (Trull et al., 2008). Furthermore, angry temperament, angry reactions to others, and the immediacy of anger expression are all common clinical features of BPD. Concerning depression, psychodynamic theories posit that depression can be conceptualized as chronic anger turned inward (Busch, 2009), and anger and irritability are often abruptly and intensely expressed by depressed individuals in their interpersonal relationships (perhaps due to high levels of rejection sensitivity) often leading to social dysfunction.

All other scales measuring personality and psychiatric problems showed more specific patterns of interrelations with the different aspects of anger. Notably, the NEO neuroticism scale, which correlates highly with borderline affect instability, was associated only with the immediacy of an anger response rather than showing a broader association with anger. This

is likely due to the NEO scale's assessment of a range of negative affects (e.g., anxiety, dysphoria) in addition to anger. ADHD was not related to anger temperament, but rather to anger reaction, and also anger immediacy. This suggests that anger among those with ADHD is most likely to manifest itself in interpersonal contexts, and that the impulsivity associated with ADHD may lead to very abrupt and intense anger expressions. The latter may reflect some deficits in cognitive control and in emotion regulation amongst those with ADHD (Shaw et al., 2014). Interestingly, alcohol problems were associated with immediacy only in the high scoring class. Clearly, this result contradicts a simple linear relation between anger and alcohol problems where increasing anger symptoms are accompanied by increasing alcohol problems. This result is noteworthy because the high scoring class differed significantly from the lower scoring class only with respect to the associations of the immediacy factor, and is interesting in the context of previous findings that in individuals with low levels of trait anger the relation between alcohol and aggression was less pronounced than in individuals with higher levels of trait anger (Parrot and Zeichner, 2002).

The differential pattern of associations of the studied comorbid behaviors with the different aspects of anger is an important finding that can inform future studies in clinical samples to investigate differential treatment strategies (e.g., anger management, cognitive-behavioral treatment, relaxation therapy; Del Vecchio and O'Leary, 2004). Our results indicate that taking into account specific patterns of associations at the level of more narrow defined factors rather than the level of broader domains is likely to be beneficial when defining symptom profiles.

4.3. Strength and Limitations

This study is the first to investigate population heterogeneity of anger and the association of the different clusters to a number of related psychiatric disorders. An important strength of the current study is the large population based sample drawn from the Netherlands Twin Register (NTR) used for the analyses (Willemsen et al., 2013). A limitation of our study is that all measures are selfreports, and do not take into consideration possible psychiatric diagnoses or treatments. In addition, cluster solutions concerning self-reports of anger may at least to some extent be culture dependent, and generalizations beyond the Dutch population should be made with caution.

Acknowledgements

GH Lubke was supported by NIH DA-018673. MHM de Moor was supported by the Netherlands Organization for Scientific Research (NWO) (VENI-016-115-035). We acknowledge funding for data collection from the Netherlands Organization for Scientific Research (NWO) and MagW/ZonMW): Twin family database for behavior genomics studies (480-04-004); Genetic determinants of risk behavior in relation to alcohol use and alcohol use disorder (ZonMw Addiction-31160008); Genotype/phenotype database for behavior genetic and genetic epidemiological studies (ZonMw Middelgroot 911-09-032); Spinozapremie (SPI 56-464-14192); CMSB: Center for Medical Systems Biology (NWO Genomics); NBIC/BioAssist (RK/2008.024); BBMRI -NL (184.021.007); Biobanking and Biomolecular Resources Research Infrastructure; the VU University: Institute for Health and Care Research (EMGO+) and Neuroscience Campus Amsterdam (NCA); the European Science Council (ERC Advanced, 230374)

References

- Achenbach, TM. Department of Psychiatry. Burlington, VT: University of Vermont; 1997. Manual for the Young Adult Self-report and Young Adult Behavior Checklist.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed.. Washington, DC: Author; 2013.
- Boomsma DI, Saviouk V, Hottenga JJ, Distel MA, de Moor MHM, Vink JM, Geels LM, Van Beek JHDA, Bartels M, de Geus EJC, Willemsen G. Genetic epidemiology of Attention Deficit Hyperactivity Disorder (ADHD Index) in adults. *PLOS ONE*. 2010
- Busch FN. Anger and depression. *Advances in Psychiatric Treatment*. 2009; 15:271–278.
- Chida Y, Steptoe A. The Association of Anger and Hostility With Future Coronary Heart Disease: A Meta-Analytic Review of Prospective Evidence. *Journal of the American College of Cardiology*. 2009; 53:936–946. [PubMed: 19281923]
- Costa PT, MacCrae RR. Revised NEO Personality Inventory (NEO PI-R) and NEO Five-Factor Inventory (NEO FFI): Professional Manual. Psychological Assessment Resources. 1992
- De Fruyt F, Mervielde I. The assessment of the big five in the Dutch language domain. *Psychologica Belgica*. 1998; 38:1–22.
- De Moor MHM, Beem AL, Stubbe JH, Boomsma DI, De Geus EJC. Regular exercise, anxiety, depression and personality: a population-based study. *Preventive Medicine*. 2006; 42:273–279. [PubMed: 16439008]
- Dhalla S, Kopec JA. The CAGE questionnaire for alcohol misuse: a review of reliability and validity studies. *Clinical & Investigative Medicine*. 2007; 30:33–41. [PubMed: 17716538]
- Del Vecchio T, O'Leary KD. Effectiveness of anger treatments for specific anger problems: A meta-analytic review. *Clinical Psychology Review*. 2004; 24:15–34. [PubMed: 14992805]
- DiGiuseppe R, McDermut W, Unger F, Fuller JR, Zimmerman M, Chelminski I. The comorbidity of anger symptoms with personality disorders in psychiatric outpatients. *Journal of Clinical Psychology*. 2012; 68:67–77. [PubMed: 21948109]
- Distel MA, Ligthart L, Willemsen G, Nyholt DR, Trull TJ, Boomsma DI. Personality, health and lifestyle in a questionnaire family study: a comparison between highly cooperative and less cooperative families. *Twin Research and Human Genetics*. 2007; 10:348–353. [PubMed: 17564524]
- Distel MA, Roeling MP, Tielbeek JJ, van Toor D, Derom CA, Trull TJ, Boomsma DI. The covariation of trait anger and borderline personality: A bivariate twin-siblings study. *Journal of Abnormal Psychology*. 2012; 121:458–466. [PubMed: 22149910]
- Distel MA, Trull TJ, Willemsen G, Vink JM, Derom CA, Lynskey M, Martin NG, Boomsma DI. The Five Factor Model of personality and borderline personality disorder: A genetic analysis of comorbidity. *Biological Psychiatry*. 2009; 66:1131–1138. [PubMed: 19748081]
- Eckhardt C, Norlander B, Deffenbacher J. The assessment of anger and hostility: A critical review. *Aggression and Violent Behavior*. 2004; 9:17–43.
- Ewing JA. Detecting alcoholism: The CAGE questionnaire. *JAMA The Journal of the American Medical Association*. 1984; 252:1905–1907. [PubMed: 6471323]
- Freedman DA. On the so-called “Huber sandwich estimator” and “robust standard errors”. *American Statistician*. 2006; 60:299–302.
- Geels LM, Vink JM, van Beek JHDA, Bartels M, Willemsen G, Boomsma DI. Increases in alcohol consumption in women and elderly groups: Evidence from an epidemiological study. *BMC Public Health*. 2013; 13:207. [PubMed: 23497391]
- Hakulinen C, Jokela M, Keltikangas-Järvinen L, Merjonen P, Raitakari OT, Hintsanen M. Longitudinal measurement invariance, stability and change of anger and cynicism. *Journal of Behavioral Medicine*. 2014
- Hallquist MN, Pilkonis PA. Refining the phenotype of borderline personality disorder: Diagnostic criteria and beyond. *Personality disorders: Theory, research, and treatment*. 2012; 3:228–246.

- He J, Qiu P, Park KY, Xu Q, Potegal M. Young Chinese children's anger and distress: Emotion category and intensity identified by the time course of behaviors. *International Journal of Behavioral Development*. 2013; 37:349–356.
- Hurvich CM, Tsai C. The Impact of Model Selection on Inference in Linear Regression. *The American Statistician*. 1990; 44:214–217.
- Joreskog, KG. Testing Structural Equation Models. In: Bollen, KA.; Long, JS., editors. *Testing Structural Equation Models*. Newbury Park, CA: Sage; 1993. p. 294–325.
- Kassinove, H.; Tafrate, RC. Anger-related disorders: Basic issues, models, and diagnostic considerations. In: Feindler, EL., editor. *Anger related disorders: A practitioner's Guide to Comprehensive Treatments*. New York, NY: Springer Publishing; 2006. p. 1–27.
- Lenzenweger MF, Clarkin JF, Yeomans FE, Kernberg OF, Levy KN. Refining the borderline personality disorder phenotype through finite mixture modeling: Implications for classification. *Journal of Personality Disorders*. 2008; 22:313–331. [PubMed: 18684047]
- Lubke GH, Laurin C, Amin N, Hottenga JJ, Willemsen G, van Grootheest G, Abdellaoui A, Karssen LC, Oostra BA, van Duijn CM, Penninx BWJH, Boomsma DI. Genome-wide analyses of borderline personality features. *Molecular Psychiatry*. 2014; 19:923–929. [PubMed: 23979607]
- Lubke GH, Muthén BO. Investigating population heterogeneity with factor mixture models. *Psychological Methods*. 2005; 10:21–39. [PubMed: 15810867]
- Lubke GH, Neale MC. Distinguishing between latent classes and continuous factors with categorical outcomes: Class invariance of parameters of factor mixture models. *Multivariate Behavioral Research*. 2008; 43:592–620. [PubMed: 20165736]
- McDermut W, Fuller JR, DiGiuseppe R, Chelminski I, Zimmerman M. Trait anger and Axis I disorders: Implications for REBT. *Journal of Rational-Emotive and Cognitive-Behavioral Therapy*. 2009; 27:121–135.
- Morey, LC. *The Personality Assessment Inventory: Professional Manual*. Odessa, FL: Psychological Assessment Resources; 1991.
- Muthén, LK.; Muthén, BO. *Mplus User's Guide Sixth Edition*. Los Angeles, CA: Muthen & Muthen; 1998–2012.
- Nyholt DR. A simple correction for multiple testing for SNPs in linkage disequilibrium with each other. *American Journal of Human Genetics*. 2004; 74:765–769. [PubMed: 14997420]
- Nylund KL, Asparouhov T, Muthén BO. Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Structural Equation Modeling*. 2007; 14:535–569.
- Okuda M, Picazo J, Olfson M, Hasin DS, Liu SM, Bernardi S, Blanco C. Prevalence and correlates of anger in the community: results from a national survey. *CNS spectrums*. 2015; 20:130–139. [PubMed: 25831968]
- Owen JM. Transdiagnostic cognitive processes in high trait anger. *Clinical Psychology Review*. 2011; 31:193–202.
- Parrott DJ, Zeichner A. Effects of alcohol and trait anger on physical aggression in men. *Journal of Studies on Alcohol*. 2002; 63:196–204. [PubMed: 12033696]
- Phillips LH, Henry JD, Hosie JA, Milne AB. Age, anger regulation and well-being. *Aging & Mental Health*. 2006; 10:250–256. [PubMed: 16777652]
- Rebollo I, Boomsma DI. Genetic analysis of anger: Genetic dominance or competitive sibling interaction. *Behavior Genetics*. 2006; 36:216–228. [PubMed: 16402285]
- Schwarz G. Estimating the dimensions of a model. *Annals of Statistics*. 1978; 6:461–464.
- Skodol AE, Gunderson JG, Pfohl B, Widiger TA, Livesley WJ, Siever LJ. The borderline diagnosis I: Psychopathology, Comorbidity, and Personality Structure. *Biological Psychiatry*. 2002; 51:936–950. [PubMed: 12062877]
- Shaw P, Stringaris A, Nigg J, Leibenluft E. Emotion dysregulation in attention deficit hyperactivity disorder. *American Journal of Psychiatry*. 2014; 171:276–293. [PubMed: 24480998]
- Smith TW, Glazer K, Ruiz JM, Gallo LC. Hostility, Anger, Aggressiveness, and Coronary Heart Disease: An Interpersonal Perspective on Personality, Emotion, and Health. *Journal of Personality*. 2004; 72:1217–1270. [PubMed: 15509282]

- Spielberger, CD.; Jacobs, G.; Russell, JS.; Crane, RS. Assessment of anger: the state-trait anger scale. In: Butcher, JN.; Spielberger, CD., editors. *Advances in Personality Assessment*. Vol. 2. Hillside, NJ: Erlbaum; 1983. p. 159-187.
- Trull TJ, Solhan MB, Tragesser SL, Jahng S, Wood PK, Piasecki TM, Watson D. Affective instability: Measuring a core feature of borderline personality disorder with ecological momentary assessment. *Journal of Abnormal Psychology*. 2008; 117:647–661. [PubMed: 18729616]
- Tueller S, Lubke GH. Evaluation of structural equation mixture models: Parameter estimates and correct class assignment. *Structural Equation Modeling*. 2010; 17:165–192. [PubMed: 20582328]
- Van der Ploeg, HM.; Defares, PB.; Spielberger, CD. Handleiding bij de Zelf-analyse Vragenlijst. The Netherlands: Swets & Zeitlinger, Lisse; 1982.
- Willemsen G, Vink JM, Abdellaoui A, den Braber A, van Beek JH, Draisma HH, van Dongen J, van 't Ent D, Geels LM, van Lien R, Ligthart L, Kattenberg M, Mbarek H, de Moor MHM, Neijts M, Pool R, Stroo N, Kluit C, Suchiman HED, Slagboom PE, de Geus EJC, Boomsma DI. The Adult Netherlands Twin Register: Twenty-Five Years of Survey and Biological Data Collection. *Twin Research and Human Genetics*. 2013; 16:271–281. [PubMed: 23298648]

Figure 1.

Response Profiles of the two classes on the 10 STAS Items

Note: The three panels depict the cumulative probabilities of scoring zero (upper panel), zero or one (middle panel), and zero, one or two (lower panel), respectively. The 10 STAS items are on the x-axis. The probability of scoring zero has a similar pattern for the two classes, but the higher scoring class endorses zero less often (see upper panel). Almost all participants in the lower scoring class have answered either zero or one on all items (see middle panel, probabilities for the low scoring class depicted in black are all close to 1). The higher scoring class also endorses higher response categories, and shows a differential

response pattern across items. The lower panel shows that the higher scoring class scores higher than two especially on items 4, 7, 8, and 10.

- Trait Anger can be modeled as a two-class three-factor structure
- Regressing the three factors on associated psychiatric behaviors shows a differential pattern
- Borderline affect instability and depression are associated with all three factors in both classes
- All other psychiatric behaviors are associated only with 1 or 2 of the factors.
- Results refute a linear relationship between anger symptoms and alcohol problems

Table 1

Response rates on the 10 STAS items for males and females

	almost never	sometimes	often	almost always	NA
1: "I am quick tempered"	0.569 / 0.611	0.381 / 0.342	0.044 / 0.041	0.003 / 0.003	0.002 / 0.003
2: "I have a fiery temper"	0.658 / 0.627	0.295 / 0.318	0.041 / 0.049	0.004 / 0.004	0.001 / 0.003
3: "I am a hotheaded person"	0.756 / 0.790	0.213 / 0.179	0.025 / 0.022	0.003 / 0.002	0.003 / 0.006
4: "I feel quickly angry"	0.365 / 0.313	0.532 / 0.573	0.094 / 0.104	0.006 / 0.007	0.003 / 0.004
5: "I am hot-tempered"	0.767 / 0.765	0.203 / 0.201	0.025 / 0.024	0.002 / 0.003	0.004 / 0.007
6: "I fly off the handle"	0.800 / 0.763	0.183 / 0.213	0.011 / 0.015	0.002 / 0.002	0.005 / 0.007
7: "when I get mad I say nasty things"	0.429 / 0.427	0.464 / 0.471	0.084 / 0.078	0.020 / 0.017	0.003 / 0.007
8: "it makes me furious when criticized in front of others"	0.465 / 0.422	0.464 / 0.501	0.056 / 0.057	0.011 / 0.013	0.003 / 0.007
9: "I am quickly irritated"	0.491 / 0.401	0.434 / 0.497	0.058 / 0.080	0.004 / 0.006	0.013 / 0.016
10: "I feel infuriated when I do a good job and get a poor evaluation"	0.335 / 0.287	0.527 / 0.572	0.104 / 0.105	0.033 / 0.032	0.002 / 0.004

Note: Male/female response rates are given in frequencies, and are separated by a slash. The content of items 4, 5, and 9 is different in the Dutch version of the STAS.

Table 2

Correlations between the 10 STAS items

	STAS1	STAS2	STAS3	STAS4	STAS5	STAS6	STAS7	STAS8	STAS9	STAS10
STAS1	1									
STAS2	0.61	1								
STAS3	0.62	0.59	1							
STAS4	0.43	0.43	0.38	1						
STAS5	0.50	0.56	0.58	0.45	1					
STAS6	0.43	0.44	0.43	0.35	0.46	1				
STAS7	0.26	0.27	0.24	0.28	0.26	0.29	1			
STAS8	0.22	0.23	0.21	0.29	0.25	0.24	0.33	1		
STAS9	0.43	0.43	0.38	0.62	0.43	0.36	0.29	0.30	1	
STAS10	0.25	0.24	0.23	0.30	0.25	0.24	0.31	0.44	0.3	1

Table 3

Data exploration: Correlations of Neuroticism, PAI-BOR subscales, ADHD index, DSM-depression, and CAGE alcohol problems for male and females prior to model fitting

	ang	neurot	bor_ai	bor_ip	bor_nr	bor_sh	adhd	depres	alcohol
ang	-	0.42	0.548	0.379	0.344	0.259	0.485	0.278	0.111
neurot	0.37	-	0.601	0.643	0.457	0.185	0.554	0.552	0.095
bor_ai	0.564	0.562	-	0.564	0.516	0.29	0.539	0.482	0.143
bor_ip	0.35	0.588	0.507	-	0.497	0.277	0.517	0.475	0.096
bor_nr	0.322	0.431	0.507	0.463	-	0.275	0.385	0.362	0.124
bor_sh	0.311	0.213	0.349	0.286	0.31	-	0.341	0.195	0.1
adhd	0.51	0.475	0.495	0.454	0.34	0.36	-	0.468	0.135
depres	0.265	0.495	0.397	0.422	0.302	0.165	0.432	-	0.103
alcohol	0.151	0.104	0.148	0.129	0.133	0.162	0.155	0.134	-

Note: Female correlations are shown in the upper triangular, and male correlations in the lower triangular.

Abbreviations are as follows: ang=anger total STAS score, neurot=NEO neuroticism, bor_ai=borderline affect instability, bor_ip=borderline identity problems, bor_nr=borderline negative relations, bor_sh=borderline self harm, adhd=CAARS ADHD index, depress=DSM depression symptoms, alcohol=CAGE alcohol problems.

Table 4

Model fit results

	log Likelihood	# parameters	BIC	saBIC
LCA C2	−18259.270	62	37003.581	36806.592
LCA C3	−17690.645	94	36116.675	35818.013
LCA C4	−17365.716	126	35717.161	35316.827
F1C1	−17376.632	41	35074.033	34943.766
F2C1	−17164.978	45	34558.756	34415.780
F2C2	−16945.583	78	34360.309	34112.483
F2C3	−16845.291	113	34603.874	34244.845
F2C4	−16784.063	148	34723.727	34253.494
F3C1	−16947.075	46	34254.038	34107.885
F3C2	−16816.382	81	34266.480	34009.122
F3C3	−16822.285	119	34482.748	34104.655
F3C4	−16818.046	157	34698.028	34199.199

Note: Models were fitted to a subsample of N=2,499 to separate model selection from the estimation of regressions on comorbid behaviors in order to avoid inflated type 1 error. LCA C# indicates results for latent class models with # latent classes. F#C# are factor mixture models where # indicates the number of factors within class and the number of classes (e.g., F2C3 is a 2-factor 3-class model). BIC stands for Bayesian Information Criterion, with smaller values indicating better model fit. SaBIC is a BIC with a different sample size adjustment.³¹

Table 5

Standardized factor loadings of the 3-factor 2-class model

	temperament					reaction			immediacy
	STAS_1	STAS_2	STAS_3	STAS_5	STAS_6	STAS_7	STAS_8	STAS_10	STAS_4
high c 1	0.835	0.841	0.860	0.809	0.710	0.671	0.526	0.649	0.880
low c 2	0.846	0.851	0.869	0.821	0.725	0.738	0.598	0.717	0.839
									0.884
									0.845

Note: The loadings for the higher scoring class are provided in row 1. The general pattern is similar for the two classes. Loadings were also very similar in the models fitted to the smaller model selection sample and the larger sample used to investigate associations (i.e., loadings correlated 0.95)

Table 6

Standardized regression coefficients of the three anger factors on comorbid behaviors within the high scoring minority class and within the low scoring majority class of the best fitting 3-factor 2-class model

	anger temperament	anger reaction	anger immediacy
<i>high scoring minority class</i>			
BOR-AI	0.484 (<0.001)	0.369 (<0.001)	0.484 (<0.001)
BOR-NR	-	0.126 (0.004)	-0.009 (0.751)
BOR-IP	0.004 (0.914)	0.082 (0.099)	-
BOR-SH	0.100 (0.001)	-	-0.028 (0.379)
neuroticism	-	-	0.221 (<0.001)
ADHD	-	0.168 (0.003)	0.107 (0.003)
alcohol problems	-	-	0.080 (0.001)
depression	0.173 (<0.001)	0.306 (<0.001)	0.394 (<0.001)
<i>low scoring majority class</i>			
BOR-AI	0.508 (<0.001)	0.238 (<0.001)	0.398 (<0.001)
BOR-NR	-	0.100 (<0.001)	-0.046 (0.012)
BOR-IP	-0.044 (0.007)	0.073 (0.006)	-
BOR-SH	0.112 (<0.001)	-	0.015 (0.420)
neuroticism	-	-	0.073 (<0.001)
ADHD	-	0.233 (<0.001)	0.209 (<0.001)
alcohol problems	-	-	0.026 (0.111)
depression	0.204 (<0.001)	0.221 (<0.001)	0.326 (<0.001)

Note: Presented are standardized regression coefficients of the regression of the factors on the comorbid behaviors and their p-values (between brackets). All regression coefficients were estimated simultaneously by fitting a 3-factor 2-class model. Blanks indicate that effects were not included in the model because coefficients were below 0.05 in initial exploratory analyses (see text). Significant effects are in bold, the significance level was determined after correction for multiple testing at $\alpha = 0.0036$ (see text).

Abbreviations are bor-ai=borderline affect instability, bor-ip=borderline identity problems, bor-nr=borderline negative relations, bor-sh=borderline self harm, adhd=CAARS ADHD index, depression=DSM depression symptoms, alcohol=dichotomized CAGE alcohol problems