Published in final edited form as:

Org Lett. 2011 December 16; 13(24): 6480–6483. doi:10.1021/ol202772k.

Iron-catalyzed Rearrangements and Cycloaddition Reactions of 2H-Chromenes

Yi Luan, Huan Sun, and Scott E. Schaus

Department of Chemistry and Center for Chemical Methodology and Library Development (CMLD-BU), Life Sciences and Engineering Building, Boston University, 24 Cummington Street, Boston, Massachusetts 02215, United States

Scott E. Schaus: seschaus@bu.edu

Abstract

Iron(III) salts catalyse the tandem rearrangement/hetero-Diels—Alder reaction of 2H-chromenes to yield tetrahydrochromeno heterocycles. The process can occur as a homodimerization and cycloaddition process using electron rich dienophiles. Deuterium labeling and mechanistic studies revealed a hydride shift and *ortho*-quinone methide cycloaddition reaction pathway.

The tetrahydrochromeno polycyclic core is a characteristically unique structure among the large number of biologically active phenolic compounds isolated from Nature.¹ Natural product examples include mulberrofuran G², australisine A³ and sorocinol E⁴, bearing a quaternary ketal carbon (Figure 1). Mulberrofuran, australisine and sorocinol natural products possess hypotensive effects and cytotoxicities against human cancer cell lines.^{2, 3} The dimeric flavonoid dependensin isolated from the antimalarial *uvarzia dependens* also contains a benzopyranobenzopyran polycyclic ring structure.⁵ We envisaged synthetic access to these natural product core structures via tetrahydro-chromeno and benzopyrano starting materials. Herein, we describe an FeCl₃·6H₂O-catalyzed rearrangement and hetero-Diels-Alder reaction of 2H-chromenes with dienophiles to access these unique heterocyclic structures.⁶

We have been interested in the synthesis and utility of 2H-chromenes (4'-methoxyflav-3-enes).⁷ Homodimerization of small molecules is a highly efficient way to construct complex dimeric structures⁸ and recently, Kumar and co-workers demonstrated the homodimerization of 2H-chromenes under Brønsted acidic conditions.⁹

The homodimerization of 2H-chromenes provides rapid access to ketal polycyclic or benzopyranobenzopyran core structures, but as reported, did not readily provide discrete

Correspondence to: Scott E. Schaus, seschaus@bu.edu.

Supporting Information Available Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

access to heterodimeric structures, similar to those found in Nature. During the course of our study on dimerization of 4'-methoxyflav-3-ene, we discovered that iron salts, especially $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ served as an efficient catalyst for the homodimerization of 2*H*-chromene **1a** (Table 1, entries 1-4). Anhydrous FeCl_3 performed poorly, suggesting the hydrate as $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ is essential for catalytic activity (entry 5). To eliminate the possibility that a trace amount of HCl was acting as the catalytic species, catalytic quantities of aqueous HCl was evaluated in the reaction to afford a low yield of **2a** in 1:1 dr (entry 6). The yield was slightly improved using anhydrous methanol as the solvent but in no diastereo-selectivity (entry 7).⁹ Dry HCl failed to provide product from the reaction (entry 8). Similar yields and diastereo-selectivities were achieved using alternative iron(III) salts (entries 9–10), while other metal chloride salts gave little or no yield (entries 11–12). We chose $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ as the catalyst to further investigate in the reaction.

The optimized reaction conditions proved general for a range of 2*H*-chromenes investigated (Table 2). The reaction proceeded smoothly with 6-methyl chromene **1b**, affording chromeno[2,3-*b*]chromene dimer **2b** in good yields (entry 2). Electron-deficient chromenes **1c-1e** underwent dimerization in a similar fashion, providing compounds **2c-2e** in moderate to good yields, but ultimately requiring longer reaction times (entries 3-5). The relative stereochemistry of both the major and minor diastereomers was determined by single crystal X-ray analysis of the dimer **2d**.¹⁰ A change in chemoselectivity was observed in the dimerization of 7-methoxy chromene **1f**. Pyran **1f** afforded isomer **3f** as the major product in good yield at room temperature; only a trace amount (<5%) of isomer **2f** was identified (entry 6). The chemoselectivity reverted to the hetero-Diels—Alder chromeno[2,3-*b*]chromene dimer product with the methoxy group in the 6-position (4',6-dimethoxy-3-flavene **1g**, Table 2, entry 7). Electron rich chromene **1h** did lead to the formation of isomer **3h** in good yield (entry 8). Alternatively, the use of the dimethyl carbamate protecting group attenuated the reactivity of 7-oxygenated arene **1i** to yield the chromeno[2,3-*b*]chromene dimer **2i** (entry 9). The more electron rich chromene reaction partners appeared to have the appropriate reactivity profile to react with the heterodiene as it was being formed in the reaction mixture.

The Fe-catalyzed reaction was utilized in the synthesis of dependensin (**5**) via the homodimerization of 5,7,8-trimethoxyflav-3-ene **4**; easily prepared in three steps from commercially available starting materials. Dependensin has been previously synthesized¹¹ in a similar fashion with our route demonstrating a reduction in reaction sequence steps in high stereocontrol (Scheme 1). Subjecting 5,7,8-trimethoxy-3-flavene **4** to the optimized FeCl_3 -dimerization conditions afforded (±)-dependensin (**5**) as a single diastereomer in 29% overall yield from 1,2,3,5-tetramethoxybenzene.

We designed experiments to gain insight into the reaction sequence leading to the dimerization product. We postulated use of 2-deutero-4'-methoxyflav-3-ene **1-D** in the dimerization reaction would illustrate the isomerization of the chromene to the dienophile. The chromene **1-D** was synthesized via boronate addition onto the deuterated 2-ethoxy-2*H*-chromene. Subjecting **1-D** to the dimerization conditions led to the formation of dimer **2-D**; epimeric at the benzylic positions (Scheme 2). Under *uv* irradiation, *oxa-6π* rearrangement of chromenes¹² results in the formation of the corresponding *ortho*-quinone methide

(*o*QM).¹³ We propose the similar *oxa*-6 π rearrangement of 2*H*-chromene could occur under Fecatalyzed conditions. Illustrated in Scheme 3, the Fecatalyst is responsible for promoting the hydride shift to yield the dienophile and the ring opening reaction.

The ring opening process can lead to the formation of *Z*-ortho-quinone methide that rapidly equilibrates to the *E*-conformer under the reaction conditions (**E**)-**7-D**.¹⁴ The second 2*H*-chromene undergoes a hydride shift to yield the 4'-methoxy-2-flavene **6-D** as the reactive dienophile.¹⁵ Cycloaddition of the *E*-configured vinyl *o*QM (**E**)-**7-D** and **6-D** in an inverse electron demand [4+2] fashion furnishes the dimer **2-D**. The chemoselectivity observed in the 4',7-dimethoxy-3-flavene **1f** dimerization (Table 2, entry 7) is rationalized by a [4+2] cycloaddition of a highly electron rich olefin with the *o*QM at a faster rate than the hydride shift.

We proposed further experiments based on our mechanistic hypothesis aimed at the heterodimerization process. We first postulated that 4'-methoxy-2-flavene **6** would react with the *o*QM generated *in situ* from 2*H*-chromene **1h** under the Fe-promoted conditions (Scheme 4, reaction 1). The reaction proceeded well yielding the corresponding heterodimer **9h** in high yield and 4:1 diastereoselectivity, with only trace amounts (<5%) of homodimer **3h** formed. We demonstrated the intermediacy of the *o*QM by making (**E**)-**8** and reacting it with dienophile **6** under the same reaction conditions (Scheme 4, reaction 2).¹⁶ The product was isolated in comparable yield and the same diastereoselectivity to provide further evidence for the intermediacy of both reaction partners in the [4+2] cycloaddition pathway.

We envisaged a cycloaddition processes to occur with electron rich dienophiles utilizing 2*H*-chromenes as *o*QM precursors based on our preliminary studies of the reaction.¹⁷ Inspired by natural products such as the mulberrofurans and australisine we first evaluated hetero-Diels-Alder reactions between 2*H*-chromenes **1** and 4'-methoxy-2-flavenes **6** under the Fe-promoted conditions.

The reactions proceeded well with 6-methyl and 1,3-benzodioxol substituted chromenes **1b** and **1j** (Table 3, entries 1 and 9). Electron-deficient 2*H*-chromenes **1c-1e** were also able to participate, but at a slower rate (entries 2-4). Electron-rich 2*H*-chromenes **1f-1h** underwent hetero-Diels—Alder reaction in good yields, providing selectively functionalized **9f-9h** with similar diastereoselectivities (entries 5-7). Attenuating the electron-donating group with a dimethyl carbamate did not affect the yield or selectivity of the reaction (entry 8).

We further explored the scope of the cycloaddition reaction pathway using the *in situ* generated *o*QM of chromene **1f** with dienophiles **10** (Table 4). Under the optimized reaction conditions, *p*-methoxystyrene and its derivatives **10a-10c** successfully afforded chromans **11a-11c** in good yields (Table 4, entries 1-3). Furthermore, dihydronaphthalene **10d** yielded the [4+2] adduct **11d** as a single diastereomer (entry 4). Electron rich indene **10e** was also a suitable dienophile (entry 5) and the reaction proceeded smoothly with dihydropyran **10f** to produce the corresponding acetal (entry 6).

In conclusion, we have developed an iron-catalyzed tandem rearrangement/hetero-Diels—Alder approach to access tetrahydrochromeno heterocycles. Our studies have revealed an

*o*QM involved cycloaddition reaction mechanism. Further mechanistic evidence, including trapping of the *in situ* generated *o*QM and use of the *o*QM (*E*)-8, supports the proposed multi-step process. Ongoing studies include expansion of the scope and synthetic utility of the reaction sequence.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This research was supported by the NIH (R01 GM078240, S.E.S. and Y.L.) and BU Undergraduate Research Opportunities Program (H.S.).

References

1. (a) Nomura, T.; Fukai, T.; Hano, Y. *Studies in Natural Products Chemistry*. Rahman, A., editor. Elsevier; 2003. (b) Bhandari P, Crombie L, Harper MF, Rossiter JT, Sanders MD, Whiting A. J. Chem. Soc. Perkin Trans. 1. 1992;1685–1697. (c) Reisch J, Adebajo AC, Kumar V, Aladesanmi AJ. *Phytochemistry*. 1994; 36:1073–1076.
2. (a) Fukai T, Hano Y, Hirakura K, Nornura T, Uzawa J, Fukushima K. *Chem. Pharm. Bull.* 1985; 33:3195. [PubMed: 4085065] (b) Fukai T, Hano Y, Hirakura K, Nornura T, Uzawa J, Fukushima K. *Heterocycles*. 1984; 22:1007–1011.
3. Zhang QJ, Tang YB, Chen RY, Yu DQ. *Chemistry & Biodiversity*. 2007; 4:1533–1540. [PubMed: 17638335]
4. Hano Y, Yamanaka J, Momose Y, Nomura T. *Heterocycles*. 1995; 41:2811–2821.
5. Nkonya MHH, Waibel R, Achenbach H. *Phytochemistry*. 1993; 34:853–856.
6. (a) Bolm C, Legros J, Paih JL, Zani L. *Chem. Rev.* 2004; 104:6217–6254. [PubMed: 15584700] (b) Rao WD, Chan PWH. *Org. Biomol. Chem.* 2010; 8:4016–4025. [PubMed: 20644887] (c) Iovel I, Mertins K, Kischel J, Zapf A, Beller M. *Angew. Chem. Int. Ed.* 2005; 44:3913–3917.
7. Moquist PN, Kodama T, Schaus SE. *Angew. Chem. Int. Ed.* 2010; 49:7096–7100.
8. (a) Ibrahim AA, Wei PH, Harzmann GD, Kerrigan NJ. *J. Org. Chem.* 2010; 75:7901–7904. [PubMed: 21033697] (b) Guidi V, Sandoval S, McGregor MA, Rosen W. *Tetrahedron Lett.* 2010; 51:5086–5090. (c) Chapman OL, Engel MR, Springer JP, Clardy JC. *J. Am. Chem. Soc.* 1971; 93:6696–6698.
9. Devakarama R, Black DStC, Kumar N. *Tetrahedron Lett.* 2010; 51:3636–3638.
10. CCDC 812201 and 812202 contains the supplementary crystallographic data for this paper.
11. Deodhar M, Black DStC, Kumar N. *Tetrahedron*. 2007; 63:5227–5235.
12. (a) Delbaere S, Micheau JC, Vermeersch G. *J. Org. Chem.* 2003; 68:8968–8973. [PubMed: 14604369] (b) Padwa A, Lee GA. *J. Chem. Soc. Chem. Commun.* 1972:795–796.
13. Pathak TP, Sigman MS. *J. Org. Chem. ASAP*.
14. (a) Bishop LM, Winkler M, Houk KN, Bergman RG, Trauner D. *Chem. Eur. J.* 2008; 14:5405–5408. [PubMed: 18449871] (b) Jurd L. *Tetrahedron*. 1977; 33:163–168.
15. For similar hydride transfer, see: McQuaid KM, Long JZ, Sames D. *Org. Lett.* 2009; 11:2972–2975. [PubMed: 19548698] McQuaid KM, Sames D. *J. Am. Chem. Soc.* 2009; 131:402–403. [PubMed: 19099474]
16. Jurd L, Roitman JN, Wong RY. *Tetrahedron*. 1979; 35:1041–1054.
17. (a) Van de Water RW, Pettus TRR. *Tetrahedron*. 2002; 58:5367–5405. (b) Wu KL, Mercado EV, Pettus TRR. *J. Am. Chem. Soc.* 2011; 133:6114–6117. [PubMed: 21452818] (c) Marsini MA, Huang Y, Lindsey C, Wu KL, Pettus TRR. *Org. Lett.* 2008; 10:1477–1480. [PubMed: 18336038] (d) Jensen KH, Webb JD, Sigman MS. *J. Am. Chem. Soc.* 2010; 132:17471–17482. [PubMed:

21082845] (e) Lumb JP, Choong KC, Trauner D. J. Am. Chem. Soc. 2008; 130:9230–9231.
[PubMed: 18582058]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.
Representative natural products.

Scheme 1.
Synthesis of (±)-Dependensin

Scheme 2.
Deuterium-labeled **1a** in the Homodimerization

Scheme 3.
Proposed Mechanism for Cycloaddition

Scheme 3.
Cycloaddition Reactions of 4'-Methoxy-2-flavene **6**

Table 1

Acid-catalyzed Homodimerization of **1a**.^a

entry	catalyst	solvent	dr ^b	yield ^c
1	FeCl ₃ ·6H ₂ O	PhCH ₃	2:1	40%
2	FeCl ₃ ·6H ₂ O	CH ₃ OH		0%
3	FeCl ₃ ·6H ₂ O	CH ₂ Cl ₂	2:1	81%
4 ^d	FeCl ₃ ·6H ₂ O	CH ₂ Cl ₂	3:1	85%
5 ^e	FeCl ₃	CH ₂ Cl ₂	2:1	22%
6	HCl(aq)	CH ₂ Cl ₂	1:1	5%
7 ^f	HCl(aq)	CH ₃ OH	1:1	34%
8	HCl in ether	CH ₂ Cl ₂		0%
9	Fe(OTf) ₃ ·6H ₂ O	CH ₂ Cl ₂	2:1	80%
10	Fe(OTf) ₃	CH ₂ Cl ₂	2:1	85%
11	ZnCl ₂	CH ₂ Cl ₂	2:1	14%
12	AlCl ₃	CH ₂ Cl ₂	2:1	12%

^a Reaction conditions: 0.25 mmol of **1a**, 10 mol % catalyst, 0.2 M in the solvent for 12 h at room temperature.

^b Ratio was determined by ¹H NMR analysis.

^c Isolated yield.

^d 15 mol % FeCl₃·6H₂O.

^e Anhydrous conditions.

^f 0.32 mmol of **1a** in 20 mL of methanol, with 10 drops of HCl, then refluxed at 65 °C for 12 h.⁹

Table 2

FeCl₃·6H₂O Catalyzed Dimerization^a

entry	2H-chromene (1)	dr ^b	2:3	yield ^c
1	R ¹ = H, R ² = H (1a)	3:1	>99:1	85% (2a)
2	R ¹ = CH ₃ , R ² = H (1b)	3:1	>99:1	89% (2b)
3	R ¹ = H, R ² = Cl (1c)	3:1	>99:1	82% (2c)
4 ^d	R ¹ = Br, R ² = H (1d)	3:1	>99:1	86% (2d)
5 ^e	R ¹ = NO ₂ , R ² = H (1e)	1.5:1	>99:1	72% (2e)
6	R ¹ = H, R ² = OCH ₃ (1f)	>99:1	5:95	86% (3f)
7	R ¹ = OCH ₃ , R ² = H (1g)	2:1	>99:1	87% (2g)
8	R ¹ , R ² = 6,7-(OCH ₂ O) (1h)	>99:1	3:97	66% (3h)
9	R ¹ = H, R ² = OCON(CH ₃) ₂ (1i)	3:1	>99:1	70% (2i)

^a 0.25 mmol of **1**, 15 mol % FeCl₃·6H₂O, 0.2 M in CH₂Cl₂ for 12 h at rt.^b ¹H NMR analysis of **2** or **3**.^c Isolated yield.^d 24 h.^e 72 h.

Table 3

Cycloaddition Reactions using 4'-Methoxyflav-2-ene as the Dienophile

entry	2H-chromene (1)	dr	yield ^a
1	R ¹ = CH ₃ , R ² = H, R ³ = 4-OCH ₃ (1b)	3:1	87% (9b)
2	R ¹ = H, R ² = Cl, R ³ = 4-OCH ₃ (1c)	3:1	87% (9c)
3	R ¹ = Br, R ² = H, R ³ = 4-OCH ₃ (1d)	2:1	72% (9d)
4 ^b	R ¹ = NO ₂ , R ² = H, R ³ = 4-OCH ₃ (1e)	3:1	65% (9e)
5	R ¹ = H, R ² = OCH ₃ , R ³ = 4-OCH ₃ (1f)	3:1	87% (9f)
6	R ¹ = OCH ₃ , R ² = H, R ³ = 4-OCH ₃ (1g)	2:1	83% (9g)
7	R ¹ , R ² = 6,7-(OCH ₂ O), R ³ = 4-OCH ₃ (1h)	4:1	81% (9h)
8	R ¹ = H, R ² = OCON(CH ₃) ₂ , R ³ = 4-OCH ₃ (1i)	4:1	89% (9i)
9	R ¹ = R ² = H, R ³ = 3,4-(OCH ₂ O) (1j)	2:1	90% (9j)

^a Isolated yield.^b 24 h, another 0.186 mmol of **6** and 40 mol % FeCl₃·6H₂O were added after first 12 h.

Table 4Cycloaddition Reactions using the *Ortho*-quinone methide Precursor 4',7-Dimethoxy-3-flavene **1f**

entry	dienophile	equiv	cycloadduct	dr	yield ^a
1	10a	2	11a	2:1	78%
2	10b	10	11b	1:1	67%
3	10c	2	11c	1.5:1	84%
4	10d	5	11d	>99:1	82%
5	10e	5	11e	2:1	74%
6	10f	20	11f	1:1	61%

^a Isolated yield. Ar = PMP (*para*-methoxyphenyl)