Published in final edited form as:

Mol Pharm. 2015 September 8; 12(9): 3502–3506. doi:10.1021/acs.molpharmaceut.5b00390.

Enhanced Peptide Stability Against Protease Digestion Induced by Intrinsic Factor Binding of a Vitamin B₁₂ Conjugate of Exendin-4

Ron L. Bonaccorso[†], Oleg G. Chepurny[‡], Christoph Becker-Pauly^{||}, George G. Holz^{‡,§}, and Robert P. Doyle^{†,‡,*}

[†]Department of Chemistry, Center for Science and Technology, Syracuse University, 111 College Place, Syracuse, New York 13244, United States

[‡]Department of Medicine, State University of New York, Upstate Medical University, Syracuse, New York 13210, United States

[§]Department of Pharmacology, State University of New York, Upstate Medical University, Syracuse, New York 13210, United States

^{||}Institute of Biochemistry, Christian-Albrechts-University, 24118 Kiel, Germany

Abstract

Peptide digestion from proteases is a significant limitation in peptide therapeutic development. It has been hypothesized that the dietary pathway of vitamin B₁₂ (B₁₂) may be exploited in this area, but an open question is whether B₁₂-peptide conjugates bound to the B₁₂ gastric uptake protein intrinsic factor (IF) can provide any stability against proteases. Herein, we describe a new conjugate of B₁₂ with the incretin peptide exendin 4 that demonstrates picomolar agonism of the glucagon-like peptide-1 receptor (GLP1-R). Stability studies reveal that Ex-4 is digested by pancreatic proteases trypsin and chymotrypsin and by the kidney endopeptidase meprin β . Prebinding the B₁₂ conjugate to IF, however, resulted in up to a 4-fold greater activity of the B₁₂-Ex-4 conjugate relative to Ex-4, when the IF-B₁₂-Ex-4 complex was exposed to 22 μ g/mL of trypsin, 2.3-fold greater activity when exposed to 1.25 μ g/mL of chymotrypsin, and there was no decrease in function at up to 5 μ g/mL of meprin β .

Keywords

vitamin B₁₂; exendin-4; intrinsic factor; trypsin; AKAR3

*Corresponding Author. rpdoyle@syr.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.molpharmaceut.5b00390.

NMR data and scatterplot analyses (PDF)

The authors declare no competing financial interest.

INTRODUCTION

The human vitamin B₁₂ (B₁₂) dietary uptake pathway is a complex process that facilitates access in humans to a vital cofactor of methionine synthase and methyl malonyl CoA mutase enzymes.¹ This pathway involves three major binders, two of which, intrinsic factor (IF) and haptocorrin (HC), being critical for oral uptake (the third, transcobalamin II (TCII), facilitates entry into cells upon enterocyte passage).² HC primarily protects B₁₂ against acid digestion in the stomach and is enzymatically digested upon entry of the HC-B₁₂ complex into the duodenum, whereupon the B₁₂ is bound by IF. While IF is produced in gastric parietal cells and can bind B₁₂ in the stomach, HC binding is preferred at the lower pH here and it is only upon digestion of HC and a rise in pH in the intestine that IF binding of B₁₂ occurs naturally.^{3,4} Concomitant with the rise in pH is the release of pancreatic proteases, and it is critical to note that IF, unlike HC, is resistant to pancreatic protease digestion.⁵ IF is critical then for delivery of B₁₂ through the intestinal tract to the ileum where cubilin-amnionless based receptor mediated enterocyte passage occurs.⁶ Employing this pathway for oral peptide delivery, for example, requires conjugation of the peptide to B₁₂ in such a way that IF recognition of B₁₂ is not critically hindered and that B₁₂ conjugated peptide can still exhibit the desired pharmacological function. Such concerns are typically readily addressed, however, and there are now several significant examples of B₁₂-peptide conjugates that meet the above criteria.^{7–11} What is not understood, but is no less important, is whether such peptide function is maintained when the conjugate is bound to IF and whether IF, so effective at protecting B₁₂, can provide any protection to a B₁₂ conjugated peptide upon exposure to a protease. To investigate these questions we decided to focus on a highly potent peptide (Ex-4) that is the basis of a pharmaceutical (exenatide) currently approved for treatment of diabetes mellitus.¹²

Ex-4 was discovered in the venom of the Gila monster in 1992 by Eng et al. and is an incretin mimetic, sharing 53% homology with glucagon-like peptide-1 (GLP-1). Like GLP-1, Ex-4 stimulates the release of insulin through agonism of the GLP-1 receptor (GLP-1R) (EC₅₀ 33 pM), effectively lowering blood glucose levels. Unlike GLP-1, Ex-4 is resistant to the enzyme dipeptidyl peptidase IV (DPP-IV), which rapidly cleaves and inactivates GLP-1 in vivo.^{13,14} Since DPP-IV cleaves any peptide with an alanine or proline at the second position from the N-terminus, substituting a glycine for the alanine in GLP-1 results in the resistance seen in Ex-4. This resistance allows Ex-4 to have a half-life of 2.4 h compared to <2 min as seen for GLP-1.¹⁵ Such resistance to DPP-IV does not, however, translate to other proteases, and exenatide therefore must be administered subcutaneously.

The hypothesis herein then is that this pancreatic degradation and general protease limitation may be overcome, at least to some degree above unmodified peptide, by conjugating B₁₂ to Ex-4 and subsequently adding IF, assuming the necessary maintenance of B₁₂ binding by IF and Ex-4 agonism are controlled. To test these hypotheses we synthesized a B₁₂-Ex-4 conjugate focusing on the lysine 12 (K12) position of Ex-4 and the ribose 5'-hydroxyl group of the B₁₂ moiety as sites of conjugation since both sites on the respective moieties had published precedent for allowable modification.^{16–18} Binding to IF was confirmed by radioassay, and agonism of the GLP-1 receptor was then established for an azido modified K12-Ex-4 (**1**), B₁₂-Ex-4 (**4**), and IF-B₁₂-Ex-4 (IF-**4**). With such establishing parameters

controlled for stability against the abundant intestinal endopeptidases, trypsin, chymotrypsin, and the kidney protease meprin β ¹⁹ were compared for **1**, **4**, and IF-**4**.

EXPERIMENTAL SECTION

(AzidoK12)-Ex-4 (**1**) was conjugated to B₁₂ at the K12 position using Huisgens/Sharpless click chemistry,²⁰ using Ex-4 modified at the lysine 12 ϵ -amine with an azido group during solid-phase synthesis. The 5' hydroxyl group of B₁₂ was also modified prior to coupling, being selectively oxidized to a carboxylic acid (**2**) using 2-iodoxybenzoic acid, as previously described by us.²¹ Subsequent coupling of 1-amino-3-butyne to **2** with 1-ethyl-3-(3-(dimethylamino)propyl)carbodiimide (EDCI) and 1-hydroxybenzo-triazole (HOBt) produced B₁₂ with a terminal alkyne at the ribose 5'-position (**3**) (see Scheme 1). Compound **3** was purified using a Shimadzu Prominence HPLC on an Eclipse XDB C18 5 μ m 4.6 mm $\times$ 150 mm column with a mobile phase of 0.1% TFA water and elution with acetonitrile on a gradient starting at 15% acetonitrile increasing to 35% over 20 min (NMR for **3** is provided as Supplementary Figure S1).

Compounds **1** and **3** were coupled using copper(II) sulfate and sodium L-ascorbate (see Scheme 1). The new B₁₂-Ex-4 conjugate (**4**) was purified with a Shimadzu HPLC using an Eclipse XDB C18 5 μ m 4.6 mm $\times$ 150 mm column with a mobile phase of 0.1% TFA water and elution with acetonitrile. A gradient run from 20% acetonitrile to 42% acetonitrile during the first 3 min and then 42 to 47.5% acetonitrile during the next 10 min was used to separate **4** from starting materials. The product was confirmed by matrix-assisted laser desorption/ionization time of flight mass spectrometer (MALDI-ToF MS) (see Figure 1, inset). Compound **4** was purified to greater than 97% purity by HPLC (see Figure 1). The tendency of Ex-4 to aggregate resulted in a small shoulder at 6.5 min.²²

DISCUSSION

Initially, IF binding of **4** was confirmed by radiometric chase assay using ⁵⁷Co-labeled B₁₂ and compared to free B₁₂, as cyanocobalamin (see Figure 2).²³ Significant IF binding of **4** (6.8 nM) was maintained, albeit reduced from unmodified B₁₂ (0.12 nM).

Once IF binding of **4** (IF-**4**) was confirmed, agonism of the GLP-1R was assayed for **1**, **4**, and IF-**4** using HEK-293 cells stably transfected with the GLP-1R (HEK-GLP-1R).²⁵ To this end, we employed a new assay that uses adenoviral transduction to express the genetically encoded FRET reporter AKAR3 that serves as a sensitive readout for cAMP production due to the fact that AKAR3 undergoes a decrease of 485/535 nm emission FRET ratio when it is phosphorylated by cAMP-dependent protein kinase A (PKA) subsequent to GLP-1R activation.^{26–28} This is the first instance to our knowledge of a FRET assay for GLP-1R using viral AKAR3 and offers a ready route to sensitive high-throughput screening of the GLP-1R. An EC₅₀ for **1**, **4**, and IF-**4** were measured at 26, 68, and 132 pM, respectively (see Figure 3). It is worth noting that the azido modification to the K12 position of Ex-4 showed no significant reduction in potency compared to unmodified Ex-4 suggesting a useful general route for selective conjugation to Ex-4 through click chemistry

approaches.²⁰ Compounds **4** and IF-**4** show that further conjugation to the K12 position effects function but still demonstrates low picomolar effective concentrations.

Compounds **1**, **4**, and IF-**4** were analyzed for stability against proteolysis by measuring remaining function at the receptor compared to undigested controls. Compounds **1**, **4**, and IF-**4** were tested for function at [100 nM], a concentration at which each had comparable percent change in FRET ratio (see Table 1). Each protease was analyzed separately so that the protective nature of B₁₂ and IF could be analyzed for their effect versus the specific protease. The pH sensitivity of the assay prevented the use of actual intestinal fluids when testing the compounds.

Digestion was conducted in a standard extracellular solution containing trypsin at 11, 22, or 50 µg/mL, chymotrypsin at 1.25, 3, or 6.25 µg/mL, or meprin β at 1 or 5 µg/mL (see Figures 4 and S2).

At the lowest concentrations of trypsin (11 µg/mL) and chymotrypsin (1.25 µg/mL) there is up to 50% greater function for IF-**4** relative to **4** alone with the highest concentration of trypsin (50 µg/mL) and chymotrypsin (6.25 µg/mL) assayed showing complete lack of function for all systems. The digestion was monitored by measuring agonism of the drugs at the GLP-1R, initially over the course of 3 h, although it was quickly noted that there was no change after 1.5 h indicating that the digestion had stopped by this time point (data not shown). Subsequent triplicate runs were then performed on digestions of 1.5 h.

Meprin β digestion revealed a 2-fold increase in function with B₁₂ conjugation and a 3-fold increase in function when prebound to IF (see Figures 5 and S3). No function was seen for **1** at concentrations greater than 3 µg/mL. The protection provided from B₁₂ conjugation and subsequent binding to IF show that key residues are being protected. Results of the AKAR3 assays show maintenance of function where otherwise none was observed or improvement of function when **4** is first bound to IF.

CONCLUSIONS

The conservation or improved relative function demonstrated herein for Ex-4 when conjugated to B₁₂, and more significantly when bound by IF, is an important first-step in addressing the use and putative role of IF in protecting an administered peptide (orally or by injected means). Protection against pancreatic protease-catalyzed hydrolytic digestion of **4** was maximal at a trypsin concentration of 22 µg/mL and 3 µg/mL of chymotrypsin when **4** was prebound to IF, providing a 4-fold and 5-fold positive increase in function, respectively, as measured by GLP-1R agonism (utilizing the AKAR3 screening assay). The digestion with metalloendoprotease meprin β showed the most significant protection when comparing **1** and IF-**4**. No reduction in function was seen at the highest concentration of meprin β tested (5 µg/mL), while **1** showed no function at concentrations greater than 1 µg/mL of meprin β. B₁₂ provided some protection against trypsin relative to the native peptide. The effect is seen at 11 µg/mL of trypsin with a relative 4-fold increase and at 1.25 and 3 µg/mL of chymotrypsin with a relative increase of 3- and 5-fold. The fact that the IF bound form IF-**4** still maintained significant function at the GLP-1R is also highly significant since many

routes to protect against intestinal degradation involve encapsulation, which prevents possible luminal function or absorption when in place. The use of IF to improve the protease stability of a peptide offers significant scope for exploitation. Even a small improvement in oral function, for example, may be sufficient to achieve the desired effect. Combining this approach with a highly potent peptide with known gut receptors that can produce a vagal afferent response (such as, but not limited to, GLP-1/Ex-4 or PYY3-36), for example, may allow for a positive clinical outcome to be achieved orally, without need even for systemic delivery. Finally, as demonstrated by the stability against meprin β , there is no suggestion that this approach is limited to oral use against gastric proteases, but could also be expanded into serum (through subcutaneous or intravenous injection of IF bound B₁₂-peptide conjugates, for instance), facilitating greatly improved pharmacokinetics (especially when combined with prior results showing B₁₂ conjugation already improved sc absorption of a PYY3-36 conjugate⁸), making this a possible platform technology for peptide drug development.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

The authors acknowledge the lab of Prof. Ebba Nexø (Department of Clinical Medicine–Clinical Biochemistry, University of Aarhus, Denmark) for performing the IF binding assays²³ and Prof. C. Becker-Pauly for providing meprin β produced with support of the German Research Foundation (DFG) Grant SFB877. R.P.D. acknowledges Xeragenx LLC (St. Louis, MO, USA; Xeragenx.com) for research funding and for providing intrinsic factor (#XGX-003).

REFERENCES

1. Banerjee, R. Chemistry and Biochemistry of B₁₂. New York: John Wiley & Sons; 1999.
2. Nielsen MJ, Rasmussen MR, Andersen CBF, Nexø E, Moestrup SK. Vitamin B₁₂ transport from food to the body's cells—a sophisticated, multistep pathway. *Nat. Rev. Gastroenterol. Hepatol.* 2012; 9(6):345–354. [PubMed: 22547309]
3. Gräsbeck R. Hooked to vitamin B₁₂ since 1955: A historical perspective. *Biochimie.* 2013; 95(5): 970–975. [PubMed: 23274132]
4. Alpers DH, Russell-Jones G. Gastric intrinsic factor: The gastric and small intestinal stages of cobalamin absorption. A personal journey. *Biochimie.* 2013; 95(5):989–994. [PubMed: 23274574]
5. Allen RH, Seetharam B, Podell E, Alpers DH. Effect of Proteolytic Enzymes on the Binding of Cobalamin to R Protein and Intrinsic Factor: In Vitro Evidence That A Failure To Partially Degrade Protein Is Responsible For Cobalamin Malabsorption In Pancreatic Insufficiency. *J. Clin. Invest.* 1978; 61(1):47–54. [PubMed: 22556]
6. Gherasim C, Lofgren M, Banerjee R. Navigating the B₁₂ Road: Assimilation, Delivery, and Disorders of Cobalamin. *J. Biol. Chem.* 2013; 288(19):13186–13193. [PubMed: 23539619]
7. Clardy-James S, Chepurny OG, Leech CA, Holz GG, Doyle RP. Synthesis, Characterization and Pharmacodynamics of Vitamin-B(12)-Conjugated Glucagon-Like Peptide-1. *ChemMedChem.* 2013; 8(4):582–586. [PubMed: 23203941]
8. Henry KE, Elfers CT, Burke RM, Chepurny OG, Holz GG, Blevins JE, Roth CL, Doyle RP. Vitamin B₁₂ Conjugation of Peptide-YY3–36 Decreases Food Intake Compared to Native Peptide-YY3–36 Upon Subcutaneous Administration in Male Rats. *Endocrinology.* 2015; 156(5):1739–1749. [PubMed: 25658456]

9. Fazen CH, Valentin D, Fairchild TJ, Doyle RP. Oral delivery of the appetite suppressing peptide hPYY(3–36) through the vitamin B12 uptake pathway. *J. Med. Chem.* 2011; 54:8707–8711. [PubMed: 22054008]
10. Chalasani KB, Russell-Jones GJ, Jain AK, Diwan PV, Jain SK. Effective oral delivery of insulin in animal models using vitamin B12-coated dextran nanoparticles. *J. Controlled Release.* 2007; 122(2):141–150.
11. Siega P, Wuerges J, Arena F, Gianolio E, Fedosov SN, Dreos R, Geremia S, Aime S, Randaccio L. Release of Toxic Gd3+ Ions to Tumour Cells by Vitamin B12 Bioconjugates. *Chem. - Eur. J.* 2009; 15(32):7980–7989. [PubMed: 19562781]
12. Eng J, Kleinman WA, Singh L, Singh G, Raufman JP. Isolation and characterization of exendin-4, an exendin-3 analogue, from *Heloderma suspectum* venom. Further evidence for an exendin receptor on dispersed acini from guinea pig pancreas. *J. Biol. Chem.* 1992; 267(11):7402–7405. [PubMed: 1313797]
13. Mentlein R, Gallwitz B, Schmidt WE. Dipeptidyl-peptidase IV hydrolyses gastric inhibitory polypeptide, glucagon-like peptide-1(7–36)amide, peptide histidine methionine and is responsible for their degradation in human serum. *Eur. J. Biochem.* 1993; 214(3):829–835. [PubMed: 8100523]
14. Nielsen LL, Young AA, Parkes DG. Pharmacology of exenatide (synthetic exendin-4): a potential therapeutic for improved glycemic control of type 2 diabetes. *Regul. Pept.* 2004; 117(2):77–88. [PubMed: 14700743]
15. Schnabel CA, Wintle M, Kolterman O. Metabolic Effects of the Incretin Mimetic Exenatide in the Treatment of Type 2 Diabetes. *Vascular Health and Risk Management.* 2006; 2(1):69–77. [PubMed: 17319471]
16. Clardy SM, Allis DG, Fairchild TJ, Doyle RP. Vitamin B12 in drug delivery: breaking through the barriers to a B12 bioconjugate pharmaceutical. *Expert Opin. Drug Delivery.* 2011; 8(1):127–140.
17. Son S, Chae SY, Kim CW, Choi YG, Jung SY, Lee S, Lee KC. Preparation and Structural, Biochemical, and Pharmaceutical Characterizations of Bile Acid-Modified Long-Acting Exendin-4 Derivatives. *J. Med. Chem.* 2009; 52(21):6889–6896. [PubMed: 19827752]
18. Jin C-H, Chae SY, Son S, Kim TH, Um KA, Youn YS, Lee S, Lee KC. A new orally available glucagon-like peptide-1 receptor agonist, biotinylated exendin-4, displays improved hypoglycemic effects in db/db mice. *J. Controlled Release.* 2009; 133(3):172–177.
19. Broder C, Becker-Pauly C. The metalloproteases meprin α and meprin β : unique enzymes in inflammation, neurodegeneration, cancer and fibrosis. *Biochem. J.* 2013; 450(2):253–264. [PubMed: 23410038]
20. Kolb HC, Finn MG, Sharpless KB. Click Chemistry: Diverse Chemical Function from a Few Good Reactions. *Angew. Chem., Int. Ed.* 2001; 40(11):2004–2021.
21. Clardy-James SM, Bernstein JL, Kerwood DJ, Doyle RP. Site-Selective Oxidation of Vitamin B12 Using 2-Iodoxybenzoic Acid. *Synlett.* 2012; 23:2363–2366.
22. Andersen NH, Brodsky Y, Neidigh JW, Prickett KS. Medium-Dependence of the secondary structure of exendin-4 and glucagon-like-peptide-1. *Bioorg. Med. Chem.* 2002; 10(1):79–85. [PubMed: 11738609]
23. Stupperich E, Nexø E. Effect of the cobalt-N coordination on the cobamide recognition by the human vitamin B12 binding proteins intrinsic factor, transcobalamin and haptocorrin. *Eur. J. Biochem.* 1991; 199(2):299–303. [PubMed: 2070790]
24. Fedosov SN, Laursen NB, Nexø E, Moestrup SK, Petersen TE, Jensen EØ, Berglund L. Human intrinsic factor expressed in the plant *Arabidopsis thaliana*. *Eur. J. Biochem.* 2003; 270(16):3362–3367. [PubMed: 12899693]
25. Gromada J, Rorsman P, Dissing S, Wulff BS. Stimulation of cloned human glucagon-like peptide 1 receptor expressed in HEK 293 cells induces cAMP-dependent activation of calcium-induced calcium release. *FEBS Lett.* 1995; 373(2):182–186. [PubMed: 7589461]
26. Allen MD, Zhang J. Subcellular dynamics of protein kinase A activity visualized by FRET-based reporters. *Biochem. Biophys. Res. Commun.* 2006; 348(2):716–721. [PubMed: 16895723]
27. Chepurny OG, Kelley GG, Dzihura I, Leech CA, Roe MW, Dzihura E, Li X, Schwede F, Genieser H-G, Holz GG. PKA-dependent potentiation of glucose-stimulated insulin secretion by Epac

- activator 8-pCPT-2-O-Me-cAMP-AM in human islets of Langerhans. *Am. J. Physiol. Endocrinol. Metabol.* 2010; 298:E622–E633.
28. Holz, GG.; Chepurny, OG.; Leech, CA.; Roe, MW. High-throughput FRET assays for fast time-dependent detection of cyclic AMP in pancreatic beta cells. In: Xiaodong, C., editor. *Cyclic Nucleotide Signaling*. Boca Raton, FL: CRC Press, Taylor & Francis Group; 2015. p. 35-59.
29. Becker C, Kruse MN, Sloty KA, Köhler D, Harris JR, Rösmann S, Sterchi EE, Stöcker W. Differences in the Activation Mechanism between the α and β Subunits of Human Meprin. *Biol. Chem.* 2003; 384:825–883. [PubMed: 12817480]

Figure 1.

LC trace showing purified **4** as a monomer (~7 min) and dimer (~6.5 min) and MALDI-ToF MS (inset) of **4** showing m/z of 5658.153 Da, which corresponds to the +1 of **4**.

Figure 2.

IF binding to B₁₂ (0.12 nM) and **4** (6.8 nM). IF used in these assays was produced in the plant *Arabidopsis* in the apo-form and of high purity.²⁴

Figure 3.
Dose–response analysis of **1**, **4**, and IF-**4** yielded EC₅₀ values of 26, 68, and 132 pM, respectively, as determined by monitoring the 485/535 nm FRET emission ratio.

Figure 4.

Digestion for 1.5 h of 100 nM **1**, **4**, and IF-4 with 50, 22, or 11 $\mu\text{g/mL}$ of trypsin or 1.25, 3, or 6.25 $\mu\text{g/mL}$ of chymotrypsin using AKAR3 to measure function. The data shows the maximum expression normalized to 100% of the conjugates done in triplicate (mean $\pm$ SEM). Basal control contained trypsin at 50 $\mu\text{g/mL}$ of trypsin. (N.C. = no change). A scatterplot analysis is provided in the Supporting Information (Figure S2).

Figure 5.

Thirty minute meprin β digestion of 100 nM **1**, **4** and IF-**4** with 1 and 5 $\mu\text{g/mL}$ of meprin β . The data shows the maximum expression normalized to 100% of the conjugates done in triplicate (mean $\pm$ SEM). Basal control contained 2 $\mu\text{g/mL}$ of meprin β . Recombinant human meprin β was produced in insect cells and purified and activated as described previously.²⁹ A scatterplot analysis is provided in the Supporting Information (Figure S3).

Scheme 1. Synthesis of B₁₂-Ex-4 Conjugate 4a

^aReagents and conditions: (i) EDCI, HOBt, 1-amino-3-butyne, rt, 16 h; (ii) **1**, CuSO₄, sodium ascorbate, 1 h.

Table 1

Percent Change in FRET at 100 nM for 1, 4, and IF-4

compd	% change in FRET at 100 nM
1	-12 ± 0.01
4	-13 ± 0.02
IF-4	-12 ± 0.01