Published in final edited form as:

Photochem Photobiol Sci. 2015 December ; 14(12): 2151–2158. doi:10.1039/c5pp00334b.

Photoactivatable, biologically-relevant phenols with sensitivity toward 2-photon excitation†

Duncan E. McLain^{a,b}, Adam. C. Rea^b, Magnus B. Widegren^a, and Timothy M. Dore^{a,b}

Timothy M. Dore: timothy.dore@nyu.edu

^aNew York University Abu Dhabi, PO Box 129188, Abu Dhabi, United Arab Emirates

^bDepartment of Chemistry, University of Georgia, Athens, Georgia, 30602, USA

Abstract

Spatio-temporal release of biologically relevant small molecules provides exquisite control over the activation of receptors and signaling pathways. This can be accomplished via a photochemical reaction that releases the desired small molecule in response to irradiation with light. A series of biologically-relevant signaling molecules (serotonin, octopamine, capsaicin, *N*-vanillyl-nonanoylamide, estradiol, and tyrosine) that contain a phenol moiety were conjugated to the 8-bromo-7-hydroxyquinolinyl (BHQ) or 8-cyano-7-hydroxyquinolinyl (CyHQ) photoremovable protecting groups (PPGs). The CyHQ caged compounds proved sensitive toward 1PE and 2PE processes with quantum efficiencies of 0.2 – 0.4 upon irradiation at 365 nm and two-photon action cross sections of 0.15 – 0.31 GM when irradiated at 740 nm. All but one compound, BHQ-estradiol, were found to be sensitive to photolysis through 1PE and 2PE with quantum efficiencies of 0.30 – 0.40 and two photon cross sections of 0.40 – 0.60 GM. Instead of releasing estradiol, BHQ-estradiol underwent debromination.

Introduction

Biological effectors containing a phenol functional group are ubiquitous in nature. Phenols are found in neurotransmitters, synthetic neural agonists and antagonists, steroids, and amino acids, among other classes of compounds (Figure 1). The physiological function of these phenols is quite broad. For example, serotonin (5-HT) plays a role in the modulation of mood, appetite, and sleep, and there is a growing body of evidence that it influences left-right patterning in the embryonic development of vertebrates.^{1–3} Octopamine is primarily an insect neurotransmitter involved in reinforcing reward encoding and certain behaviors.⁴ Capsaicin, the active ingredient in hot peppers, is a TRPV1 ion channel agonist, and heterologously expressed TRPV1 channels have been activated by in vivo application of capsaicin to effect behavioral changes.⁵ *N*-Vanillyl-nonanoylamide (VNA) is an equipotent TRPV1 agonist.⁶ Steroids are involved in myriad signaling pathways⁷ and have been recently implicated in neurogenesis.⁸ Amino acids, tyrosine in particular, are involved in the

†Electronic Supplementary Information (ESI) available: Synthetic procedures for the preparations of **1b**, **2–7**, **10b**, and **11–16**; ¹H NMR and ¹³C NMR spectra; experimental procedures for determining ϵ , Q_u , and δ_u ; and individual photolysis reaction progress curves. See DOI: 10.1039/b000000x/

Correspondence to: Timothy M. Dore, timothy.dore@nyu.edu.

regulation of MAPK and MEK/ERK through phosphorylation and dephosphorylation.⁹ Developing ways to artificially activate phenols in model biological systems would enable the conduct of experiments aimed at understanding the role that they play in physiology at the molecular level.

One method of activating biological effectors is through the use of a photoremovable protecting group (PPG). This strategy typically employs a covalent linkage between the PPG and a biologically relevant functional group on the desired biological effector; it has been used successfully to “cage” carboxylates, phosphates, alcohols, aldehydes and ketones, diols, thiols, amines, and phenols.¹⁰ A major limitation faced when using a PPG on a phenol is that most PPGs require a carbonate linkage between the PPG and the phenol to effect release. A common way around this design problem has been to instead protect a different functional group, typically a primary or secondary amine. In either case, this limitation necessitates a decarboxylation step from either the phenol (~ 4 ms)¹¹ or the amine (~ 0.4 s)^{12, 13} to return the biological effector to full activity. Direct release of phenols enables the study of biological processes faster than the rate(s) of decarboxylation; an example of this is the *ortho*-carboxynitrobenzyl (o-CNB) PPG, which has demonstrated the ability to directly release phenols with a time constant of 16 μ s, albeit with a low quantum efficiency (0.03).¹⁴

The kinetics of photolysis are important in design considerations from two different standpoints. The first is in regard to spatial resolution, and the second is in regard to the kinetics of the biological process under investigation. Spatial resolution in 3-dimensions is primarily achieved through two-photon excitation (2PE): the nearly simultaneous absorbance of two photons of light to reach an excited state restricts the volume of excitation to the focal point of the laser, which can provide spatial resolution down to one femtoliter.¹⁰ The exquisite spatial resolution provided by 2PE can only be achieved when the kinetics of photolysis are faster than the rate of diffusion ($\sim 3.3 \times 10^{-6}$ cm²/s),¹⁵ otherwise diffusional spread results in a release volume larger than the excitation volume. When photolysis kinetics are slower than the rate of diffusion, the sensitivity of the PPG toward 2PE must raise accordingly, and a reported threshold of 0.10 GM¹⁶ for biological application may be insufficient for certain PPGs and a revised threshold of >3 GM¹⁷ may be warranted. The photorelease of the effector must be faster than the physiological process under study. This is akin to watching a video of sprinters in a 50-m dash at a low vs. a high frame rate. If the frame rate is 0.1 frames/s, one would see the runners lined up for the start and the next image would show the racers post finish. Using a frame rate of 10 frames/s would capture nearly all of the action in between the start and finish. Likewise, to investigate a biological process, the photolysis reaction needs to be faster than the action of the biological effector on the system.

The kinetics of biological processes involving the phenols shown in Figure 1 have been studied and occur on a range of timescales. For example, the estrogen receptor activation rate constant is $\sim 1.7 \times 10^6$ M⁻¹ s⁻¹,¹⁸ tyrosine phosphorylation by the tyrosine kinase Lck occurs with an apparent Michaelis rate constant k_{cat} of 2.8×10^{-3} s⁻¹,¹⁹ and octopamine binds to its receptor with an on-rate constant of $\sim 3.2 \times 10^5$ M⁻¹ s⁻¹.²⁰ Capsaicin induced channel opening of the TRPV1 channel has three phases with the fastest occurring in 0.46 ms.²¹ There are two major types of 5-HT receptors: the 5-HT₃ receptor, a ligand gated ion

channel with kinetics ($4 \times 10^2 \text{ s}^{-1}$) measured by Hess and co-workers using a caged 5-HT,²² and six G-protein coupled receptor subtypes (5-HT_{1,2,4-7}), which display binding on timescales in the 100–500 ms range.²³

Photoactivatable versions of each of the biologically-relevant phenols shown in Figure 1 have been previously reported (Figure 2). The neurotransmitter 5-HT has been protected at the phenol by the α -carboxynitrobenzyl (CNB)¹⁴ and the nitrophenethyl (NPE)²⁴ PPGs and at the amine with CNB,¹⁴ through a carbamate linkage with NPE (Tocris cat. #3991), and as a complex with the ruthenium bipyridyl PPG.²⁵ More recently, we reported the synthesis and photochemistry of an 8-bromo-7-hydroxyquinoline (BHQ)-protected 5-HT (**1a** Figure 3).³ Octopamine has not been protected on its phenol. Rather, a carbamate linkage between the amine and several 7-aminocoumarinyl derivatives²⁶ and a pyrenylmethyl PPG²⁷ have been reported. Several strategies for protecting capsaicin and VNA have been developed. It has been protected at the phenol using CNB and 7-bromo-8-hydroxycoumarinyl derivatives,²⁸ whereas VNA has been protected at the phenol with NB,²⁹ DMNB,^{29, 30} NVOC,^{11, 29} and MeOPA^{29, 30} or the amide nitrogen with NB or DMNB.¹¹ There are two previously reported photoactivatable forms of estradiol, one protected by DMNB at the phenol and one protected by NPEC at the C-17 hydroxyl group.³¹ Tyrosine has been protected at the phenol with CNV³² and the amine with PMOC.²⁷

Most of the photoactivatable phenols described in the previous paragraph photolyze with low quantum efficiencies ($Q_u = 0.003\text{--}0.05$)^{11, 14, 25, 27, 28, 32} or the quantum efficiencies are not reported.^{24, 29–31} The value of λ_{max} is less than 300 nm for the NB-,¹¹ CNB-,¹⁴ and NPEC-conjugated³¹ effectors, which raises the potential for damaging biological tissues during photoactivation experiments. The coumarinyl-protected compounds have the highest quantum efficiencies at biologically useful wavelengths ($\lambda_{\text{max}} > 365 \text{ nm}$),^{26, 28} but they are linked to the respective biological effectors through carbonate or carbamate linkages, which significantly slows the release kinetics,^{12, 13} because the resulting carbonate or carbamate must decarboxylate to reveal the active biological effector. Nitrobenzyl PPGs also show slow release kinetics, especially with poor leaving groups.³³ Only the coumarinyl derivatives,^{26, 28} NV-VNA,¹¹ and the $[\text{Ru}(\text{bpy})_2(5\text{HT})_2]^{2+}$ complex²⁵ photolyze readily by 2PE.

Hydroxyquinoline derivatives have demonstrated the ability to efficiently release a number of biologically relevant functional groups, including carboxylates, phosphates, diols, and thiols, through 1PE and 2PE.^{34–37} The hydroxyquinoline PPG has also been shown to directly release phenols without the use of a carbonate linker,³ and the timescale, measured for BHQ-OAc, is on the order of nanoseconds,³⁸ thus making this PPG suitable for use in studying temporally resolved signaling events. By directly releasing phenols, photoactivatable versions of relevant phenol-containing biological effectors with potentially rapid release kinetics are now possible. Indeed, 5-HT was recently caged and subsequently released in *Xenopus* embryos to provide temporal resolution in the examination of left-right patterning defects caused by abnormal distribution of the monoamine in the developing embryo.^{3, 39}

We describe the synthesis of representative biologically active photactivatable phenols (Figure 3), their sensitivity toward both 1PE and 2PE with subsequent release of the phenol, and their hydrolytic stability in the dark. The phenols are comprised of 5-HT, octopamine, capsaicin, VNA, estradiol, and tyrosine.

Results and discussion

We protected the biologically relevant phenols using two hydroxyquinoline-based PPGs, BHQ and CyHQ (Scheme 1). Compound **1a** was prepared as previously reported.³ The other compounds were synthesized from MOM-protected BHQ-OH (**8a**) and CyHQ-OH (**8b**), which were prepared by published protocols.^{36, 38} The hydroxyl group on **8a** or **8b** was activated with methanesulfonyl chloride to produce mesylate **9a**, which has been previously reported,³ or **9b**, respectively, in high yield. Displacement of the methanesulfonate with the desired phenol using cesium carbonate in acetone produced each of the protected caged phenols **10–16**. Deprotection conditions depended upon the protecting groups involved, but generally, trimethylsilyl chloride in methanol effected global deprotection, which was followed by HPLC purification to provide the photoactivatable phenols **1–7**.

The caged capsaicinoids **3a** and **4a** were too insoluble in neutral aqueous buffer, so BHQ-VAA (**5a**), which is sufficiently soluble, was used as a surrogate for assessing the photophysical and photochemical properties of the photoactivatable capsaicin and VNA. The CyHQ-protected capsaicinoids were much more water-soluble, but for direct comparison, CyHQ-VAA (**5b**) was photolyzed as a surrogate for **3b** and **4b**.

Both BHQ- (**a** series) and CyHQ-protected (**b** series) phenols had a maximum wavelength of absorption near 370 nm, although comparing the average values of ϵ (2480 vs 5660 M⁻¹ cm⁻¹ for BHQ and CyHQ, respectively) shows that the CyHQ series had around twice the molar absorptivity than the BHQ-protected phenols (Table 1).

Irradiation of the **a** and **b** series of caged compounds **1**, **2**, **5**, and **7** plus **6b** with 365-nm light in KMOPS buffer (pH 7.2) released the corresponding phenols in 55–65% yield and with average quantum efficiencies of 0.29 for the BHQ series and 0.36 for the CyHQ series (Figure 4). The chemical yields reported are similar to those of a variety of other caged compounds.¹⁰ Several factors can reduce the chemical yield of the photolysis: the caged compounds might undergo undesired photoreactions,³⁸ the effectors might possess some photosensitivity and react further to form secondary photoproducts,⁴⁰ or undesired secondary photoproducts might be generated through the interaction of the spent PPG and the biological effector, or a combination of all of these. The BHQ- and CyHQ-protected phenols have similar quantum efficiencies, but the sensitivity, which is quantified by multiplying the quantum efficiency by the molar absorptivity ($Q_u \times \epsilon$) was much greater for the CyHQ-protected compounds, owing to their much larger extinction coefficients.

BHQ-protected estradiol **6a** did not photolyze to the expected products of **17a** and estradiol. Instead, debromination of **6a** was the primary reaction pathway, a phenomenon that has been observed previously.⁴¹ To further investigate the debromination reaction, we irradiated at 365 nm a model compound, BHQ-OH (**17a**), which did not undergo photoinduced cleavage

of the C–O bond, but lost bromine instead with a quantum efficiency of 0.12 in pH 7.2 KMOPS buffer. The extent of debromination in the BHQ series of photoactivatable phenols can be significant. A factor that contributes to the extent of debromination is the leaving group ability, and effectors that are poor leaving groups (high pK_a) conjugated to PPGs tend to be inert.¹⁰ The calculated pK_a of estradiol was found to be 10.27 ± 0.60 (ACD/Labs version 11.02 accessed through SciFinder). All of the other phenols have calculated pK_a values lower than this value (10.09 ± 0.40 for 5-HT, 9.97 ± 0.26 for octopamine, 10.02 ± 0.31 for VAA, and 9.09 ± 0.10 for tyrosine), which might be the threshold for leaving group ability for BHQ-protected effectors. The ACD/Labs-calculated values were used to provide an estimated pK_a of the phenols under identical conditions and avoid issues arising from the variations in conditions found in the literature reporting pK_a values from experimental titrations.

Sufficient hydrolytic stability of the PPG-effector conjugate is required, so that there is no problem with leakage of the activated effector prior to photoactivation. Experiments where the caged compound is subjected to aqueous environments for hours or days require robust resistance to hydrolysis in the dark. Such experiments, might include those that use CyHQ-estradiol for neurogenesis or gene regulation studies or BHQ/CyHQ-tyrosine when controlling kinase cascades via phosphorylation of tyrosine residues. All of the compounds were examined for their ability to resist hydrolysis under simulated physiological conditions. Solutions of the relevant compound (100 μ M) in KMOPS pH 7.2 were prepared, stored in the dark at room temperature, analyzed periodically over a period of 1 week, and in all cases showed minimal (<3%) hydrolysis over the measurement time frame.

To assess the sensitivity of the photoactivatable phenols to 2PE, **1**, **2**, **5**, **6b**, and **7** were photolyzed at 740 nm using a femtosecond-pulsed, mode-locked Ti:sapphire laser. HPLC or uHPLC was used to monitor the reaction progress and from the decay curves (Figure 5) and the 2PE-generated fluorescence output of fluorescein as an external standard, the 2-photon uncaging action cross-sections (δ_u) were calculated as previously described (Table 1).^{3, 16} The BHQ derivatives were more sensitive to 2PE than the CyHQ derivatives, having average values of δ_u of 0.55 GM compared to 0.25 GM for the CyHQ derivatives. The cross-sections reported are similar to that of previously reported quinoline- and coumarin-based PPGs and significantly higher than most nitrobenzyl and nitroindoline derivatives.^{10, 42–44} BHQ and CyHQ are sufficiently sensitive to be applied to the photoactivation of biologically relevant phenols in vivo through 2PE.

Conclusion

The sensitivity of the hydroxyquinoline PPGs BHQ and CyHQ toward 2PE, combined with sub-microsecond release kinetics provided an opportunity to create a variety of photoactivatable phenolic biological effectors (5-HT, octopamine, capsaicin, VNA, estradiol, and tyrosine). The CyHQ-protected phenols were more sensitive to 1PE owing to their larger molar absorptivities, whereas the BHQ-derivatives were more sensitive to 2PE. Instead of releasing estradiol, BHQ-estradiol was efficiently debrominated upon exposure to 365-nm light. This photochemical reaction pathway dominates when the attached effector is a poor leaving group. Alternative reaction pathways were not observed with the CyHQ-protected

phenols. The synthesis of the XHQ-phenol conjugates is modular and can be applied to any biologically relevant phenol, not just the ones described here. The photoactivatable phenols reported will enable the examination of their associated biological processes at the cellular and molecular level. BHQ-*O*-5HT (**1a**) has already been used to study the role of 5-HT in left/right asymmetry in the developing vertebrate embryo.^{3, 39} Biological applications of the other photoactivatable phenols will be reported in due course.

Experimental

Synthesis

General procedures—MOM-protected BHQ-OH (**8a**)³⁸ and CyHQ-OH (**8b**)³⁶ were prepared as previously described, as were **1a**, **9a**, and **10a**.³ A representative experimental procedure for preparing a photoactivatable phenol is given below. Experimental procedures for preparing each caged phenol can be found in the Electronic Supplementary Information. uHPLC and HPLC were carried out on Agilent 1290 and 1260 systems, respectively, with Zorbax Eclipse C-18 reverse phase columns.

(8-Cyano-7-(methoxymethoxy)quinolin-2-yl)methyl methanesulfonate (9b)—Alcohol **8b** (0.31 g, 1.2 mmol) was stirred in THF (5 mL). Methanesulfonyl chloride (0.189 mL, 2.44 mmol) and triethylamine (0.51 mL, 3.7 mmol) were added. The reaction progress was monitored by uHPLC, and upon completion, celite was added to the reaction mixture, which was subsequently dried in vacuo, loaded onto a column and purified via flash chromatography using EtOAc/hexane (1:2) to provide **9b** as a white powder (0.132 g, 0.408 mmol, 34%): ¹H NMR (400 MHz, CDCl₃) δ 8.23 (d, *J* = 8.4 Hz, 1H), 8.02 (d, *J* = 9.2 Hz, 1H), 7.58 (dd, *J* = 12.6, 8.8 Hz, 2H), 5.55 (s, 2H), 5.47 (s, 2H), 3.59 (s, 3H), 3.29 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ 162.5, 157.0, 148.2, 137.5, 133.7, 122.9, 119.1, 116.3, 114.5, 99.5, 95.1, 71.6, 56.9, 38.5; HRMS-ESI (*m/z*) [M+H]⁺ calcd for C₁₄H₁₄N₂O₅S 323.06962; found, 323.06966.

tert-Butyl (2-(4-((8-cyano-7-(methoxymethoxy)quinolin-2-yl)methoxy)phenyl)-2-hydroxyethyl)carbamate (11b)—*tert*-Butyl (2-hydroxy-2-(4-hydroxyphenyl)ethyl)carbamate (0.043 g, 0.17 mmol) and **9b** (0.050 g, 0.16 mmol) were stirred in acetone (1 mL). Cs₂CO₃ (0.10 g, 0.33 mmol) was added, and the reaction progress was monitored by uHPLC. Upon completion, brine was added and the resulting mixture was extracted with EtOAc. The extract was dried over sodium sulfate, concentrated in vacuo and purified via flash chromatography eluting with EtOAc/hexane (1:2), which yielded **11b** as a yellow residue upon drying (0.025 g, 0.052 mmol, 31%): ¹H NMR (600 MHz, CD₃OD) δ 9.05 (d, *J* = 8.3 Hz, 1H), 8.26 (d, *J* = 9.1 Hz, 1H), 7.90 (d, *J* = 8.3 Hz, 1H), 7.64 (d, *J* = 8.9 Hz, 1H), 7.27 (d, *J* = 7.9 Hz, 2H), 6.82 (d, *J* = 7.9 Hz, 2H), 5.27 (s, 2H), 4.82 (dd, *J* = 11.9, 3.7 Hz, 1H), 3.35 (s, 3H), 3.59–3.20 (m, 2H), 1.53 (s, 9H); ¹³C NMR (151 MHz, CD₃OD) δ 167.6, 162.0, 157.3, 147.0, 145.5, 140.8, 135.5, 131.6, 126.9, 122.6, 120.0, 117.6, 115.0, 111.9, 97.1, 88.9, 85.0, 69.3, 60.7, 53.9, 46.0, 26.2; HRMS-ESI (*m/z*) [M+H]⁺ calcd for C₂₆H₂₉N₃O₆ 480.21291; found, 480.21438.

2-((4-(2-amino-1-hydroxyethyl)phenoxy)methyl)-7-hydroxyquinoline-8-carbonitrile (2b)—Compound **11b** (0.025 g, 0.052 mmol) was dissolved in methanol (1 mL). TMSCl (0.115 mL, 0.918 mmol) was added. The resulting solution was stirred in the dark and monitored by uHPLC. Upon completion, the reaction was concentrated in vacuo and purified by HPLC using a 10 minute gradient from 5% CH₃CN/95% H₂O (0.1% TFA) to 100% CH₃CN. Fractions containing only one peak were combined and concentrated in vacuo to provide **2b** as a pale yellow solid (0.014 g, 0.042 mmol, 80% yield): ¹H NMR (600 MHz, CD₃OD) δ 8.27 (d, *J* = 8.4 Hz, 1H), 8.01 (d, *J* = 9.1 Hz, 1H), 7.58 (d, *J* = 8.3 Hz, 1H), 7.26 (d, *J* = 8.5 Hz, 2H), 6.82 (d, *J* = 8.4 Hz, 2H), 5.03 (s, 2H), 4.81 (d, *J* = 6.5 Hz, 1H), 4.81 (d, *J* = 12.7 Hz, 1H), 3.12 – 3.02 (m, 2H); ¹³CNMR (600 MHz, CD₃OD) δ 163.6, 157.3, 148.4, 137.0, 133.7, 131.6, 126.8, 121.3, 117.7, 116.8, 115.1, 115.0, 94.0, 69.3, 64.7, 47.1, 45.9; HRMS-ESI (*m/z*) [M+H]⁺ calcd for C₁₉H₁₇N₃O₃ 336.13427; found, 336.13471.

Photochemistry

Determination of the Quantum Efficiency for One-Photon Photolysis (*Q_u*)—

Briefly, solutions (100 μM) of the relevant substrate in KMOPS pH 7.2 were irradiated in quartz cuvettes (21-Q-10, Starna, Atascadero, CA) with a 365 ± 10 nm LED (Cairn OptoLED Lite). Aliquots (25 μL) were removed at various time points and analyzed by uHPLC (MeCN/H₂O 0.1% TFA: 6-min gradient 5% CH₃CN to 100% CH₃CN, 2 min gradient 100% CH₃CN to 5% CH₃CN), using an external standard to determine concentrations. The reaction progress was plotted and the data fit to a single exponential decay function and *Q_u* values were calculated according to previously reported protocols.³⁴ See Electronic Supplementary Information for detailed experimental procedure.

Determination of the Two-Photon Uncaging Action Cross-Section (*δ_u*)—

Solutions (100 μM) of the substrates in KMOPS were prepared and stored in the dark. Aliquots (25 μL) of this solution were placed in a microcuvette (10 × 1 × 1 mm illuminated dimensions, 25 μL effective filling volume) and irradiated with a fs-pulsed and mode-locked Ti:Sapphire laser (Chameleon Ultra II, Coherent or a Mai Tai HP DeepSee, Spectra-Physics) with 740-nm light at an average power of 220–300 mW measured after passing through the cuvette. Three samples of each substrate were irradiated for various time periods and analyzed by HPLC or uHPLC as described for one-photon experiments. The reaction progress was plotted and the data fit to a single exponential decay function and *δ_u* values were calculated according to previously reported methods using fluorescein as an external standard.³⁴ See Electronic Supplementary Information for detailed experimental procedure.

Hydrolysis in the Dark

Substrates (100 μM) were dissolved in KMOPS and stored in the dark at room temperature. Aliquots (20 μL or 5 μL) were removed and analyzed periodically by HPLC or uHPLC, respectively, as described for one-photon photolysis. None of the compounds synthesized displayed a measurable rate of dark hydrolysis over an approximately 100-h time period.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by grants from the National Science Foundation (CHE-1012412, CHE-1317760) and the National Institutes of Health (R01NS070159). The research was carried out using Core Technology Platform resources at New York University Abu Dhabi and mass spectrometry resources at the Proteomics and Mass Spectrometry Core Facility at the University of Georgia.

Notes and references

1. Levin M, Buznikov GA, Lauder JM. Of Minds and Embryos: Left-Right Asymmetry and the Serotonergic Controls of Pre-Neural Morphogenesis. *Dev. Neurosci.* 2006; 28:171–185. [PubMed: 16679764]
2. Vandenberg N L, Levin M. Far from solved: a perspective on what we know about early mechanisms of left-right asymmetry. *Dev. Dyn.* 2010; 239:3131–3146. [PubMed: 21031419]
3. Rea AC, Vandenberg LN, Ball RE, Snouffer AA, Hudson AG, Zhu Y, McLain DE, Johnston LL, Lauderdale JD, Levin M, Dore TM. Light Activated Serotonin for Exploring Its Action in Biological Systems. *Chem. Biol.* 2013; 20:1536–1546. [PubMed: 24333002]
4. Roeder T. Tyramine and octopamine: ruling behavior and metabolism. *Annu. Rev. Entomol.* 2005; 50:447–477. [PubMed: 15355245]
5. Guler AD, Rainwater A, Parker JG, Jones GL, Argilli E, Arenkiel BR, Ehlers MD, Bonci A, Zweifel LS, Palmiter RD. Transient activation of specific neurons in mice by selective expression of the capsaicin receptor. *Nat. Commun.* 2012; 3:1749/1741–1749/1710.
6. Walpole, CSJ.; Wrigglesworth, R. Capsaicin in the Study of Pain. Wood, JN., editor. London: Academic Press; 1993. p. 63-81.
7. Prossnitz ER, Arterburn JB, Smith HO, Oprea TI, Sklar LA, Hathaway HJ. Estrogen signaling through the transmembrane G protein-coupled receptor GPR30. *Annu. Rev. Physiol.* 2008; 70:165–190. [PubMed: 18271749]
8. Suzuki S, Gerhold LM, Bottner M, Rau SW, Dela Cruz C, Yang E, Zhu H, Yu J, Cashion AB, Kindy MS, Merchenthaler I, Gage FH, Wise PM. Estradiol enhances neurogenesis following ischemic stroke through estrogen receptors α and β . *J Comp. Neurol.* 2007; 500:1064–1075. [PubMed: 17183542]
9. De Luca A, Maiello MR, D'Alessio A, Pergameno M, Normanno N. The RAS/RAF/MEK/ERK and the PI3K/AKT signalling pathways: role in cancer pathogenesis and implications for therapeutic approaches. *Expert Opin. Ther. Targets.* 2012; 16:S17–S27. [PubMed: 22443084]
10. Klan P, Solomek T, Bochet CG, Blanc A, Givens R, Rubina M, Popik V, Kostikov A, Wirz J. Photoremovable Protecting Groups in Chemistry and Biology: Reaction Mechanisms and Efficacy. *Chem. Rev.* 2013; 113:119–191. [PubMed: 23256727]
11. Zhao J, Gover TD, Muralidharan S, Auston DA, Weinreich D, Kao JPY. Caged Vanilloid Ligands for Activation of TRPV1 Receptors by 1- and 2-Photon Excitation. *Biochemistry.* 2006; 45:4915–4926. [PubMed: 16605259]
12. Papageorgiou G, Corrie JET. Synthesis and properties of carbamoyl derivatives of photolabile benzoin. *Tetrahedron.* 1997; 53:3917–3932.
13. Papageorgiou G, Barth A, Corrie JET. Flash photolytic release of alcohols from photolabile carbamates or carbonates is rate-limited by decarboxylation of the photoproduct. *Photochem. Photobiol. Sci.* 2005; 4:216–220. [PubMed: 15696240]
14. Breiting H-GA, Wieboldt R, Ramesh D, Carpenter BK, Hess GP. Synthesis and Characterization of Photolabile Derivatives of Serotonin for Chemical Kinetic Investigations of the Serotonin 5-HT₃ Receptor. *Biochemistry.* 2000; 39:5500–5508. [PubMed: 10820023]
15. Xia P, Bungay PM, Gibson CC, Kovbasnjuk ON, Spring KR. Diffusion coefficients in the lateral intercellular spaces of Madin-Darby canine kidney cell epithelium determined with caged compounds. *Biophys. J.* 1998; 74:3302–3312. [PubMed: 9635784]
16. Furuta T, Wang SSH, Dantzker JL, Dore TM, Bybee WJ, Callaway EM, Denk W, Tsien RY. Brominated 7-hydroxycoumarin-4-ylmethyls: photolabile protecting groups with biologically

- useful cross-sections for two photon photolysis. *Proc. Natl. Acad. Sci. U.S.A.* 1999; 96:1193–1200. [PubMed: 9990000]
17. Kiskin NI, Chillingworth R, McCray JA, Piston D, Ogden D. The efficiency of two-photon photolysis of a "caged" fluorophore *o*-1-(2-nitrophenyl)ethylpyranine, in relation to photodamage of synaptic terminals. *Eur. Biophys. J.* 2002; 30:588–604. [PubMed: 11908850]
18. Weichman BM, Notides AC. Analysis of estrogen receptor activation by its [3H]estradiol dissociation kinetics. *Biochemistry.* 1979; 18:220–225. [PubMed: 420773]
19. Housden HR, Skipp PJS, Crump MP, Broadbridge RJ, Crabbe T, Perry MJ, Gore MG. Investigation of the kinetics and order of tyrosine phosphorylation in the T-cell receptor ζ chain by the protein tyrosine kinase Lck. *Eur. J. Biochem.* 2003; 270:2369–2376. [PubMed: 12755691]
20. Dudai Y. High-affinity octopamine receptors revealed in *Drosophila* by binding of [3H]octopamine. *Neurosci. Lett.* 1982; 28:163–167. [PubMed: 6803195]
21. Raisinghani M, Pabbidi RM, Premkumar LS. Activation of transient receptor potential vanilloid 1 (TRPV1) by resiniferatoxin. *J Physiol.* 2005; 567:771–786. [PubMed: 16037081]
22. Solt K, Ruesch D, Forman SA, Davies PA, Raines DE. Differential effects of serotonin and dopamine on human 5-HT3A receptor kinetics: interpretation within an allosteric kinetic model. *J Neurosci.* 2007; 27:13151–13160. [PubMed: 18045909]
23. Vilardaga J-P, Romero G, Feinstein TN, Wehbi VL. Kinetics and dynamics in the G protein-coupled receptor signaling cascade. *Methods Enzymol.* 2013; 522:337–363. [PubMed: 23374192]
24. Boahen YO, MacDonald GM. A concise approach to caged serotonin for Fourier transform infrared (FT-IR) difference photolysis studies. *J Ghana Sci. Assoc.* 2005; 7:54–59.
25. Zayat L, Salierno M, Etchenique R. Ruthenium(II) Bipyridyl Complexes as Photolabile Caging Groups for Amines. *Inorg. Chem.* 2006; 45:1728–1731. [PubMed: 16471986]
26. Schaal J, Dekowski B, Wiesner B, Eichhorst J, Marter K, Vargas C, Keller S, Eremina N, Barth A, Baumann A, Eisenhardt D, Hagen V. Coumarin-Based Octopamine Phototriggers and their Effects on an Insect Octopamine Receptor. *Chem Bio Chem.* 2012; 13:1458–1464.
27. Fernandes MJG, Costa SPG, Goncalves MST. Synthesis and light triggered release of catecholamines from pyrenylmethyl carbamate cages. *New J. Chem.* 2013; 37:2369–2376.
28. Gilbert D, Funk K, Dekowski B, Lechler R, Keller S, Moehrlen F, Frings S, Hagen V. Caged capsaicins: new tools for the examination of TRPV1 channels in somatosensory neurons. *Chem Bio Chem.* 2007; 8:89–97.
29. Van Ryssen MP, Avlonitis N, Giniatullin R, McDougall C, Carr JL, Stanton-Humphreys MN, Borgstroem ELA, Brown CTA, Fayuk D, Surin A, Niittykoski M, Khiroug L, Conway SJ. Synthesis, photolysis studies and in vitro photorelease of caged TRPV1 agonists and antagonists. *Org. Biomol. Chem.* 2009; 7:4695–4707. [PubMed: 19865707]
30. Carr JL, Wease KN, Van Ryssen MP, Paterson S, Agate B, Gallagher KA, Brown CTA, Scott RH, Conway SJ. In vitro photo-release of a TRPV1 agonist. *Bioorg. Med. Chem. Lett.* 2006; 16:208–212. [PubMed: 16213723]
31. Hayashi, K-i; Hashimoto, K.; Kusaka, N.; Yamazoe, A.; Fukaki, H.; Tasaka, M.; Nozaki, H. Caged gene-inducer spatially and temporally controls gene expression and plant development in transgenic Arabidopsis plant. *Bioorg. Med. Chem. Lett.* 2006; 16:2470–2474. [PubMed: 16480868]
32. Russell AG, Sadler MJ, Laidlaw HJ, Gutierrez-Loriente A, Wharton CW, Carteau D, Bassani DM, Snaith JS. Photorelease of tyrosine from α -carboxy-6-nitroveratryl (α CNV) derivatives. *Photochem. Photobiol. Sci.* 2012; 11:556–563. [PubMed: 22249211]
33. Il'ichev YV, Schwoerer MA, Wirz J. Photochemical Reaction Mechanisms of 2-Nitrobenzyl Compounds: Methyl Ethers and Caged ATP. *J Am. Chem. Soc.* 2004; 126:4581–4595. [PubMed: 15070376]
34. Fedoryak OD, Dore TM. Brominated hydroxyquinoline as a photolabile protecting group with sensitivity to multiphoton excitation. *Org. Lett.* 2002; 4:3419–3422. [PubMed: 12323033]
35. Zhu Y, Pavlos CM, Toscano JP, Dore TM. 8-Bromo-7-hydroxyquinoline as a Photoremovable Protecting Group for Physiological Use: Mechanism and Scope. *J Am. Chem. Soc.* 2006; 128:4267–4276. [PubMed: 16569001]

36. Davis MJ, Kragor CH, Reddie KG, Wilson HC, Zhu Y, Dore TM. Substituent Effects on the Sensitivity of a Quinoline Photoremovable Protecting Group to One- and Two-Photon Excitation. *J Org. Chem.* 2009; 74:1721–1729. [PubMed: 19140722]
37. Abate-Pella D, Zeliadt NA, Ochocki JD, Warmka JK, Dore TM, Blank DA, Wattenberg EV, Distefano MD. Photochemical Modulation of Ras-Mediated Signal Transduction Using Caged Farnesyltransferase Inhibitors: Activation via One- and Two-Photon Excitation. *Chem Bio Chem.* 2012; 13:1009–1016.
38. Ma J, Rea AC, An H, Ma C, Guan X, Li M-D, Su T, Yeung CS, Harris KT, Zhu Y, Nganga JL, Fedoryak OD, Dore TM, Phillips DL. Unraveling the Mechanism of the Photodeprotection Reaction of 8-Bromo- and 8-Chloro-7-hydroxyquinoline Caged Acetates. *Chem.—Eur. J.* 2012; 18:6854–6865. [PubMed: 22511356]
39. Vandenberg LN, Blackiston DJ, Rea AC, Dore TM, Levin M. Left-right patterning in *Xenopus* conjoined twin embryos requires serotonin signaling and gap junctions. *Int. J. Dev. Biol.* 2014; 58:799–809. [PubMed: 25896280]
40. Gostkowski ML, Allen R, Plenert ML, Okerberg E, Gordon MJ, Shear JB. Multiphoton-excited serotonin photochemistry. *Biophys. J.* 2004; 86:3223–3229. [PubMed: 15111435]
41. Ma J, Cheng SC, An H, Li M-D, Ma C, Rea AC, Zhu Y, Nganga JL, Dore TM, Phillips DL. Comparison of the Absorption, Emission, and Resonance Raman Spectra of 7-Hydroxyquinoline and 8-Bromo-7-Hydroxyquinoline Caged Acetate. *J Phys. Chem. A.* 2011; 115:11632–11640. [PubMed: 21905734]
42. Warther D, Gug S, Specht A, Bolze F, Nicoud JF, Mourot A, Goeldner M. Two-photon uncaging: New prospects in neuroscience and cellular biology. *Bioorg. Med. Chem.* 2010; 18:7753–7758. [PubMed: 20554207]
43. Dore, TM.; Wilson, HC. Photosensitive Molecules for Controlling Biological Function. Chambers, JJ.; Kramer, RH., editors. New York: Humana Press; 2011. p. 57-92.
44. Bort G, Gallavardin T, Ogden D, Dalko PI. From One-Photon to Two-Photon Probes: "Caged" Compounds, Actuators, and Photoswitches. *Angew. Chem., Int. Ed.* 2013; 52:4526–4537.

Figure 1.
Some biologically active phenols.

Figure 2.
Photoactivatable, biologically relevant phenols from the literature.

1a: BHQ-O-5HT (X = Br)
1b: CyHQ-O-5HT (X = CN)

2a: BHQ-octopamine (X = Br)
2b: CyHQ-octopamine (X = CN)

3a: BHQ-capsaicin (X = Br)
3b: CyHQ-capsaicin (X = CN) } R =

4a: BHQ-VNA (X = Br)
4b: CyHQ-VNA (X = CN) } R = (CH₂)₇CH₃
5a: BHQ-VAA (X = Br)
5b: CyHQ-VAA (X = CN) } R = CH₃

6a: BHQ-estradiol (X = Br)
6b: CyHQ-estradiol (X = CN)

7a: BHQ-tyrosine (X = Br)
7b: CyHQ-tyrosine (X = CN)

Figure 3.
Photoactivatable, biologically relevant phenols prepared.

Figure 4.

Time courses for the photolysis of the photoactivatable phenols at 365 nm (1PE) and the rise of the biologically active phenols. Top: BHQ series, Bottom: CyHQ series. The percentage remaining is the average of at least 3 runs, and the error bars represent the standard deviation of the measurement. Lines are least-squares fits of a single exponential decay or a single exponential rise to max.

Figure 5.

Reaction progress curves for the photolysis of the photoactivatable phenols at 740 nm (2PE). Top: BHQ series, Bottom: CyHQ series. The percentage remaining is the average of at least 3 runs, and the error bars represent the standard deviation of the measurement. Lines are least-squares fits of a single exponential decay or a single exponential rise to max.

Scheme 1.

Synthesis and photolysis of hydroxyquinoline-protected phenols.

Table 1

Photophysical and photochemical data for the photoactivatable phenols

Photoactivatable Phenol	λ_{max} (nm)	Molar Absorptivity (ϵ) ^a	Quantum Efficiency (Q_u)	Sensitivity ($Q_u \times \epsilon$)	2-Photon Uncaging Action Cross-Section (δ_u) ^b
BHQ- <i>O</i> -5HT 1a ³	368	2000	0.30	600	0.50
CyHQ- <i>O</i> -5HT 1b	372	5700	0.39	2223	0.23
BHQ-Octopamine 2a	368	2700	0.36	972	0.49
CyHQ-Octopamine 2b	372	5400	0.34	1944	0.18
BHQ-VAA 5a	370	2700	0.18	486	0.61
CyHQ-VAA 5b	368	6100	0.41	2501	0.24
BHQ-Estradiol 6a	369	2400	—	—	—
CyHQ-Estradiol 6b	373	6200	0.29	1798	0.24
BHQ-Tyrosine 7a	366	2600	0.32	832	0.61
CyHQ-tyrosine 7b	371	4900	0.38	1862	0.36

^a $\text{M}^{-1} \text{cm}^{-1}$, $\lambda = 370 \text{ nm}$,^b $\text{GM} = 10^{-50} \text{ cm}^4 \text{ s/photon}$