Published in final edited form as:

Prostate. 2010 September 1; 70(12): 1265–1273. doi:10.1002/pros.21162.

Activation of FOXO1 Is Critical for the Anticancer Effect of Methylseleninic Acid in Prostate Cancer Cells

Haitao Zhang^{1,2,3,*}, Jian Fang^{1,2}, Dian Yao⁴, Yue Wu⁴, Clement Ip⁴, and Yan Dong^{2,3,5}

¹Department of Pathology and Laboratory Medicine, Tulane University School of Medicine, New Orleans, Louisiana

²Tulane Cancer Center, Tulane University School of Medicine, New Orleans, Louisiana

³School of Basic Medicine, Jilin University, Changchun, Jilin, PR China

⁴Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, New York

⁵Department of Structural and Cellular Biology, Tulane University School of Medicine, New Orleans, Louisiana

Abstract

BACKGROUND—Previous studies have demonstrated that physiological concentrations of methylseleninic acid (MSA) inhibits the growth of prostate cancer cells. The growth inhibitory effect could be attributed to cell cycle block and apoptosis induction. The current study was designed to investigate the involvement of forkhead box O1 (FOXO1) in the anticancer effect of MSA.

METHODS—LNCaP and LAPC-4 cells were treated with 10 μ M MSA for various time points, and the expression of FOXO1 was analyzed by qRT-PCR and Western blotting. FOXO1 activity was determined by a luciferase construct containing FOXO binding sites. The *trans*-activation activity of the androgen receptor (AR) was determined by the ARE-luciferase assay. FOXO1 gene silencing was achieved by using a small interfering RNA (siRNA).

RESULTS—MSA treatment led to a rapid and robust increase of FOXO1 expression, as well as an increase of the FOXO1 transcriptional activity. Blocking FOXO1 activation by gene silencing abolished apoptosis induction by MSA, suggesting FOXO1 plays a critical role in mediating the apoptotic effect of MSA. Recent studies have shown that FOXO1 and AR antagonize the actions of each other. We examined the consequence of FOXO1 induction on AR activity. Consistent with previous reports, we found that ectopic expression of FOXO1 suppressed the transcriptional activity of AR. Furthermore, FOXO1 silencing attenuated MSA suppression of AR activity, suggesting that FOXO1 induction contributes to suppression of AR signaling by MSA.

CONCLUSIONS—In prostate cancer cells, MSA activates the FOXO1 signaling pathway. FOXO1 activation is critical for the anticancer effects of MSA.

*Correspondence to: Dr. Haitao Zhang, Department of Pathology and Laboratory Medicine, Tulane University School of Medicine, 1430 Tulane Avenue SL-79, New Orleans, LA 70112. hzhang@tulane.edu.

Keywords

methylseleninic acid; FOXO1; apoptosis; androgen receptor

INTRODUCTION

Prostate cancer (PCa) is a significant public health problem that engenders huge medical care and human suffering costs in the United States. A number of case-control studies have demonstrated an inverse relationship between selenium status and prostate cancer risk [1–5]. One of the more important studies of selenium as a chemopreventive agent is the Nutritional Prevention of Cancer (NPC) trial initiated by Larry Clark [6,7]. The study was a randomized, double-blind, placebo-controlled trial involving 1,312 patients (mostly men) who were recruited initially because of a history of basal cell or squamous cell carcinoma of the skin. Individuals in the treatment arm were given 200 µg selenized yeast per day for a mean of 4.5 years. After a total follow-up of 8,271 person-years, selenium treatment did not decrease the recurrence of these non-melanoma skin cancers. However, patients receiving the supplement showed a much lower risk of developing total (HR = 0.75) or prostate cancer (HR = 0.48) [6,7].

Encouraged by the prostate cancer results of the NPC trial, the National Cancer Institute launched the Selenium and Vitamin E Cancer Prevention Trial (SELECT) in 2001. An interim data analysis after a median follow-up of 5.46 years suggested selenium, either alone or in combination with vitamin E, did not lower prostate cancer risk in this study population [8]. The trial was halted in October 2008, but the follow-up will continue for 3 more years.

The results of the SELECT have caused much controversy. One hot topic of debate is the formulation of selenium used in the SELECT. Selenomethionine, which was used in the SELECT, can be incorporated non-specifically into proteins in place of methionine [9]. Compartmentation into tissue proteins limits selenomethionine from being further metabolized. On the other hand, monomethylated forms of selenium, including methylseleninic acid (MSA) and methylselenocysteine (MSC), can be easily metabolized to methylselenol, which is considered to be the critical metabolite for the anticancer activity of selenium [9,10]. By itself, methylselenol is highly reactive and difficult to prepare. Therefore, the proximal precursors including MSA and MSC are superior to selenomethionine with regard to providing a steady stream of methylselenol. In fact, studies published before and after the launch of the SELECT have showed that MSA and MSC have stronger anticancer activities than selenomethionine [11–14]. While the conversion of MSC to methylselenol requires the action of β -lyase, MSA can be easily reduced to methylselenol through non-enzymatic reactions involving glutathione (GSH) or NADPH [15]. Due to the fact that epithelial cells express low level of β -lyase, MSA is 10 times more potent than MSC in affecting biological processes *in vitro* [13]. MSA is widely accepted to be the best reagent for delineating the molecular action of selenium in cell culture studies [16–19]. It also has excellent anticancer activity in animals [13,20,21].

In view of the above information, we believe that the potential of selenium compounds as chemopreventive agents for prostate cancer should not be dismissed. Unraveling the

mechanisms of action for these agents is urgent and will no doubt be helpful in rational design of future intervention trials. We and others have previously profiled selenium-induced gene expression changes in prostate cancer cells [17,22,23]. Based on the datasets generated from the microarray studies, we conducted a systematic data mining analysis, taking advantage of several publicly available clinical prostate cancer datasets, in order to gain new insights into novel molecular targets that may be relevant to the anticancer activity of MSA [24]. The analysis drew our attention to forkhead box O1 (FOXO1). We found that the expression of FOXO1 is consistently decreased in a large number of prostate cancer specimens, and the microarray analyses showed MSA up-regulates the expression of FOXO1 [24]. FOXO1 is a member of the FOXO family of transcription factors that induces the expression of pro-apoptotic genes including Fas ligand [25,26], bcl-2 family proteins [27–29], and TRAIL [30]. FOXO1 is also involved in cell cycle regulation [31]. FOXO1 is phosphorylated and suppressed by AKT [32,33], which is an important survival molecule for prostate cancer. In prostate cancer cells, androgen receptor (AR) interacts with FOXO1 and inhibits its activation of downstream targets [34]. The current study was designed to examine the role of FOXO1 in mediating the anticancer effect of MSA.

MATERIALS AND METHODS

Materials

MSA was purchased from PharmaSe (Lubbock, TX). Fetal bovine serum, RPMI 1640, and the Lipofectamine PLUS transfection reagents were purchased from Invitrogen (Carlsbad, CA). Immobilon PVDF membrane was purchased from Millipore (Bedford, MA) and ECL Western blotting detection reagent from Amersham Pharmacia Biotech (Arlington Heights, IL). For Western blotting analysis, the antiglyceraldehyde-3-phosphate dehydrogenase (GAPDH) antibody was from Chemicon (Temecula, CA) and anti-FOXO1 was from Cell Signaling (Danvers, MA). The Cell Death Detection ELISA kit was purchased from Roche Applied Science (Indianapolis, IN). The p3XIRS-luc reporter construct was kindly provided by Dr. Kun-Liang Guan at the University of Michigan, and the pcDNA3-FKHR expression vector was obtained from Dr. Frederic G. Barr at the University of Pennsylvania. The pcDNA3-AR-FL expression vector was a gift from Dr. Shuyun Yeh at the University of Rochester.

Cell Culture and Treatment

The human LNCaP prostate cancer cell line was obtained from the American Type Culture Collection (Manassas, VA). The LAPC-4 cell line was provided by Dr. Charles L. Sawyers at the University of California at Los Angeles Jonsson Comprehensive Cancer Center. Both LNCaP and LAPC-4 express AR and require androgen for their growth. Cells were cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum, 100 unit/ml of penicillin, 100 µg/ml of streptomycin, and 2mM of glutamine. In some experiments, cells were cultured in an androgen-defined condition by using charcoal-stripped FBS in the presence of 1 nM R1881 (a potent synthetic androgen). Treatment with MSA usually began at 72 hr after seeding, when the cultures were 60–80% confluent.

Quantitative Reverse transcription-Polymerase Chain Reaction (qRT-PCR)

The PCR primers and Taqman probes for β -actin, FOXO1, and AR were Assays-on-Demand products from Applied Biosystems (Foster City, CA). The PCR conditions were as follows: an initial incubation at 50°C for 2min, then a denaturation at 95°C for 10 min, followed by 40 cycles of 95°C for 15 sec and 60°C for 1 min. The relative quantitation of gene expression was done using the comparative CT (C_T) method [35]. Details of the procedure were described in our previous publication [22].

Transient Transfection and Reporter Gene Assay

Supercoiled plasmid DNAs were prepared by the Qiagen column procedure (Qiagen, Valencia, CA). Twenty-four hours before transfection, cells were trypsinized and seeded at a density to reach 90–95% confluency at the time of transfection. Transient transfection was carried out by using the Lipofectamine™ and Plus™ reagents (Invitrogen) per instruction of the manufacturer. After incubating with the transfection mixture for 3 hr, the cells were trypsinized and replated in triplicate into 6-well plates to achieve equal transfection efficiency. The cells were allowed to attach overnight before 10 μ M MSA was added to the culture medium. At 6 or 16 hr following treatment, cells were lysed with 1X Passive Lysis Buffer (Promega, Madison, WI), and the luciferase activity was assayed by using the Luciferase Assay System (Promega). Protein concentration in the cell extract was determined by using the bicinchoninic acid protein assay kit (Pierce, Rockford, IL). Luciferase activities were normalized to the protein concentration of the same sample. The transfection experiments were repeated three times.

Gene Silencing With siRNA

A small interfering RNA (siRNA) designed to target FOXO1 (Cat. # HSS103719) and a matching negative control oligonucleotide were purchased from Invitrogen. These oligonucleotides were transiently transfected into LNCaP cells by using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. At 48 hr posttransfection, 10 μ M MSA was added to the culture medium and the cells were treated for an additional 24 hr. RNA was prepared from the cells and qRT-PCR was performed to determine the efficiency of gene silencing.

Statistical Analysis

The Student's *t*-test was used to determine significant differences between different groups. Unless otherwise indicated, $P < 0.05$ was considered statistically significant. All analyses were two-tailed.

RESULTS

MSA Induces FOXO1 Expression

We first performed qRT-PCR and Western blotting to confirm the modulation of FOXO1 by MSA in LNCaP cells, as first noted from our microarray analysis [24]. Cells were treated with 10 μ M MSA for various lengths of time before they were lysed for RNA and protein purification. The qRT-PCR results are shown in Figure 1a. Induction of FOXO1 mRNA was

observed as early as 1 hr after exposure to MSA, suggesting that FOXO1 is a proximal target of MSA. The mRNA level peaked at 2 hr, then declined gradually with time, but still remained elevated at 24 hr. Western blotting of FOXO1 was carried out in LNCaP and LAPC-4 cells (Fig. 1b). No change in protein level was detected until at least after 3 hr. Thus the increases of FOXO1 protein appeared to lag behind the increases of the message, although the protein signal was decidedly stronger by 6 hr in cells treated with MSA.

MSA Induces the Transcriptional Activity of FOXO

As mentioned in the Introduction section, FOXO1 is a transcription factor. In order to study the effect of MSA on the activity of FOXO1 as a transcription factor, we transiently transfected LNCaP and LAPC-4 cells with a luciferase reporter construct, p3XIRS-luc. This construct has three tandem repeats of a FOXO1 binding element, the insulin-responsive sequence (IRS), inserted upstream of the minimal thymidine kinase promoter [33]. It is widely used as an indicator of the transcriptional activity of FOXO proteins. As shown in Figure 2a, the transcriptional activity of this reporter construct was induced by approximately two-fold in LNCaP cells after 6 hr of treatment with 10 μ M MSA. A pronounced induction (>5-fold) was observed in LNCaP after 16 hr of treatment ($P < 0.01$). Nearly identical results were obtained in LAPC-4 cells (Fig. 2b).

FOXO1 Gene Silencing Blocks MSA-Induced Apoptosis

MSA has been shown to induce apoptosis in prostate cancer cells by several groups, including ours [17–19,36]. The experiments described above suggested that MSA induces the FOXO1 signaling pathway, which is known to positively regulate apoptosis. To establish the role of FOXO1 in MSA-induced apoptosis, we employed the RNA interference technique to knockdown the expression of FOXO1. A commercially available siRNA targeting FOXO1 was obtained. To confirm the specificity of the siRNA, we performed a Local Alignment Search Tool (BLAST) against the entire human transcriptome using the sequence provided by the manufacturer. With the exception of FOXO1, the search identified no other homology with the siRNA sequence, including other FOXO members. When introduced into LNCaP cells, the FOXO1 siRNA, named siFOXO1 hereafter, was able to decrease the baseline expression of FOXO1 by approximately 50% (Fig. 3a). Consistent with our previous finding, a two-fold induction of FOXO1 was observed when the cells were treated with 10 μ M MSA for 24 hr (comparing columns 1 and 3). siFOXO1 was able to abolish this induction by MSA (comparing columns 3 and 4).

Apoptosis was quantitated in siRNA-transfected and MSA-treated cells by using an ELISA-based method. The result is shown in Figure 3b. In general, the level of apoptosis in these cells correlated well with the expression level of FOXO1 (Fig. 3a), confirming that FOXO1 plays an important role in apoptosis regulation. More importantly, when the induction of FOXO1 was blocked by the addition of siFOXO1, no induction of apoptosis was observed (Fig. 3b, comparing columns 1 and 4). These results suggest that FOXO1 is a key mediator of apoptosis induction by MSA.

FOXO1 Activation Suppresses AR *trans*-Activation

To examine the effect of FOXO1 activation on the transcriptional activity of AR, we transiently co-transfected LNCaP cells with a reporter construct containing three repeats of the androgen response element (ARE) ligated in tandem to the luciferase reporter, together with a FOXO1 expression vector, pcDNA3-FKHR, or the empty vector. The ARE-luciferase reporter assay is commonly used to assess the *trans*-activating activity of AR. Following transfection, cells were exposed to 1 nM R1881, a synthetic androgen, for 6 or 16 hr before they were lysed for luciferase assay. As shown in Figure 4a, the AR transcriptional activity was greatly stimulated by the addition of the ligand. In the presence of ectopically expressed FOXO1, the induction was significantly diminished (Fig. 4a, comparing columns 2 and 4 for both time points). Therefore, our results confirmed published studies showing that FOXO1 activation suppresses AR signaling [37–40].

FOXO1 Induction Contributes to AR Suppression by MSA

It has been found previously that MSA is a potent suppressor of AR signaling [22,23,41]. The mechanisms involved in suppression of AR signaling by MSA include reduction in AR mRNA transcription [22,41] and stability, increase in AR protein turnover reduction in AR translocation, inhibition of co-activator recruitment, and increased co-repressor recruitment to the promoters of AR-regulated genes [22,41,42]. The result from the previous section prompted us to investigate whether FOXO1 induction is a contributing factor for AR suppression by MSA. Once again, we employed the gene knockdown approach. LNCaP cells were co-transfected with the ARE-luciferase construct and siFOXO1, and treated with 0 or 10 μ M MSA. In the presence of the scrambled oligo, MSA suppressed AR activity by approximately 70% (Fig. 4b). This is in line with our previous observations [22]. However, when FOXO1 was silenced, the suppression was attenuated to about 60% ($P < 0.01$). This was further confirmed when we examined the modulation of prostate-specific antigen (PSA) expression by MSA in the presence or absence of siFOXO1 (data not shown). These results are in agreement with previous studies showing that MSA suppresses AR signaling through a multitude of mechanisms and identified FOXO1 activation as a novel mechanism contributing to the inhibition of AR *trans*-activation by MSA.

DISCUSSION

Despite the protective effect of selenium against prostate cancer demonstrated by the NPC study and several studies which showed selenium is very effective in switching off androgen signaling, recent results from the SELECT showed that selenium, alone or in combination with vitamin E, did not prevent prostate cancer in a randomized trial of 33,000 men at average risk [8]. Several potential reasons have been discussed to explain the discrepancy of the findings in SELECT and the NPC trial. In addition to the dose and formulation of selenium used in the trial, one important consideration is the baseline selenium level. The NPC trial showed that the protective effect of selenium was limited to patients with baseline serum selenium in the lower two tertiles [7]. The average baseline selenium level of the participants in SELECT was much higher than that observed in the NPC study. In fact, 78% of men in SELECT had baseline selenium above the range that selenium provided protection in the NPC trial (<121.6 ng/ml) [8]. Another important consideration is how selenium exerts

its anticancer activity. The Physicians' Health Study demonstrated an inverse association of plasma selenium level with risk of advanced prostate cancer, not localized prostate cancer, suggesting selenium might function by slowing down tumor progression [5]. In view of the above information, we believe that the negative finding by SELECT should not be simply interpreted as selenium is ineffective against prostate cancer. Instead, the outcome of this trial, as well as those of several recently published clinical trials [43–45], may indicate that it is difficult to find a single chemoprevention strategy which can benefit the general population. There is an urgent need to re-evaluate all the pre-clinical and clinical evidence to identify the subset of patients that are most likely to benefit from selenium supplementation.

This report is the first to show that MSA induces the expression of FOXO1. The elevated expression is accompanied by an increase of the FOXO transcriptional activity. We further demonstrated that FOXO1 is a key mediator of apoptosis induction by MSA. The above conclusion is supported by the following observations. First, FOXO1 induction occurred very early following MSA treatment, suggesting that FOXO1 is a proximal target of MSA. Second, MSA failed to induce apoptosis when FOXO1 stimulation was abolished by the addition of a FOXO1-specific siRNA. There are two major cell death signaling pathways, one triggered through death receptors (the extrinsic pathway), and the other through the mitochondria (the intrinsic pathway). A signature of the intrinsic pathway is the release of cytochrome *c* from the mitochondria, which is regulated by the Bcl-2 family of proteins. As a pro-apoptotic member of the Bcl-2 family, Bim functions by antagonizing the actions of the anti-apoptotic Bcl-2 and Bcl-XL. Both TRAIL and TRADD are associated with the extrinsic pathway. MSA has been shown to activate caspases that are involved in both the intrinsic and extrinsic apoptosis signaling pathways [18,36]. We are currently working on identifying the pro-apoptotic targets of FOXO1 that are induced by MSA. In addition to its role in regulating apoptosis, FOXO1 also plays an important role in cell cycle control. It up-regulates the expression of p27 [31,46] and down-regulates the expression of cyclins D1 and D2 [47,48], a pattern consistent with the G1 cell cycle block by MSA [17,49]. Therefore, it is possible that FOXO1 also mediates the cell cycle effects of MSA. Research along this line is currently ongoing in our laboratories.

Several mechanisms could account for the induction of FOXO1 signaling by MSA. One is through the induction of FOXO1 expression, as evidenced by the increased transcript and protein levels following MSA treatment. It has been shown that AR interacts and suppresses the activity of FOXO1 in prostate cancer cells [34,50]. Another potential mechanism of MSA activation of FOXO1 is through decreasing AR expression and thereby relieving the inhibition of FOXO1 by AR. This is supported by the fact that ectopic expression of AR could attenuate the induction of FOXO1 activity by MSA (data not shown). Yet, there might be a third mechanism by which MSA induces FOXO1. A key regulator of cellular FOXO1 activity is Akt, an important survival molecule for many cancer types, including prostate cancer. Akt phosphorylates FOXO1, which leads to nuclear exclusion and proteosomal degradation of FOXO1 [33]. MSA has been shown to suppress the PI3K-Akt signaling pathway [19,51–53]. Therefore, it is possible that relieving the suppression by Akt may contribute to MSA induction of FOXO1. Further experimental evidence is needed to support this hypothesis.

In agreement with previous reports [37,38,40,54], our data showed increased abundance of FOXO1 leads to decreased AR activity. Together with the well-documented AR inhibition of FOXO1 activity, it appears that in prostate cancer cells, the AR and FOXO1 signaling pathways antagonize the action of each other. The outcome is likely determined by the relative abundance of AR and FOXO1 proteins. When AR signaling dominates, the growth inhibitory signals conveyed by FOXO1 are muted, and the cells undergo proliferation. On the other hand, when FOXO1 signaling dominates, the antiproliferative and pro-apoptotic signaling prevail. When prostate cancer cells are exposed to MSA, AR signaling is suppressed whilst FOXO1 signaling is stimulated. By doing so, MSA could shift the balance heavily in favor of FOXO1, leading to cell cycle arrest and apoptosis. Therefore, modulating the crosstalk between AR and FOXO1 could be the key mechanism underlining the anticancer effect of MSA in the prostate.

CONCLUSIONS

The work described herein demonstrates that MSA activates FOXO1 signaling pathway. FOXO1 plays a critical role in mediating the apoptotic activity of MSA, and also contributes to the suppression of androgen signaling by MSA. This study enhances our understanding of the molecular mechanisms of the anticancer activity of MSA, which will be critical for designing future prostate cancer intervention studies with MSA.

ACKNOWLEDGMENTS

This work was supported by DOD New Investigator grant no. W81XWH-05-1-0598 (H. Zhang), American Cancer Society grant no. RSG-07-218-01-TBE (Y. Dong), National Cancer Institute grant nos. K01 CA114252 (Y. Dong), P01 CA126804 (C. Ip, H. Zhang, and Y. Dong), start-up funds from the Louisiana Cancer Research Consortium (H. Zhang and Y. Dong).

Abbreviations

AR	androgen receptor
ARE	androgen response element
FOXO1	forkhead box O1
MSA	methylseleninic acid
NPC	Nutritional Prevention of Cancer
PCa	prostate cancer
PSA	prostate-specific antigen
qRT-PCR	quantitative reverse transcription-polymerase chain reaction
SELECT	Selenium and Vitamin E Chemoprevention Trial
siRNA	small interfering RNA

REFERENCES

1. Yoshizawa K, Willett WC, Stampfer MJ, Spiegelman D, Rimm EB, Giovannucci E. Study of prediagnostic selenium level in toenails and the risk of advanced prostate cancer. *J Natl Cancer Inst.* 1998; 90:1219–1224. [PubMed: 9719083]
2. Helzlsouer KJ, Huang HY, Alberg AJ, Hoffman S, Burke A, Norkus EP, Morris JS, Comstock GW. Association between α -tocopherol, γ -tocopherol, selenium, and subsequent prostate cancer. *J Natl Cancer Inst.* 2000; 92:2018–2023. [PubMed: 11121464]
3. Nomura AMY, Lee J, Stemmermann GN, Combs GF. Serum selenium and subsequent risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev.* 2000; 9:883–887. [PubMed: 11008904]
4. Brooks JD, Jeffrey ME, Chan DW, Sokoll LJ, Landis P, Nelson WG, Muller D, Andres R, Carter HB. Plasma selenium level before diagnosis and the risk of prostate cancer development. *J Urol.* 2001; 166:2034–2038. [PubMed: 11696701]
5. Li H, Stampfer MJ, Giovannucci EL, Morris JS, Willett WC, Gaziano JM, Ma J. A prospective study of plasma selenium levels and prostate cancer risk. *J Natl Cancer Inst.* 2004; 96:696–703. [PubMed: 15126606]
6. Clark LC, Combs GF Jr, Turnbull BW, Slate EH, Chalker DK, Chow J, Davis LS, Glover RA, Graham GF, Gross EG, Krongrad A, Leshner JL Jr, Park HK, Sanders BB Jr, Smith CL, Taylor JR. Effects of selenium supplementation for cancer prevention in patients with carcinoma of the skin. A randomized controlled trial. Nutritional Prevention of Cancer Study Group. *JAMA.* 1996; 276(24): 1957–1963. [PubMed: 8971064]
7. Duffield-Lillico AJ, Reid ME, Turnbull BW, Combs GF Jr, Slate EH, Fischbach LA, Marshall JR, Clark LC. Baseline characteristics and the effect of selenium supplementation on cancer incidence in a randomized clinical trial: A summary report of the nutritional prevention of cancer trial. *Cancer Epidemiol Biomarkers Prev.* 2002; 11:630–639. [PubMed: 12101110]
8. Lippman SM, Klein EA, Goodman PJ, Lucia MS, Thompson IM, Ford LG, Parnes HL, Minasian LM, Gaziano JM, Hartline JA, Parsons JK, Bearden JD III, Crawford ED, Goodman GE, Claudio J, Winquist E, Cook ED, Karp DD, Walther P, Lieber MM, Kristal AR, Darke AK, Arnold KB, Ganz PA, Santella RM, Albanes D, Taylor PR, Probstfield JL, Jagpal TJ, Crowley JJ, Meyskens FL Jr, Baker LH, Coltman CA Jr. Effect of selenium and vitamin E on risk of prostate cancer and other cancers: The Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA.* 2009; 301(1): 39–51. [PubMed: 19066370]
9. Ip C. Lessons from basic research in selenium and cancer prevention. *J Nutr.* 1998; 128:1845–1854. [PubMed: 9808633]
10. Ip C, Ganther HE. Activity of methylated forms of selenium in cancer prevention. *Cancer Res.* 1990; 50:1206–1211. [PubMed: 2105164]
11. Ip C, Birringer M, Block E, Kotreba M, Tyson JF, Uden PC, Lisk DJ. Chemical speciation influences comparative activity of selenium-enriched garlic and yeast in mammary cancer prevention. *J Agric Food Chem.* 2000; 48(6):2062–2070. [PubMed: 10888499]
12. Medina D, Thompson H, Ganther H, Ip C. Se-methylselenocysteine: A new compound for chemoprevention of breast cancer. *Nutr Cancer.* 2001; 40(1):12–17. [PubMed: 11799917]
13. Ip C, Thompson HJ, Zhu Z, Ganther HE. In vitro and in vivo studies of methylseleninic acid: Evidence that a monomethylated selenium metabolite is critical for cancer chemoprevention. *Cancer Res.* 2000; 60:2882–2886. [PubMed: 10850432]
14. Li G, Lee H, Wang Z, Hu H, Liao J, Watts J, Combs GJ, Lü J. Superior in vivo inhibitory efficacy of methylseleninic acid against human prostate cancer over selenomethionine or selenite. *Carcinogenesis.* 2008; 29(5):1005–1012. [PubMed: 18310093]
15. Ganther HE, Lawrence JR. Chemical transformations of selenium in living organisms. Improved forms of selenium for cancer prevention. *Tetrahedron.* 1997; 53(36):12299–12310.
16. Dong Y, Ganther HE, Stewart C, Ip C. Identification of molecular targets associated with selenium-induced growth inhibition in human breast cells using cDNA microarrays. *Cancer Res.* 2002; 62(3):708–714. [PubMed: 11830524]

17. Dong Y, Zhang H, Hawthorn L, Ganther HE, Ip C. Delineation of the molecular basis for selenium-induced growth arrest in human prostate cancer cells by oligonucleotide array. *Cancer Res.* 2003; 63(1):52–59. [PubMed: 12517777]
18. Jiang C, Wang Z, Ganther H, Lu J. Caspases as key executors of methyl selenium-induced apoptosis (anoikis) of DU-145 prostate cancer cells. *Cancer Res.* 2001; 61(7):3062–3070. [PubMed: 11306488]
19. Jiang C, Wang Z, Ganther H, Lu J. Distinct effects of methylseleninic acid versus selenite on apoptosis, cell cycle, and protein kinase pathways in DU145 human prostate cancer cells. *Mol Cancer Ther.* 2002; 1(12):1059–1066. [PubMed: 12481429]
20. Dong Y, Ip C, Ganther H. Evidence of a field effect associated with mammary cancer chemoprevention by methylseleninic acid. *Anticancer Res.* 2002; 22(1A):27–32. [PubMed: 12017302]
21. Ip C, Dong Y, Ganther HE. New concepts in selenium chemoprevention. *Cancer Metastasis Rev.* 2002; 21:281–289. [PubMed: 12549766]
22. Dong Y, Lee SO, Zhang H, Marshall J, Gao AC, Ip C. Prostate specific antigen expression is down-regulated by selenium through disruption of androgen receptor signaling. *Cancer Res.* 2004; 64(1):19–22. [PubMed: 14729601]
23. Zhao H, Whitfield ML, Xu T, Botstein D, Brooks JD. Diverse effects of methylseleninic acid on the transcriptional program of human prostate cancer cells. *Mol Biol Cell.* 2004; 15(2):506–519. [PubMed: 14617803]
24. Zhang H, Dong Y, Marshall J, Hawthorn L, Nowak N, Gao AC, Ip C. Microarray data mining for potential selenium targets in chemoprevention of prostate cancer. *Cancer Genomics Proteomics.* 2005; 2:97–114. [PubMed: 18548127]
25. Kavurma MM, Khachigian LM. Signaling and transcriptional control of Fas ligand gene expression. *Cell Death Differ.* 2003; 10(1):36–44. [PubMed: 12655294]
26. Suhara T, Kim HS, Kirshenbaum LA, Walsh K. Suppression of Akt signaling induces Fas ligand expression: Involvement of caspase and Jun kinase activation in Akt-mediated Fas ligand regulation. *Mol Cell Biol.* 2002; 22(2):680–691. [PubMed: 11756562]
27. Dijkers PF, Medema RH, Lammers JW, Koenderman L, Coffey PJ. Expression of the pro-apoptotic Bcl-2 family member Bim is regulated by the forkhead transcription factor FKHR-L1. *Curr Biol.* 2000; 10(19):1201–1204. [PubMed: 11050388]
28. Gilley J, Coffey PJ, Ham J. FOXO transcription factors directly activate bim gene expression and promote apoptosis in sympathetic neurons. *J Cell Biol.* 2003; 162(4):613–622. [PubMed: 12913110]
29. Tang TT, Dowbenko D, Jackson A, Toney L, Lewin DA, Dent AL, Lasky LA. The forkhead transcription factor AFX activates apoptosis by induction of the BCL-6 transcriptional repressor. *J Biol Chem.* 2002; 277(16):14255–14265. [PubMed: 11777915]
30. Modur V, Nagarajan R, Evers BM, Milbrandt J. FOXO proteins regulate tumor necrosis factor-related apoptosis inducing ligand expression. Implications for PTEN mutation in prostate cancer. *J Biol Chem.* 2002; 277(49):47928–47937. [PubMed: 12351634]
31. Nakamura N, Ramaswamy S, Vazquez F, Signoretti S, Loda M, Sellers WR. Forkhead transcription factors are critical effectors of cell death and cell cycle arrest downstream of PTEN. *Mol Cell Biol.* 2000; 20(23):8969–8982. [PubMed: 11073996]
32. Guo S, Rena G, Cichy S, He X, Cohen P, Unterman T. Phosphorylation of serine 256 by protein kinase B disrupts transactivation by FKHR and mediates effects of insulin on insulin-like growth factor-binding protein-1 promoter activity through a conserved insulin response sequence. *J Biol Chem.* 1999; 274(24):17184–17192. [PubMed: 10358076]
33. Tang ED, Nunez G, Barr FG, Guan KL. Negative regulation of the forkhead transcription factor FKHR by Akt. *J Biol Chem.* 1999; 274(24):16741–16746. [PubMed: 10358014]
34. Li P, Lee H, Guo S, Unterman TG, Jenster G, Bai W. AKT-independent protection of prostate cancer cells from apoptosis mediated through complex formation between the androgen receptor and FKHR. *Mol Cell Biol.* 2003; 23(1):104–118. [PubMed: 12482965]
35. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-CT} method. *Methods.* 2001; 25(4):402–408. [PubMed: 11846609]

36. Zu K, Ip C. Synergy between selenium and vitamin E in apoptosis induction is associated with activation of distinctive initiator caspases in human prostate cancer cells. *Cancer Res.* 2003; 63:6988–6995. [PubMed: 14583501]
37. Dong XY, Chen C, Sun X, Guo P, Vessella RL, Wang RX, Chung LW, Zhou W, Dong JT. FOXO1A is a candidate for the 13q14 tumor suppressor gene inhibiting androgen receptor signaling in prostate cancer. *Cancer Res.* 2006; 66(14):6998–7006. [PubMed: 16849544]
38. Fan W, Yanase T, Morinaga H, Okabe T, Nomura M, Daitoku H, Fukamizu A, Kato S, Takayanagi R, Nawata H. Insulin-like growth factor 1/insulin signaling activates androgen signaling through direct interactions of FOXO1 with androgen receptor. *J Biol Chem.* 2007; 282(10):7329–7338. [PubMed: 17202144]
39. Liu P, Li S, Gan L, Kao TP, Huang H. A transcription-independent function of FOXO1 in inhibition of androgen-independent activation of the androgen receptor in prostate cancer cells. *Cancer Res.* 2008; 68(24):10290–10299. [PubMed: 19074897]
40. Ma Q, Fu W, Li P, Nicosia SV, Jenster G, Zhang X, Bai W. FoxO1 mediates PTEN suppression of androgen receptor N- and C-terminal interactions and coactivator recruitment. *Mol Endocrinol.* 2009; 23(2):213–225. [PubMed: 19074551]
41. Dong Y, Zhang H, Gao AC, Marshall JR, Ip C. Androgen receptor signaling intensity is a key factor in determining the sensitivity of prostate cancer cells to selenium inhibition of growth and cancer-specific biomarkers. *Mol Cancer Ther.* 2005; 4:1047–1055. [PubMed: 16020662]
42. Chun JY, Nadiminty N, Lee SO, Onate SA, Lou W, Gao AC. Mechanisms of selenium down-regulation of androgen receptor signaling in prostate cancer. *Mol Cancer Ther.* 2006; 5(4):913–918. [PubMed: 16648561]
43. Zhang SM, Cook NR, Albert CM, Gaziano JM, Buring JE, Manson JE. Effect of combined folic acid, vitamin B6, and vitamin B12 on cancer risk in women: A randomized trial. *JAMA.* 2008; 300(17):2012–2021. [PubMed: 18984888]
44. Gaziano JM, Glynn RJ, Christen WG, Kurth T, Belanger C, MacFadyen J, Bubes V, Manson JE, Sesso HD, Buring JE. Vitamins E and C in the prevention of prostate and total cancer in men: The Physicians' Health Study II randomized controlled trial. *JAMA.* 2009; 301(1):52–62. [PubMed: 19066368]
45. Lin J, Cook NR, Albert C, Zaharris E, Gaziano JM, Van DM, Buring JE, Manson JE. Vitamins C and E and beta carotene supplementation and cancer risk: A randomized controlled trial. *J Natl Cancer Inst.* 2009; 101(1):14–23. [PubMed: 19116389]
46. Medema RH, Kops GJ, Bos JL, Burgering BM. AFX-like Forkhead transcription factors mediate cell-cycle regulation by Ras and PKB through p27kip1. *Nature.* 2000; 404(6779):782–787. [PubMed: 10783894]
47. Ramaswamy S, Nakamura N, Sansal I, Bergeron L, Sellers WR. A novel mechanism of gene regulation and tumor suppression by the transcription factor FKHR. *Cancer Cell.* 2002; 2(1):81–91. [PubMed: 12150827]
48. Schmidt M, Fernandez dM, van der HA, Klompmaker R, Kops GJ, Lam EW, Burgering BM, Medema RH. Cell cycle inhibition by FoxO forkhead transcription factors involves downregulation of cyclin D. *Mol Cell Biol.* 2002; 22(22):7842–7852. [PubMed: 12391153]
49. Wang Z, Jiang C, Lu J. Induction of caspase-mediated apoptosis and cell-cycle G1 arrest by selenium metabolite methylselenol. *Mol Carcinog.* 2002; 34(3):113–120. [PubMed: 12112305]
50. Huang H, Muddiman DC, Tindall DJ. Androgens negatively regulate forkhead transcription factor FKHR (FOXO1) through a proteolytic mechanism in prostate cancer cells. *J Biol Chem.* 2004; 279(14):13866–13877. [PubMed: 14726521]
51. Hu H, Jiang C, Li G, Lu J. PKB/AKT and ERK regulation of caspase-mediated apoptosis by methylseleninic acid in LNCaP prostate cancer cells. *Carcinogenesis.* 2005; 26(8):1374–1381. [PubMed: 15845651]
52. Wu Y, Zu K, Warren MA, Wallace PK, Ip C. Delineating the mechanism by which selenium deactivates Akt in prostate cancer cells. *Mol Cancer Ther.* 2006; 5(2):246–252. [PubMed: 16505097]
53. Lee JH, Shin SH, Kang S, Lee YS, Bae S. A novel activation-induced suicidal degradation mechanism for Akt by selenium. *Int J Mol Med.* 2008; 21(1):91–97. [PubMed: 18097621]

54. Liu P, Kao TP, Huang H. CDK1 promotes cell proliferation and survival via phosphorylation and inhibition of FOXO1 transcription factor. *Oncogene*. 2008; 27(34):4733–4744. [PubMed: 18408765]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 1. Effect of methylseleninic acid (MSA) on FOXO1 expression. **a:** Change of FOXO1 mRNA in LNCaP cells as a function of time of MSA treatment, determined by qRT-PCR. The results are shown as mean $\pm$ standard error of mean (SEM). **b:** Western analysis of FOXO1 protein level as a function of time of MSA treatment, in both LNCaP and LAPC-4 cells. The band intensity was quantified by volume densitometry and normalized to that of GAPDH. The results were expressed as fold induction over untreated.

Fig. 2.

Induction of FOXO transcriptional activity by MSA. LNCaP (a) and LAPC-4 (b) cells were transfected with the p3XIRS-luc construct and treated with 10 μ M MSA for the indicated times. At the end of treatment, cells were lysed for luciferase assay. Total protein concentration was also determined and used to normalize the luciferase reading. The results were expressed as mean $\pm$ SEM; *, $P < 0.05$.

Fig. 3. Effect of FOXO1 gene silencing on MSA-induced apoptosis. **a:** qRT-PCR analysis of FOXO1 expression in cells transfected with small interference RNAs (siRNAs) and treated with or without MSA. The data were expressed as fold relative to the scrambled, untreated control. **b:** Quantitation of apoptotic cell death by an ELISA method.

a FOXO1 overexpression on AR activity

b FOXO1 knockdown on suppression of AR activity by MSA

Fig. 4.

Induction of FOXO1 contributes to AR suppression by MSA. **a:** Increased expression of FOXO1 reduced the transcriptional activity of AR. LNCaP cells were co-transfected with the ARE-luciferase reporter construct and either the pcDNA3-FKHR or the pcDNA3 vector, and treated with 1nM R1881 for the indicated times. **b:** FOXO1 knockdown attenuated the suppression of AR trans-activation by MSA. LNCaP cells were co-transfected with the ARE-luciferase construct and either the scrambled control or siFOXO1, and treated with 10 μ M MSA for 24 hr. The luciferase reading was normalized by protein concentration. The

experiment was done three times and the results were expressed as mean percent inhibition $\pm$ SEM.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript