Published in final edited form as:

Angew Chem Int Ed Engl. 2011 August 22; 50(35): 8172–8175. doi:10.1002/anie.201103271.

Catalytic Diastereoselective Petasis Reactions**

Dr. Giovanni Muncipinto,

Broad Institute of Harvard and MIT, Howard Hughes Medical Institute, Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge, Massachusetts 02138 (USA)

Dr. Philip N. Moquist,

Department of Chemistry, Center for Chemical Methodology and Library Development at Boston University (CMLD-BU), Life Science and Engineering Building, Boston University, 24 Cummings Street, Boston, Massachusetts 02215 (USA)

Dr. Stuart L. Schreiber, and

Broad Institute of Harvard and MIT, Howard Hughes Medical Institute, Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge, Massachusetts 02138 (USA)

Dr. Scott E. Schaus* [Prof.]

Department of Chemistry, Center for Chemical Methodology and Library Development at Boston University (CMLD-BU), Life Science and Engineering Building, Boston University, 24 Cummings Street, Boston, Massachusetts 02215 (USA)

The Petasis boronic acid Mannich reaction is a versatile multicomponent reaction of boronic acids, amines and aldehydes that generates highly functionalized α -amino acids and β -amino alcohols.^[1] When enantiopure α -hydroxy aldehyde derivatives are used as the carbonyl component in the reaction, enantiopure β -amino alcohols are produced with exclusively *anti* diastereoselectivity.^[2] This motif has proven useful in the synthesis of stereodefined, biologically active molecules including sialic acids,^[3] iminocyclitols,^[4] and pyrrolizidine alkaloids.^[5] The characteristic of the diastereoselective boronic acid Mannich reaction that makes it valuable, namely its predictable sense of *anti* diastereoselectivity, is also its limitation as *syn* β -amino alcohols are unattainable under these conditions [Eq. 1].^[6] Previous attempts to obtain *syn* β -amino alcohols via the Petasis reaction have been unsuccessful and underscore the difficulty in overriding the intrinsic selectivity of the reaction.^[7] Herein, we report the first diastereoselective Petasis reaction catalyzed by chiral biphenols that enables the synthesis of *anti* and *syn* β -amino alcohols in pure form.

**This research was supported by the NIH (R01 GM078240, S.E.S. and P.N.M.). The NIGMS-sponsored Center of Excellence in Chemical Methodology and Library Development (P50-GM069721, S.L.S and G.M.) enabled this research.

*Fax: (+1) 617-353-6466, seschaus@bu.edu.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

(1)

This collaborative project was undertaken with the goal of developing a catalyst-controlled diastereoselective Petasis reaction. We recently developed the first enantioselective Petasis reaction between alkenyl boronates, secondary amines and ethyl glyoxylates catalyzed by chiral biphenols (Scheme 1a) and anticipated this type of ligand-exchange catalysis would be applicable to the diastereoselective variant.^[8,9] An immediate application of this methodology is a synthetic route to the full matrix of stereoisomeric products of a pathway conceived for use in small-molecule screening (Scheme 1b).^[10] This type of library development continues to represent a substantial challenge given current limitations in synthetic methodology. The synthesis of compounds having stereogenic carbons in diversity-oriented synthesis (DOS) appears to be useful based on one study showing a correlation between compounds with intermediate stereochemical complexity and improved binding selectivity.^[11] In addition, stereochemistry-based structure activity relationship (SSAR) can provide important clues that facilitate optimization and modification studies following the discovery of a small-molecule lead.^[12]

Initial development of the *syn* selective Petasis reaction focused on (*S*)-5-benzyl-2,2-dimethyl-1,3-dioxolan-4-ol **5aL**-phenylalanine methyl ester **6a** and (*E*)-diethyl styrylboronate **7a** modified reaction from our previous library synthesis.^[10] The uncatalyzed reaction of these components produced exclusively the *anti* β -amino alcohol **8** (Table 1, entry 1). Catalysts (*S*)-VAPOL **1**(*S*)-H₈-BINOLs **2a** and **2b**, (*S*)-BINOLs **3a** and **3b** were tested in the reaction, and although *syn* β -amino alcohol **8** was observed in the product mixture, these catalysts primarily gave the *anti* diastereomer (Table 1, entries 2–6). A breakthrough occurred with catalyst (*S*)-3,3'-Br₂-BINOL **4** which produced the *syn* diastereomer **8** as the major product in 4:1 d.r. (Table 1, entry 7). Attempts to optimize the diastereoselectivity via solvent effects (Table 1, entries 9–11) and boronate ligation were unsuccessful (Table 1, entries 12 & 13); however, an increase in *syn* selectivity to 5.5:1 d.r. was found with the addition 4 Å molecular sieves (Table 1, entry 8). In addition, the two diastereomers are separable on normal-phase chromatography allowing for isolation of the *syn* product in 54% yield.

This result shows for the first time it is possible to overcome the inherent selectivity of the diastereomeric Petasis reaction and obtain *syn* β -amino alcohols in their pure form. The ability of the catalyst to control the diastereoselectivity from >99:1 d.r. *anti* to 5.5:1 d.r. *syn* represents remarkable kinetic control of the reaction. Interestingly, the enantiomers of the

catalysts form a matched and a mismatched relationship with other components of the reaction. When enantiomeric catalyst (**R**)-**4** is used in the reaction the *anti* product is formed exclusively in 86% yield (Table 1, entry 14), while the racemic catalyst ($\pm$)-**4** gave the *anti* product in 9:1 d.r. (Table 1, entry 15). Therefore, the (*S*)-catalysts give mismatched selectivity and form the *syn* product, while the (*R*)-catalysts are matched and reinforce the *anti* pathway.

With optimized conditions in hand, we explored the scope of the *syn*-selective reaction. In all cases, the uncatalyzed reaction gave only the *anti* β -amino alcohol. Alkene addition with boronate **7a** formed primarily the *syn* diastereomer with lactol **5b** and L-phenylalanine-derived amines **6a** and **6b** (Table 2, entries 1 & 2). The Cbz-protected amino lactol **5c** worked as well indicating high functional group tolerance in the reaction (Table 2, entries 3 & 4). Isolation of the pure *syn* β -amino alcohol **9–12** was possible in these reactions in yields up to 80%. Aryl addition is also possible in the reaction using 4-methoxyphenylboronate **7d** which afforded the *syn* product with amino ester **6a** (Table 2, entries 5–7). However, the uses of amino acetal **6b** in the aryl addition reaction led to poor diastereoselectivity and the products were inseparable by silica gel chromatography (Table 2, entries 8–10). Alkynylboronate **7e** in combination with lactol **5a** and amine **6a** afforded the *syn* product in 5:1 d.r. with 62% yield of the pure *syn* diastereomer (Table 2, entry 11).

Next, the achiral amino ester **6c** and D-amino ester **6d** were tested under the catalyzed conditions. The reaction of amine **6c**alkenylboronate **7a** and lactol **5a** gave the *syn* diastereomer in 2:1 d.r. in 70% yield (Table 2, entry 12). Changing the aldehyde component afforded the *syn* product in 1.5:1 with lactol **6b** and 1:1 d.r. with lactol **6c** (Table 2, entries 13 and 14). The reaction with D-phenylalanine methyl ester **6d** with boronate **7a** and lactol **5a** with (**S**)-**4** forms primarily the *anti* product **27** in 3:1 d.r. *anti:syn* (Table 2, entry 15). Attempts to optimize the reaction through the use of other biphenol catalysts failed to generate better diastereoselectivity.^[13] These results demonstrate the influence of the amine stereocenter as a third element of diastereocontrol is dramatic. It becomes apparent that the D-amine is a matched pair with (*S*)-lactol and reinforces the *anti* selectivity of the reaction, while the L-amine has matched selectivity with (*S*)-catalyst to form the *syn* product.

Due to the significance of the amino acid stereochemistry on product diastereoselectivity, we compared the catalyzed reaction of boronate **7a** and lactol **5a** with the various D- and L-amino acid derivatives. Whereas the L-phenylglycine amino ester **6e** gave the *syn* product in 2:1 d.r., the D-enantiomer **6j** produced the *anti* product in 20:1 d.r. (Table 3, entries 1 & 5). Similar results appeared in the reaction with enantiomers of leucine-**6f** and **6k** and valine-derived amino esters **6g** and **6l** (Table 3, entries 2, 3, 5 & 6), in which the L-amine provides the *syn* and the D-amine provides *anti* product. The size of the substituents also affects the selectivity of the reaction, with smaller groups affording higher levels of *syn* product. Interestingly, the D-phenylalanine dimethyl acetal **6m** gave the *syn* product in 4:1 d.r. under the catalyzed conditions and was isolated as the pure *syn* diastereomer in 38% yield (Table 3, entry 10). Therefore, phenylalanine dimethyl acetals **6b** and **6m** are able to form the full matrix of stereoisomeric products in the library.

As an extension of this methodology, glycolaldehyde dimer **34** was used in the reaction to synthesize primary β -amino alcohols. The uncatalyzed reaction of amine **6**boronate **7a** and glycolaldehyde **34** produced a 4:1 mixture of the (*S,S*)-amino alcohol **35** and (*S,R*)-amino alcohol **36** (Scheme 3). Use of 20 mol% (*S*)-**4** gave >20:1 d.r. of (*S,S*)-amino alcohol **35**. This indicates both the catalyst and amine direct the boronate addition to the form of the (*S*)-stereogenic center of the amino alcohol. The (*R*)-catalyst (*R*)-**4** produced the opposite diastereomers **36** in 10:1 d.r. and easily overcomes the inherent selectivity of the amine component. These results are further evidence of the matched selectivity of catalyst (*S*)-**4** with L-amino acids and catalyst (*R*)-**4** with D-amino acids.

Next, we undertook a preliminary mechanistic investigation of the catalyzed reaction. The stereochemical model for the *anti* diastereoselective Petasis reaction involves an α -hydroxy-directed boronate addition to the imine, which proceeds through a Felkin-Anh-type transition state.^[14] Unsurprisingly, the use of 3-phenylpropanal and (*S*)-2-methoxy-3-phenylpropanal in the catalyzed reaction gave only trace amounts of product (<1% yield). This indicates the boronate coordination to the α -hydroxy group remains critical for reactivity and explains the immense stereochemical influence of the lactol. The boronate also appears to undergo a single-ligand exchange with the catalyst as evidenced by ¹H NMR and ESI-MS analysis.^[13] Therefore, it appears the boronate undergoes both ligand exchange with the catalyst and coordination with the α -hydroxy aldehyde during the course of the reaction. This type of activation is in line with the current mechanistic model of diastereoselective Petasis reaction, as well as previous models for catalytic activation of boronates with chiral diols.^[8,9]

Because this reaction relies on three stereocontrolling elements, it can be classified as a catalytic triple diastereoselective reaction. Although catalytic reactions involving double diastereoselectivity are well studied, catalyzed reactions involving three elements of diastereocontrol are rare. To this end, Masamune and Kishi have pioneered and produced elegant work in this area.^[15] However, since these initial reports have not investigated all enantiomers of the three stereocontrolling elements, this is the first report for which all enantiomers of the components have been examined. What we learned from this reaction can be simplified into a few generalized rules. (1) The uncatalyzed reaction maintains exclusively *anti* diastereoselectivity regardless of the amine, lactol, or boronate components. (2) While the amine is unable to overcome the diastereocontrol of the lactol, the structure and stereochemistry of the amine play a large role in the diastereoselectivity of the catalyzed reaction. (3) The matched combination of an L-amine and (*S*)-catalyst usually produces predominantly the *syn* diastereomers. (4) In general, the matched combination of D-amine and (*S*)-lactol using the catalyst-promoted conditions leads to the *anti* β -amino alcohol with the exception of amino acetal **6ma** characteristic of this particular reaction we are continuing to explore.

In conclusion, we have reported the first diastereoselective Petasis reaction of boronates, α -hydroxyl aldehydes, and amines to produce *syn* β -amino alcohols. In many cases the *syn* product can be obtained in isomerically pure form for further elaboration. Furthermore, the full matrix of stereoisomers for use in library synthesis was achieved using phenylalanine methyl acetal, and we believe other amines will be successful at this task. Although these

preliminary results indicate a number of challenges have yet to be solved, this work represents a substantial improvement in the utility and scope of the reaction. Our current efforts are focused on the improvement of this catalyst system and further investigations are ongoing to understand mechanism and activity of this reaction.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. a) Petasis NA, Akritopoulou I. *Tetrahedron Lett.* 1993; 34:583–586. b) Petasis, NA. *Multicomponent Reactions*. Zhu, J.; Bienayme, H., editors. Weinheim: Wiley-VCH; 2005. p. 199–222. c) Kaiser PF, Churches QI, Hutton CA. *Aust. J. Chem.* 2007; 60:799–810. d) Candeias NR, Montalbano F, Cal PMSD, Gois PMP. *Chem. Rev.* 2010; 110:6169–6193. [PubMed: 20677749]
2. a) Petasis NA, Zavialov IA. *J. Am. Chem. Soc.* 1997; 119:445–446. b) Petasis NA, Zavialov IA. *J. Am. Chem. Soc.* 1998; 120:11798–11799. c) Prakash GKS, Mandal M, Schweizer S, Petasis NA, Olah GA. *Org. Lett.* 2000; 2:3173–3176. [PubMed: 11009374] d) Prakash GKS, Mandal M, Schweizer S, Petasis NA, Olah GA. *J. Org. Chem.* 2002; 67:3718–3723. [PubMed: 12027685]
3. Hong Z, Liu L, Hsu C-C, Wong C-H. *Angew. Chem.* 2006; 118:7577–7581. *Angew. Chem. Int. Ed.* 2006; 45:7417–7421.
4. Hong Z, Liu L, Sugiyama M, Fu Y, Wong C-H. *J. Am. Chem. Soc.* 2009; 131:8352–8353. [PubMed: 19530721]
5. a) Davis AS, Pyne SG, Skelton BW, White AH. *J. Org. Chem.* 2004; 69:3139–3143. [PubMed: 15104453] b) Au CWG, Pyne SG. *J. Org. Chem.* 2006; 71:7097–7099. [PubMed: 16930074] c) Ritthiwigrom T, Pyne SG. *Org. Lett.* 2008; 10:2769–2771. [PubMed: 18517213] d) Ritthiwigrom T, Willis AC, Pyne SG. *J. Org. Chem.* 2010; 75:815–824. [PubMed: 20028000] e) Au CWG, Nash RJ, Pyne SG. *Chem. Commun.* 2010; 46:713–715.
6. For a review on the synthesis of 1,2-amino alcohols: Bergmeier SC. *Tetrahedron.* 2000; 56:2561–2576.
7. a) Koolmeister T, Sodergren M, Scobie M. *Tetrahedron Lett.* 2002; 43:5969–5970. b) Southwood TJ, Curry MC, Hutton CA. *Tetrahedron.* 2006; 62:236–242.
8. Lou S, Schaus SE. *J. Am. Chem. Soc.* 2008; 130:6922–6923. [PubMed: 18459782]
9. a) Wu TR, Chong JM. *J. Am. Chem. Soc.* 2005; 127:3244–3245. [PubMed: 15755118] b) Wu TR, Chong JM. *J. Am. Chem. Soc.* 2007; 129:4908–4909. [PubMed: 17402741] c) Barnett DB, Moquist PN, Schaus SE. *Angew. Chem.* 2009; 128:8835–8838. Barnett DB, Moquist PN, Schaus SE. *Angew. Chem. Int. Ed.* 2009; 48:8679–8682.
10. a) Kumagai N, Municipinto G, Schreiber SL. *Angew. Chem.* 2006; 118, 3717–3720. *Angew. Chem. Int. Ed.* 2006; 45:3635–3638. b) Municipinto G, Kaya T, Wilson JA, Kumagai N, Clemons PA, Schreiber SL. *Org. Lett.* 2010; 12:5230–5233. [PubMed: 20977261]
11. a) Nielsen TE, Schreiber SL. *Angew. Chem.* 2008; 120:52–61. *Angew. Chem. Int. Ed.* 2008; 47:48–56. b) Schreiber SL. *Nature.* 2009; 457:153–154. [PubMed: 19129834] c) Clemons PA, Bodycombe NE, Carrinski HA, Wilson JA, Wagner BK, Koehler AN, Schreiber SL. *Proc. Natl. Acad. Sci. U.S.A.* 2010; 107:18787–18792. [PubMed: 20956335]
12. For examples of SSAR studies: Zhang QS, Lu HJ, Curran DP. *J. Am. Chem. Soc.* 2004; 126:36–37. [PubMed: 14709047] Curran DP, Zhang Q, Richard C, Lu H, Gudipathi V, Wilcox CS. *J. Am. Chem. Soc.* 2006; 128:9561–9573. [PubMed: 16848495] Wrona IE, Lowe JT, Turbyville TJ, Johnson TR, Beignet J, Beutler JA, Panek JS. *J. Org. Chem.* 2008; 74:1897–1916. [PubMed: 19191575] Macaurelle LA, et al. *J. Am. Chem. Soc.* 2010; 132:16962–16976. [PubMed: 21067169]
13. See supporting information for details.
14. a) Schlienger N, Bryce MR, Hansen TK. *Tetrahedron.* 2000; 56:10023–10030. b) Petasis NA, Boral S. *Tetrahedron Lett.* 2001; 42:539–542. c) Voisin AS, Bouillon A, Lancelot J-C, Lesnard A,

- Oulyadi H, Rault S. *Tetrahedron Lett.* 2006; 47:2165–2169.d) Candeias NR, Veiros LF, Afonso CAM, Gois PMP. *Eur. J. Org. Chem.* 2009:1859–1867.e) Candeias NR, Cal PMSD, Andre V, Duarte MT, Veiros LF, Gois PMP. *Tetrahedron.* 2010; 66:2736–2745.f) Tao JC, Li SH. *Chin. J. Chem.* 2010; 28:41–49.
15. a) Duplantier AJ, Nantz MH, Roberts JC, Short RP, Somfai P, Masamune S. *Tetrahedron Lett.* 1989; 30:7357–7360.b) Namba K, Kishi Y. *Org. Lett.* 2004; 6:5031–5033. [PubMed: 15606128] c) Choi H-W, Nakajima K, Demeke D, Kang F-A, Jun H-S, Wan Z-K, Kishi Y. *Org. Lett.* 2002; 4:4435–4438. [PubMed: 12465906] d) Wan Z-K, Choi H-W, Kang F-A, Nakajima K, Demeke D, Kishi Y. *Org. Lett.* 2002; 4:4431–4434. [PubMed: 12465905]

Figure 1.
Chiral Diol Catalysts.

Table 1

Diastereoselective Petasis Reaction.^[a]

Entry	Catalyst	Boronate	Solvent	Yield ^[b] [%]	d.r. <i>anti:syn</i> ^[c]
1	–	7a	PhCF_3	24	<i>anti</i> only
2	1	7a	PhCF_3	69	10:1
3	2a	7a	PhCF_3	73	6:1
4	2b	7a	PhCF_3	66	8:1
5	3a	7a	PhCF_3	41	4:1
6	3b	7a	PhCF_3	81	3:2
7	(<i>S</i>)- 4	7a	PhCF_3	70	1:4
8	(<i>S</i>)- 4	7a	PhCF_3	68 (54)	1:5.5 (<i>dl</i>)
9	(<i>S</i>)- 4	7a	PhCH_3	67	1:1
10	(<i>S</i>)- 4	7a	CH_2Cl_2	57	3:1
11	(<i>S</i>)- 4	7a	THF	<5	n.d.
12	(<i>S</i>)- 4	7b	PhCF_3	48	1.5:1
13	(<i>S</i>)- 4	7c	PhCF_3	<5	n.d.
14	(<i>R</i>)- 4	7a	PhCF_3	86	<i>anti</i> only
15	<i>rac</i> - 4	7a	PhCF_3	84	9:1

^[a] Reactions were run with 0.4 mmol boronate, 0.2 mmol lactol, 0.2 mmol amine, and 20 mol% catalyst in organic solvent (0.2 M) for 24 h under Ar, followed by flash chromatography on silica gel.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

[b] Isolated yield of diastereomeric mixture. Yield in parenthesis is isolated yield of the syn diastereomer (>20:1 d.r.).

[c] Determined by ^1H NMR.

[d] Run with 4Å molecular sieves.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Diastereoselective Petasis Reaction^[a]

Entry	Lactol	Amine	Boronate	Product	Yield [%] ^[b]	d.r. <i>antisyn</i> ^[c]
1	5b	6a	7a	9	95 (80)	1:5.5
2	5b	6b	7a	10	96 (84)	1:7

Entry	Lactol	Amine	Boronate	Product	Yield [%] ^[b]	d.r. <i>antisyn</i> ^[c]
9	5b	6b	7d	17	70 (n.d.)	10:1
10	5c	6b	7d	18	71 (n.d.)	4:1

Entry	Lactol	Amine	Boronate	Product	Yield [%] ^[b]	d.r. <i>antisyn</i> ^[c]
15	5a	6d	7a	23	61 (n.d.)	3:1

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

^[a] Reactions were run with 0.2 mmol boronate, 0.1 mmol lactol, 0.1 mmol amine, 20 mol% catalyst, and 4Å molecular sieves in PhCF₃ (0.2M) for 16-60 h under Ar, followed by flash chromatography on silica gel.

^[b] Isolated yield of diastereomeric mixture. Yield in parenthesis is isolated yield of the *syn* diastereomer (>20:1 d.r.).

^[c] Determined by ¹H NMR. The *anti* products were synthesized using the matched catalyst (*R*)-**4** and same reaction conditions.

Table 3

Diastereoselective Petasis Reaction with L- and D-amines.^[a]

Entry	L-Amine	Yield [%] ^[b]	d.r. <i>anti:syn</i>	Entry	D-Amine	Yield [%] ^[b]	d.r. <i>anti:syn</i> ^[c]
1		55 (n.d.)	1:2	6		57 (n.d.)	20:1
2		80 (56)	1:7.5	7		71 (n.d.)	2:1
3		65 (n.d.)	1:2.5	8		68 (n.d.)	6:1
4		72 (48)	1:2	9		74 (n.d.)	10:1
5		94 (80)	1:6	10		83 (38)	1:4

^[a] Reactions were run with 0.2 mmol boronate, 0.1 mmol lactol, 0.1 mmol amine, 20 mol% catalyst, and 4 Å molecular sieves in PhCF₃ (0.2M) for 16-60 h under Ar, followed by flash chromatography on silica gel.

^[b] Isolated yield of diastereomeric mixture. Yield in parenthesis is isolated yield of only the syn diastereomer.

^[c] Determined by ¹H NMR. The *anti* products were synthesized using the matched catalyst (*R*)-**4** and same reaction conditions.