Published in final edited form as:

J Psychopathol Behav Assess. 2015 December 1; 37(4): 634–644. doi:10.1007/s10862-015-9492-z.

An fMRI Pilot Study of Cognitive Reappraisal in Children: Divergent Effects on Brain and Behavior

Lea R. Dougherty, Ph.D., Sarah L. Blankenship, M.S., Philip A. Spechler, B.A., Srikanth Padmala, M.S., and Luiz Pessoa, Ph.D.

University of Maryland College Park

Abstract

Although neuroimaging studies in adults demonstrate that cognitive reappraisal effectively down-regulates negative affect and results in increased prefrontal and decreased amygdala activity, very limited empirical data exist on the neural basis of cognitive reappraisal in children. This study aimed to pilot test a developmentally-appropriate guided cognitive reappraisal task in order to examine the effects of cognitive reappraisal on children's self-reports of affect and brain responses. Study 1 ($N=19$, 4–10 years-old) found that children successfully employed guided cognitive reappraisal to decrease subjective ratings of negative affect, supporting the effectiveness of the guided cognitive reappraisal task. Study 2 ($N=15$, ages 6–10 years-old) investigated the neural responses to guided cognitive reappraisal and found that the neural responses showed increased activation in the amygdala and ventromedial prefrontal cortex during the cognitive reappraisal condition compared to the no regulation condition. In addition, amygdala activity was positively correlated with ventromedial prefrontal cortex activation during cognitive reappraisal. Findings suggest that the neural networks supporting cognitive reappraisal in children involve similar brain regions but brain responses deviate from findings in adults. Our findings suggest that the neural networks supporting emotion regulation are still developing during middle childhood, and future research is necessary to delineate age-related development of the neural network involved in cognitive reappraisal.

Keywords

emotion regulation; cognitive reappraisal; children; fMRI; amygdala; prefrontal cortex

Emotion regulation is broadly defined as the ability to modify one's emotional experiences and expressions (Gross, 1998). Emotion regulation strategies are adaptive in that they enable individuals to respond appropriately to emotional content in their environments and social relationships. Deficits in the ability to regulate one's emotions are linked to a wide range of cognitive, academic, interpersonal, and physical health problems (Graziano et al., 2007; Gross, 1998; Gross & John, 2003). Furthermore, maladaptive emotion regulation is a core feature of numerous psychiatric disorders and has been implicated in the etiology and

instructed 7–17 year-olds to attend to, increase, or decrease feelings of disgust before viewing neutral and disgusting images selected from the International Affective Picture System (IAPS; Lang et al., 2008). Children were provided with specific strategies to increase (e.g., “pretend it’s in front of you”) and decrease (e.g., “pretend its fake”) their affective response. Similar to findings in adults (Urry et al., 2006), the authors reported that down-regulation of disgust resulted in an inverse relationship between left amygdala and medial PFC activity, reinforcing the idea of a down-regulating effect driven by prefrontal control regions. In addition, left amygdala activation decreased with age, suggesting that the network supporting reappraisal undergoes developmental change.

In a recent developmental study, McRae and colleagues (2012) investigated reappraisal networks in children (10–13 years), adolescents (14–17 years), and young adults (18–22 years) and found developmental changes in the neural mechanisms supporting cognitive reappraisal. Similar to the paradigm used by Pitskel and colleagues (2011), at the beginning of each trial, children were provided with a reappraisal instruction to attend to or decrease negative affect followed by the presentation of a negative or neutral IAPS image. During reappraisal, activation in the left inferior frontal gyrus and a left ventrolateral PFC region increased linearly with age while activation in the posterior cingulate, medial PFC, and anterior temporal cortex showed quadratic changes with age, with adolescents showing greater activation than adults or children during reappraisal. Interestingly, down-regulation of the amygdala was not observed in any age group during reappraisal; however, several prefrontal regions did show significant activation during reappraisal, but this effect was not observed in children ages 10–13 years. These findings suggest that cognitive reappraisal mechanisms may differ across development.

While there have been recent attempts to investigate the development of reappraisal networks in younger individuals, the existing literature is inconclusive. Some studies suggest children are capable of successfully down-regulating amygdala activity (Pitskel et al., 2011), while others indicate they are not (McRae et al., 2012). Even more, there is still a dearth of research investigating emotion regulation networks during early to middle childhood (6–10 years), a period of ongoing neurodevelopment and reorganization in regions implicated in emotion regulation. The prefrontal cortex undergoes developmental changes into adulthood (Gogtay et al., 2004), and the amygdala reaches its peak volume around age 10 (Uematsu, et al., 2010). Recent evidence also suggests that during fearful face processing, amygdala-prefrontal functional connectivity undergoes a developmental change around age 10 where a shift from positive to negative connectivity between these regions occurs (Gee et al., 2013). Taken together, evidence of the protracted development of regions implicated in emotion regulation and the connectivity between these regions suggests that early to middle childhood is a period of interest in understanding the development of emotion regulation networks.

The current study aimed to pilot a developmentally-appropriate guided cognitive reappraisal task (Dennis & Hajcak, 2009) modified for functional magnetic resonance imaging (fMRI) use in order to examine the neural basis of cognitive reappraisal during early to middle childhood (6–10 years). Given the current state of the literature, we tentatively hypothesized that (1) guided cognitive reappraisal would effectively down-regulate negative affect, as

assessed via children's self-reports; and (2) guided cognitive reappraisal would be associated with increased prefrontal and decreased amygdala activation. We also hypothesized that ventromedial PFC would show inverse coupling with amygdala activity during reappraisal. However, these hypotheses are tentative as recent findings by McRae and colleagues (2012) suggest that children at this age, in contrast to adults, may not have the control mechanisms necessary to down-regulate amygdala activation using reappraisal strategies.

Method

Overview

Study 1 examined the behavioral effects of cognitive reappraisal on regulating negative affect. Using a self-report measure of affect, we assessed the degree to which directed reappraisal stories up-regulated or down-regulated children's emotional responses to negatively valenced pictures. Study 2 examined changes in functional brain activity associated with the cognitive reappraisal paradigm used in study 1.

Study 1: Behavioral Study of Cognitive Reappraisal

Participants

Participants were recruited via a commercial mailing list and print advertisements distributed throughout local schools, community centers, daycare and healthcare providers in the greater Washington, DC area. Interested parents of child participants were interviewed on the telephone to determine study eligibility. Eligible participants were between the ages of 4 and 10 years old, native English speakers, and free from any cognitive or developmental disorders based on parent-report. A total of 19 children were recruited for the behavioral study (12 females; Age $M = 6.7$ years, $SD = 1.1$, range=4–10). The study was approved by the human subjects review committee, and informed consent was obtained from parents. Child assent was obtained from children 7 years and older.

Emotion regulation task

The emotion regulation task was based on a directed reappraisal paradigm developed for children ages 5–10 years-old that has been shown to modulate brain responses to emotional stimuli as measured by event related potentials (see Dennis & Hajcak, 2009). In the current study, participants were presented with 45 negative pictures (e.g. frightening animals, negative facial expressions) and 15 neutral pictures¹. Developmentally appropriate negative- and neutral-valenced images were selected from the International Affective Picture System (IAPS; Lang et al., 2008) and the Emotional Picture Set (EmoPicS; Wessa et al., 2010). The negative pictures were divided into three conditions: negative-attend, negative-decrease, and negative-increase. All neutral pictures were presented in the neutral-attend condition. Each picture was preceded by a brief story containing an implicit instruction to attend, decrease, or increase their affective response to the upcoming picture. Participants were instructed to “match the story to the picture” and were trained in a practice session before the main

¹IAPS:

1050,1120,1201,1300,1302,1321,1525,1930,1932,2120,2130,2457,2458,2703,2780,2810,2900,3022,3230,3280,6300,6312,6370,7380, 9050,9250,9321,9421,9440,9480,9490,9561,9582,9594;EmoPicS:208,209,210,211,212, 216,225,235,246,251,329

experiment. The decrease stories provided a directed reappraisal of the negative-valenced picture to down-regulate negative affective responses. For example, prior to the presentation of a picture of a crying father and son, the participant heard the following directed reappraisal: “This family just won a big prize and is so happy they are crying”. Conversely, the increase stories provided a directed reappraisal to up-regulate the negative affective response from the picture (e.g., “This family just lost their home in a fire and is sad and crying”). The attend conditions (negative-attend and neutral-attend) provided an objective description of the picture, absent of any regulatory cues or affective language (e.g., “This is a picture of a man holding a small boy in his arms”). See Online Resource 1 for a complete list of stimuli and associated stories.

The stories paired with negative pictures were counterbalanced across participants such that all negative pictures were balanced across each negative condition (i.e., decrease, increase, attend). Neutral pictures were always paired with an “attend” story. None of the images or stories was repeated during the session. The stories were recorded by a female staff member using an external microphone attached to a computer using the open-source program Audacity (<http://audacity.sourceforge.net/>). Each story had a spoken duration of approximately 5 seconds (s) ($M=4.52$, $SD=0.50$). Our final stimuli set consisted of 60 story-picture pairs (15 per condition). Both the behavioral and fMRI versions (described below) of the experiments were programmed using Presentation software (Neurobehavioral Systems, Albany, CA, USA).

We assessed children’s self-reported affect following the presentation of each picture using an abbreviated version of the self-assessment manikin (SAM) valence rating system (Lang et al., 2008). The 5-point valence scale depicts five characters who range from unhappy (1) to happy (5) with 3 being neutral. Participants were told to “point to the picture of SAM that best shows how you feel”. Children’s responses were hand recorded by the experimenter. The abbreviated SAM has been shown to assess self-reported affective responses in children as young as 5 years (Hajcak & Dennis, 2009).

The experiment was divided into three “runs” of approximately 5–6 minutes each in duration. Each run consisted of 20 trials (5 per trial type). All four experimental conditions were intermixed in a pseudorandom fashion. Each picture was presented for 2.5 s (visual angle = $9.5^\circ \times 9.5^\circ$), preceded by the story lasting approximately 5 s during which a fixation mark was presented on a blank screen. A 2 s delay period was included between each story and picture (see Figure 1). The picture disappeared to a blank screen for 500 ms followed by a SAM rating prompt. After recording the participant’s response, the experimenter initiated the next trial.

Before the start of the experimental runs, participants were introduced to the task in a practice run. The practice run consisted of 8 trials (2 per trial type) using a separate set of stimuli and stories. As in the main experimental runs, all four experimental conditions were intermixed in a pseudorandom fashion. None of the stories or pictures from the practice run was included in the main runs. The participants were instructed to match the story to the picture, and the experimenter probed the child to confirm that s/he understood the task and the rating scale before proceeding to the experimental task.

Data Analysis Plan—First, to assess self-reported emotional reactivity to negative stimuli, self-reported SAM ratings of negative-attend vs. neutral-attend conditions were compared using a paired *t*-test. Second, we investigated the effects of directed reappraisal to up- and down-regulate negative-valenced pictures by comparing SAM ratings of negative-increase vs. negative-attend and negative-decrease vs. negative-attend conditions, respectively, using separate paired *t*-tests.

Results

Self-Report Ratings—As seen in Figure 2, self-reported affect was sensitive to condition. Lower SAM ratings are indicative of stronger negative affect. Self-reported affect was rated higher in negative affect in the negative-attend condition ($M=2.53$, $SD=0.80$, range: 3.87–1.20) than the neutral-attend condition ($M=3.74$, $SD=0.73$, range: 5.00–2.13), $t_{18} = -4.60$, $p < .001$, indicating an increase in negative affect in response to negative vs. neutral stimuli. As evidence for the down-regulation of negative affect as a function of cognitive reappraisal, self-reported affect was rated lower in negative affect in the negative-decrease condition ($M=3.50$, $SD=0.74$, range: 4.67–1.47) than the negative-attend condition, $t_{18} = 4.76$, $p < .001$. In contrast, self-reported affect was rated higher in negative affect in the negative-increase condition ($M=1.55$, $SD=0.50$, range: 2.93–1.00) than the negative-attend condition, $t_{18} = -7.30$, $p < .001$.

Study 2: fMRI Study of Cognitive Reappraisal

Methods

Participants—Novel participants were recruited in the same manner as the behavioral study. Potential participants were excluded if they were left handed; below 6 or above 10 years of age; non-native English speakers; had a current or past diagnosis of neurological or psychiatric disorder based on parent-report; had a history of head trauma; currently used psychotropic medication; or had any MRI contraindication (e.g., metal in the body, orthodontic braces). From a total of 21 participants, 15 children were included in the final analysis. Two children were excluded due to excessive motion inside the scanner (cumulative motion relative to reference volume exceeding 6 mm in any direction); two children were excluded due to an inability to tolerate the scanning environment; one child was excluded due to technical difficulties; and one child's scan was terminated prematurely due to too much motion. Of the 15 children, children's mean age was 7.94 years ($SD=1.21$, range = 6–9 years), and 8 children were female (53%). Lastly, two participants included in the analysis only provided usable data for 2 of the 3 runs due to excessive head motion in one of the runs. The study was approved by the human subjects review committee, and informed consent was obtained from parents. Child assent was obtained from children 7 years and older.

Emotion regulation task—The fMRI paradigm used the same stories and pictures from the behavioral paradigm described in Study 1 with the addition of 6 new neutral stimuli² to balance the number of trials per condition. Three conditions were included in the fMRI task:

²IAPS: 2250,2342,2411;EmoPicS:268,302,339

neutral-attend, negative-attend, and negative-decrease. The negative-increase condition was not included to minimize the time children spent in the scanner and increase the number of trials per condition. In addition, self-report SAM ratings were not obtained in the fMRI version given concerns about excessive head movement while providing behavioral responses.

Prior to scanning, eligible participants prepared for the MRI and practiced the behavioral task outside and inside a mock scanner 1 hour before the MRI scan. The practice sessions consisted of 9 trials (3 per trial type presented in pseudorandom fashion). None of the stories or pictures from the practice run was included in the fMRI task. After successful completion in the mock scanner session, participants participated in the MRI session. The fMRI task consisted of 3 runs, and each run included 21 trials (7 per trial type presented in pseudorandom fashion). Each trial was nearly identical to the version used in the behavioral experiment with the exception of no rating period after picture presentation and a longer inter trial interval of 4 – 8 s used to jitter the stimulus onset times to aid in the estimation of fMRI responses.

MRI data acquisition—MRI data were collected using a 3 Tesla Siemens TRIO scanner (Siemens Medical Systems, Erlangen, Germany) with a 12-channel head coil (without parallel imaging). Each scanning session began with a high-resolution MPRAGE anatomical scan (TR = 1900 ms, TE = 2.52 ms, TI = 900 ms, 1 mm cubic voxels, 250 mm field of view). Subsequently, during each functional run, 170 EPI volumes were acquired with a TR of 2000 ms and TE of 24 ms. Each volume consisted of 36 oblique slices with a thickness of 3 mm and an in-plane resolution of 3 mm x 3 mm (FOV = 192 mm). Slices were positioned approximately 30 degrees relative to the plane defined by the line connecting the anterior and posterior commissures, helping to decrease susceptibility artifacts in regions such as the ventromedial prefrontal cortex, temporal cortex, and amygdala.

Imaging analysis—Preprocessing of the data was conducted using the AFNI software package (Cox, 1996), with the exception of the skull-stripping of the anatomical data, which was processed using statistical parametric mapping (SPM8; Wellcome Trust Centre for Neuroimaging, London, UK). The first 3 volumes of each functional run were discarded to account for equilibration effects. The remaining volumes were slice-time corrected using a Fourier interpolation, such that all slices were realigned to the first acquisition slice to account for timing differences. Six-parameter rigid-body motion correction within and across runs was performed using Fourier interpolation (Cox & Jesmanowicz, 1999) so that all volumes were spatially registered to the first volume. Prior to spatial normalization to standard space, the anatomical and functional data were coregistered. We spatially normalized the anatomical data to the MNI anatomical template for children ages 4.5–8.5 years using a 12-parameter affine transformation. The rationale and production of this atlas can be found in Fonov et al. (2011). The same transformation was then applied to the functional data. All volumes were spatially smoothed using a Gaussian filter with a full-width at half maximum of 6 mm (i.e., two times the voxel dimension). Finally, the signal intensity of each voxel was scaled to a mean of 100 (on a per run basis), which allowed the interpretation of the estimated regression coefficients in terms of percent signal change.

Voxelwise analysis—The central goal of the voxelwise analysis was to define regions of interest (ROIs). Each participant's fMRI data were analyzed using a linear model framework with AFNI. Constant, linear, and quadratic terms were included for each run separately (as covariates of no interest) to model baseline and drifts of the MRI signal. To account for the signal variance related to head motion, six estimated motion parameters were included as nuisance regressors in the model. Further, we excluded the current and previous TR data whenever frame-to-frame movement within a run was more than half a voxel size (i.e., 1.5 mm) in any direction. Specifically, collapsing across conditions, there was an average of .99%/4.87 TRs (range: 0–5%; 0–24 TRs) of data lost per participant due to motion, and the number of excluded volumes (i.e., TR) did not differ across condition, $F(2,28) = .08$, $p = .92$; hence, minimal data was lost due to motion. No assumptions were made about the shape of the hemodynamic response function. All trials from three conditions were pooled together resulting in a total of one main event type in the design matrix. Trials were pooled together in order to define ROIs in an unbiased manner (see below). Responses were estimated on each trial starting from trial onset to 24 s post-onset using cubic spline basis functions. This method is closely related to the use of finite impulses (“stick functions”), the commonly employed technique that can be considered the simplest form of basis expansion. Cubic splines allow a smoother approximation of the underlying responses, instead of the discrete approximation obtained by finite impulses. As an index of picture related activation, we averaged the estimated responses at time points 12, 14, and 16 s post-trial onset (i.e., approximately 4, 6, 8 s post-picture onset). We used the average of these time points, as picture-related response would be maximal at these time points, while the effect of story-related response would be minimal. We further confirmed that these three time points had maximal picture-related responses in early visual cortex and minimal story-related responses in early auditory cortex. Nevertheless, given our design, we cannot rule out some contribution of story-related response on subsequent picture related activation.

Group analysis—Whole-brain voxelwise random-effects analyses were restricted to graymatter voxels based on the MNI pediatric gray matter template (Fonov et al., 2011). The template was resampled to match the resolution of the functional data. To assess picture related processing (vs. baseline), a one sample *t*-test (vs. zero) was run on the estimated response strength index of picture related activation. We did not analyze auditory story related responses because our primary goal was to investigate how the responses to the pictures were modulated by the preceding auditory stories.

ROI analysis—To enhance statistical power, we focused our analysis on *a priori* ROIs commonly implicated in the context of emotion regulation: the ventromedial PFC, dorsolateral PFC, and amygdala (Table 1). The left and right amygdala ROI (Figure 3A–3B) were anatomically defined initially in the talairach space using the maximum-probabilistic atlas containing subcortical regions (“DD_Desai_MPM”) provided in the AFNI package (Destrieux et al., 2010). The ROIs were transformed into the MNI space using the 3dWarp command and then aligned to the MNI template for children using the Nudge tool in the AFNI package. The ROIs were then resampled to match the resolution of the functional data. We then selected those voxels activated during the voxelwise analysis with a threshold set at $p = .05$ (uncorrected). The other ROIs were functionally defined based on the

voxelwise analysis with a threshold set at $p = .05$ (uncorrected) and a minimum cluster extent of 10 voxels (Figure 3C–3E). We adopted the selection criterion of picture-related processing (pooled over all three conditions) to determine ROIs, because it was statistically independent of the central goal of our analysis: to examine the emotion regulation (i.e., negative-decrease vs. negative-attend) and the emotional reactivity (i.e., negative-attend vs. neutral-attend) contrasts (Kriegeskorte et al., 2009; Vul et al., 2009). A representative time series for each ROI was then created by averaging the time series of all gray matter voxels inside the ROI. Then, for each ROI, we ran the analysis on the representative time series data to estimate the hemodynamic response function of three main event types: neutral-attend, negative-attend, and negative-decrease. For each condition, as an index of picture related activation, we averaged the estimated responses at time points 12, 14, and 16 s post-trial onset (i.e., approximately 4, 6, 8 s post-picture onset).

First, to assess emotional reactivity, in each ROI we contrasted the response indices of negative-attend and neutral-attend conditions using a paired t -test. Second, to assess emotional down-regulation, in each ROI we contrasted the response indices of negative-decrease and negative-attend conditions using a paired t -test. For the sake of completeness, additional voxelwise analyses were also performed and followed the same emotional reactivity and emotional regulation contrasts as in the ROI analysis. Results from these supplementary voxelwise analyses are reported in Online Resource 2.

Association between brain responses in amygdala and ventromedial PFC—

Because previous studies have highlighted the importance of the functional relationship between the amygdala and ventromedial PFC during emotion regulation (e.g., Banks et al., 2007; Urry et al., 2006), we investigated the association between brain responses in the amygdala ROIs and the ventromedial PFC ROI. For each participant, we created the emotion down-regulation index (negative-decrease minus negative-attend) for each ROI, separately. Then, across participants, we investigated the linear relation of the down-regulation index between each amygdala ROI and ventromedial PFC ROI using robust regression. We employed iterative reweighted least squares, the robust fit function from Matlab (Mathworks, Natick, MA, USA), given that standard Pearson correlation is sensitive to even a few influential data points (Wilcox, 2005).

Results

ROI results—We focused our analysis on *a priori* regions that were previously implicated in the cognitive reappraisal literature. First, we compared the negative-attend vs. neutral-attend pictures to investigate children's emotional reactivity to negative stimuli. We observed significant differential activation in the right amygdala and right dorsolateral PFC and marginally significant activation in the left dorsolateral PFC (Table 2). In these regions, as expected, activation during negative-attend pictures was greater than during the neutral-attend pictures (Figures 3A–3D). To investigate children's emotion regulation, we contrasted negative-decrease vs. negative-attend pictures. We observed significant differential activation in the right amygdala and ventromedial PFC (Table 2). In the right amygdala, responses during negative-decrease pictures were *higher* compared to negative-

attend pictures (Figure 3A). Negative-decrease condition showed stronger responses compared to negative-attend condition in ventromedial PFC (Figure 3E).

Associations between brain regions—Given the importance of the functional link between amygdala and ventromedial PFC during emotion regulation, particularly cognitive reappraisal, we evaluated the association between responses in these regions. A robust regression analysis using the emotion regulation index revealed a positive correlation between responses in the right amygdala and ventromedial PFC (robust $R^2 = 0.54$, $p = .036$; Figure 4A). The association between responses in the left amygdala and ventromedial PFC was not detected (robust $R^2 = 0.40$, $p = .144$; Figure 4B).

Discussion

The current pilot study examined the neural basis of cognitive reappraisal during middle childhood (6–10 years) using a developmentally-appropriate directed reappraisal task. We found that children can successfully use guided cognitive reappraisal to decrease subjective ratings of negative affect. In contrast to the behavioral data, children's brain responses demonstrated *increased* activation in the amygdala and ventromedial PFC during the reappraisal condition compared to the no regulation condition. These findings suggest that the neural networks supporting reappraisal at this age involve similar brain regions identified in adults but brain responses seem to deviate from findings in adults.

Our behavioral results indicate that children are capable of using guided reappraisal to decrease subjective measurements of negative affect, confirming our hypothesis that negative- decrease (reappraised) images would receive lower self-reported ratings of negative affect than negative-attend images. These results are consistent with previous research that indicates that reappraisal is effective at down-regulating negative affect in children, adolescents, and adults (Banks, et al., 2007; McRae et al., 2010; McRae et al., 2012; Ray et al., 2010).

Our neuroimaging results revealed that similar to adults, children's emotional reactivity (negative-attend – neutral-attend) engaged bilateral amygdala and dorsolateral PFC, supporting children's increased neural responses to negative stimuli. In addition, both the ROI and supplemental whole brain voxelwise analyses showed that the brain regions involved in cognitive reappraisal in adults (Buhle et al., in 2014; Kohn et al., 2014; Ochsner, Silvers, & Buhle, 2012), which include the amygdala and PFC, were also activated during cognitive reappraisal in children ages 6–10 years. However, contrary to the extant adult literature that suggests that prefrontal cortices down-regulate amygdala activity during reappraisal, children's brain responses did not show decreased activation in the amygdala. In the current study, reappraisal (negative-decrease – negative-attend) elicited a significant *increase* in activation in the right amygdala and ventromedial PFC. Interestingly, Levesque and colleagues (2003; 2004) reported that during reappraisal children displayed more distributed regions of prefrontal activation in comparison to adults. In addition, McRae et al. (2012) showed that while adolescents and adults displayed a reappraisal effect in regions of PFC, 10–13 year-old children did not show this effect. Similar to these neuroimaging findings, DeCicco, Solomon, and Dennis (2012) found that the amplitude of the late positive

potential (LPP), an event-related potential responsive to reappraisal in adults, was not reduced during reappraisal in 5–7 year-old children. Further, there is evidence that reappraisal is also ineffective at down-regulating physiological responses to negative stimuli at age 10 (de Veld, Riksen-Walraven & de Weerth, 2012), while in adults the opposite is found (Ray et al., 2010). Taken together, these findings suggest that the neural networks underlying cognitive reappraisal undergo significant developmental change across childhood.

We observed *greater* amygdala activity following reappraisal (negative-decrease condition) compared to the negative-attend condition. This effect was, of course, unexpected and further experiments are needed to clarify it. However, it is intriguing to note that, during reappraisal, amygdala activity was *positively* correlated with ventromedial PFC activation. This positive relationship may be related to the observation of increased amygdala responses following reappraisal. It is possible that the greater recruitment of prefrontal regions during cognitive reappraisal simultaneously recruits subcortical areas, such as the amygdala. Naturally, our finding is in contrast to a robust body of literature in adults that shows an inverse relationship between ventromedial PFC and amygdala during reappraisal (for review, see Kohn et al., 2014). But our findings may be related to evidence that amygdala-prefrontal functional connectivity during fearful face processing undergoes a developmental shift around age 10 from positive to negative connectivity (Gee et al., 2013).

Of particular interest is the apparent discrepancy between our behavioral and neuroimaging findings: while 6–10 year-old children were capable of using reappraisal to decrease subjective ratings of image valence, the observed decrease in subjective ratings of negative affect was not reflected in decreased amygdala activity. Other investigations have reported a discontinuity between subjective ratings of affect and neural responses in children. For example, McRae et al. (2012) reported that reappraisal elicited a reduction in negative affect based on behavioral ratings in children ages 10–13 years old, while there were no effects of reappraisal on amygdala and prefrontal cortex responses.

As the first fMRI investigation of reappraisal in this age range, our findings suggest that the neural circuitry supporting emotion regulation has not fully developed to a mature, adult-like network. By extending the current literature to a younger age group, we were able to begin identifying the developmental trajectory of emotion regulation networks and shed light on the reason for potential inconsistencies in the literature. Given methodological limitations of such a young sample, we employed an age-appropriate paradigm utilized in an ERP investigation of reappraisal in this age whereby reappraisals were orally presented to the children instead of requiring participants to self-generate their own reappraisals (Dennis & Hajcak, 2009). This paradigm limited the cognitive load for our young participants, reduced the risk for motion artifacts, and provides consistency between two modes of reappraisal investigations at this age.

The study also had limitations. First, given that this study was a pilot study, our sample size was small, which limited our statistical power and ability to use a whole-brain search space. As suggested by Levesque and colleagues (2004; 2003), children show more distributed loci of activation in PFC and whole-brain analyses may be more sensitive to these differences,

which may be overlooked in a ROI-based analysis. Second, given evidence of gender differences in emotion regulation (e.g., Dennis & Hajcak, 2009; McRae et al., 2008) and brain changes occurring during early to middle childhood, larger samples would allow for investigations of gender and age differences in reappraisal circuitry. Third, due to concerns about excessive head motion and task length for children in this age range, children did not complete a SAM rating following each trial while in the scanner. It will be important for future research to examine whether child affect ratings vary as a function of children's brain responses. Fourth, even though the guided reappraisal task allowed us to investigate cognitive reappraisal in young children, the paradigm deviates from the self-generated reappraisals adult studies employ. It is possible that self-generated reappraisals would engage different neural networks. Future research will need to compare the neural processes underlying guided reappraisals and self-generated reappraisals across development. Lastly, our small sample size and lack of correction for multiple comparisons in ROI analysis makes it difficult to draw firm conclusions from this study alone; nevertheless, future research can build upon the present study to investigate the neural basis of emotion regulation in children and its role in developmental psychopathology.

In sum, our findings suggest that while children are capable of using reappraisal to decrease subjective ratings of negative affect, neural responses in the amygdala revealed a divergent pattern with increased activation during the reappraisal condition. In light of the developmental literature, the discrepancy between behavioral and neuroimaging findings may be attributable to immature connectivity between PFC and amygdala which purportedly subserves top-down regulation of the amygdala during reappraisal. This literature, in accordance with our current findings, suggest that neurodevelopmental changes occur during middle childhood through adolescence, which equip children with the ability to employ reappraisal to down-regulate neural responses to negative content. Future research is necessary to delineate age-related development of the neural network involved in cognitive reappraisal. Our findings suggest that the neural networks supporting emotion regulation are still developing during middle childhood. Failure to employ top-down regulation of amygdala activity during this age may result in prolonged neurophysiological responses and could indicate a mechanism by which childhood is marked by a particular vulnerability to emotional events.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by the University of Maryland (UMD) Maryland Neuroimaging Center Seed Grant Program (LRD, LP), UMD College of Behavioral and Social Sciences Dean's Research Initiative Award (LRD), the UMD Research and Scholars Award (LRD) and in part by the National Institute of Mental Health R01 MH071589 (LP). The authors wish to thank the children and families for their participation in the study and the Maryland Neuroimaging Center.

References

- Banks SJ, Eddy KT, Angstadt M, Nathan PJ, Phan KL. Amygdala-frontal connectivity during emotion regulation. *Social Cognitive and Affective Neuroscience*. 2007; 2:303–312. [PubMed: 18985136]
- Beauregard M, Paquette V, Lévesque J. Dysfunction in the neural circuitry of emotional self-regulation in major depressive disorder. *Neuroreport*. 2006; 17(8):843–6. [PubMed: 16708026]
- Buhle JT, Silvers JA, Wager TD, Lopez R, Onyemekwu C, Kober H, Weber J, Ochsner KN. Cognitive reappraisal of emotion: A meta-analysis of human neuroimaging studies. *Cerebral Cortex*. 2014; 24:2981–2990. [PubMed: 23765157]
- Carthy T, Horesh N, Apter A, Gross JJ. Patterns of emotional reactivity and regulation in children with anxiety disorders. *Journal of Psychopathology and Behavioral Assessment*. 2009; 32:23–36.
- Cousineau D. Confidence intervals in within-subject designs: A simpler solution to Loftus and Masson's method. *Tutorial in Quantitative Methods for Psychology*. 2005; 1:4–45.
- Cox RW, Jesmanowicz A. Real-time 3D image registration for functional MRI. *Magnetic Resonance in Medicine*. 1999; 42:1014–1018. [PubMed: 10571921]
- DeCicco JM, Solomon B, Dennis TA. Neural correlates of cognitive reappraisal in children: An ERP study. *Developmental Cognitive Neuroscience*. 2012; 2:70–80. [PubMed: 22163262]
- Dennis TA, Cole PM, Wiggins CN, Cohen LH, Zalewski M. The functional organization of pre-school age children's emotion expressions and actions in challenging situations. *Emotion*. 2009; 9:520–530. [PubMed: 19653775]
- Dennis TA, Hajcak G. The late positive potential: a neurophysiological marker for emotion regulation in children. *Journal of Child Psychiatry*. 2009; 50:1373–1383.
- Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *Neuroimage*. 2010; 53:1–15. [PubMed: 20547229]
- de Veld DMJ, Riksen-Walraven JM, de Weerth C. The relation between emotion regulation strategies and physiological stress responses in middle childhood. *Psychoneuroendocrinology*. 2012; 37:1309–1319. [PubMed: 22309825]
- Fonov V, Evans AC, Botteron K, Almli CR, McKinstry RC, Collins DL. Unbiased average age-appropriate atlases for pediatric studies. *Neuroimage*. 2011; 54:313–327. [PubMed: 20656036]
- Gee DG, Humphreys KL, Flannery J, Goff B, Telzer EH, Shapiro M, Hare TA, Bookheimer SY, Tottenham N. A developmental shift from positive to negative connectivity in human amygdala-prefrontal circuitry. *The Journal of Neuroscience*. 2013; 33:4584–4593. [PubMed: 23467374]
- Gogtay N, Giedd JN, Lusk L, Hayashi KM, Greenstein D, Vaituzis AC, Nugent TF, Herman DH, Clasen LS, Toga AW, Rapoport JL, Thompson PM. Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences*. 2004; 101:8174.8179.
- Graziano PA, Reavis RD, Keane SP, Calkins SD. The role of emotion regulation and children's early academic success. *Journal of School Psychology*. 2007; 45:3–19. [PubMed: 21179384]
- Gross JJ. Antecedent- and response-focused emotion regulation: Divergent consequences for experience, expression, and physiology. *Journal of Personality and Social Psychology*. 1998; 74:224–237. [PubMed: 9457784]
- Gross JJ, John OP. Individual differences in two emotion regulation processes: Implications for affect, relationships, and well-being. *Journal of Personality and Social Psychology*. 2003; 85:348–362. [PubMed: 12916575]
- Gullone E, Hughes EK, King NJ, Tong B. The normative development of emotion regulation strategy use in children and adolescents: a 2-year follow-up study. *Journal of Child Psychology and Psychiatry*. 2010; 51:567–574. [PubMed: 19845818]
- Kim P, Evans GW, Angstadt M, Ho SS, Sripada CS, Swain JE, Liberzon I, Phan KL. Effects of childhood poverty and chronic stress on emotion regulatory brain function in childhood. *Proceedings of the National Academy of the Sciences*. 2012; 110:18442–18447.
- Kriegeskorte N, Simmons WK, Bellgowan PS, Baker CI. Circular analysis in systems neuroscience: the dangers of double dipping. *Nature Neuroscience*. 2009; 12:535–540. [PubMed: 19396166]

- Kohn N, Eickhoff SB, Scheller M, Laird AR, Fox PT, Habel U. Neural network of cognitive emotion regulation - an ALE meta-analysis and MACM analysis. *Neuroimage*. 2014; 87:345–355. [PubMed: 24220041]
- Lang, PJ.; Bradley, MM.; Cuthbert, BN. Technical Report A-8. Gainesville, FL: University of Florida; 2008. International affective picture system (IAPS): Affective ratings of pictures and instruction manual.
- Levesque J, Joanette Y, Mensour B, Beaudoin G, Leroux JM, Bourgouin P, Beauregard M. Neural basis of emotional self-regulation in childhood. *Neuroscience*. 2004; 129:361–369. [PubMed: 15501593]
- Levesque J, Eugene F, Joanette Y, Paquette V, Mensour B, Beaudoin G, Leroux J-M, Bourgouin P, Beauregard M. Neural circuitry underlying voluntary self-regulation of sadness. *Biological Psychiatry*. 2003; 53:502–510. [PubMed: 12644355]
- Manian N, Bornstein MH. Dynamics of emotion regulation in infants of clinically depressed and nondepressed mothers. *Journal of Child Psychology and Psychiatry*. 2009; 50:1410–1418. [PubMed: 19788549]
- McRae K, Gross JJ, Weber J, Robertson ER, Sokol-Hessner P, Ray RD, Gabrieli JDE, Ochsner KN. The development of emotion regulation: an fMRI study of cognitive reappraisal in children, adolescent and young adults. *Social Cognitive and Affective Neuroscience*. 2012; 7:11–22. [PubMed: 22228751]
- McRae K, Hughes B, Chopra S, Gabrieli JDE, Gross JJ, Ochsner KN. The neural bases of distraction and reappraisal. *Journal of Cognitive Neuroscience*. 2010; 22:248–262. [PubMed: 19400679]
- McRae K, Ochsner KN, Mauss IB, Gabrieli JD, Gross JJ. Gender differences in emotion regulation: an fMRI study of cognitive reappraisal. *Group Processes and Intergroup Relations*. 2008; 11:143–162.
- Monk CS. The development of emotion-related neural circuitry in health and psychopathology. *Development and Psychopathology*. 2008; 20:1231–1250. [PubMed: 18838040]
- Opitz PC, Rauch LC, Terry DP, Urry HL. Prefrontal mediation of age differences in cognitive reappraisal. *Neurobiology of Aging*. 2010; 33:645–655. [PubMed: 20674090]
- Ochsner KN, Gross JJ. The cognitive control of emotion. *TRENDS in Cognitive Science*. 2005; 9:242–249.
- Ochsner KN, Gross JJ. Cognitive emotion regulation: Insights from social cognitive and affective neuroscience. *Current Directions in Psychological Science*. 2008; 17:152–158.
- Ochsner KN, Silvers JA, Buhle JT. Functional imaging studies of emotion regulation: A synthetic review and evolving model of the cognitive control of emotion. *Annals of the New York Academy of Sciences*. 2012:E1–E24. [PubMed: 23025352]
- Payer DE, Baicy K, Lieberman MD, London ED. Overlapping neural substrates between intentional and incidental down-regulation of negative emotions. *Emotion*. 2012; 12:229–235. [PubMed: 22468617]
- Pitskel NB, Bolling DZ, Kaiser MD, Crowley MJ, Pelphrey KA. How grossed out are you?. The neural bases of emotion regulation from childhood to adolescence. *Developmental Cognitive Neuroscience*. 2011; 1(3):324–337. [PubMed: 21686071]
- Ray RD, McRae K, Ochsner KN, Gross JJ. Cognitive reappraisal of negative affect: Converging evidence from EMG and self-report. *Emotion*. 2010; 10:587–592. [PubMed: 20677875]
- Schaefer SM, Jackson DC, Davidson RJ, Aguirre GK, Kimberg DY, Thompson-Schill SL. Modulation of amygdalar activity by the conscious regulation of negative emotion. *Journal of Cognitive Neuroscience*. 2002; 14:913–921. [PubMed: 12191458]
- Southam-Gerow MA, Kendall PC. Emotion regulation and understanding implications for child psychopathology and therapy. *Clinical Psychology Review*. 2002; 22:189–222. [PubMed: 11806019]
- Stegge, H.; Meerum Terwogt, M. Awareness and regulation of emotion in typical and atypical development. In: Gross, JJ., editor. *Handbook of emotion regulation*. New York: Guilford Press; 2007. p. 249–268.

- Uematsu A, Matsui M, Tanaka C, Takahashi T, Noguchi K, Suzuki M, Nishijo H. Developmental trajectories of amygdala and hippocampus from infancy to early adulthood in healthy individuals. *PLOS ONE*. 2012; 7(10):1–10.
- Urry HL, van Reekum CM, Johnstone T, Kalin NH, Thurow ME, Schaefer HS, Jackson CA, Frye CJ, Greischar LL, Alexander AL, Davidson RJ. Amygdala and ventromedial prefrontal cortex are inversely coupled during regulation of negative affect and predict the diurnal pattern of cortisol secretion among older adults. *Journal of Neuroscience*. 2006; 26(16):4415–4425. [PubMed: 16624961]
- Vul E, Harris C, Winkielman P, Pashler H. Puzzlingly high correlations in fMRI studies of emotion, personality, and social cognition. *Perspectives on Psychological Science*. 2009; 4:274–290. [PubMed: 26158964]
- Wager TD, Keller MC, Lacey SC, Jonides J. Increased sensitivity in neuroimaging analyses using robust regression. *NeuroImage*. 2005; 26:99–113. [PubMed: 15862210]
- Wessa M, Kanske P, Neumeister P, Bode K, Heissler J, Schonfelder S. EmoPics: Subjektive und psychophysiologische Evaluation neuen Bildmaterials für die klinisch-biopsychologische Forschung. *Zeitschrift für Klinische Psychologie und Psychotherapie*. 2010; 11(Supplementum 1): 77.
- Wilcox RR. Outlier detection. *Encyclopedia of statistics in behavioral science*. 2005
- Wolgast M, Lundh L, Viborg G. Cognitive reappraisal and acceptance: An experimental comparison of two emotion regulation strategies. *Behaviour Research and Therapy*. 2011; 49:858–866. [PubMed: 21999941]

Figure 1.

Experimental paradigm: Each trial began with an auditory story lasting 5.3 s, followed by a delay lasting 2 s, and followed by a negative or neutral-valenced picture lasting 2.5 s. Study 1 concluded each picture display with a subjective report of affect prompt. Study 2 did not collect subjective reports; each picture was followed by a 4 – 8 s jittered inter-trial interval (ITI).

Figure 2.

Mean affect ratings as a function of condition in Study 1. Affect ratings ranged from from 1 (negative) to 5 (positive). Error bars represent standard error of the mean. Asterisks represent significance at a threshold of $p < .001$.

Figure 3.

Visualization of the ROIs on pediatric MNI atlas. Graphs display the average deconvolved responses (left) and response indices (right) as a function of trial type in bilateral amygdala (A,B), bilateral dorsolateral PFC (C,D), and ventromedial PFC (E). Shaded regions indicate the time points used to index responses to pictures. Error bars denote the standard within subject error term (Cousineau, 2005). PFC=prefrontal cortex. Arrows in Figure 3E indicate the onset times of the auditory story and picture phases.

Figure 4.

Relationship between brain regions during down-regulation using robust regression. Across participants a significant linear relationship was observed involving fMRI interactions scores ($[\text{negative-decrease vs. negative-attend}]_{\text{Amygdala}} - [\text{negative-decrease vs. negative-attend}]_{\text{ventromedialPFC}}$) between Right amygdala and ventromedial PFC (A); a linear relationship was not significant between Left amygdala and ventromedial PFC (B). PFC=prefrontal cortex.

Table 1

Peak coordinates in MNI space and number of voxels [k] in each ROI

ROI	x	Y	z	K
Left Dorsolateral PFC	56	-31	13	75
Right Amygdala	-16	5	-14	72
Left Amygdala	-20	-2	-17	52
Ventromedial PFC	2	-22	-26	18
Right Dorsolateral PFC	-52	-19	28	12

Note. PFC = prefrontal cortex.

Table 2

ROI paired t-test results

ROI	Negative-attend vs Neutral-attend	Negative-decrease vs Negative-attend
Right Amygdala	$t_{14}=2.23$, $p=.043$	$t_{14}=2.98$, $p=.010$
Left Amygdala	$t_{14}=1.59$, $p=.134$	$t_{14}=1.81$, $p=.092$
Left Dorsolateral PFC	$t_{14}=2.14$, $p=.051$	$t_{14}=0.99$, $p=.341$
Right Dorsolateral PFC	$t_{14}=2.34$, $p=.035$	$t_{14}=1.52$, $p=.151$
Ventromedial PFC	$t_{14}=0.10$, $p=.924$	$t_{14}=2.54$, $p=.024$

Note. PFC = prefrontal cortex.